Bresenham's mid-point line drawing algorithm (Generalized version):

A generalized version of the Bresenham's mid-point line drawing algorithm will work for any straight line (its starting and ending points can be in any octant).

Pseudocode:

Procedure Bresenham_line (x_S, y_S, x_E, y_E) //Starting and ending points of line are passed as arguments Begin

```
x = x_S, y = y_S // Initialization of starting point
dx = Abs(x_E - x_S), dy = Abs(y_E - y_S)
s1=Sign(x_E - x_{S}), s2=Sign(y_E - y_{S})
If(dy>dx)
       Temp=dx
       dx=dy
       dy=Temp
       Swap=1
Else
       Swap=0
EndIf
d = 2dy - dx // Initialization of decision variable d
Steps=1
SetPixel(x,y)
While(Steps\leqdx)
Begin
       If(d>0)
                x=x+s1
                y=y+s2
                d=d+2(dy-dx)
       Else
                If(Swap=1)
                        y=y+s2
                Else
                        x=x+s1
                EndIf
                d=d+2dy
       EndIf
       SetPixel(x,y)
       Steps=Steps+1
EndWhile
```

Functions used:

End

SetPixel(x, y) is used to plot the corresponding pixel defined by co-ordinate (x, y) on the screen.

Abs(k) returns absolute value of its argument k.

Sign() returns -1, 0, 1 as its argument (say, k) <0, =0 or >0 i.e. Sign(k)=k/Abs(k).

Abhishek Dey
Assistant Professor
Department of Computer Science
Bethune College, Kolkata

Q. Find plotted pixels or plotted points of straight line P(4, 2), Q(6, 14) using Bresenham's midpoint line drawing algorithm.

Answer:

$$dx = 2$$
, $dy = 12$

$$s1=Sign(6-4)=Sign(2)=1$$
, $s1=Sign(14-2)=Sign(12)=1$

d = -8

As, dy>dx, so swap them. Hence, dx=12, dy=2

Steps=1

Initialize, x = 4, y = 2

Now 1st point (4, 2) is plotted

Next points will be:

Steps	d	X	y	Plotted
				point
1	-4	4	3	(4,3)
2	0	4	4	(4,4)
3	4	4	5	(4,5)
4	-16	5	6	(5,6)
5	-12	5	7	(5,7)
6	-8	5	8	(5,8)
7	-4	5	9	(5,9)
8	0	5	10	(5,10)
9	4	5	11	(5,11)
10	-16	6	12	(6,12)
11	-12	6	13	(6,13)
12	-8	6	14	(6,14)

Hence, the plotted points / pixels are: (4,3), (4,4), (4,5), (5,6), (5,7), (5,8), (5,9), (5,10), (5,11), (6,12), (6,13), (6,14).

HW. Find plotted pixels or plotted points of straight line P(4, -6), Q(2, 1) using Bresenham's mid-point line drawing algorithm.