

EE23010 Assignment

Sayyam Palrecha* EE22BTECH11047

Question 10.13.3.19

Two dice are thrown at the same time. Find the probability of getting

- (i) same number on both dice.
- (ii) different numbers on both dice.

Solution: Let the random variables:

parameters	value	description
X	$1 \leq X \leq 6$	outcome of the first die
Y	$1 \leq Y \leq 6$	outcome of the second die

Consider a random variable Z :

$$Z = X - Y$$

Z can take values ranging from $\{-5$ to $5\}$.

We need to find the PMF of Z

We know that,

$$p_X(k) = \begin{cases} \frac{1}{6}, & 1 \leq k \leq 6 \\ 0, & \text{otherwise} \end{cases} \quad (2)$$

$p_Y(k)$ is same as $p_X(k)$.

PMF of Z using z -transform:

$$\Pr(k = Z) = \Pr(k = X - Y)$$

applying the z -transform on both the sides

$$z\{\Pr(k = Z)\} = z\{\Pr(k = X - Y)\} \quad (4)$$

This gives us the z -transform of convolution of sequences

$$z\{p_Z(k)\} = z\{p_X(k) * p_{-Y}(k)\} \quad (5)$$

Let $X(z)$ and $Y(-z)$ represent $z\{p_X(k)\}$ and $z\{p_{-Y}(k)\}$ respectively,

$$= X(z) \cdot Y(-z) \quad (6)$$

$$= \left(\sum_{i=1}^6 p_X(i) \cdot z^i \right) \cdot \left(\sum_{j=1}^6 p_Y(j) \cdot z^{-j} \right) \quad (7)$$

$$= \left(\frac{1}{6} \sum_{i=1}^6 z^i \right) \left(\frac{1}{6} \sum_{j=1}^6 z^{-j} \right) \quad (8)$$

$$= \frac{1}{36} \sum_{i=1}^6 \sum_{j=1}^6 (z^i \cdot z^{-j}) \quad (9)$$

$$= \frac{1}{36} \sum_{i=1}^6 \sum_{j=1}^6 z^{i-j} \quad (10)$$

- (1) Here i and j represents outcomes on the dice X and dice Y respectively, so $i - j$ represents the value k .

$$\Rightarrow p_Z(k) = \frac{1}{36} \sum_{i=1}^6 \sum_{j=1}^6 z^{i-j} \Big|_{i-j=k} \quad (11)$$

Here i and j should satisfy the condition: $i - j = k$

$$= \frac{1}{36} \sum_{j=1, i=1+k}^{6, 6+k} z^k \Big|_{i-j=k} \quad (12)$$

- (i) Finding the probability for $Z = 0$

From the result (12)

$$p_Z(0) = \frac{1}{36} \sum_{j=1, i=1}^{6, 6} z^0 \Big|_{i=j} \quad (13)$$

- (3) The coefficient of z^k (z^0 here), is the probability $P(Z = 0)$

$$\Rightarrow \Pr(Z = 0) = \frac{1}{36}(6) \quad (14)$$

$$= \frac{1}{6} \quad (15)$$

- (ii) Finding the probability for $Z \neq 0$

$$\Pr(Z \neq 0) = 1 - \Pr(Z = 0) \quad (16)$$

$$= 1 - \frac{1}{6} \quad (17)$$

$$= \frac{5}{6} \quad (18)$$




Fig. 0. Theoretical PMF of Z ($p_Z(k)$)