

Honors Discrete Mathematics:

Lecture 7 Notes

Professor Gerandy Brito Spring 2022

Sarthak Mohanty

Cardinality

There is an easy way to test if two numbers have the same cardinality, which will become useful when we deal with infinite sets.

Theorem. Two sets A and B are said to have the same cardinality if there exists a bijection (surjection?) from A to B . The proof of this theorem is outside the scope of this course.

Now consider the collection of all sets (i.e.: the power set of the universe!) and define the relation

$$A\mathcal{R}B \text{ iff } A \text{ and } B \text{ have the same cardinality.}$$

Using the above theorem, we will now show that this relation is an equivalence relation.

1. Reflexivity: There exists a bijection from A to B $f : A \rightarrow B$.
2. Symmetry: Since $a\mathcal{R}b$, we have defined a bijection from A to B . As discussed before, bijections have inverses, so we can define an inverse $f^{-1}B \rightarrow A$, so $b\mathcal{R}a$.
3. Transitivity: By definition, there exists a bijection from A to B . Furthermore, there exists a bijection from B to C . The composition $h : A \rightarrow C = g \circ f$ is a bijection as well (proof shown below), so $a\mathcal{R}c$.

Proposition. Let $f : A \rightarrow B$ and $g : B \rightarrow C$ be bijections. Then $h : A \rightarrow C$ is a bijection as well.

Proof. Let's start by proving that $g \circ f$ is one-to-one. Suppose $g \circ f(a) = g \circ f(b)$. Then $g(f(a)) = g(f(b))$. Since g is one-to-one, this implies $f(a) = f(b)$. Since f is one-to-one, $a = b$. Therefore $g \circ f$ is one-to-one.

Let's now prove that $g \circ f$ is onto. Suppose $c \in C$. Since $c \in C$ and g is onto, there exists $b \in B$ such that $g(b) = c$. Since $b \in B$ and f is onto, there exists an $a \in A$ such that $f(a) = b$. Therefore $g(f(a)) = c$, i.e., $g \circ f(a) = c$. Since $g \circ f$ is one-to-one and onto, it is a bijection.

Definition: A set is said to be *countable* if it is finite or has the same cardinality as the natural numbers. Examples: Show the following are countable sets by defining bijections for each of the following sets.

- From \mathbb{N} to the set of positive even numbers.
- From \mathbb{N} to the set of negative integers.
- From \mathbb{N} to the set of odd positive numbers.
-

Solution:

- $f(n) = 2n$
- $f(n) = -n$
- Let A denote the set of odd natural numbers. Let $f(x) = 2x - 1$ for all $x \in \mathbb{N}$. Then $f : \mathbb{N} \rightarrow A$ is a bijection.
-

Lemma. Let A, B be countable sets. Then $A \cup B$ is also countable.

Proof. Suppose both $|A|$ and $|B|$ is finite. Then the proof is trivial.

Now suppose at least one of $|A|$, $|B|$ is infinite. WLOG, suppose $|A|$ is infinite and the size of B is k . Define a series of elements with the element at index i equal to the i -th element of A . Then shift every element k spaces to the right, and fill the first k spaces in the series with the elements in B . Since series are functions, we have shown there exists a bijection from $|A|$ to $|B|$

Now suppose both $|A|$ and $|B|$ are infinite. Let $f(n) = 2n$ and $g(n) = 2n - 1$. Define a function $h(n)$ as

$$h(n) = \begin{cases} f(n) & \text{if } n \text{ is even} \\ g(n) & \text{if } n \text{ is odd} \end{cases}$$

We now have another tool to show sets are countable: The finite union of countable sets is also countable.

Lemma. Let A, B be countable sets. If $A \subseteq B$ and B is countable, then so is A . Proof is left as an exercise for the reader.

Lemma. Let A, B be countable sets. Then $A \times B$ is countable.

Proof. (TBC) We know $A \times B = \{a_i, b_j : a_i \in A, b_j \in B\}$.

Define a function f as follows:

$$\begin{aligned} f(a_1, b_1) &= 1, \\ f(a_2, b_1) &= 2, \quad f(a_1, b_2) = 3, \\ f(a_3, b_1) &= 4, \quad f(a_3, b_2) = 5, \quad f(a_3, b_3) = 6, \\ f(1, 4) &= 7, \quad f(2, 3) = 8, \quad f(3, 2) = 9, \quad f(4, 1) = 10, \end{aligned}$$

and so on. Then $f : \mathbb{A} \times \mathbb{B} \rightarrow \mathbb{N}$ is a bijection.

Example. Show that \mathbb{N} to $A = \{x \in \mathbb{Q} : x > 0\}$ is a bijection.

We have many options, we can use the previous lemma and define the Cartesian product as $A \times B$, where A is the set of all numerators and B is the set of all denominators.

Let A denote the set of positive rational numbers. Define a function f as follows:

$$\begin{aligned} f(1) &= \frac{1}{1}, \\ f(2) &= \frac{1}{2}, \quad f(3) = \frac{2}{1}, \\ f(4) &= \frac{1}{3}, \quad f(5) = \frac{3}{1}, \\ f(6) &= \frac{1}{4}, \quad f(7) = \frac{2}{3}, \quad f(8) = \frac{3}{2}, \quad f(9) = \frac{4}{1}, \end{aligned}$$

Cantor's Diagonal Lemma

Let f be a function from \mathbb{N} to $(0, 1)$. Prove that there exists $y \in (0, 1)$ such that y does not belong to the range of f . (in other words, prove the set of real numbers is not countable.)

Solution

We are given a function $f : \mathbb{N} \rightarrow (0, 1)$. We wish to find a number $y \in (0, 1)$ such that

$$y \notin \{f(1), f(2), f(3), f(4), \dots\}.$$

For each $n \in \mathbb{N}$ and each $k \in \mathbb{N}$, let x_{nk} be the k -th digit in the *standard* decimal expansion of $f(n)$. Then

$$\begin{aligned} f(1) &= 0.\overline{x_{11}}x_{12}x_{13}x_{14}\dots, \\ f(2) &= 0.x_{21}\overline{x_{22}}x_{23}x_{24}\dots, \\ f(3) &= 0.x_{31}x_{32}\overline{x_{33}}x_{34}\dots, \\ f(4) &= 0.x_{41}x_{42}x_{43}\overline{x_{44}}\dots, \\ &\text{and so on.} \end{aligned}$$

We shall define the number y by defining the digits in its decimal expansion so that they are different from the “diagonal” entries $x_{11}, x_{22}, x_{33}, x_{44}, \dots$ that are highlighted in the equations above. For each $n \in \mathbb{N}$, let

$$y_n = \begin{cases} 5 & \text{if } x_{nn} \neq 5, \\ 4 & \text{if } x_{nn} = 5. \end{cases}$$

Then for each $n \in \mathbb{N}$, $y_n \neq x_{nn}$. Now let y be the number whose standard decimal expansion is

$$y = 0.y_1y_2y_3y_4\dots$$

Then $y \in (0, 1)$. In fact, $0.444\dots \leq y \leq 0.555\dots$. To see that y is not in the range of f , note that for each $n \in \mathbb{N}$, $y \neq f(x)$ (because the numbers y and $f(n)$ differ in their n -th decimal place; in other words, $y_n \neq x_{nn}$).

Post Lecture

Question 3

Describe bijections (without justifications): Whenever the bijection is defined by a single formula, also provide its inverse.

- (a) from \mathbb{Z} to \mathbb{N} .
- (b) from $(-\frac{\pi}{2}, \frac{\pi}{2})$ to \mathbb{R} . [A suitable trigonometric function will do.]
- (c) from $(0, 1)$ to \mathbb{R} . [Compose a linear map with the map in part (b).]

Solution

- (a) Let $n \in \mathbb{Z}$. Define $f(n)$ by

$$f(n) = \begin{cases} 2n + 1 & \text{if } n \geq 0, \\ -2n & \text{if } n < 0. \end{cases}$$

Then $f : \mathbb{Z} \rightarrow \mathbb{N}$ is a bijection.

- (b) Let $f(x) = \tan(x)$ for all $x \in (-\frac{\pi}{2}, \frac{\pi}{2})$. Let $g(y) = \tan^{-1}(y)$ for all $y \in \mathbb{R}$. Then $f^{-1} = g$ and $f : (-\frac{\pi}{2}, \frac{\pi}{2}) \rightarrow \mathbb{R}$ is a bijection.
- (c) Let $f(x) = \tan(x)$ and $g(x) = \sin^{-1}(2x - 1)$. Now f is a bijection from $(-\frac{\pi}{2}, \frac{\pi}{2})$ to \mathbb{R} and g is a bijection from $(0, 1)$ to $(-\frac{\pi}{2}, \frac{\pi}{2})$. Then $(f \circ g)(x) = \tan(\sin^{-1}(2x - 1))$ is a bijection from $(0, 1)$ to \mathbb{R} . The inverse of $f \circ g$ is $(g^{-1} \circ f^{-1})(y) = \frac{\sin(\tan^{-1}(y)) + 1}{2}$ for all $y \in \mathbb{R}$.