

**Problem 1**

- a) 0
- b) No
- c) 0

**Notes on SPD Matrices, Inner Products, Norms, and Metrics**

**Problem 1:** prove Fact 3 – If  $A$  is SPD, then  $A$  is invertible, and  $A^{-1}$  is SPD too.

Proof: Let  $A$  be a SPD matrix. Assume  $A$  is not invertible, thus there exist vector  $x \neq 0$  such that  $Ax = 0$ .  
 $x^T Ax = x^T 0 = 0$  in contradiction to the fact that  $A$  is SPD. ( $x^T Ax > 0 \quad \forall x \neq 0$ )

Therefore  $A$  is invertible, meaning  $A^{-1}$  exists.

Let  $\lambda$  be an eigenvalue of  $A$ , hence:

$$Av = \lambda v \rightarrow A^{-1}Av = A^{-1}\lambda v \rightarrow A^{-1}Av = \lambda A^{-1}v \rightarrow Iv = \lambda A^{-1}v \rightarrow \frac{1}{\lambda}v = A^{-1}v$$

According to fact 1, all eigenvalues  $\lambda$  of  $A$  are positive. Therefore, all  $\frac{1}{\lambda}$  (the eigenvalues of  $A^{-1}$ ) are positive, and again from fact 1 we obtain that  $A^{-1}$  is SPD.

**Problem 2:** prove Fact 4 – Let  $Q$  be an  $n \times n$  SPD matrix. Then  $\langle \cdot, \cdot \rangle_Q: R^n \times R^n \rightarrow R$ ,  $\langle \cdot, \cdot \rangle_Q: (x, y) \rightarrow x^T Q y$  is an inner product.

Solution: Let  $Q$  be an  $n \times n$  SPD matrix and  $x, y, z \in R^n$ ,  $\alpha, \beta \in R$

We will show that the properties of definition 6 hold.

- 1)  $\langle x, y \rangle_Q = x^T Q y \underset{x^T Q y \in R}{=} (x^T Q y)^T = y^T Q^T x \underset{Q \text{ is symmetric}}{=} y^T Q x = \langle y, x \rangle_Q$
- 2)  $\langle \alpha x + \beta y, z \rangle_Q = (\alpha x + \beta y)^T Q z = (\alpha x^T + \beta y^T) Q z = \alpha x^T Q z + \beta y^T Q z = \alpha \langle x, z \rangle_Q + \beta \langle y, z \rangle_Q$
- 3)  $\langle x, x \rangle_Q = x^T Q x \underset{Q \text{ is SPD}}{\geq} 0$
- 4)  $\Leftarrow$  Let  $x = 0_v$ , then  $x^T Q x = 0_v^T Q 0_v = 0$   
 $\Rightarrow$  Let  $x^T Q x = 0$ , then from SPD definition,  $x = 0_v$

**Problem 3:** prove Fact 8 – Every norm induces a metric:  $d(x, y) = \|x - y\|$ .

Proof: Let  $x, y, z \in V (= M)$ , we will show that the properties of definition 11 hold.

- 1)  $\|x - y\| \underset{\text{from norm definition}}{\geq} 0$
- 2)  $\Rightarrow$  Let  $\|x - y\| = 0$  then, from definition,  $x - y = 0_v \rightarrow x = y$   
 $\Leftarrow$  Let  $x = y$ , then  $x - y = 0_v$  and from definition  $\|0_v\| = 0$
- 3)  $\|x - y\| = \|-1(y - x)\| \underset{\text{from norm definition}}{=} |-1| \cdot \|y - x\| = \|y - x\|$
- 4)  $\|x - z\| = \|(x - y) + (y - z)\| \underset{\text{from norm definition}}{\leq} \|x - y\| + \|y - z\|$

**Problem 4:** prove Fact 10 – Let  $Q$  be an SPD matrix and  $LL^T$  denote its Cholesky decomposition.


$$\text{Then } \|x\|_Q = \|L^T x\|_{l_2}.$$

Proof: Let  $Q$  be an SPD matrix and  $LL^T$  denote its Cholesky decomposition.


$$\|x\|_Q = x^T Q x = x^T L L^T x = (L^T x)^T L^T x = \|L^T x\|_{l_2}$$

**Computer Exercise 1:**


→ The relevant code will be added separately in the submission page.




a:




b:



c:



d:



e:

## Notes on Convexity

### Problem 1:

- 1)  $f: R \rightarrow R, f(x) = x^2$  is convex:

Let assume that  $x_1, x_2 \in R$ , so:

$$\begin{aligned}
 f(tx_1 + (1-t)x_2) &\stackrel{?}{\leq} tf(x_1) + (1-t)f(x_2) \\
 (tx_1 + (1-t)x_2)^2 &\stackrel{?}{\leq} tx_1^2 + (1-t)x_2^2 \\
 t^2x_1^2 + 2t(1-t)x_1x_2 + (1-t)^2x_2^2 &\stackrel{?}{\leq} tx_1^2 + x_2^2 - tx_2^2 \\
 t^2x_1^2 + 2(t-t^2)x_1x_2 + x_2^2 - 2tx_2^2 + t^2x_2^2 &\stackrel{?}{\leq} tx_1^2 + x_2^2 - tx_2^2 \\
 t^2x_1^2 - tx_1^2 + 2(t-t^2)x_1x_2 + tx_2^2 - 2tx_2^2 + t^2x_2^2 &\stackrel{?}{\leq} 0 \\
 (t^2-t)x_1^2 + 2(t-t^2)x_1x_2 + (t-2t+t^2)x_2^2 &\stackrel{?}{\leq} 0 \\
 (t^2-t)x_1^2 - 2(t^2-t)x_1x_2 + (t^2-t)x_2^2 &\stackrel{?}{\leq} 0 \\
 (t^2-t)(x_1-x_2)^2 &\stackrel{?}{\leq} 0
 \end{aligned}$$

This is true because  $(x_1 - x_2)^2 \geq 0$  by definition and because  $t \in (0,1)$  we will get that  $t > t^2$ .

- 2)  $f: R \rightarrow R, f(x) = |x|$  is convex:

Let assume that  $x_1, x_2 \in R$ , so:

$$\begin{aligned}
 f(tx_1 + (1-t)x_2) &= |tx_1 + (1-t)x_2| = \\
 &= |tx_1 + (1-t)x_2| \stackrel{\text{by the triangle inequality}}{\leq} |tx_1| + |(1-t)x_2| \stackrel{\text{since } t \in (0,1)}{=} t|x_1| + (1-t)|x_2|
 \end{aligned}$$

- 3)  $f: R \rightarrow R, f(x) = ax + b$  is convex:

Let assume that  $x_1, x_2 \in R$ , so:

$$\begin{aligned}
 f(tx_1 + (1-t)x_2) &\stackrel{?}{\leq} tf(x_1) + (1-t)f(x_2) \\
 a(tx_1 + (1-t)x_2) + b &\stackrel{?}{\leq} t(ax_1 + b) + (1-t)(ax_2 + b) \\
 atx_1 + a(1-t)x_2 + b &\stackrel{?}{\leq} tax_1 + tb + a(1-t)x_2 + (1-t)b \\
 b &\stackrel{?}{\leq} tb + (1-t)b \\
 b &\stackrel{?}{\leq} b(t(1-t)) \\
 t - t^2 &\stackrel{?}{\geq} 0 \\
 &\text{since } t \in (0,1)
 \end{aligned}$$

- 4)  $f: R \rightarrow R, f(x) = \sin(x)$  is not convex:

For  $x_1 = 0, x_2 = \pi, t = 0.5$ :

$$f(tx_1 + (1-t)x_2) = \sin(0.5 * 0 + 0.5 * \pi) = \sin\left(\frac{\pi}{2}\right) = 1$$

$$tf(x_1) + (1-t)f(x_2) = 0.5 * \sin(0) + 0.5 * \sin(\pi) = 0 + 0 = 0$$

and because  $1 \not\leq 0$ ,  $f(x) = \sin(x)$  is not convex.

- 5)  $f: R \rightarrow R, f(x) = -x^2$  is not convex:

For  $x_1 = -1, x_2 = 1, t = 0.5$ :

$$f(tx_1 + (1-t)x_2) = -(0.5 * -1 + 0.5 * 1)^2 = -(0)^2 = 0$$

$$tf(x_1) + (1-t)f(x_2) = 0.5 * -(-1)^2 + 0.5 * -(1)^2 = -0.5 + -0.5 = -1$$

and because  $0 \not\geq -1$ ,  $f(x) = -x^2$  is not convex.

### Notes on Argmin and Argmax

**Problem 1:** Let  $f : S \rightarrow R$  be a function from some set  $S$  into  $R$ . Then  $\operatorname{argmin}_{x \in S} f(x) = \operatorname{argmax}_{x \in S} -f(x)$ .

Proof: Let  $f : S \rightarrow R$  be a function from some set  $S$  into  $R$ .

$$\operatorname{argmax}_{x \in S} -f(x) = \{x : -f(x) \geq -f(x') \ \forall x' \in S\} = \{x : f(x) \leq f(x') \ \forall x' \in S\} = \operatorname{argmin}_{x \in S} f(x)$$

**Problem 2:** For some  $g : R \rightarrow R$  monotonically non-increasing function, and some  $f : S \rightarrow R$  it holds that:

$$\operatorname{argmax}_{x \in S} f(x) \notin \operatorname{argmax}_{x \in S} g(f(x))$$

Proof: set  $f$  as  $f(x) = -x^2$  and  $g$  as  $g(x) = -x$ .

It is easy to see that  $\operatorname{argmax}_{x \in R} f(x) = \operatorname{argmax}_{x \in R} (-x^2) = 0$ .

But now:  $g(f(0)) = g(0) = -0 = 0$ , while for value  $x = 1$ ,  $g(f(x)) = g(-1) = 1$ , and  $1 > 0$ .

Therefore  $\operatorname{argmax}_{x \in R} f(x) = 0 \notin \operatorname{argmax}_{x \in R} g(f(x))$ .

**Problem 3:** Let  $g : R \rightarrow R$  be a monotonically non-decreasing function. Let  $f : S \rightarrow R$  be a function we seek to maximize. Then,  $\operatorname{argmax}_{x \in S} f(x) \subset \operatorname{argmax}_{x \in S} g(f(x))$ .

Proof: By the definition of argmax:

$$x_0 \in \operatorname{argmax}_{x \in S} f(x) \rightarrow f(x_0) \geq f(x) \ \forall x \rightarrow g(f(x_0)) \geq g(f(x)) \ \forall x \rightarrow x_0 \in \operatorname{argmax}_{x \in S} (g \circ f)(x)$$

**Problem 4:** Let  $r \in R^n$  depend on  $\theta \in R^k$ . Show that  $\operatorname{argmin}_{\theta \in R^k} \|r\|_{l_2} = \operatorname{argmin}_{\theta \in R^k} \|r\|_{l_2}^2$ .

Proof: By using Fact 5 for  $g(x) = x^2$ , and  $f(x) = \|r(\theta)\|$ , we directly deduce that:

$$\operatorname{argmin}_{\theta \in R^k} \|r\|_{l_2} = \operatorname{argmin}_{\theta \in R^k} \|r\|_{l_2}^2$$

that is because that  $f(x) \geq 0 \ \forall x$ , and  $g(x) = x^2$  is monotonically increasing on  $[0, \infty]$ .

**Problem 5:** Let  $\sigma \in R_{>0}$  and let  $r \in R^n$  depend on  $\theta \in R^k$ . Show that:

$$\operatorname{argmax}_{\theta \in R^k} \frac{1}{(2\pi\sigma)^{\frac{n}{2}}} \exp\left(-\frac{1}{2} \frac{\|r\|_{l_2}^2}{\sigma^2}\right) = \operatorname{argmin}_{\theta \in R^k} \|r\|_{l_2}$$

Proof: From Problem 4 we get that:

$$\operatorname{argmin}_{\theta \in R^k} \|r\|_{l_2} = \operatorname{argmin}_{\theta \in R^k} \|r\|_{l_2}^2$$

From Fact 1 we get that  $\operatorname{argmin}_{\theta \in R^k} \|r\|_{l_2}^2 = \operatorname{argmax}_{\theta \in R^k} (-\|r\|_{l_2}^2)$

Now, again from Fact 5, for monotonically increasing function  $g(x) = \frac{1}{2} \frac{x}{\sigma^2}$  we get that:

$$\operatorname{argmax}_{\theta \in R^k} (-\|r\|_{l_2}^2) = \operatorname{argmax}_{\theta \in R^k} \left(-\frac{1}{2} \frac{\|r\|_{l_2}^2}{\sigma^2}\right)$$

And again for monotonically increasing function  $g^*(x) = \exp(x)$  we get that

$$\operatorname{argmax}_{\theta \in R^k} \left(-\frac{1}{2} \frac{\|r\|_{l_2}^2}{\sigma^2}\right) = \operatorname{argmax}_{\theta \in R^k} \left(\exp\left(-\frac{1}{2} \frac{\|r\|_{l_2}^2}{\sigma^2}\right)\right)$$

And one last time, for monotonically increasing function  $g^{**}(x) = \frac{1}{(2\pi\sigma)^{\frac{n}{2}}} x$  we get that :

$$\operatorname{argmax}_{\theta \in R^k} \left(\exp\left(-\frac{1}{2} \frac{\|r\|_{l_2}^2}{\sigma^2}\right)\right) = \operatorname{argmax}_{\theta \in R^k} \left(\frac{1}{(2\pi\sigma)^{\frac{n}{2}}} \exp\left(-\frac{1}{2} \frac{\|r\|_{l_2}^2}{\sigma^2}\right)\right)$$

So overall we achieved that:  $\operatorname{argmax}_{\theta \in R^k} \frac{1}{(2\pi\sigma)^{\frac{n}{2}}} \exp\left(-\frac{1}{2} \frac{\|r\|_{l_2}^2}{\sigma^2}\right) = \operatorname{argmin}_{\theta \in R^k} \|r\|_{l_2}$

### Notes on Linear Least Squares

**Problem 1:** Find  $\operatorname{argmin}_{\theta \in \mathbb{R}^k} \|H\theta - y\|_{l_2}^2 + \lambda_1 \theta_1^2 + \lambda_3 \theta_3^2$

where  $H \in \mathbb{R}^{N \times 5}$ ,  $y \in \mathbb{R}^N$ ,  $\theta = [\theta_1 \ \theta_2 \ \theta_3 \ \theta_4 \ \theta_5]^T \in \mathbb{R}^5$ ,  $\lambda_1, \lambda_3 > 0$

Proof: We will bring our problem to a least squares manner:

$$\|H\theta - y\|_{l_2}^2 + \lambda_1 \theta_1^2 + \lambda_3 \theta_3^2 = \left\| \begin{bmatrix} \sqrt{\lambda_1} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \sqrt{\lambda_3} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}^T \theta - \begin{bmatrix} y \\ 0_{5 \times 1} \end{bmatrix} \right\|^2$$

Thus, from linear least squares, the minimizer satisfies the following equation:

$$\begin{aligned} \left[ \begin{bmatrix} \sqrt{\lambda_1} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \sqrt{\lambda_3} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}^T \right] \left[ \begin{bmatrix} \sqrt{\lambda_1} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \sqrt{\lambda_3} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \theta \right] &= \left[ \begin{bmatrix} \sqrt{\lambda_1} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \sqrt{\lambda_3} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}^T \right] \begin{bmatrix} y \\ 0_{5 \times 1} \end{bmatrix} \\ &\rightarrow \left( H^T H + \begin{bmatrix} \lambda_1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \lambda_3 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \right) \theta = H^T y \end{aligned}$$

$H^T H$  is always SPD, and the addition of positive values on the diagonal won't change that so the new matrix is invertible.

$$\rightarrow \theta = \left( H^T H + \begin{bmatrix} \lambda_1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \lambda_3 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \right)^{-1} H^T y$$

**Problem 2:** Find  $\operatorname{argmin}_{\hat{x} \in \operatorname{span}(v_1, \dots, v_k)} \|x - \hat{x}\|_{l_2}$

Proof: as mentioned at the notes,  $\hat{x} \in \operatorname{span}(v_1, \dots, v_k) \rightarrow \hat{x} = \sum_{j=1}^k \theta_j v_j$ , when  $\theta = \begin{bmatrix} \theta_1 \\ \vdots \\ \theta_k \end{bmatrix} \in \mathbb{R}^k$

and  $V = [v_1, \dots, v_k] \in \mathbb{R}^{d \times k}$  and therefore :

$$\hat{x} = V * \theta$$

Meaning the two following problems are equal:

$$\operatorname{argmin}_{\hat{x} \in \operatorname{span}(v_1, \dots, v_k)} \|x - \hat{x}\|_{l_2} \underset{\square^2 \text{ is } \uparrow}{=} \operatorname{argmin}_{\hat{x} \in \operatorname{span}(v_1, \dots, v_k)} \|x - \hat{x}\|_{l_2}^2 = \operatorname{argmin}_{\theta \in \mathbb{R}^k} \|x - V\theta\|_{l_2}^2$$

We know how to solve it by LS and the normal equations:

$$\operatorname{argmin}_{\theta \in \mathbb{R}^k} \|x - V\theta\|_{l_2}^2 = \operatorname{argmin}_{\theta \in \mathbb{R}^k} \|V\theta - x\|_{l_2}^2 \rightarrow V^T V \theta = V^T x$$

Because  $V$  is orthogonal we get:

$$I\theta = V^T x \rightarrow \theta = V^T x$$

### Notes on Random Vectors

**Problem 1:**  $P$  satisfies finite additivity; namely, if  $A_1, A_2, \dots, A_n$  is a finite collection of pairwise disjoint events then  $P(\bigcup_{i=1}^n A_i) = \sum_{i=1}^n P(A_i)$ .

Proof:

$$\sum_{i=1}^n P(A_i) = \sum_{i=1}^n P(A_i) + \sum_{i=n+1}^{\infty} 0 \stackrel{\substack{\text{by infromal} \\ \text{definition 2} \\ \text{(point 1)}}}{=} \sum_{i=1}^n P(A_i) + \sum_{i=n+1}^{\infty} P(\emptyset)$$

Let us denote  $A_i = \emptyset$  for every  $i > n$ , and now:

$$\begin{aligned} \sum_{i=1}^n P(A_i) + \sum_{i=n+1}^{\infty} P(A_i) &= \sum_{i=1}^{\infty} P(A_i) \stackrel{\substack{\text{by infromal} \\ \text{definition 2} \\ \text{(point 3)}}}{=} P\left(\bigcup_{i=1}^{\infty} A_i\right) \\ &= P\left(\bigcup_{i=1}^n A_i \cup \bigcup_{i=n+1}^{\infty} A_i\right) \stackrel{\substack{\text{from empty} \\ \text{set property}}}{=} P\left(\bigcup_{i=1}^n A_i \cup \emptyset\right) = P\left(\bigcup_{i=1}^n A_i\right) \end{aligned}$$

**Problem 2:** Prove that  $P(A^c) = 1 - P(A)$

Proof:

$$\begin{aligned} 1 &\stackrel{\substack{\text{by infromal} \\ \text{definition 2} \\ \text{(point 2)}}}{=} P(\Omega) \stackrel{\substack{\text{definition of} \\ \text{compliment}}}{=} P(A^c \cup A) \stackrel{\substack{A^c \cap A = \emptyset \\ \text{from} \\ \text{fact 1}}}{=} P(A^c) + P(A) \\ &\rightarrow P(A^c) = 1 - P(A) \end{aligned}$$

**Problem 3:** Let  $X$  be a RV whose codomain is  $R$ , Find  $X^{-1}(B)$  and  $P(X^{-1}(B))$  where

- 1)  $B = R$
- 2)  $B = \emptyset$

Proof:

- 1)  $X^{-1}(B) = \{\omega \in \Omega : X(\omega) \in B\} = \{\omega \in \Omega : X(\omega) \in R\}$   
And because  $X: \Omega \rightarrow R$ , for every  $\omega \in \Omega$ ,  $X(\omega) \in R$ , and therefore:  
 $\{\omega \in \Omega : X(\omega) \in R\} = \Omega$

$$\text{And } P(X^{-1}(B)) = P(\Omega) \stackrel{\substack{\text{by infromal} \\ \text{definition 2} \\ \text{(point 2)}}}{=} 1$$

- 2)  $X^{-1}(B) = \{\omega \in \Omega : X(\omega) \in B\} = \{\omega \in \Omega : X(\omega) \in \emptyset\}$   
And because  $X: \Omega \rightarrow R$ , there is no  $\omega \in \Omega$  that  $X(\omega) \in \emptyset$ , and therefore:  
 $\{\omega \in \Omega : X(\omega) \in \emptyset\} = \emptyset$

$$\text{And } P(X^{-1}(B)) = P(\emptyset) \stackrel{\substack{\text{by infromal} \\ \text{definition 2} \\ \text{(point 1)}}}{=} 0$$

**Problem 4:** Is the following function,  $F: R \rightarrow [0,1]$ , a CDF of some RV?  $F(x) = \begin{cases} 0 & x \leq 42 \\ 1 & x > 42 \end{cases}$

Proof:

No, we will show that  $F(x)$  is not right-continuous.

A function is right-continuous at  $c$  if  $\lim_{x \rightarrow c^+} F(x) = F(c)$

$$\text{Therefore, } \lim_{x \rightarrow 42^+} F(x) = \lim_{x \rightarrow 42^+} \begin{cases} 0 & x \leq 42 \\ 1 & x > 42 \end{cases} = 1 \neq 0 = F(42)$$

Means  $F(x)$  does not holds right-continuous property of CDF  $\rightarrow F$  is not CDF.

**Problem 5:** Let  $X$  be a continuous  $n$ -dimensional RV.

- 1) Let  $x \in R^n$ . Find  $P(X = x) = P(X^{-1}(x))$ .
- 2) Give an example for  $B \subseteq R^n$  such that  $B$  contains at least one element of  $R^n$  and  $P(X^{-1}(B)) = 0$ .
- 3) Let  $\{B_i\}_{i=1}^{\infty}$  be a countable collection of nonempty pairwise disjoint subsets of  $R^n$  (i.e.,  $B_i \neq \emptyset$  for every  $i$  and  $B_i \cap B_j = \emptyset$  whenever  $i \neq j$ ). Give an example for such a collection where, in addition,

$$P(X \in \bigcup_{i=1}^{\infty} B_i) = P(X^{-1}(\bigcup_{i=1}^{\infty} B_i)) \text{ Is zero.}$$

Proof:

- 1)  $P(X = x) = P(x \leq X \leq x)$

Let us denote  $x = [x_1 \dots x_n]$ ,  $X = [X_1 \dots X_n]$  now:

$$\begin{aligned} P(x \leq X \leq x) &= P(x_1 \leq X_1 \leq x_1, \dots, x_n \leq X_n \leq x_n) = \lim_{\forall i: x'_i \rightarrow x_i^-} (P(x'_1 < X_1 \leq x_1, \dots, x'_n < X_n \leq x_n)) \\ &= \lim_{\forall i: x'_i \rightarrow x_i^-} (P(X_1 \leq x_1, \dots, X_n \leq x_n) - P(X_1 \leq x'_1, \dots, X_n \leq x'_n)) \\ &= P(X_1 \leq x_1, \dots, X_n \leq x_n) - \lim_{\forall i: x'_i \rightarrow x_i^-} (P(X_1 \leq x'_1, \dots, X_n \leq x'_n)) \\ &= P(X \leq x) - \lim_{x' \rightarrow x^-} (P(X \leq x')) = F(x) - \lim_{x' \rightarrow x^-} F(x') \end{aligned}$$

Because  $X$  is continuous RV then the CDF  $F(x)$  is continuous (definition 9). Thus,  $\lim_{x' \rightarrow x^-} F(x') = F(x)$

Now,  $F(x) - \lim_{x' \rightarrow x^-} F(x') = F(x) - F(x) = 0$ .


- 2) For  $B = \{[1 \dots n]\} \in R^n$ , denote  $x = [1 \dots n]$ , therefore:  $P(X^{-1}(B)) = P(X^{-1}(x)) = P(X = x) \stackrel{\text{from section 1}}{=} 0$

- 3) For  $B_i = \left\{ \left[ \underbrace{i \dots i}_{n \text{ entries}} \right] \right\} \subset R^n$ , denote  $x_i = \left[ \underbrace{i \dots i}_{n \text{ entries}} \right]$ , therefore:  
 $(*) P(X^{-1}(B_i)) = P(X^{-1}(x_i)) = P(X = x_i) \stackrel{\text{from section 1}}{=} 0$

We can notice that for every  $i \neq j$ :  $B_i \neq \emptyset$  and  $B_i \cap B_j = \emptyset$ . Now, for:  $P(X \in \bigcup_{i=1}^{\infty} B_i)$  there exist  $k$  such that  $X \in B_k$ , and therefore  $X = x_k = [k, \dots, k]$ , so:  $P(X \in \bigcup_{i=1}^{\infty} B_i) = P(X = x_k) \stackrel{\text{from (*)}}{=} 0$

**Problem 6:** Let  $X: \Omega \rightarrow R$  stand for the result of rolling a fair die. Thus, for  $i = 1, \dots, 6$ ,  $a_i = i$  and  $p_i = \frac{1}{6}$  and Draw  $F(X)$  and  $p(X)$ .

Proof:



**Problem 7:** Write a similar expression for  $p(x_3, x_4, x_5)$ .

Proof:

$$\begin{aligned} p(x_3, x_4, x_5) &= \sum_{x_1, x_2} p(x) = \sum_{x_1} \sum_{x_2} p(x) = \sum_{x_1} (p(x_1, -1, x_3, x_4, x_5) + p(x_1, 1, x_3, x_4, x_5)) \\ &= p(-1, -1, x_3, x_4, x_5) + p(-1, 1, x_3, x_4, x_5) + p(1, -1, x_3, x_4, x_5) + p(1, 1, x_3, x_4, x_5) \end{aligned}$$

**Problem 8:** Let  $X$  stand for the result of rolling a fair die (i.e.,  $X$  is distributed uniformly on  $\{1, 2, 3, 4, 5, 6\}$ ). Show that  $E(X) = 3.5$ .

Proof:

$$\sum_x xp(x) = 1 * \frac{1}{6} + 2 * \frac{1}{6} + 3 * \frac{1}{6} + 4 * \frac{1}{6} + 5 * \frac{1}{6} + 6 * \frac{1}{6} = 3.5$$

**Problem 9:** let  $x \in R^n$ . Then the  $n \times n$  matrix  $xx^T$  is symmetric; namely,  $xx^T = (xx^T)^T$ .

Proof:  $(xx^T)^T = (x^T)^T x^T = xx^T$

**Problem 10:** Show that  $E(X^T) = (E(X))^T$ .

Proof:

$$(E(X))^T = \left( E \left( \begin{bmatrix} X_1 \\ \vdots \\ X_n \end{bmatrix} \right) \right)^T = \left( \begin{bmatrix} E(X_1) \\ \vdots \\ E(X_n) \end{bmatrix} \right)^T = [E(X_1) \dots E(X_n)] = E([X_1 \dots X_n]) = E \left( \left( \begin{bmatrix} X_1 \\ \vdots \\ X_n \end{bmatrix} \right)^T \right) = E(X^T)$$

**Problem 11:** Show that  $(R_x)_{i,j}$  (i.e., the element in the  $i$ -th row and  $j$ -th column of  $R_x$ ) is  $E(X_i X_j)$

Proof:

Denote  $X = \begin{bmatrix} X_1 \\ X_2 \\ \vdots \\ X_n \end{bmatrix} \in R^n$  RV (discrete or continuous – we will see it results in the same ☺)

(\*)  $g: R^n \rightarrow R^{n \times n}$  such that  $g(X) = XX^T$

$$\begin{aligned} R_x = E(XX^T) &\stackrel{(*)}{=} E(g(X)) = E \left( \begin{bmatrix} X_1^2 & \dots & X_1 X_n \\ \vdots & \ddots & \vdots \\ X_n X_1 & \dots & X_n^2 \end{bmatrix} \right) \stackrel{\text{Definition 23,24-}}{\equiv} \begin{bmatrix} E(X_1^2) & \dots & E(X_1 X_n) \\ \vdots & \ddots & \vdots \\ E(X_n X_1) & \dots & E(X_n^2) \end{bmatrix} \\ &\rightarrow (R_x)_{i,j} = E(X_i X_j) \end{aligned}$$

**Problem 12:** Show that  $R_x$  is symmetric, i.e., show that  $R_x = R_x^T$ .

Proof: We will show that  $E(X_i X_j) = E(X_j X_i)$

For  $X_i, X_j \in R^n$  discrete RVs:

$$E(X_i X_j) = \sum_{x_i} \sum_{x_j} x_i x_j p(x_i, x_j) = \sum_{x_j} \sum_{x_i} x_j x_i p(x_j, x_i) = E(X_j X_i)$$

For  $X_i, X_j \in R^n$  continuous RVs:

$$E(X_i X_j) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} x_i x_j p(x_i, x_j) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} x_j x_i p(x_j, x_i) = E(X_j X_i)$$

Now, from as we saw in problem 11:

$$R_x = \begin{bmatrix} E(X_1^2) & \dots & E(X_1 X_n) \\ \vdots & \ddots & \vdots \\ E(X_n X_1) & \dots & E(X_n^2) \end{bmatrix} \stackrel{E(X_i X_j) = E(X_j X_i)}{\equiv} \begin{bmatrix} E(X_1^2) & \dots & E(X_n X_1) \\ \vdots & \ddots & \vdots \\ E(X_1 X_n) & \dots & E(X_n^2) \end{bmatrix} = R_x^T$$



**Problem 13:** Show that  $(\Sigma_x)_{i,j}$  (i.e., the element in the  $i$ -th row and  $j$ -th column of  $\Sigma_x$ ) is

$$E\left((X_i - E(X_i))(X_j - E(X_j))\right)$$

Proof:

Similar to problem 11, denote  $X = \begin{bmatrix} X_1 \\ X_2 \\ \vdots \\ X_n \end{bmatrix} \in R^n$  RV.

(\*)  $g: R^n \rightarrow R^{n \times n}$  such that  $g(X) = (X - E(X))(X - E(X))^T$

$$\Sigma_x = E\left((X - E(X))(X - E(X))^T\right) \stackrel{(*)}{=} E(g(X))$$

$$\begin{aligned} &= E\left(\begin{bmatrix} (X_1 - E(X_1))^2 & \dots & (X_1 - E(X_1))(X_n - E(X_n)) \\ \vdots & \ddots & \vdots \\ (X_n - E(X_n))(X_1 - E(X_1)) & \dots & (X_n - E(X_n))^2 \end{bmatrix}\right) \\ &\stackrel{\text{Definition 23,24-}}{=} \begin{bmatrix} E((X_1 - E(X_1))^2) & \dots & E((X_1 - E(X_1))(X_n - E(X_n))) \\ \vdots & \ddots & \vdots \\ E((X_n - E(X_n))(X_1 - E(X_1))) & \dots & E((X_n - E(X_n))^2) \end{bmatrix} \\ &\rightarrow (\Sigma_x)_{i,j} = E\left((X_i - E(X_i))(X_j - E(X_j))\right) \end{aligned}$$

**Problem 14:** Show that  $(\Sigma_x)_{i,i}$  is the variance of  $X_i$ .

Proof: From definition,  $V(X_i) = E((X_i - E(X_i))^2)$

From problem 13, we obtain:

$$(\Sigma_x)_{i,i} = E((X_i - E(X_i))(X_i - E(X_i))) = E((X_i - E(X_i))^2) = V(X_i)$$

**Problem 15:** Show that  $\Sigma_x$  is symmetric, i.e., show that  $\Sigma_x = \Sigma_x^T$ .

Proof: We will show that  $E((X_i - E(X_i))(X_j - E(X_j))) = E((X_j - E(X_j))(X_i - E(X_i)))$

Denote  $X_i - E(X_i) = Y_i$ ,  $X_j - E(X_j) = Y_j$ , hence we need to prove:

$$E(Y_i Y_j) = E(Y_j Y_i)$$

But we already proved it in problem 12 for discrete and continuous RVs

$$\rightarrow (*) E((X_i - E(X_i))(X_j - E(X_j))) = E((X_j - E(X_j))(X_i - E(X_i)))$$

Now from problem 13:

$$\begin{aligned} \Sigma_x &= \begin{bmatrix} E((X_1 - E(X_1))^2) & \dots & E((X_1 - E(X_1))(X_n - E(X_n))) \\ \vdots & \ddots & \vdots \\ E((X_n - E(X_n))(X_1 - E(X_1))) & \dots & E((X_n - E(X_n))^2) \end{bmatrix} \stackrel{(*)}{=} \\ &\begin{bmatrix} E((X_1 - E(X_1))^2) & \dots & E((X_n - E(X_n))(X_1 - E(X_1))) \\ \vdots & \ddots & \vdots \\ E((X_1 - E(X_1))(X_n - E(X_n))) & \dots & E((X_n - E(X_n))^2) \end{bmatrix} = \Sigma_x^T \end{aligned}$$

**Problem 16:** Prove:

1.  $\Sigma = E(XX^T) - \mu\mu^T$
2.  $\Sigma = E((X - \mu)X^T)$
3.  $\Sigma = E(X(X - \mu)^T)$

Proof:

1.  $\Sigma \stackrel{\text{Definition}}{=} E\left((X - E(X))(X - E(X))^T\right) \stackrel{E(X)=\mu}{=} E((X - \mu)(X - \mu)^T) = E((X - \mu)(X^T - \mu^T))$   
 $= E(XX^T - X\mu^T - \mu X^T + \mu\mu^T) \stackrel{\text{Linearity of Expectation}}{=} E(XX^T) - E(X\mu^T) - E(\mu X^T) + E(\mu\mu^T)$   
 $\stackrel{\mu, \mu^T \text{ are constant vectors}}{=} E(XX^T) - \mu^T E(X) - \mu E(X^T) + \mu\mu^T \stackrel{E(X)=\mu}{=} E(XX^T) - \mu^T \mu - \mu E(X)^T + \mu\mu^T$   
 $\stackrel{E(X^T)=E(X)^T}{=} E(XX^T) - \mu^T \mu - \mu\mu^T + \mu\mu^T = E(XX^T) - \mu^T \mu$
2.  $\Sigma \stackrel{\text{Definition}}{=} E\left((X - E(X))(X - E(X))^T\right) \stackrel{E(X)=\mu}{=} E((X - \mu)(X - \mu)^T) = E((X - \mu)(X^T - \mu^T)) =$   
 $E((X - \mu)X^T - (X - \mu)\mu^T) \stackrel{\text{Linearity of Expectation}}{=} E((X - \mu)X^T) - E((X - \mu)\mu^T)$   
 $= E((X - \mu)X^T) - \mu^T E(X - \mu) \stackrel{\text{Linearity of Expectation}}{=} E((X - \mu)X^T) - \mu^T (E(X) - E(\mu))$   
 $= E((X - \mu)X^T) - \mu^T (\mu - \mu) = E((X - \mu)X^T)$
3.  $\Sigma \stackrel{\text{Definition}}{=} E\left((X - E(X))(X - E(X))^T\right) \stackrel{E(X)=\mu}{=} E((X - \mu)(X - \mu)^T) = E((X - \mu)(X^T - \mu^T)) =$   
 $E(X(X^T - \mu^T) - \mu(X^T - \mu^T)) \stackrel{\text{Linearity of Expectation}}{=} E(X(X^T - \mu^T)) - E(\mu(X^T - \mu^T))$   
 $\stackrel{\mu \text{ is a constant vector}}{=} E(X(X^T - \mu^T)) - \mu E(X^T - \mu^T) = E(X(X^T - \mu^T)) - \mu(E(X^T) - \mu^T) \stackrel{E(X^T)=E(X)^T}{=} E(X(X^T - \mu^T)) - \mu(E(X)^T - \mu^T)$   
 $= E(X(X^T - \mu^T)) - \mu(\mu^T - \mu^T) = E(X(X^T - \mu^T))$

**Problem 17:** Prove:

1.  $R_{YX} = R_{XY}^T$
2.  $\Sigma_{YX} = \Sigma_{XY}^T$
3.  $E\left((X - E(X))(Y - E(Y))^T\right) = R_{XY} - \mu_X \mu_Y^T$

Proof:

1.  $R_{XY}^T = E(XY^T)^T = E((XY^T)^T) = E(YX^T) = R_{YX}$
2.  $\Sigma_{XY}^T = E\left((X - E(X))(Y - E(Y))^T\right)^T = E\left(\left((X - E(X))(Y - E(Y))^T\right)^T\right)$   
 $= E\left((Y - E(Y))(X - E(X))^T\right) = \Sigma_{YX}$
3.  $E\left((X - E(X))(Y - E(Y))^T\right) = E((X - \mu_X)(Y - \mu_Y)^T) = E((X - \mu_X)(Y^T - \mu_Y^T))$   
 $= E(XY^T - X\mu_Y^T - \mu_X Y^T + \mu_X \mu_Y^T) = E(XY^T) - E(X\mu_Y^T) - E(\mu_X Y^T) + E(\mu_X \mu_Y^T)$   
 $= R_{XY} - \mu_X \mu_Y^T - \mu_X \mu_Y^T + \mu_X \mu_Y^T = R_{XY} - \mu_X \mu_Y^T$

**Problem 18:** Let  $Y = AX + b$  affine transformation. Prove:

1.  $\mu_Y = A\mu_X + b$
2.  $\Sigma_Y = A\Sigma_X A^T$

Proof:

$$\begin{aligned}
 1. \quad \mu_Y &= E(AX + b) = E(AX) + E(b) \stackrel{\substack{A, b \text{ are constants}}}{=} AE(X) + b = A\mu_X + b \\
 2. \quad \Sigma_Y &= E((Y - \mu_Y)(Y - \mu_Y)^T) = E((AX + b - \mu_Y)(AX + b - \mu_Y)^T) \stackrel{1.}{=} \\
 &= E((AX + b - A\mu_X - b)(AX + b - A\mu_X - b)^T) = E(A(X - \mu_X)(X - \mu_X)^T) \\
 &= E(A(X - \mu_X)(X - \mu_X)^T A^T) \stackrel{A \text{ is a constant}}{=} AE((X - \mu_X)(X - \mu_X)^T)A^T = A\Sigma_X A^T
 \end{aligned}$$

**Problem 19:** Let  $X = [X_1 \dots X_n]^T$  be an  $n$ -dimensional RV with mean  $\mu$  and covariance  $\Sigma = \sigma^2 I$ .  
Find:  $V(\sum_{i=1}^n X_i)$

Proof:

Denote:  $A = 1^T, b = 0$

$\rightarrow Y = AX + b = 1^T X$  the affine transformation of  $X$ .

From fact 10:

$$\begin{aligned}
 \Sigma_Y &= A\Sigma_X A^T = 1^T \sigma^2 I (1^T)^T = 1^T \sigma^2 I 1 = 1^T \sigma^2 1 = \sigma^2 1^T 1 \\
 &= \sigma^2 \sum_{i=1}^n 1 = \sigma^2 \mathbf{n} \in R^{1 \times 1} = V(Y) = V(1^T X) = V\left(\sum_{i=1}^n X_i\right)
 \end{aligned}$$

**Problem 20:**  $A \perp\!\!\!\perp B|C \leftrightarrow p(A|B, C) = p(A|C) \leftrightarrow p(B|A, C) = p(B|C)$

Proof:

We'll prove:

- 1)  $A \perp\!\!\!\perp B|C \leftrightarrow p(A|B, C) = p(A|C)$
- 2)  $A \perp\!\!\!\perp B|C \leftrightarrow p(B|A, C) = p(B|C)$

$$\begin{aligned}
 1) \quad A \perp\!\!\!\perp B|C &\stackrel{\text{definition 31}}{\Leftrightarrow} P(A \cap B|C) = P(A|C)P(B|C) \stackrel{\substack{\text{Bayes Theorem on} \\ P(A \cap B|C), P(B|C)}}{\Leftrightarrow} \frac{P(A \cap B \cap C)}{P(C)} = P(A|C) \frac{P(B \cap C)}{P(C)} \\
 &\stackrel{\text{multiply by } P(C)}{\Leftrightarrow} P(A \cap B \cap C) = P(A|C)P(B \cap C) \\
 &\stackrel{\substack{\text{divide by } P(B \cap C) > 0}}{\Leftrightarrow} \frac{P(A \cap B \cap C)}{P(B \cap C)} = P(A|C) \stackrel{\substack{\text{Bayes Theorem on} \\ P(A|B, C)}}{\Leftrightarrow} P(A|B, C) = P(A|C)
 \end{aligned}$$

$$\begin{aligned}
 2) \quad A \perp\!\!\!\perp B|C &\stackrel{\text{definition 31}}{\Leftrightarrow} P(A \cap B|C) = P(A|C)P(B|C) \stackrel{\substack{\text{Bayes Theorem on} \\ P(A \cap B|C), P(A|C)}}{\Leftrightarrow} \frac{P(A \cap B \cap C)}{P(C)} = \frac{P(A \cap C)}{P(C)} P(B|C) \\
 &\stackrel{\text{multiply by } P(C)}{\Leftrightarrow} P(A \cap B \cap C) = P(A \cap C)P(B|C) \\
 &\stackrel{\substack{\text{divide by } P(A \cap C) > 0}}{\Leftrightarrow} \frac{P(B \cap A \cap C)}{P(A \cap C)} = P(B|C) \stackrel{\substack{\text{Bayes Theorem on} \\ P(B|A, C)}}{\Leftrightarrow} P(B|A, C) = P(B|C)
 \end{aligned}$$

**Problem 21:** Show that  $E(\mathbb{1}_A) = P(A)$

Proof:

$$E(\mathbb{1}_A) = 1 * P(\mathbb{1}_A = 1) + 0 * p(\mathbb{1}_A = 0) = 1 * P(\omega \in A) + 0 * p(\omega \notin A) = P(\omega \in A) = P(A)$$

**Problem 22:** prove Fact 13:

- 1) If  $X \in R^n$  and  $Y \in R^m$  are orthogonal RVs, then the correlation matrix of  $Z = [X^T Y^T]^T$  is  $\begin{bmatrix} R_X & 0_{n \times m} \\ 0_{m \times n} & R_Y \end{bmatrix}$
- 2) If  $X \in R^n$  and  $Y \in R^m$  are uncorrelated RVs, then the covariance matrix of  $Z = [X^T Y^T]^T$  is  $\begin{bmatrix} \Sigma_X & 0_{n \times m} \\ 0_{m \times n} & \Sigma_Y \end{bmatrix}$

Proof:

- 1)  $X \in R^n$  and  $Y \in R^m$  are orthogonal RVs, hence from definition 37:

$$E(XY^T) = 0_{n \times m}$$

Recall from definition 28 that

$$\begin{aligned} E(XY^T) &= R_{XY}, & R_{YX} &= R_{XY}^T \\ \rightarrow R_{XY} &= 0_{n \times m}, & R_{YX} &= 0_{m \times n} \end{aligned}$$

$$\begin{aligned} \text{Finally, we saw in fact 9 that the correlation matrix of } Z \text{ is } R_Z &= \begin{bmatrix} R_X & R_{XY} \\ R_{YX} & R_Y \end{bmatrix} \\ \rightarrow R_Z &= \begin{bmatrix} R_X & R_{XY} \\ R_{YX} & R_Y \end{bmatrix} = \begin{bmatrix} R_X & 0_{n \times m} \\ 0_{m \times n} & R_Y \end{bmatrix} \end{aligned}$$

- 2)  $X \in R^n$  and  $Y \in R^m$  are uncorrelated RVs, hence from definition 38:  $X - \mu_X$  and  $Y - \mu_Y$  are orthogonal, and again from definition 37 we obtain:

$$E((X - \mu_X)(Y - \mu_Y)^T) = 0_{n \times m}$$

Recall from definition 28 and problem 17 that

$$\begin{aligned} E((X - \mu_X)(Y - \mu_Y)^T) &= \Sigma_{XY}, & \Sigma_{YX} &= \Sigma_{XY}^T \\ \rightarrow \Sigma_{XY} &= 0_{n \times m}, & \Sigma_{YX} &= 0_{m \times n} \end{aligned}$$

$$\begin{aligned} \text{Finally, we saw in fact 9 that the covariance matrix of } Z \text{ is } \Sigma_Z &= \begin{bmatrix} \Sigma_X & \Sigma_{XY} \\ \Sigma_{YX} & \Sigma_Y \end{bmatrix} \\ \rightarrow \Sigma_Z &= \begin{bmatrix} \Sigma_X & \Sigma_{XY} \\ \Sigma_{YX} & \Sigma_Y \end{bmatrix} = \begin{bmatrix} \Sigma_X & 0_{n \times m} \\ 0_{m \times n} & \Sigma_Y \end{bmatrix} \end{aligned}$$

**Problem 23:** prove Fact 14:

$$X \perp\!\!\!\perp Y \rightarrow X \perp Y$$

Proof:

X and Y are independent RVs, hence by definition  $p(x, y) = p(x)p(y) \forall x, y$ .

We need to show that X and Y are uncorrelated, meaning:  $E(XY^T) = \mu_X \mu_Y^T$

For  $X, Y \in R^n$  discrete RVs:

$$E(XY^T) = \sum_x \sum_y xy^T p(x, y) = \sum_x \sum_y xy^T p(x)p(y) = \sum_x xp(x) \sum_y y^T p(y) = \mu_X \mu_Y^T$$

For  $X, Y \in R^n$  continuous RVs:

$$E(XY^T) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xy^T p(x, y) dx dy = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xy^T p(x)p(y) dx dy = \int_{-\infty}^{\infty} xp(x) dx \int_{-\infty}^{\infty} y^T p(y) dy = \mu_X \mu_Y^T$$

**Problem 24:** Let  $X = [X_1 \ X_2]^T$  be a two-dimensional random vector taking values in  $S = \{0,1\}^2$ . The probability mass function (pmf) of  $X$  is given by:

$$\begin{aligned} p(0,0) &= 0.5 \\ p(0,1) &= 0.1 \\ p(1,0) &= 0.3 \\ p(1,1) &= 0.1 \end{aligned}$$

- (a) Find the marginals,  $p(X_1), p(X_2)$ .
- (b) Find the 2D mean vector,  $E(X)$ .
- (c) Find the 2-by-2 correlation matrix,  $E(XX^T)$ .
- (d) Find the 2-by-2 covariance matrix,  $E((X - \mu)(X - \mu)^T)$ ,  $\mu = E(X)$ .
- (e) Are  $X_1$  and  $X_2$  independent?
- (f) Are  $X_1$  and  $X_2$  correlated?

Proof:

(a)

$$p(X_1 = 0) = \sum_{x_2 \in \{0,1\}} p(0, x_2) = p(0,0) + p(0,1) = 0.5 + 0.1 = 0.6$$

$$p(X_1 = 1) = \sum_{x_2 \in \{0,1\}} p(1, x_2) = p(1,0) + p(1,1) = 0.3 + 0.1 = 0.4$$

$$\rightarrow p(x_1) = \begin{cases} 0.6, & x_1 = 0 \\ 0.4, & x_1 = 1 \\ 0, & \text{otherwise} \end{cases}$$

$$p(X_2 = 0) = \sum_{x_1 \in \{0,1\}} p(x_1, 0) = p(0,0) + p(1,0) = 0.5 + 0.3 = 0.8$$

$$p(X_2 = 1) = \sum_{x_1 \in \{0,1\}} p(x_1, 1) = p(0,1) + p(1,1) = 0.1 + 0.1 = 0.2$$

$$\rightarrow p(x_2) = \begin{cases} 0.8, & x_2 = 0 \\ 0.2, & x_2 = 1 \\ 0, & \text{otherwise} \end{cases}$$

$$(b) \ E(X) = E([X_1 \ X_2]^T) = \begin{bmatrix} E(X_1) \\ E(X_2) \end{bmatrix} = \begin{bmatrix} 0 * p(X_1 = 0) + 1 * p(X_1 = 1) \\ 0 * p(X_2 = 0) + 1 * p(X_2 = 1) \end{bmatrix} = \begin{bmatrix} p(X_1 = 1) \\ p(X_2 = 1) \end{bmatrix} = \begin{bmatrix} 0.4 \\ 0.2 \end{bmatrix}$$

$$(c) \ E(XX^T) = \sum_x x x^T p(x) = 0.5 * \begin{bmatrix} 0 \\ 0 \end{bmatrix} \begin{bmatrix} 0 & 0 \end{bmatrix} + 0.1 * \begin{bmatrix} 0 \\ 1 \end{bmatrix} \begin{bmatrix} 0 & 1 \end{bmatrix} + 0.3 * \begin{bmatrix} 1 \\ 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \end{bmatrix} + 0.1 * \begin{bmatrix} 1 \\ 1 \end{bmatrix} \begin{bmatrix} 1 & 1 \end{bmatrix} = \\ 0.5 * \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} + 0.1 * \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} + 0.3 * \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + 0.1 * \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 0.3 + 0.1 & 0.1 \\ 0.1 & 0.1 + 0.1 \end{bmatrix} = \begin{bmatrix} 0.4 & 0.1 \\ 0.1 & 0.2 \end{bmatrix}$$

$$(d) \ E((X - \mu)(X - \mu)^T) = E(XX^T) - \mu\mu^T = \begin{bmatrix} 0.4 & 0.1 \\ 0.1 & 0.2 \end{bmatrix} - \begin{bmatrix} 0.4 \\ 0.2 \end{bmatrix} \begin{bmatrix} 0.4 & 0.2 \end{bmatrix} = \begin{bmatrix} 0.4 & 0.1 \\ 0.1 & 0.2 \end{bmatrix} - \begin{bmatrix} 0.16 & 0.08 \\ 0.08 & 0.04 \end{bmatrix} = \\ \begin{bmatrix} 0.24 & -0.7 \\ -0.7 & -0.2 \end{bmatrix}$$

$$(e) \ X_1 \text{ and } X_2 \text{ are not independent since } p(0,0) = 0.5 \neq 0.48 = 0.3 * 0.8 = p(X_1 = 0) * p(X_2 = 0)$$

$$(f) \ E(X_1 X_2) = 0 * 0 * p(0,0) + 0 * 1 * p(0,1) + 1 * 0 * p(1,0) + 1 * 1 * p(1,1) = p(1,1) = 0.1$$

$$E(X_1 X_2) - E(X_1)E(X_2) = 0.1 - 0.4 * 0.2 = 0.02 \neq 0$$

Meaning,  $X_1$  and  $X_2$  are not uncorrelated  $\rightarrow X_1$  and  $X_2$  are correlated.

**Problem 25:** Let  $X = [X_1 \ X_2]^T$  be a two-dimensional random vector taking values in  $S = \{0,1\}^2$ . The probability mass function (pmf) of  $X$  is given by:

$$\begin{aligned} p(0,0) &= 0.25 \\ p(0,1) &= 0.25 \\ p(1,0) &= 0.25 \\ p(1,1) &= 0.25 \end{aligned}$$

- (a) Find the marginals,  $p(X_1), p(X_2)$ .
- (b) Find the 2D mean vector,  $E(X)$ .
- (c) Find the 2-by-2 covariance matrix,  $E((X - \mu)(X - \mu)^T)$ ,  $\mu = E(X)$ .
- (d) Are  $X_1$  and  $X_2$  independent?
- (e) Are  $X_1$  and  $X_2$  uncorrelated?

Let  $Y_1 = X_1 + X_2$  and let  $Y_2 = |X_1 - X_2|$

- (f) Find the pmf of  $Y = [Y_1 \ Y_2]^T$ .
- (g) Find the marginal pmf of  $Y_1$  and the marginal pmf of  $Y_2$ .
- (h) Find the 2D mean vector of  $Y$ ,  $E(Y)$ .
- (i) Find the 2-by-2 covariance matrix of  $Y$ ,  $E((Y - \mu_Y)(Y - \mu_Y)^T)$ ,  $\mu_Y = E(Y)$ .
- (j) Are  $Y_1$  and  $Y_2$  independent?
- (k) Are  $Y_1$  and  $Y_2$  uncorrelated?

Proof:

(a)

$$\begin{aligned} p(X_1 = 0) &= \sum_{x_2 \in \{0,1\}} p(0, x_2) = p(0,0) + p(0,1) = 0.25 + 0.25 = 0.5 \\ p(X_1 = 1) &= \sum_{x_2 \in \{0,1\}} p(1, x_2) = p(1,0) + p(1,1) = 0.25 + 0.25 = 0.5 \end{aligned}$$

$$\rightarrow p(x_1) = \begin{cases} 0.5, & x_1 = 0 \\ 0.5, & x_1 = 1 \\ 0, & \text{otherwise} \end{cases}$$

Similarly, we will obtain:

$$\rightarrow p(x_2) = \begin{cases} 0.5, & x_2 = 0 \\ 0.5, & x_2 = 1 \\ 0, & \text{otherwise} \end{cases}$$

$$(b) \ E(X) = E([X_1 \ X_2]^T) = \begin{bmatrix} E(X_1) \\ E(X_2) \end{bmatrix} = \begin{bmatrix} 0 * p(X_1 = 0) + 1 * p(X_1 = 1) \\ 0 * p(X_2 = 0) + 1 * p(X_2 = 1) \end{bmatrix} = \begin{bmatrix} p(X_1 = 1) \\ p(X_2 = 1) \end{bmatrix} = \begin{bmatrix} 0.5 \\ 0.5 \end{bmatrix}$$

$$\begin{aligned} (c) \ \Sigma_X &= E(XX^T) - E(X)E(X^T) = \sum_x xx^T p(x) - (\sum_x xp(x)) * (\sum_x x^T p(x)) = 0.25 * \begin{bmatrix} 0 \\ 0 \end{bmatrix} \begin{bmatrix} 0 & 0 \end{bmatrix} + 0.25 * \\ &\begin{bmatrix} 0 \\ 1 \end{bmatrix} \begin{bmatrix} 0 & 1 \end{bmatrix} + 0.25 * \begin{bmatrix} 1 \\ 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \end{bmatrix} + 0.25 * \begin{bmatrix} 1 \\ 1 \end{bmatrix} \begin{bmatrix} 1 & 1 \end{bmatrix} - (0.25 * \begin{bmatrix} 0 \\ 0 \end{bmatrix} + 0.25 * \begin{bmatrix} 0 \\ 1 \end{bmatrix} + 0.25 * \begin{bmatrix} 1 \\ 0 \end{bmatrix} + 0.25 * \\ &\begin{bmatrix} 1 \\ 1 \end{bmatrix}) * (0.25 * \begin{bmatrix} 0 & 0 \end{bmatrix} + 0.25 * \begin{bmatrix} 0 & 1 \end{bmatrix} + 0.25 * \begin{bmatrix} 1 & 0 \end{bmatrix} + 0.25 * \begin{bmatrix} 1 & 1 \end{bmatrix}) = 0.25 * \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} + 0.25 * \\ &\begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} + 0.25 * \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + 0.25 * \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} - (\begin{bmatrix} 0 \\ 0.25 \end{bmatrix} + \begin{bmatrix} 0.25 \\ 0 \end{bmatrix} + \begin{bmatrix} 0.25 \\ 0.25 \end{bmatrix}) * (\begin{bmatrix} 0 & 0.25 \end{bmatrix} + \begin{bmatrix} 0.25 & 0 \end{bmatrix} + \\ &\begin{bmatrix} 0.25 & 0.25 \end{bmatrix}) = 0.25 * \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix} - \begin{bmatrix} 0.5 \\ 0.5 \end{bmatrix} * \begin{bmatrix} 0.5 & 0.5 \end{bmatrix} = \begin{bmatrix} 0.5 & 0.25 \\ 0.25 & 0.5 \end{bmatrix} - \begin{bmatrix} 0.25 & 0.25 \\ 0.25 & 0.25 \end{bmatrix} = \begin{bmatrix} 0.25 & 0 \\ 0 & 0.25 \end{bmatrix} \end{aligned}$$

(d)  $X_1$  and  $X_2$  are independent since

$$\begin{aligned} p(0,0) &= 0.25 = 0.5 * 0.5 = p(X_1 = 0) * p(X_2 = 0) \\ p(0,1) &= 0.25 = 0.5 * 0.5 = p(X_1 = 0) * p(X_2 = 1) \\ p(1,0) &= 0.25 = 0.5 * 0.5 = p(X_1 = 1) * p(X_2 = 0) \\ p(1,1) &= 0.25 = 0.5 * 0.5 = p(X_1 = 1) * p(X_2 = 1) \end{aligned}$$

$$(e) \ E(X_1 X_2) = 0 * 0 * p(0,0) + 0 * 1 * p(0,1) + 1 * 0 * p(1,0) + 1 * 1 * p(1,1) = p(1,1) = 0.25$$

$$E(X_1 X_2) - E(X_1)E(X_2) = 0.25 - 0.5 * 0.5 = 0$$

Meaning,  $X_1$  and  $X_2$  are uncorrelated.

(f) We can notice that  $Y_1 \in \{0,1,2\}$  and  $Y_2 \in \{0,1\}$

$$\begin{aligned}
 p(0,0) &= p(Y_1 = 0, Y_2 = 0) = p(X_1 + X_2 = 0, |X_1 - X_2| = 0) = p(X_1 = 0, X_2 = 0) = 0.25 \\
 p(0,1) &= p(Y_1 = 0, Y_2 = 1) = p(X_1 + X_2 = 0, |X_1 - X_2| = 1) = 0 \\
 p(1,0) &= p(Y_1 = 1, Y_2 = 0) = p(X_1 + X_2 = 1, |X_1 - X_2| = 0) = 0 \\
 p(1,1) &= p(Y_1 = 1, Y_2 = 1) = p(X_1 + X_2 = 1, |X_1 - X_2| = 1) \\
 &= p(X_1 = 1, X_2 = 0) + p(X_1 = 0, X_2 = 1) = 0.25 + 0.25 = 0.5 \\
 p(2,0) &= p(Y_1 = 2, Y_2 = 0) = p(X_1 + X_2 = 2, |X_1 - X_2| = 0) = p(X_1 = 1, X_2 = 1) = 0.25 \\
 p(2,1) &= p(Y_1 = 2, Y_2 = 1) = p(X_1 + X_2 = 2, |X_1 - X_2| = 1) = 0
 \end{aligned}$$

(g)

$$\begin{aligned}
 p(Y_1 = 0) &= \sum_{y_2 \in \{0,1\}} p(0, y_2) = p(0,0) + p(0,1) = 0.25 + 0 = 0.25 \\
 p(Y_1 = 1) &= \sum_{y_2 \in \{0,1\}} p(1, y_2) = p(1,0) + p(1,1) = 0 + 0.5 = 0.5 \\
 p(Y_1 = 2) &= \sum_{y_2 \in \{0,1\}} p(2, y_2) = p(2,0) + p(2,1) = 0.25 + 0 = 0.25 \\
 &\rightarrow p(y_1) = \begin{cases} 0.25, & y_1 = 0 \\ 0.5, & y_1 = 1 \\ 0.25, & y_1 = 2 \\ 0, & \text{otherwise} \end{cases}
 \end{aligned}$$

$$\begin{aligned}
 p(Y_2 = 0) &= \sum_{y_1 \in \{0,1,2\}} p(y_1, 0) = p(0,0) + p(1,0) + p(2,0) = 0.25 + 0 + 0.25 = 0.5 \\
 p(Y_2 = 1) &= \sum_{y_1 \in \{0,1,2\}} p(y_1, 1) = p(0,1) + p(1,1) + p(2,1) = 0 + 0.5 + 0 = 0.5 \\
 &\rightarrow p(y_2) = \begin{cases} 0.5, & y_2 = 0 \\ 0.5, & y_2 = 1 \\ 0, & \text{otherwise} \end{cases}
 \end{aligned}$$

(h)  $E(Y) = E([Y_1 \ Y_2]^T) = \begin{bmatrix} E(Y_1) \\ E(Y_2) \end{bmatrix} = \begin{bmatrix} 0 * p(Y_1 = 0) + 1 * p(Y_1 = 1) + 2 * p(Y_1 = 2) \\ 0 * p(Y_2 = 0) + 1 * p(Y_2 = 1) \end{bmatrix} =$

$$\begin{bmatrix} p(Y_1 = 1) + p(Y_1 = 2) \\ p(Y_2 = 1) \end{bmatrix} = \begin{bmatrix} 0.5 + 0.25 \\ 0.5 \end{bmatrix} = \begin{bmatrix} 0.75 \\ 0.5 \end{bmatrix}$$

(i)  $E(YY^T) - E(Y)E(Y^T) = \sum_y y y^T p(y) - (\sum_y y p(y)) * (\sum_y y^T p(y)) = 0.25 * \begin{bmatrix} 0 \\ 0 \end{bmatrix} [0 \ 0] + 0 * \begin{bmatrix} 0 \\ 1 \end{bmatrix} [0 \ 1] + 0 * \begin{bmatrix} 1 \\ 0 \end{bmatrix} [1 \ 0] + 0.5 * \begin{bmatrix} 1 \\ 1 \end{bmatrix} [1 \ 1] + 0.25 * \begin{bmatrix} 2 \\ 0 \end{bmatrix} [2 \ 0] + 0 * \begin{bmatrix} 2 \\ 1 \end{bmatrix} [2 \ 1] -$

$$\begin{aligned}
 & (0.25 * \begin{bmatrix} 0 \\ 0 \end{bmatrix} + 0 * \begin{bmatrix} 0 \\ 1 \end{bmatrix} + 0 * \begin{bmatrix} 1 \\ 0 \end{bmatrix} + 0.5 * \begin{bmatrix} 1 \\ 1 \end{bmatrix} + 0.25 * \begin{bmatrix} 2 \\ 0 \end{bmatrix} + 0 * \begin{bmatrix} 2 \\ 1 \end{bmatrix}) (0.25 * [0 \ 0] + 0 * [0 \ 1] + 0 * [1 \ 0] + 0.5 * [1 \ 1] + 0.25 * [2 \ 0] + 0 * [2 \ 1]) = 0.25 * \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} + 0.5 * \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} + 0.25 * \begin{bmatrix} 4 & 0 \\ 0 & 0 \end{bmatrix} - \\
 & \left( \begin{bmatrix} 0.5 \\ 0.5 \end{bmatrix} + \begin{bmatrix} 0.5 \\ 0 \end{bmatrix} \right) * ([0.5 \ 0.5] + [0.5 \ 0]) = \begin{bmatrix} 0.5 + 4 * 0.25 & 0.5 \\ 0.5 & 0.5 \end{bmatrix} - \left( \begin{bmatrix} 1 \\ 0.5 \end{bmatrix} * [1 \ 0.5] \right) = \begin{bmatrix} 1.5 & 0.5 \\ 0.5 & 0.5 \end{bmatrix} - \\
 & \begin{bmatrix} 1 & 0.5 \\ 0.5 & 0.25 \end{bmatrix} = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.25 \end{bmatrix}
 \end{aligned}$$

(j)  $Y_1$  and  $Y_2$  are not independent since  $p(0,0) = 0.25 \neq 0.125 = 0.25 * 0.5 = p(Y_1 = 0) * p(Y_2 = 0)$

(k)  $E(Y_1 Y_2) = 0 * 0 * p(0,0) + 1 * 0 * p(1,0) + 2 * 0 * p(2,0) + 0 * 1 * p(0,1) + 1 * 1 * p(1,1) + 2 * 1 * p(2,1) = p(1,1) + 2 * p(2,1) = 0.5 + 2 * 0 = 0.5$

$$E(Y_1 Y_2) - E(Y_1)E(Y_2) = 0.5 - 0.75 * 0.5 = 0.5 - 0.375 = 0.125 \neq 0$$

Meaning,  $X_1$  and  $X_2$  are not uncorrelated  $\rightarrow X_1$  and  $X_2$  are correlated.

**Problem 26:**  $X \perp\!\!\!\perp Y|Z \leftrightarrow p(x, y, z) = p(x|z)p(y|z)$

Proof: We'll prove:

- 1)  $X \perp\!\!\!\perp Y|Z \leftrightarrow p(x|y, z) = p(x|z)$
- 2)  $X \perp\!\!\!\perp Y|Z \leftrightarrow p(y|x, z) = p(y|z)$

Let us notice that:

$$\begin{aligned}
 (*) \quad p(x, y|z) &\stackrel{\text{Bayes Theorem}}{=} \frac{p(x, y, z)}{p(z)} \stackrel{\text{Definition 40}}{=} p(x|z)p(y|z) \stackrel{\text{multiply by } p(z)}{\leftrightarrow} p(x, y, z) = p(x|z)p(y|z)p(z) \\
 1) \quad X \perp\!\!\!\perp Y|Z \leftrightarrow p(x|y, z) &\stackrel{\text{Bayes Theorem}}{=} \frac{p(x, y, z)}{p(y, z)} \stackrel{\text{from } (*)}{=} \frac{p(x|z)p(y|z)p(z)}{p(y, z)} \stackrel{\text{Bayes Theorem}}{=} \frac{p(x|z)\frac{p(y, z)}{p(z)}p(z)}{p(y, z)} = p(x|z) \\
 2) \quad X \perp\!\!\!\perp Y|Z \leftrightarrow p(y|x, z) &\stackrel{\text{Bayes Theorem}}{=} \frac{p(x, y, z)}{p(x, z)} \stackrel{\text{from } (*)}{=} \frac{p(x|z)p(y|z)p(z)}{p(x, z)} \stackrel{\text{Bayes Theorem}}{=} \frac{\frac{p(x, z)}{p(z)}p(y|z)p(z)}{p(x, z)} = p(y|z)
 \end{aligned}$$

**Problem 27:** Let  $B$  be some bounded hyper-rectangle in  $R^3$ .

- 1) Let  $p(x, y, z) \propto \begin{cases} \exp(x + xz + yz) & \text{if } (x, y, z) \in B \\ 0 & \text{otherwise} \end{cases}$

Show that  $X \perp\!\!\!\perp Y|Z$

- 2) Let  $p(x, y, z) \propto \begin{cases} \exp(xyz) & \text{if } (x, y, z) \in B \\ 0 & \text{otherwise} \end{cases}$

Show that  $X \not\perp\!\!\!\perp Y|Z$

Proof:

- 1) For  $f(x, z) = \begin{cases} \exp(x + xz) & \text{if } (x, y, z) \in B \text{ for some } y \\ 0 & \text{otherwise} \end{cases}$   
and  $g(y, z) = \begin{cases} \exp(yz) & \text{if } (x, y, z) \in B \text{ for some } x \\ 0 & \text{otherwise} \end{cases}$   
we get that:

$$p(x, y, z) \propto \begin{cases} \exp(x + xz + yz) & \text{if } (x, y, z) \in B \\ 0 & \text{otherwise} \end{cases} = \begin{cases} f(x, z) * g(y, z) & \text{if } (x, y, z) \in B \\ 0 & \text{otherwise} \end{cases}$$

and now from Fact 18 we get that:  $X \perp\!\!\!\perp Y|Z$ .

- 2) For bounded hyper-rectangle  $B = \{(i, j, k): 0 \leq i, j, k \leq 1\}$  we can see that for  $x = 1, y = 0.5, z = 1$ :

$$\begin{aligned}
 p(x, y|z) &= \frac{p(x, y, z)}{p(z)} = \frac{p(x, y, z)}{\iint p(xyz) dx dy} \propto \frac{\exp(1 * 0.5 * 1)}{\iint \exp(xy * 1) dx dy} = \frac{\exp(0.5)}{\iint \exp(xy) dx dy} \approx \frac{1.6487}{1.3179} \approx 1.251 \\
 p(x|z)p(y|z) &= \frac{p(x, z)}{p(z)} * \frac{p(y, z)}{p(z)} = \frac{\int p(xyz) dy}{\iint p(xyz) dx dy} * \frac{\int p(xyz) dx}{\iint p(xyz) dx dy} \\
 &\propto \frac{\int \exp(1 * y * 1) dy}{\iint \exp(xy * 1) dx dy} * \frac{\int \exp(x * 0.5 * 1) dx}{\iint \exp(xy * 1) dx dy} = \frac{\int \exp(y) dy}{\iint \exp(xy) dx dy} * \frac{\int \exp(0.5x) dx}{\iint \exp(xy) dx dy} \\
 &= \frac{\int \exp(y) dy}{\iint \exp(xy) dx dy} * \frac{\int \exp(0.5x) dx}{\iint \exp(xy) dx dy} \approx \frac{(e - 1)}{1.3179} * \frac{2\sqrt{e} - 2}{1.3179} \approx 1.283
 \end{aligned}$$

And:  $1.251 \neq 1.283$

Therefore:  $X \not\perp\!\!\!\perp Y|Z$ .



**Problem 28:** Show that  $\mathcal{P}(\Omega)$  is  $\sigma$ -field

Proof:

We will show that:

- 1)  $\Omega \in \mathcal{P}(\Omega)$
- 2)  $\mathcal{P}(\Omega)$  is closed under complementation
- 3)  $\mathcal{P}(\Omega)$  is closed under countable unions

1) From definition of power-set:  $\Omega \in \mathcal{P}(\Omega)$

2) Mark  $\{\sigma_1, \sigma_2, \dots\} \in \Omega$ . For  $A = \{\sigma_i: \text{for some } i's\} \in \Omega$ ,  $A^c = \Omega \setminus A = \{\sigma_j: j \neq i\}$ .

Now, from power-set definition we get that:  $\{\sigma_j: j \neq i\} \in \mathcal{P}(\Omega)$ .

3) Mark  $\{\sigma_1, \sigma_2, \dots\} \in \Omega$ . For  $A_i = \{\sigma_j^{(i)}: \text{for some } j's\} \in \Omega$

$\bigcup_{n=1}^{\infty} A_n = \bigcup \{\sigma_j: \sigma_j^{(i)} \in A_i\}$  Now, from power-set definition we get that:  $\bigcup \{\sigma_j: \sigma_j^{(i)} \in A_i\} \in \mathcal{P}(\Omega)$ .

**Problem 29:** Let  $\mathcal{F}$  be a  $\sigma$ -field.

- 1) Show that  $\emptyset \in \mathcal{F}$ .
- 2) Using De Morgan's laws, show that  $\mathcal{F}$  is closed under countable intersections. Namely, if  $A_1, A_2, \dots$  are in  $\mathcal{F}$  then  $\bigcap_{n=1}^{\infty} A_n \in \mathcal{F}$

Proof:

1) Because  $\Omega \in \mathcal{F}$ , we get from property (2) of  $\sigma$ -field that:  $\Omega^c = \emptyset \in \mathcal{F}$

2) De Morgan's laws state that  $(A \cup B)^c = A^c \cap B^c$ . Therefore:  $\bigcup_i A_i = \bigcap_i A_i^c$

For all  $A_i \in \mathcal{F}$ , from property (2) of  $\sigma$ -field:  $A_i^c \in \mathcal{F}$

from property (3) of  $\sigma$ -field:  $\bigcup_i A_i^c \in \mathcal{F}$ , and therefore:  $\bigcup_i A_i^c = \bigcap_i (A_i^c)^c = \bigcap_i A_i \in \mathcal{F}$ .