Supplemental Material for "Fast and efficient inference for flexible spatial extremes models"

Peng Zhong, Scott A. Sisson, and Boris Beranger
UNSW Data Science Hub, School of Mathematics and Statistics,
University of New South Wales, Sydney 2052, Australia.
{peng.zhong, scott.sisson, b.beranger}@unsw.edu.au

March 28, 2025

In the following, starred equation numbers (*) refer to equations in the main manuscript.

A Proof of Theorem 1

For any $k \in \{1, \ldots, D-1\}$, we have

$$\begin{aligned}\lim_{\boldsymbol{x}_{\overline{B}_k} \downarrow \mathbf{0}_{D-k}} -V_{B_k}(\boldsymbol{x}) &= \int_0^\infty r^k f_{\boldsymbol{W}_{B_k}}(r\boldsymbol{x}_{B_k}) \lim_{\boldsymbol{x}_{\overline{B}_k} \downarrow \mathbf{0}_{D-k}} \Pr(\boldsymbol{W}_{\overline{B}_k} \in [\mathbf{0}, r\boldsymbol{x}_{\overline{B}_k}] \mid \boldsymbol{W}_{B_k} = r\boldsymbol{x}_{B_k}) \mathrm{d}r \\ &= \int_0^\infty r^k f_{\boldsymbol{W}_{B_k}}(r\boldsymbol{x}_{B_k}) \Pr(\boldsymbol{W}_{\overline{B}_k} = \mathbf{0}_{D-k} \mid \boldsymbol{W}_{B_k} = r\boldsymbol{x}_{B_k}) \mathrm{d}r,\end{aligned}$$

which implies that if $\Pr(\boldsymbol{W}_{\overline{B}_k} = \mathbf{0}_{D-k} \mid \boldsymbol{W}_{B_k} = r\boldsymbol{x}_{B_k}) = 0$ then $\lim_{\boldsymbol{x}_{\overline{B}_k} \downarrow \mathbf{0}_{D-k}} -V_{1:k}(\boldsymbol{x}) = 0$ and consequently by $(6^\star)$, $h_{B_k}\left(\boldsymbol{x}_{B_k}/\|_1 \boldsymbol{x}_{B_k}\|\right) = 0$. Moreover, if the angular density on $\partial\Omega$ is zero everywhere, then we must have

$$\lim_{x_{\overline{B}_k} \downarrow \mathbf{0}_{D-k}} -V_{B_k}(\boldsymbol{x}) = \int_0^\infty r^k f_{\boldsymbol{W}_{B_k}}(r\boldsymbol{x}_{B_k}) \Pr(W_{\overline{B}_k} = \mathbf{0}_{D-k} \mid \boldsymbol{W}_{Bk} = r\boldsymbol{x}_{B_k}) \mathrm{d}r = 0,$$

and since the integrand is non-negative, we conclude that the integrand must be zero almost everywhere. Therefore, we have $\Pr(W_{\overline{B}_k} = \mathbf{0}_{D-k} \mid W_{B_k} = r\boldsymbol{x}_{B_k}) = 0$ for any $r > 0$ and $\boldsymbol{x}_{B_k} > \mathbf{0}_k$ almost everywhere because r^k is positive and W is a non-negative continuous random process, which implies the condition $(16^\star)$ must hold.

B Useful distributions

Definition 1 (Extended skew-normal distribution Azzalini (2013, Chapter 5))**.** *A random vector* $\boldsymbol{Y} = (Y_1, \ldots, Y_D) \in \mathbb{R}^D$ *is said to follow the extended skew-normal distribution with location vector* $\boldsymbol{\mu} \in \mathbb{R}^D$*,* $D \times D$ *scale matrix* Σ*, slant parameter* $\boldsymbol{\alpha} \in \mathbb{R}^D$ *and extension parameter* $\tau \in \mathbb{R}$ *if its probability density function (pdf) can be written as*

$$\psi(\boldsymbol{x}; \boldsymbol{\mu}, \Sigma, \boldsymbol{\alpha}, \tau) = \phi(\boldsymbol{x}; \boldsymbol{\mu}, \Sigma) \frac{\Phi\left(\alpha_0 + \boldsymbol{\alpha}^\top \boldsymbol{\omega}^{-1}(\boldsymbol{x} - \boldsymbol{\mu}); 0, 1\right)}{\Phi(\tau; 0, 1)},$$

where $\alpha_0 = \tau \left(1 + \boldsymbol{\alpha}^\top \overline{\Sigma} \boldsymbol{\alpha}\right)^{1/2}$*,* $\overline{\Sigma} = \boldsymbol{\omega}^{-1} \Sigma \boldsymbol{\omega}^{-1}$*,* $\boldsymbol{\omega} = \sqrt{\operatorname{diag}(\Sigma)}$*. The corresponding cumulative distribution function (cdf) is given by*

$$\Psi(\boldsymbol{x}; \mu, \Sigma, \alpha, \alpha_0) = \frac{\Phi\left(\begin{bmatrix} \boldsymbol{\omega}^{-1}(\boldsymbol{x} - \boldsymbol{\mu}) \\ \tau \end{bmatrix}; \begin{bmatrix} 0 \\ 0 \end{bmatrix}, \begin{bmatrix} \overline{\Sigma} & -\frac{\overline{\Sigma}\boldsymbol{\alpha}}{(1+\boldsymbol{\alpha}^\top \overline{\Sigma}\boldsymbol{\alpha})^{1/2}} \\ -\frac{\boldsymbol{\alpha}^\top \overline{\Sigma}}{(1+\boldsymbol{\alpha}^\top \overline{\Sigma}\boldsymbol{\alpha})^{1/2}} & 1 \end{bmatrix}\right)}{\Phi(\tau; 0, 1)}.$$

In short, we can write $\boldsymbol{Y} \sim ESN(\boldsymbol{\mu}, \Sigma, \boldsymbol{\alpha}, \alpha_0)$*.*

Definition 2 (Multivariate truncated normal distribution)**.** *Let* $\boldsymbol{Y}$ *be a* D*-dimensional Gaussian random vector with mean* $\mu \in \mathbb{R}^D$ *and covariance matrix* Σ*, with pdf and cdf represented by* $\phi(\cdot; \mu, \Sigma)$ *and* $\Phi(\cdot; \mu, \Sigma)$*. The random vector* $\widehat{\boldsymbol{Y}} = \boldsymbol{Y} | \boldsymbol{Y} \geq \boldsymbol{0}$ *defines a truncated Gaussian random vector with pdf*

$$f_{\widehat{Y}}(\boldsymbol{y}) = c^{-1} \phi(\boldsymbol{y}; \mu, \Sigma), \quad \boldsymbol{y} \in \mathbb{R}^D_+,$$

where $c = \int_{\boldsymbol{0}}^{\infty} \phi(\boldsymbol{y}; \mu, \Sigma) d\boldsymbol{y}$*. In short, we can write* $\widehat{\boldsymbol{Y}} \sim \mathcal{N}_{[\boldsymbol{0}, \infty]}(\boldsymbol{\mu}, \Sigma)$*.*

Note that while the conditional distributions of $\widehat{\boldsymbol{Y}} \sim \mathcal{N}_{[\boldsymbol{0}, \infty]}(\boldsymbol{\mu}, \Sigma)$ remain within the truncated Gaussian family, the marginal distributions do not observe such a property (Horrace, 2005). Denote the set $A = [-\boldsymbol{\infty}, \boldsymbol{y}] = [-\infty, y_1] \times \cdots \times [-\infty, y_D]$ and $A_{B_k} = \cup_{b \in B_k} A_b$, where $A_b = [-\infty, y_1] \times \cdots \times [-\infty, y_{b-1}] \times [-\infty, 0] \times [-\infty, y_{b+1}] \times \cdots \times [-\infty, y_D]$. This implies that $[\boldsymbol{0}, \boldsymbol{y}] = A \cap \overline{A_{B_D}} =$

$A \cap_{b\in B_D} \overline{A}_b$, and by the inclusion and exclusion principle we have

$$\int_{\mathbf{0}}^{\boldsymbol{y}} \phi(\boldsymbol{x};\mu,\Sigma)\mathrm{d}\boldsymbol{x} = \int_A \phi(\boldsymbol{x};\mu,\Sigma)\mathrm{d}\boldsymbol{x} + \sum_{k=1}^{D}\sum_{B_k}(-1)^k \int_{A_{B_k}} \phi(\boldsymbol{x};\mu,\Sigma)\mathrm{d}\boldsymbol{x}$$
$$= \Phi(\boldsymbol{y};\mu,\Sigma) + \sum_{k=1}^{D}\sum_{B_k}(-1)^k \Phi(\boldsymbol{y}^{B_k};\mu,\Sigma) := \Phi^{\circ}(\boldsymbol{y};\mu,\Sigma), \tag{1}$$

where $\boldsymbol{y}^{B_k}$ corresponds to the vector $\boldsymbol{y}$ where the b-th component, $b \in B_k$, is replaced by zero.

Definition 3 (Cdf of the multivariate truncated normal distribution)**.** *Let $\widehat{\boldsymbol{Y}} \sim \mathcal{N}_{[\mathbf{0},\infty]}(\boldsymbol{\mu},\Sigma)$. The cdf of the truncated Gaussian is given by*

$$F_{\widehat{Y}}(\boldsymbol{y}) = \frac{\Phi^{\circ}(\boldsymbol{y};\mu,\Sigma)}{\Phi^{\circ}(\boldsymbol{\infty};\mu,\Sigma)}, \quad \boldsymbol{y} \in \mathbb{R}_+^D,$$

where the denominator is given by (1)*.*

Definition 4. *Let $t_\nu(\boldsymbol{y};\mu,\Sigma)$ and $T_\nu(\boldsymbol{y};\mu,\Sigma)$ be the pdf and cdf of the multivariate t distribution with mean $\mu \in \mathbb{R}^D$, positive scale matrix Σ and degree of freedom $\nu > 0$. Following a similar construction as in Definition 2, a random vector $\boldsymbol{Y}$ is said to follow the multivariate truncated student-t distribution on $\mathbb{R}_+^D$ if its pdf can be written as*

$$\frac{t_\nu(\boldsymbol{y};\mu,\Sigma)}{T_\nu^{\circ}(\boldsymbol{\infty};\mu,\Sigma)}, \quad \boldsymbol{y} \in \mathbb{R}_+^D.$$

Additionally, its cdf is given by $T_\nu^{\circ}(\boldsymbol{y};\mu,\Sigma)/T_\nu^{\circ}(\boldsymbol{\infty};\mu,\Sigma)$. In short, we can write $\boldsymbol{Y} \sim t_{[\mathbf{0},\infty]}(\boldsymbol{\mu},\Sigma,\nu)$.

C Skew Brown-Resnick processes

C.1 Proof of Theorem 2

Let $\boldsymbol{W} = \exp\{\boldsymbol{Y} - \boldsymbol{a}\}$, where $\boldsymbol{Y}$ is the random vector associated with the process $Y(\boldsymbol{s})$, i.e., $\boldsymbol{Y} \sim \mathrm{ESN}(\mathbf{0},\Sigma,\boldsymbol{\alpha},0)$ with scale matrix Σ and slant vector $\boldsymbol{\alpha}$, and $\boldsymbol{a} = (a_1,\ldots,a_D)$, $a_k = \log \mathbb{E}\left[\exp\{Y_k\}\right]$, $k = 1,\ldots,D$.

By applying the first half of (12*), we have

$$\begin{aligned} V(\boldsymbol{x}) &= \sum_{k=1}^{D} \tfrac{1}{x_k} \int_{-\infty}^{\infty} \int_{-\infty}^{y_k+\boldsymbol{a}_{-k}-a_k\mathbf{1}_{D-1}+\log(\boldsymbol{x}_{-k}/x_k)} e^{y_k-a_k}\psi(\boldsymbol{y};\mathbf{0},\Sigma,\boldsymbol{\alpha},0)\mathrm{d}\boldsymbol{y}_{-k}\mathrm{d}y_k \\ &= \sum_{k=1}^{D} \tfrac{\Phi(\tau_k;0,1)}{2} \tfrac{\exp\{\tilde{\omega}_k^2/2-a_k\}}{x_k} \int_{-\infty}^{\infty} \int_{-\infty}^{y_k+\boldsymbol{a}_{-k}-a_k\mathbf{1}_{D-1}+\log(\boldsymbol{x}_{-k}/x_k)} \psi(\boldsymbol{y};\Sigma_{:,k},\Sigma,\boldsymbol{\alpha},\tau_k)\mathrm{d}\boldsymbol{y}_{-k}\mathrm{d}y_k \end{aligned} \tag{2}$$

where $\tau_k = \boldsymbol{\alpha}^\top\omega^{-1}\Sigma_{:,k}\left(1+\boldsymbol{\alpha}^\top\overline{\Sigma}\boldsymbol{\alpha}\right)^{-1/2}$, $\overline{\Sigma} = \omega^{-1}\Sigma\omega^{-1}$, $\omega = \sqrt{\mathrm{diag}(\Sigma)}$ and $\tilde{\boldsymbol{\omega}} = (\tilde{\omega}_1,\ldots,\tilde{\omega}_D) = \omega\mathbf{1}$. The above integrals represent $\Pr[\boldsymbol{X}_{-k} - X_k \le \boldsymbol{a}_{-k} - a_k\mathbf{1}_{D-1} + \log(\boldsymbol{x}_{-k}/x_k)]$, with $\boldsymbol{X} \sim \mathrm{ESN}(\Sigma_{:,k},\Sigma,\boldsymbol{\alpha},\tau_k)$. The random vector $\boldsymbol{X}_{-k} - X_k$ can be derived from $\boldsymbol{X}$ by applying the affine transformation $A_k\boldsymbol{X}$ where $A_k = (\boldsymbol{e}_1,\cdots,\boldsymbol{e}_{k-1},-\mathbf{1}_{D-1},\boldsymbol{e}_k,\ldots,\boldsymbol{e}_{D-1})$ and $\boldsymbol{e}_k, k = 1,\ldots,D-1$ are standard basis vectors of dimension $(D-1)$. From Azzalini (2013, Chapter 5) we have

$$\boldsymbol{X}_{-k} - X_k \sim \mathrm{ESN}\left(\boldsymbol{\mu}_k,\Sigma_k,\widehat{\boldsymbol{\alpha}}_k,\tau_k\right)$$

where $\boldsymbol{\mu}_k = A_k\Sigma_{:,k}, \Sigma_k = A_k\Sigma A_k^\top, \widehat{\boldsymbol{\alpha}}_k = (1-\boldsymbol{\delta}^\top\omega A_k^\top\Sigma_k^{-1}A_k\omega\boldsymbol{\delta})^{-1/2}\widehat{\omega}_k\Sigma_k^{-1}A_k\omega\boldsymbol{\delta}, \boldsymbol{\delta} = \left(1+\boldsymbol{\alpha}^\top\overline{\Sigma}\boldsymbol{\alpha}\right)^{-1/2}\overline{\Sigma}\boldsymbol{\alpha}$, $\widehat{\omega}_k = \sqrt{\mathrm{diag}(\Sigma_k)}$ and therefore, (2) becomes

$$V(\boldsymbol{x}) = \sum_{k=1}^{D} c\tfrac{\exp\{\tilde{\omega}_k^2/2-a_k\}}{x_k}\Psi\left(\boldsymbol{a}_{-k} - a_k\mathbf{1}_{D-1} + \log(\boldsymbol{x}_{-k}/x_k);\boldsymbol{\mu}_k,\Sigma_k,\widehat{\boldsymbol{\alpha}}_k,\tau_k\right). \tag{3}$$

The normalizing constants are derived by solving $a_k = \log\left(\mathbb{E}\left[\exp\{Y_k\}\right]\right)$ for $k = 1,\ldots,D$. The moment generating function of $Y \sim \mathrm{ESN}(\mathbf{0},\Sigma,\boldsymbol{\alpha},0)$ is given by

$$M(\boldsymbol{t}) = 2\exp\left\{\tfrac{1}{2}\boldsymbol{t}^\top\Sigma\boldsymbol{t}\right\}\Phi\left(\boldsymbol{\delta}^\top\omega\boldsymbol{t}\right),$$

where $\boldsymbol{\delta}$ and ω are as defined above and considering $\boldsymbol{t} = \boldsymbol{e}_k$ yields

$$a_k = \log(2) + \tfrac{1}{2}\tilde{\omega}_k^2 + \log\left(\Phi\left(\tilde{\omega}_k\delta_k\right)\right).$$

Noting that $\tau_k = \tilde{\omega}_k\delta_k$ and

$$\boldsymbol{a}_{-k} - a_k\mathbf{1}_{D-1} + \log\left(\tfrac{\boldsymbol{x}_{-k}}{x_k}\right) = \log\left(\tfrac{\boldsymbol{x}_{-k}^{\circ\circ}}{x_k^{\circ\circ}}\right) + \tfrac{\tilde{\boldsymbol{\omega}}_{-k}^2 - \tilde{\omega}_k^2\mathbf{1}_{D-1}}{2},$$

where $\boldsymbol{x}^{\circ\circ} = \left(x_1^{\circ\circ}, \ldots, x_D^{\circ\circ}\right)$, $x_k^{\circ\circ} = 2x_k\Phi\left(\tau_k; 0, 1\right)$, $k = 1, \ldots, D$, we can therefore re-write (3) as

$$
\begin{aligned}
V(\boldsymbol{x}) &= \sum_{k=1}^{D} \tfrac{1}{x_k} \Psi\left(\log\left(\tfrac{\boldsymbol{x}_{-k}^{\circ\circ}}{x_k^{\circ\circ}}\right) + \tfrac{\tilde{\boldsymbol{\omega}}_{-k}^2 - \tilde{\omega}_k^2 \mathbf{1}_{D-1}}{2}; \boldsymbol{\mu}_k, \Sigma_k, \widehat{\boldsymbol{\alpha}}_k, \tau_k\right) \\
&= \sum_{k=1}^{D} \tfrac{1}{x_k} \frac{\Phi\left(\begin{bmatrix} \widehat{\omega}_k^{-1}\left(\log\left(\tfrac{\boldsymbol{x}_{-k}^{\circ\circ}}{x_k^{\circ\circ}}\right) + \tfrac{\tilde{\boldsymbol{\omega}}_{-k}^2 - \tilde{\omega}_k^2 \mathbf{1}_{D-1}}{2} - \boldsymbol{\mu}_k\right) \\ \tau_k \end{bmatrix}; \begin{bmatrix} \mathbf{0} \\ 0 \end{bmatrix}, \begin{bmatrix} \overline{\Sigma}_k & -\frac{\overline{\Sigma}_k \widehat{\boldsymbol{\alpha}}_k}{(1+\widehat{\boldsymbol{\alpha}}_k^\top \overline{\Sigma}_k \widehat{\boldsymbol{\alpha}}_k)^{1/2}} \\ -\frac{\widehat{\boldsymbol{\alpha}}_k^\top \overline{\Sigma}_k}{\left(1+\widehat{\boldsymbol{\alpha}}_k^\top \overline{\Sigma}_k \widehat{\boldsymbol{\alpha}}_k\right)^{1/2}} & 1 \end{bmatrix}\right)}{\Phi(\tau_k; 0, 1)} \\
&= \sum_{k=1}^{D} \tfrac{1}{x_k} \frac{\Phi\left(\begin{bmatrix} \widehat{\omega}_k^{-1}\left(\log\left(\tfrac{\boldsymbol{x}_{-k}^{\circ\circ}}{x_k^{\circ\circ}}\right) + \tfrac{\tilde{\boldsymbol{\omega}}_{-k}^2 - \tilde{\omega}_k^2 \mathbf{1}_{D-1}}{2} - \boldsymbol{\mu}_k\right) \\ \tau_k \end{bmatrix}; \begin{bmatrix} \mathbf{0} \\ 0 \end{bmatrix}, \begin{bmatrix} \overline{\Sigma}_k & -\widehat{\omega}_k^{-1} A_k \omega \boldsymbol{\delta} \\ -\delta^\top \omega A_k^\top \widehat{\omega}_k^{-1} & 1 \end{bmatrix}\right)}{\Phi(\tau_k; 0, 1)}.
\end{aligned}
$$

which completes the proof of Theorem 2.

Notice that the expression above can be much simplified by re-parametrising the slant parameter $\boldsymbol{\alpha}$ by $\boldsymbol{\eta} = \omega^{-1}\boldsymbol{\alpha}$ which implies that $\omega\boldsymbol{\delta} = \Sigma\boldsymbol{\eta}\left(\sqrt{1 + \boldsymbol{\eta}^\top \Sigma \boldsymbol{\eta}}\right)^{-1/2} \equiv \boldsymbol{\xi}$. As such, we have $\tau_k = \xi_k$ and $\widehat{\boldsymbol{\alpha}}_k = \left(1 - \boldsymbol{\xi}^\top A_k^\top \Sigma_k^{-1} A_k \boldsymbol{\xi}\right)^{-1/2} \widehat{\omega}_k \Sigma_k^{-1} A_k \boldsymbol{\xi}$, and the exponent function can be written as

$$
\begin{aligned}
V(\boldsymbol{x}) &= \sum_{k=1}^{D} \tfrac{1}{x_k} \frac{\Phi\left(\begin{bmatrix} \widehat{\omega}_k^{-1}\left(\log\left(\tfrac{\boldsymbol{x}_{-k}^{\circ\circ}}{x_k^{\circ\circ}}\right) + \tfrac{\tilde{\boldsymbol{\omega}}_{-k}^2 - \tilde{\omega}_k^2 \mathbf{1}_{D-1}}{2} - \boldsymbol{\mu}_k\right) \\ \xi_k \end{bmatrix}; \begin{bmatrix} \mathbf{0} \\ 0 \end{bmatrix}, \begin{bmatrix} \overline{\Sigma}_k & -\widehat{\omega}_k^{-1} A_k \boldsymbol{\xi} \\ -\boldsymbol{\xi}^\top A_k^\top \widehat{\omega}_k^{-1} & 1 \end{bmatrix}\right)}{\Phi(\xi_k; 0, 1)} \\
&= \sum_{k=1}^{D} \tfrac{1}{x_k} \frac{\Phi\left(\begin{bmatrix} \log\left(\tfrac{\boldsymbol{x}_{-k}^{\circ\circ}}{x_k^{\circ\circ}}\right) + \tfrac{\tilde{\boldsymbol{\omega}}_{-k}^2 - \tilde{\omega}_k^2 \mathbf{1}_{D-1}}{2} - \boldsymbol{\mu}_k \\ \xi_k \end{bmatrix}; \begin{bmatrix} \mathbf{0} \\ 0 \end{bmatrix}, \begin{bmatrix} \Sigma_k & -A_k \boldsymbol{\xi} \\ -\boldsymbol{\xi}^\top A_k^\top & 1 \end{bmatrix}\right)}{\Phi(\xi_k; 0, 1)}.
\end{aligned}
$$

C.2 Proof of Proposition 1

The conditional probability $(16^\star)$ is written as

$$
\Pr(\boldsymbol{W}_{\overline{B}_k} = 0_{D-k} \mid \boldsymbol{W}_{B_k} = r\boldsymbol{x}_{B_k}) = \Pr(\boldsymbol{Y}_{\overline{B}_k} = -\boldsymbol{\infty}_{D-k} \mid \boldsymbol{Y}_{B_k} = \log(r\boldsymbol{x}_{B_k}) + a_{B_k}),
$$

and since $\boldsymbol{Y}_{\overline{B}_k} \mid \boldsymbol{Y}_{B_k}$ is distributed according to the extended skew-Normal distribution, it directly implies that the above probability is zero and therefore the skewed Brown-Resnick process has no mass on $\partial\Omega$. It now remains to derive its densty on Ω°.

Take $\boldsymbol{w} = (w_1, \ldots, w_D) \in \mathbb{R}_+^D$ and let $w_k^{\circ\circ} = 2w_k \Phi(\tau_k; 0, 1)$, the density of $\boldsymbol{W}$ is given by

$$\begin{aligned} f_W(\boldsymbol{w}) &= f_{\tilde{\boldsymbol{Y}}}\left(\frac{\tilde{\boldsymbol{w}}^2}{2} + \log\left(\boldsymbol{w}^{\circ\circ}\right)\right) \times \prod_{k=1}^{D} \frac{1}{w_k} \\ &= \psi\left(\frac{\tilde{\boldsymbol{\omega}}^2}{2} + \log\left(\boldsymbol{w}^{\circ\circ}\right); \mathbf{0}, \Sigma, \boldsymbol{\alpha}, \mathbf{0}\right) \times \prod_{k=1}^{D} \frac{1}{w_k}. \end{aligned} \tag{4}$$

In order to complete the proof we need the following lemma.

Lemma 1. *The intensity function for the skewed Brown-Resnick process is*

$$\kappa(\boldsymbol{x}) = \frac{2\Phi(\tilde{\tau}; 0, 1)|\Sigma|^{-1/2}\left(\mathbf{1}^\top \boldsymbol{q}\right)^{-1/2}}{(2\pi)^{(D-1)/2} \prod_{k=1}^{D} x_k} \exp\left\{-\frac{1}{2}\left[\log \boldsymbol{x}^{\circ\circ\top} \mathcal{M} \log \boldsymbol{x}^{\circ\circ} + \log \boldsymbol{x}^{\circ\circ\top}\left(\frac{2\boldsymbol{q}}{\mathbf{1}^\top \boldsymbol{q}} + \mathcal{M}\tilde{\boldsymbol{\omega}}^2\right) + \frac{\boldsymbol{q}^\top \tilde{\boldsymbol{\omega}}^2 - 1}{\mathbf{1}^\top \boldsymbol{q}} + \frac{1}{4}\tilde{\boldsymbol{\omega}}^{2,\top} \mathcal{M} \tilde{\boldsymbol{\omega}}^2\right]\right\},$$

where $\boldsymbol{q} = \Sigma^{-1}\mathbf{1}$, $\mathcal{M} = \Sigma^{-1} - \boldsymbol{q}\boldsymbol{q}^\top / \mathbf{1}^\top \boldsymbol{q}$ *and*

$$\tilde{\tau} = \left(1 + \frac{\left(\boldsymbol{\alpha}^\top \omega^{-1} \mathbf{1}\right)^2}{\mathbf{1}^\top q}\right)^{-1/2} \boldsymbol{\alpha}^\top \omega^{-1}\left[\left(\mathbb{I} - \frac{\mathbf{1} q^\top}{\mathbf{1}^\top q}\right)\left(\log \boldsymbol{x}^{\circ\circ} + \frac{\tilde{\boldsymbol{\omega}}^2}{2}\right) + \frac{\mathbf{1}}{\mathbf{1}^\top q}\right].$$

Proof. By plugging (4) into (13*), we get

$$\begin{aligned} \kappa(\boldsymbol{x}) &= \int_0^\infty r^D \psi\left(\frac{\tilde{\boldsymbol{\omega}}^2}{2} + \log\left(\boldsymbol{x}^{\circ\circ}\right) + \log(r); \mathbf{0}, \Sigma, \boldsymbol{\alpha}, \mathbf{0}\right) \times \prod_{k=1}^{D} \frac{1}{r x_k} \mathrm{d}r \\ &= \int_0^\infty \psi\left(\frac{\tilde{\boldsymbol{\omega}}^2}{2} + \log\left(\boldsymbol{x}^{\circ\circ}\right) + \log(r); \mathbf{0}, \Sigma, \boldsymbol{\alpha}, \mathbf{0}\right) \times \prod_{k=1}^{D} \frac{1}{x_k} \mathrm{d}r \\ &= \int_{\mathbb{R}} e^u \psi\left(\frac{\tilde{\boldsymbol{\omega}}^2}{2} + \log\left(\boldsymbol{x}^{\circ\circ}\right) + u; \mathbf{0}, \Sigma, \boldsymbol{\alpha}, \mathbf{0}\right) \times \prod_{k=1}^{D} \frac{1}{x_k} \mathrm{d}u \\ &= \int_{\mathbb{R}} \frac{e^u 2}{\prod_{k=1}^{D} x_k} \phi\left(\frac{\tilde{\boldsymbol{\omega}}^2}{2} + \log\left(\boldsymbol{x}^{\circ\circ}\right) + u; \mathbf{0}, \Sigma\right) \Phi\left(\boldsymbol{\alpha}^\top \omega^{-1}\left(\frac{\tilde{\boldsymbol{\omega}}^2}{2} + \log\left(\boldsymbol{x}^{\circ\circ}\right) + u\right); 0, 1\right) \mathrm{d}u \end{aligned}$$

Following Wadsworth and Tawn (2014), we have

$$\begin{aligned} e^u \phi\left(\frac{\tilde{\boldsymbol{\omega}}^2}{2} + \log\left(\boldsymbol{x}^{\circ\circ}\right) + u; \mathbf{0}, \Sigma\right) = \phi\left(u; \mu_u, \omega_u^2\right) & \frac{|\Sigma|^{-1/2}\left(\mathbf{1}^\top \boldsymbol{q}\right)^{-1/2}}{(2\pi)^{(D-1)/2}} \\ & \times \exp\left\{-\frac{1}{2}\left[\log \boldsymbol{x}^{\circ\circ\top} \mathcal{M} \log \boldsymbol{x}^{\circ\circ} + \log \boldsymbol{x}^{\circ\circ\top}\left(\frac{2\boldsymbol{q}}{\mathbf{1}^\top \boldsymbol{q}} + \mathcal{M}\tilde{\boldsymbol{\omega}}^2\right)\right]\right\} \\ & \times \exp\left\{-\frac{1}{2}\left(\frac{\boldsymbol{q}^\top \tilde{\boldsymbol{\omega}}^2 - 1}{\mathbf{1}^\top \boldsymbol{q}} + \frac{1}{4}\tilde{\boldsymbol{\omega}}^{2,\top} \mathcal{M} \tilde{\boldsymbol{\omega}}^2\right)\right\} \end{aligned}$$

where $\boldsymbol{q} = \Sigma^{-1}\mathbf{1}$, $\mu_u = -\frac{\log \boldsymbol{x}^{\circ\circ}\boldsymbol{q} + \frac{1}{2}(\tilde{\boldsymbol{\omega}}^2)^\top \boldsymbol{q} - 1}{\mathbf{1}^\top \boldsymbol{q}}$, $\omega_u = (\mathbf{1}^\top \boldsymbol{q})^{-1/2}$, and $\mathcal{M} = \Sigma^{-1} - \boldsymbol{q}\boldsymbol{q}^\top / \mathbf{1}^\top \boldsymbol{q}$. In addition, note that

$$\Phi\left(\boldsymbol{\alpha}^\top \omega^{-1}\left(\frac{\tilde{\boldsymbol{\omega}}^2}{2} + \log\left(\boldsymbol{x}^{\circ\circ}\right) + u\right); 0, 1\right) = \Phi\left(\alpha_u\left(\frac{u-\mu_u}{\omega_u}\right) + \alpha_{0,u}; 0, 1\right) \frac{\Phi(\tilde{\tau}; 0, 1)}{\Phi(\tilde{\tau}; 0, 1)}$$

where $\alpha_u = \omega_u \boldsymbol{\alpha}^\top \omega^{-1}\mathbf{1}$, $\alpha_{0,u} = \boldsymbol{\alpha}^\top \omega^{-1}\left(I - \omega_u^2 \mathbf{1}\mathbf{1}^\top \Sigma^{-1}\right)\left(\log \boldsymbol{x}^{\circ\circ} + \tilde{\boldsymbol{\omega}}^2/2\right) - \omega_u^2 \boldsymbol{\alpha}^\top \omega^{-1}\mathbf{1}$ and

$$\tilde{\tau} = \alpha_{0,u}(1 + \alpha_u^2)^{-1/2} = \left(1 + \frac{(\boldsymbol{\alpha}^\top \omega^{-1}\mathbf{1})^2}{\mathbf{1}^\top \boldsymbol{q}}\right)^{-1/2} \boldsymbol{\alpha}^\top \omega^{-1}\left[\left(\mathbb{I} - \frac{\mathbf{1}\boldsymbol{q}^\top}{\mathbf{1}^\top \boldsymbol{q}}\right)\left(\log \boldsymbol{x}^{\circ\circ} + \frac{\tilde{\boldsymbol{\omega}}^2}{2}\right) + \frac{\mathbf{1}}{\mathbf{1}^\top \boldsymbol{q}}\right]. \quad (5)$$

Therefore, by matching the integrand with a univariate extended skew normal density function with respect to u, we have,

$$\begin{aligned}
\kappa(\boldsymbol{x}) = &\int_{\mathbb{R}} \phi\left(u; \mu_u, \omega_u^2\right) \Phi\left(\alpha_u\left(\tfrac{u-\mu_u}{\omega_u}\right) + \alpha_{0,u}; 0, 1\right) \tfrac{\Phi(\tilde{\tau}; 0, 1)}{\Phi(\tilde{\tau}; 0, 1)} \tfrac{2|\Sigma|^{-1/2}\left(\mathbf{1}^\top \boldsymbol{q}\right)^{-1/2}}{(2\pi)^{(D-1)/2} \prod_{k=1}^D x_k} \\
&\times \exp\left\{-\tfrac{1}{2}\left[\log \boldsymbol{x}^{\circ\circ\top} \mathcal{M} \log \boldsymbol{x}^{\circ\circ} + \log \boldsymbol{x}^{\circ\circ\top}\left(\tfrac{2\boldsymbol{q}}{\mathbf{1}^\top \boldsymbol{q}} + \mathcal{M}\tilde{\boldsymbol{\omega}}^2\right)\right]\right\} \\
&\times \exp\left\{-\tfrac{1}{2}\left(\tfrac{\boldsymbol{q}^\top \tilde{\boldsymbol{\omega}}^2 - 1}{\mathbf{1}^\top \boldsymbol{q}} + \tfrac{1}{4}\tilde{\boldsymbol{\omega}}^{2,\top} \mathcal{M} \tilde{\boldsymbol{\omega}}^2\right)\right\} \mathrm{d}u \\
= &\int_{\mathbb{R}} \phi\left(u; \mu_u, \omega_u^2\right) \Phi\left(\alpha_u\left(\tfrac{u-\mu_u}{\omega_u}\right) + \alpha_{0,u}; 0, 1\right) \tfrac{1}{\Phi(\tilde{\tau}; 0, 1)} \mathrm{d}u \\
&\times \tfrac{2\Phi(\tilde{\tau}; 0, 1)|\Sigma|^{-1/2}\left(\mathbf{1}^\top \boldsymbol{q}\right)^{-1/2}}{(2\pi)^{(D-1)/2} \prod_{k=1}^D x_k} \\
&\times \exp\left\{-\tfrac{1}{2}\left[\log \boldsymbol{x}^{\circ\circ\top} \mathcal{M} \log \boldsymbol{x}^{\circ\circ} + \log \boldsymbol{x}^{\circ\circ\top}\left(\tfrac{2\boldsymbol{q}}{\mathbf{1}^\top \boldsymbol{q}} + \mathcal{M}\tilde{\boldsymbol{\omega}}^2\right)\right]\right\} \\
&\times \exp\left\{-\tfrac{1}{2}\left(\tfrac{\boldsymbol{q}^\top \tilde{\boldsymbol{\omega}}^2 - 1}{\mathbf{1}^\top \boldsymbol{q}} + \tfrac{1}{4}\tilde{\boldsymbol{\omega}}^{2,\top} \mathcal{M} \tilde{\boldsymbol{\omega}}^2\right)\right\} \\
= &\tfrac{2\Phi(\tilde{\tau}; 0, 1)|\Sigma|^{-1/2}\left(\mathbf{1}^\top \boldsymbol{q}\right)^{-1/2}}{(2\pi)^{(D-1)/2} \prod_{k=1}^D x_k} \exp\left\{-\tfrac{1}{2}\left[\log \boldsymbol{x}^{\circ\circ\top} \mathcal{M} \log \boldsymbol{x}^{\circ\circ} + \log \boldsymbol{x}^{\circ\circ\top}\left(\tfrac{2\boldsymbol{q}}{\mathbf{1}^\top \boldsymbol{q}} + \mathcal{M}\tilde{\boldsymbol{\omega}}^2\right) + \tfrac{\boldsymbol{q}^\top \tilde{\boldsymbol{\omega}}^2 - 1}{\mathbf{1}^\top \boldsymbol{q}} + \tfrac{1}{4}\tilde{\boldsymbol{\omega}}^{2,\top} \mathcal{M} \tilde{\boldsymbol{\omega}}^2\right]\right\},
\end{aligned}$$

which gives the desired expression of the intensity function. □

C.3 Partial derivatives of the exponent function

Lemma 2. *The partial derivatives of the exponent function for the skewed Brown-Resnick process are given as*

$$\begin{aligned}
-V_{B_k}(\boldsymbol{x}) = &-\frac{V_{B_k}(\boldsymbol{x}_{B_k}, \boldsymbol{\infty}_{D-k})}{\Phi\left(\tilde{\alpha}_0\left(1 + \tilde{\boldsymbol{\alpha}}^\top \overline{\tilde{\Sigma}} \tilde{\boldsymbol{\alpha}}\right)^{-1/2}; 0, 1\right)} \\
&\times \Phi\left(\begin{bmatrix} \tilde{\omega}_{(\overline{B}_k)}^{-1}\left(\log \boldsymbol{x}_{\overline{B}_k}^{\circ\circ} - \tilde{\boldsymbol{\mu}}\right) \\ \tilde{\alpha}_0(1 + \tilde{\boldsymbol{\alpha}}^\top \overline{\tilde{\Sigma}} \tilde{\boldsymbol{\alpha}})^{-1/2} \end{bmatrix}; \begin{bmatrix} \mathbf{0} \\ 0 \end{bmatrix} \begin{bmatrix} \overline{\tilde{\Sigma}} & -(1 + \tilde{\boldsymbol{\alpha}}^\top \overline{\tilde{\Sigma}} \tilde{\boldsymbol{\alpha}})^{-1/2} \overline{\tilde{\Sigma}} \tilde{\boldsymbol{\alpha}} \\ -(1 + \tilde{\boldsymbol{\alpha}}^\top \overline{\tilde{\Sigma}} \tilde{\boldsymbol{\alpha}})^{-1/2} \overline{\tilde{\Sigma}} \tilde{\boldsymbol{\alpha}} & 1 \end{bmatrix}\right),
\end{aligned}$$

where $-V_{B_k}(\boldsymbol{x}_{B_k}, \boldsymbol{\infty}_{D-k})$ *is the marginal intensity function, which equals the intensity function in Lemma 1 with parameters replaced by the marginal covariance matrix* $\Sigma_{B_kB_k}$ *and the marginal slant parameter,* $\boldsymbol{\alpha}_{(B_k)}$*, defined as,*

$$\boldsymbol{\alpha}_{(B_k)} = \left(1 + \boldsymbol{\alpha}_{\overline{B}_k}^\top \left(\overline{\Sigma}_{\overline{B}_k\overline{B}_k} - \overline{\Sigma}_{\overline{B}_kB_k}\overline{\Sigma}_{B_kB_k}^{-1}\overline{\Sigma}_{B_k\overline{B}_k}\right)\boldsymbol{\alpha}_{\overline{B}_k}\right)^{-1/2}\left(\boldsymbol{\alpha}_{B_k} + \overline{\Sigma}_{B_kB_k}^{-1}\overline{\Sigma}_{B_k\overline{B}_k}\boldsymbol{\alpha}_{\overline{B}_k}\right),$$

and $\tilde{\Sigma} = \left(\mathcal{M}_{\overline{B}_k\overline{B}_k}\right)^{-1}$, $\overline{\tilde{\Sigma}} = \omega_{(\overline{B}_k)}^{-1}\tilde{\Sigma}\omega_{(\overline{B}_k)}^{-1}$, $\omega_{(\overline{B}_k)} = \sqrt{\text{diag}(\overline{\tilde{\Sigma}})}$, $\tilde{\boldsymbol{\alpha}} = \left(1 + \frac{(\alpha^\top\omega^{-1}\mathbf{1})^2}{\mathbf{1}^\top q}\right)^{-1/2}\omega_{(\overline{B}_k)}K_{01}\left(\mathbb{I} - \frac{\mathbf{1}q^\top}{\mathbf{1}^\top q}\right)\omega^{-1}\boldsymbol{\alpha}$,

$\tilde{\alpha}_0 = \boldsymbol{\beta}^\top K_{10}^\top(\log \boldsymbol{x}_{B_k}^{\circ\circ} + K_{10}\tilde{\boldsymbol{\omega}}^2/2) + \boldsymbol{\beta}^\top K_{01}^\top(\tilde{\boldsymbol{\mu}} + K_{01}\tilde{\boldsymbol{\omega}}^2/2) + \left(1 + \frac{(\boldsymbol{\alpha}^\top\boldsymbol{\omega}^{-1}\mathbf{1})^2}{\mathbf{1}^\top q}\right)^{-1/2}\frac{\boldsymbol{\alpha}^\top\boldsymbol{\omega}^{-1}\mathbf{1}}{\mathbf{1}^\top q}$, $K_{10} =$

$(\mathbb{I}_k, \mathbf{0}_{k,D-k})$, $K_{01} = (\mathbf{0}_{D-k,k}, \mathbb{I}_{D-k})$, $\boldsymbol{\beta} = \left(1 + \frac{(\boldsymbol{\alpha}^\top\omega^{-1}\mathbf{1})^2}{\mathbf{1}^\top q}\right)^{-1/2}\left(\mathbb{I} - \frac{q\mathbf{1}^\top}{\mathbf{1}^\top q}\right)\omega^{-1}\boldsymbol{\alpha}$, $\tilde{\boldsymbol{\mu}} = -\tilde{\Sigma}\left(\mathcal{M}_{\overline{B}_kB_k}\log \boldsymbol{x}_{B_k}^{\circ\circ} + K_{01}q/(\mathbf{1}^\top$

Proof. We can write

$$-V_{B_k}(\boldsymbol{x}) = -V_{B_k}(x_{B_k}, \boldsymbol{\infty}_{D-k})\int_{\mathbf{0}}^{\boldsymbol{x}_{\overline{B}_k}} \frac{V_{B_D}(\boldsymbol{x}_{B_k}, \boldsymbol{y})}{V_{B_k}(x_{B_k}, \boldsymbol{\infty}_{D-k})}\mathrm{d}\boldsymbol{y},$$

which involves two parts, the marginal intensity function $-V_{B_k}(x_{B_k}, \infty_{D-k})$ and the integral of the conditional intensity function over $\boldsymbol{x}_{\overline{B}_k}$. As the marginal distribution of the constructive spectral component $\boldsymbol{Y}_{B_k}$ in Theorem 3 from the main manuscript follows the distribution $\text{ESN}(\mathbf{0}, \Sigma_{B_kB_k}, \boldsymbol{\alpha}_{(B_k)}, \mathbf{0})$ (Azzalini, 2013, Chapter 5.1.4), where $\boldsymbol{\alpha}_{(B_k)}$ is defined above, the marginal intensity function $-V_{B_k}(x_{B_k}, \infty_{D-k})$ has the same expression as in Lemma 1 but with parameters, Σ and $\boldsymbol{\alpha}$, replaced by their marginal counterparts, $\Sigma_{B_kB_k}$ and $\boldsymbol{\alpha}_{(B_k)}$.

The conditional intensity function $V_{B_D}(\boldsymbol{x})/V_{B_k}(\boldsymbol{x}_{B_k}, \boldsymbol{\infty}_{D-k})$ is indeed a valid density function (Wadsworth and Tawn, 2013) with respect to $\boldsymbol{x}_{\overline{B}_k}$. Treating $\boldsymbol{x}_{B_k}$ as constant, the conditional intensity function can be expressed as

$$\begin{aligned}
&\frac{V_{B_D}(\boldsymbol{x})}{V_{B_k}(\boldsymbol{x}_{B_k}, \boldsymbol{\infty}_{D-k})} \\
&\propto \kappa(\boldsymbol{x}) \\
&= \frac{2\Phi(\tilde{\tau}; 0, 1)|\Sigma|^{-1/2}(\mathbf{1}^\top\boldsymbol{q})^{-1/2}}{(2\pi)^{(D-1)/2}\prod_{k=1}^D x_k}\exp\left\{-\frac{1}{2}\left[\log \boldsymbol{x}^{\circ\circ\top}\mathcal{M}\log \boldsymbol{x}^{\circ\circ} + \log \boldsymbol{x}^{\circ\circ\top}\left(\frac{2\boldsymbol{q}}{\mathbf{1}^\top\boldsymbol{q}} + \mathcal{M}\tilde{\boldsymbol{\omega}}^2\right) + \frac{\boldsymbol{q}^\top\tilde{\boldsymbol{\omega}}^{2,-1}}{\mathbf{1}^\top\boldsymbol{q}} + \frac{1}{4}\tilde{\boldsymbol{\omega}}^{2,\top}\mathcal{M}\tilde{\boldsymbol{\omega}}^2\right]\right\}.
\end{aligned}$$

By comparing above equation with the multivariate extended skew normal density function, we can show that the conditional intensity function with respect to $\log \boldsymbol{x}_{\overline{B}_k}^{\circ\circ}$ is a density function for a multi-

variate extended skew normal random vector, $\log \boldsymbol{X}^{\circ\circ}_{\overline{B}_k}$, with scale matrix $\tilde{\Sigma} = \left(\mathcal{M}_{\overline{B}_k\overline{B}_k}\right)^{-1}$, location parameter $\tilde{\boldsymbol{\mu}} = -\tilde{\Sigma}\left(\mathcal{M}_{\overline{B}_k B_k}\log(\boldsymbol{x}^{\circ\circ}_{B_k}) + K_{01}q/(\mathbf{1}^\top q) + K_{01}\mathcal{M}\tilde{\boldsymbol{\omega}}^2/2\right)$, where $K_{01} = (\mathbf{0}_{D-k,k}, \mathbb{I}_{D-k})$, slant parameter $\tilde{\boldsymbol{\alpha}} = \left(1 + \frac{(\alpha^\top\omega^{-1}\mathbf{1})^2}{\mathbf{1}^\top q}\right)^{-1/2}\omega_{(\overline{B}_k)}K_{01}\left(\mathbb{I} - \frac{\mathbf{1}q^\top}{\mathbf{1}^\top q}\right)\omega^{-1}\boldsymbol{\alpha}$, where $\omega_{(\overline{B}_k)} = \sqrt{\mathrm{diag}(\tilde{\Sigma})}$, and parameter

$$\tilde{\alpha}_0 = \boldsymbol{\beta}^\top K_{10}^\top(\log \boldsymbol{x}^{\circ\circ}_{B_k} + K_{10}\tilde{\boldsymbol{\omega}}^2/2) + \boldsymbol{\beta}^\top K_{01}^\top(\tilde{\boldsymbol{\mu}} + K_{01}\tilde{\boldsymbol{\omega}}^2/2) + \left(1 + \tfrac{(\alpha^\top\omega^{-1}\mathbf{1})^2}{\mathbf{1}^\top q}\right)^{-1/2}\tfrac{\boldsymbol{\alpha}^\top\boldsymbol{\omega}^{-1}\mathbf{1}}{\mathbf{1}^\top q},$$

where $K_{10} = (\mathbb{I}_k, \mathbf{0}_{k,D-k})$ and $\boldsymbol{\beta} = \left(1 + \frac{(\alpha^\top\omega^{-1}\mathbf{1})^2}{\mathbf{1}^\top q}\right)^{-1/2}\left(\mathbb{I} - \frac{q\mathbf{1}^\top}{\mathbf{1}^\top q}\right)\omega^{-1}\boldsymbol{\alpha}$. Therefore, we have

$$\begin{aligned}\frac{V_{B_D}(\boldsymbol{x})}{V_{B_k}(\boldsymbol{x}_{B_k}, \boldsymbol{\infty}_{D-k})} = &\frac{|\tilde{\Sigma}|^{-1/2}(2\pi)^{-(D-k)/2}}{\Phi\left(\tilde{\alpha}_0\left(1+\tilde{\boldsymbol{\alpha}}^\top\omega^{-1}_{(\overline{B}_k)}\tilde{\Sigma}\omega^{-1}_{(\overline{B}_k)}\tilde{\boldsymbol{\alpha}}\right)^{-1/2};0,1\right)\prod_{i\in\overline{B}_k}x_i^{\circ\circ}}\Phi\left(\tilde{\alpha}_0 + \tilde{\boldsymbol{\alpha}}^\top\omega^{-1}_{(\overline{B}_k)}\left(\log \boldsymbol{x}^{\circ\circ}_{\overline{B}_k} - \tilde{\boldsymbol{\mu}}\right);0,1\right)\\ &\times\exp\left\{-\tfrac{1}{2}(\log \boldsymbol{x}^{\circ\circ}_{\overline{B}_k} - \tilde{\boldsymbol{\mu}})^\top\mathcal{M}_{\overline{B}_k\overline{B}_k}(\log \boldsymbol{x}^{\circ\circ}_{\overline{B}_k} - \tilde{\boldsymbol{\mu}})\right\}.\end{aligned}$$

Then, by using the cumulative distribution function of $\log \boldsymbol{X}^{\circ\circ}_{\overline{B}_k}$, the integral can be computed as

$$\begin{aligned}\int_{\mathbf{0}}^{\boldsymbol{x}_{\overline{B}_k}}\frac{V_{B_D}(\boldsymbol{x}_{B_k},\boldsymbol{y})}{V_{B_k}(x_{B_k},\boldsymbol{\infty}_{D-k})}\mathrm{d}\boldsymbol{y} &= \Pr[\boldsymbol{X}_{\overline{B}_k} \le \boldsymbol{x}_{\overline{B}_k}]\\ &= \Pr[\log \boldsymbol{X}_{\overline{B}_k} \le \log \boldsymbol{x}_{\overline{B}_k}]\\ &= \Pr[\log \boldsymbol{X}^{\circ\circ}_{\overline{B}_k} \le \log \boldsymbol{x}^{\circ\circ}_{\overline{B}_k}]\\ &= \frac{\Phi\left(\begin{bmatrix}\tilde{\omega}^{-1}_{(\overline{B}_k)}(\log \boldsymbol{x}^{\circ\circ}_{\overline{B}_k} - \tilde{\boldsymbol{\mu}})\\ \tilde{\alpha}_0(1+\tilde{\boldsymbol{\alpha}}^\top\overline{\tilde{\Sigma}}\tilde{\boldsymbol{\alpha}})^{-1/2}\end{bmatrix};\begin{bmatrix}\mathbf{0}\\ 0\end{bmatrix},\begin{bmatrix}\overline{\tilde{\Sigma}} & -(1+\tilde{\boldsymbol{\alpha}}^\top\overline{\tilde{\Sigma}}\tilde{\boldsymbol{\alpha}})^{-1/2}\overline{\tilde{\Sigma}}\tilde{\boldsymbol{\alpha}}\\ -(1+\tilde{\boldsymbol{\alpha}}^\top\overline{\tilde{\Sigma}}\tilde{\boldsymbol{\alpha}})^{-1/2}\overline{\tilde{\Sigma}}\tilde{\boldsymbol{\alpha}} & 1\end{bmatrix}\right)}{\Phi(\tilde{\alpha}_0(1+\tilde{\boldsymbol{\alpha}}^\top\overline{\tilde{\Sigma}}\tilde{\boldsymbol{\alpha}})^{-1/2})}\end{aligned}$$

where $\overline{\tilde{\Sigma}} = \omega^{-1}_{(\overline{B}_k)}\tilde{\Sigma}\omega^{-1}_{(\overline{B}_k)}$. Plugging the above results, the partial derivative of the exponent function are given by

$$\begin{aligned}-V_{B_k}(\boldsymbol{x}) = &\frac{-V_{B_k}(\boldsymbol{x}_{B_k},\boldsymbol{\infty}_{D-k})}{\Phi(\tilde{\alpha}_0(1+\tilde{\boldsymbol{\alpha}}^\top\overline{\tilde{\Sigma}}\tilde{\boldsymbol{\alpha}})^{-1/2})}\\ &\times\Phi\left(\begin{bmatrix}\tilde{\omega}^{-1}_{(\overline{B}_k)}(\log \boldsymbol{x}^{\circ\circ}_{\overline{B}_k} - \tilde{\boldsymbol{\mu}})\\ \tilde{\alpha}_0(1+\tilde{\boldsymbol{\alpha}}^\top\overline{\tilde{\Sigma}}\tilde{\boldsymbol{\alpha}})^{-1/2}\end{bmatrix};\begin{bmatrix}\mathbf{0}\\ 0\end{bmatrix},\begin{bmatrix}\overline{\tilde{\Sigma}} & -(1+\tilde{\boldsymbol{\alpha}}^\top\overline{\tilde{\Sigma}}\tilde{\boldsymbol{\alpha}})^{-1/2}\overline{\tilde{\Sigma}}\tilde{\boldsymbol{\alpha}}\\ -\left(1+\tilde{\boldsymbol{\alpha}}^\top\overline{\tilde{\Sigma}}\tilde{\boldsymbol{\alpha}}\right)^{-1/2}\overline{\tilde{\Sigma}}\tilde{\boldsymbol{\alpha}} & 1\end{bmatrix}\right)\end{aligned}$$

which concludes the proof. □

From Lemma 2, the limit of the partial derivative $-V_{B_k}(\boldsymbol{x})$ for any index set $B_k \neq \emptyset$ and $B_k \subsetneq \{1,\ldots,D\}$ as $\boldsymbol{x}_{\overline{B}_k} \to \mathbf{0}$ is indeed zero.

C.4 Exact simulation algorithm

The exact simulation algorithm for skewed Brown-Resnick process arises from Algorithm 2 in Dombry *et al.* (2016), which requires simulating a process $\boldsymbol{W}/W(s_0)$ with distribution $P_{s_0}(A) = \mathrm{E}_{\boldsymbol{W}}[\mathbb{1}_{\{\boldsymbol{W}/W(s_0)\in A\}}W(s_0)]$ and where $\boldsymbol{W} = (W_0, W_1, \ldots, W_D) \equiv (W(s_0), W(s_1), \ldots, W(s_D))$ is defined as in Theorem 2 to the exception of being $D+1$ dimensional. The distribution P_{s_0} defines the joint distribution of $W_i/W_0, i = 1, \ldots, D$ under the transformed probability measure $\widehat{\Pr} = W(s_0)\mathrm{dPr}$. The moment generating function of $Y_i \equiv Y(s_i), i = 0, \ldots, D$ evaluated at $\boldsymbol{t} = (t_0, t_1, \ldots, t_D)$ can be calculated under the transformed probability measure $\widehat{\Pr}$ as

$$\begin{aligned}\mathrm{E}\left[\exp\left\{Y_0 - a_0 + \sum_{i=0}^{D} t_i Y_i\right\}\right] &= 2\exp\left\{\tfrac{1}{2}(1+t_0, \boldsymbol{t}_{B_D})^\top \Sigma (1+t_0, \boldsymbol{t}_{B_D}) - a_0\right\}\Phi(\delta^\top \omega(1+t_0, \boldsymbol{t}_{B_D})) \\ &= 2\exp\{\tfrac{1}{2}(\Sigma_{0;0} + \boldsymbol{t}^\top\Sigma\boldsymbol{t}) + \Sigma_{0;\cdot}\boldsymbol{t} - a_0\}\Phi\left(\delta_0\omega_{0;0} + \delta^\top\omega\boldsymbol{t}\right)\end{aligned}$$

where the matrices ω and Σ can be decomposed as $\Sigma = \begin{bmatrix}\Sigma_{00} & \Sigma_{B_D;0} \\ \Sigma_{B_D;0}^\top & \Sigma_{B_D;B_D}\end{bmatrix}$. Plugging in the expression for a_0 given in Theorem 2 leads to

$$\mathrm{E}\left[\exp\left\{Y_0 - a_0 + \sum_{i=0}^{D} t_i Y_i\right\}\right] = \exp\{\tfrac{1}{2}\boldsymbol{t}^\top\Sigma\boldsymbol{t} + \Sigma_{\cdot;0}\boldsymbol{t}^\top\}\frac{\Phi(\delta_0\omega_{00} + \delta^\top\omega\boldsymbol{t})}{\Phi(\delta_0\omega_{00})},$$

which corresponds to the moment generating function of the extended skew-normal random vector $(Y^*(s_0), \ldots, Y^*(s_D))$ with location parameter $\Sigma_{\cdot;0}$, scale parameter Σ, slant parameter $\boldsymbol{\alpha}$, and the extended parameter $\tau_0 = \delta_0\omega_{00}$. The distribution of P_{s_0} therefore becomes

$$\exp\{Y^*(s_i) - Y^*(s_0) - a_i + a_0\}, i = 0, \ldots, D.$$

Details about random generation from the multivariate extended skew-normal can be found in Azzalini (2013, Chapter 5.3.3). The pseudo-code of the exact simulation algorithms is summarised in Algorithm 1.

Algorithm 1: Exact simulation from the skewed Brown-Resnick process at location $\boldsymbol{s}_i, i = 1, \ldots, D$.

Input: slant parameter $\boldsymbol{\alpha} \in \mathbb{R}^D$, $D \times D$ scale matrix Σ
Output: $Z(\boldsymbol{s}_j), j = 1, \ldots, D$.

Compute $\boldsymbol{a}$ and δ as defined in Theorem 2.
Simulate $E_0 \sim \text{Exp}(1)$ and set $z = 1/E_0$.
Simulate $Y^*(\boldsymbol{s})$ from an extended skew normal with location parameter $\Sigma_{\cdot;1}$, scale matrix Σ, slant parameter $\boldsymbol{\alpha}$ and the extended parameter $\tau_1 = \delta_1 \omega_{11}$.
Set $W(\boldsymbol{s}) = \exp(Y(\boldsymbol{s}) - Y(s_1) + a_1 - \boldsymbol{a})$
Set $Z(\boldsymbol{s}_j) = W(s_j) * z$ for $j = 1, \ldots, D$.
for $i = 2, \ldots, D$ **do**
    Simulate $E_0 \sim \text{Exp}(1)$ and set $z = 1/E_0$.
    while $z > Z(\boldsymbol{s}_i)$ **do**
        Simulate $Y(\boldsymbol{s})$ from an extended skew normal with location parameter $\Sigma_{\cdot;i}$, scale matrix Σ, slant parameter $\boldsymbol{\alpha}$ and extended parameter $\tau_i = \delta_i \omega_{ii}$
        Set $W(\boldsymbol{s}) = \exp(\boldsymbol{Y}(s) - Y(\boldsymbol{s}_i) + a_i - \boldsymbol{a})$
        if $W(\boldsymbol{s}_j) \times z < Z(\boldsymbol{s}_j), \forall j = 1, \ldots, i-1$ **then**
            Set $Z(\boldsymbol{s}_j) = \max(Z(\boldsymbol{s}_j), W(\boldsymbol{s}_j) \times z), j = 1, \ldots, D$.
        Simulate $E_1 \sim \text{Exp}(1)$, Set $E_0 = E_0 + E_1$ and $z = 1/E_0$.

D Truncated extremal-t processes

D.1 Proof of Theorem 2

Let $\boldsymbol{W} = \tilde{\boldsymbol{Y}}^\nu / \boldsymbol{a}$, where $\tilde{\boldsymbol{Y}}$ is the random vector associated with the process $\tilde{Y}(\boldsymbol{s})$, i.e., $\tilde{\boldsymbol{Y}} \sim \mathcal{N}_{[\mathbf{0},\infty]}(\mathbf{0}, \Sigma)$ with unit variance and therefore Σ represent the correlation matrix, and $\boldsymbol{a} = (a_1, \ldots, a_D)$, $a_k = \mathbb{E}[\tilde{Y}_k^\nu]$, $k = 1, \ldots, D$, and $\nu > 0$.

By applying the first half of $(12^\star)$, we have

$$
\begin{aligned}
V(\boldsymbol{x}) &= \sum_{k=1}^{D} \frac{1}{a_k x_k} \int_0^\infty u_k^\nu \int_{\mathbf{0}}^{u_k \left(\frac{\boldsymbol{x}^\circ_{-k}}{x^\circ_k}\right)^{1/\nu}} \frac{\phi(\boldsymbol{u};\mathbf{0},\Sigma)}{\Phi^\circ(\boldsymbol{\infty};\mathbf{0},\Sigma)} \mathrm{d}\boldsymbol{u}_{-k} \mathrm{d}u_k \\
&= \sum_{k=1}^{D} \frac{1}{a_k x_k \Phi^\circ(\boldsymbol{\infty};\mathbf{0},\Sigma)} \int_0^\infty u_k^\nu \phi(u_k;0,1) \int_{\mathbf{0}}^{u_k \left(\frac{\boldsymbol{x}^\circ_{-k}}{x^\circ_k}\right)^{1/\nu}} \phi(\boldsymbol{u}_{-k}; \Sigma_{-k,k} u_k, \mathbb{I}_{D-1} - \Sigma_{-k,k}\Sigma_{k,-k}) \mathrm{d}\boldsymbol{u}_{-k} \mathrm{d}u_k \\
&= \sum_{k=1}^{D} \frac{1}{a_k x_k \Phi^\circ(\boldsymbol{\infty};\mathbf{0},\Sigma)} \int_0^\infty u_k^\nu \phi(u_k;0,1) \Phi^\circ\left(u_k \left(\frac{\boldsymbol{x}^\circ_{-k}}{x^\circ_k}\right)^{1/\nu}; \Sigma_{-k,k} u_k, \mathbb{I}_{D-1} - \Sigma_{-k,k}\Sigma_{k,-k}\right) \mathrm{d}u_k \\
&= \sum_{k=1}^{D} \frac{2^{(\nu-2)/2}\Gamma((\nu+1)/2)}{\Phi^\circ(\boldsymbol{\infty};\mathbf{0},\Sigma) a_j x_j \sqrt{\pi}} T^\circ_{\nu+1}\left(\left(\frac{\boldsymbol{x}^\circ_{-k}}{x^\circ_k}\right)^{1/\nu}; \Sigma_{-k,k}, \frac{\mathbb{I}_{D-1} - \Sigma_{-k,k}\Sigma_{k,-k}}{\nu+1}\right), \qquad (6)
\end{aligned}
$$

where $\boldsymbol{x}^\circ = \boldsymbol{x}\boldsymbol{a}$, $\mathbb{I}_{D-1}$ is the $(D-1)$-dimensional identity matrix, and the index $-k$ represents the set $B_D \setminus \{k\}$.

The normalising constants a_k can be either derived from the fact that $V(\infty, \ldots, \infty, x_k, \infty, \ldots, \infty) = 1/x_k$ or by comparing the integral above to the following integral

$$
a_k = E[\tilde{Y}_k^\nu] = \int_0^\infty u_k^\nu \int_{\mathbf{0}}^{\boldsymbol{\infty}} \frac{\phi(\boldsymbol{u};\mathbf{0},\Sigma)}{\Phi^\circ(\boldsymbol{\infty};\mathbf{0},\Sigma)} \mathrm{d}\boldsymbol{u}_{-k} \mathrm{d}u_k .
$$

The use of either way will lead us to

$$
a_k = \frac{2^{(\nu-2)/2}\Gamma((\nu+1)/2)}{\Phi^\circ(\boldsymbol{\infty};\mathbf{0},\Sigma)\sqrt{\pi}} T^\circ_{\nu+1}\left(\boldsymbol{\infty}_{D-1}; \Sigma_{-k,k}, \frac{\mathbb{I}_{D-1} - \Sigma_{-k,k}\Sigma_{k,-k}}{\nu+1}\right).
$$

Plugging the expression for a_k in (6), we have

$$
V(\boldsymbol{x}) = \sum_{k=1}^{D} \frac{T^\circ_{\nu+1}\left(\left(\frac{\boldsymbol{x}^\circ_{-k}}{x^\circ_k}\right)^{1/\nu}; \Sigma_{-k,k}, \frac{\mathbb{I}_{D-1} - \Sigma_{-k,k}\Sigma_{k,-k}}{\nu+1}\right)}{x_k T^\circ_{\nu+1}\left(\boldsymbol{\infty}_{D-1}; \Sigma_{-k,k}, \frac{\mathbb{I}_{D-1} - \Sigma_{-k,k}\Sigma_{k,-k}}{\nu+1}\right)}.
$$

D.2 Proof of Proposition 2

From the distribution of $\tilde{\boldsymbol{Y}}$ and therefore $\boldsymbol{W}$, the conditional probability in $(16^\star)$ is written as

$$
\begin{aligned}
\Pr(\boldsymbol{W}_{\overline{B}_k} = \mathbf{0}_{D-k} \mid \boldsymbol{W}_{B_k} = r\boldsymbol{x}_{B_k}) &= \Phi^\circ\left(\mathbf{0}_{D-k}; \Sigma_{\overline{B}_k} \Sigma^{-1}_{B_k B_k} \left(r\boldsymbol{x}^\circ_{B_k}\right)^{1/\nu}, \Sigma_{\overline{B}_k \overline{B}_k} - \Sigma_{\overline{B}_k B_k} \Sigma^{-1}_{B_k B_k} \Sigma_{\overline{B}_k B_k}\right) \\
&= 0,
\end{aligned}
$$

which, by Theorem 1, implies that the truncated extramal-t process has no mass on $\partial\Omega$. It now remains to derive its density on Ω°.

Take $\boldsymbol{w} = (w_1, \ldots, w_D) \in \mathbb{R}_+^D$ and let $\boldsymbol{w}^\circ = \boldsymbol{w}\boldsymbol{a}$, the density of $\boldsymbol{W}$ is given by

$$\begin{aligned} f_W(\boldsymbol{w}) &= f_{\tilde{\boldsymbol{Y}}}\left((\boldsymbol{w}^\circ)^{1/\nu}\right) \times \prod_{k=1}^{D} \left\{ \tfrac{a_k}{\nu} \left(w_k^\circ\right)^{1/\nu - 1} \right\} \\ &= \tfrac{\phi\left((\boldsymbol{w}^\circ)^{1/\nu}; \boldsymbol{0}, \Sigma\right)}{\Phi^\circ(\boldsymbol{\infty}; \boldsymbol{0}, \Sigma)} \times \prod_{k=1}^{D} \left\{ \tfrac{a_k}{\nu} \left(w_k^\circ\right)^{1/\nu - 1} \right\}. \end{aligned} \tag{7}$$

In order to complete the proof we need the following lemma.

Lemma 3. *The intensity function for the truncated extremal-t process is*

$$\kappa(\boldsymbol{x}) = \tfrac{\prod_{k=1}^{D}\left(a_k x_k^{1-\nu}\right)^{1/\nu}}{\Phi^\circ(\boldsymbol{\infty}; \boldsymbol{0}, \Sigma)\nu^{D-1}|\Sigma|^{1/2}\pi^{D/2}2^{(2-\nu)/2}} \left\{ \left[(\boldsymbol{x}^\circ)^{1/\nu}\right]^\top \Sigma^{-1} \left[(\boldsymbol{x}^\circ)^{1/\nu}\right] \right\}^{-(\nu+D)/2} \Gamma\left(\tfrac{\nu+D}{2}\right).$$

Proof. By plugging (7) into (13*), we get

$$\begin{aligned} \kappa(\boldsymbol{x}) &= \int_0^\infty r^D \tfrac{\phi((r\boldsymbol{x}^\circ)^{1/\nu}; \boldsymbol{0}, \Sigma)}{\Phi^\circ(\boldsymbol{\infty}; \boldsymbol{0}, \Sigma)} \prod_{k=1}^{D} \left\{ \tfrac{a_k}{\nu} (r x_k^\circ)^{1/\nu - 1} \right\} \mathrm{d}r \\ &= \tfrac{\prod_{k=1}^{D}\left\{a_k^{1/\nu} x_k^{1/\nu - 1}\right\}}{\Phi^\circ(\boldsymbol{\infty}; \boldsymbol{0}, \Sigma)\nu^D} \int_0^\infty r^{D/\nu} \phi((r\boldsymbol{x}^\circ)^{1/\nu}; \boldsymbol{0}, \Sigma) \mathrm{d}r \\ &= \tfrac{\prod_{k=1}^{D}\left(a_k x_k^{\nu-1}\right)^{1/\nu}}{\Phi^\circ(\boldsymbol{\infty}; \boldsymbol{0}, \Sigma)\nu^D|\Sigma|^{1/2}\pi^{D/2}2^{(2-\nu)/2}} \int_0^\infty r^{D/\nu} \exp\left\{ -\tfrac{r^{2/\nu}}{2} \left[(\boldsymbol{x}^\circ)^{1/\nu}\right]^\top \Sigma^{-1} \left[(\boldsymbol{x}^\circ)^{1/\nu}\right] \right\} \mathrm{d}r \\ &= \tfrac{\prod_{k=1}^{D}\left(a_k x_k^{1-\nu}\right)^{1/\nu}}{\Phi^\circ(\boldsymbol{\infty}; \boldsymbol{0}, \Sigma)\nu^{D-1}|\Sigma|^{1/2}\pi^{D/2}2^{(2-\nu)/2}} \left\{ \left[(\boldsymbol{x}^\circ)^{1/\nu}\right]^\top \Sigma^{-1} \left[(\boldsymbol{x}^\circ)^{1/\nu}\right] \right\}^{-(\nu+D)/2} \Gamma\left(\tfrac{\nu+D}{2}\right), \end{aligned}$$

which completes the proof. □

By Lemma 3, the density on the interior of the simplex, Ω° is therefore given by

$$h(\boldsymbol{w}) = \tfrac{\prod_{k=1}^{D}\left(a_k w_k^{1-\nu}\right)^{1/\nu}}{\Phi^\circ(\boldsymbol{\infty}; \boldsymbol{0}, \Sigma)\nu^{D-1}|\Sigma|^{1/2}\pi^{D/2}2^{(2-\nu)/2}} \left\{ \left[(\boldsymbol{w}^\circ)^{1/\nu}\right]^\top \Sigma^{-1} \left[(\boldsymbol{w}^\circ)^{1/\nu}\right] \right\}^{-(\nu+D)/2} \Gamma\left(\tfrac{\nu+D}{2}\right), \quad \boldsymbol{w} = \tfrac{\boldsymbol{x}}{\|\boldsymbol{x}\|}.$$

D.3 Partial derivatives of the exponent function for the truncated extremal-t processes

Lemma 4. *The partial derivatives of the exponent function of the truncated extremal-t process are given by*

$$-V_{B_k}(\boldsymbol{x}) = \frac{\prod_{i\in B_k}\left\{a_i x_i^{1-\nu}\right\}^{1/\nu} 2^{(\nu-2)/2}\Gamma\left(\frac{k+\nu}{2}\right)\nu^{1-k}}{\Phi^\circ(\boldsymbol{\infty};\mathbf{0},\Sigma)\pi^{k/2}|\Sigma_{B_kB_k}|^{1/2}Q_{\Sigma_{B_kB_k}}\left(\boldsymbol{x}^\circ_{B_k}\right)^{k+\nu}} \times T^\circ_{\nu+k}\left(\frac{\left(\boldsymbol{x}^\circ_{\overline{B}_k}\right)^{1/\nu}}{Q_{\Sigma_{B_kB_k}}\left(\boldsymbol{x}^\circ_{B_k}\right)}; \frac{\Sigma_{\overline{B}_kB_k}\Sigma^{-1}_{B_kB_k}\left(\boldsymbol{x}^\circ_{B_k}\right)^{1/\nu}}{Q_{\Sigma_{B_kB_k}}\left(\boldsymbol{x}^\circ_{B_k}\right)}, \frac{\Sigma_{\overline{B}_k\overline{B}_k}-\Sigma_{\overline{B}_kB_k}\Sigma^{-1}_{B_kB_k}\Sigma_{\overline{B}_kB_k}}{k+\nu}\right),$$

where $Q_\Sigma(\boldsymbol{x}) = \left\{\boldsymbol{x}^\top\Sigma^{-1}\boldsymbol{x}\right\}^{1/2}$.

Proof. Since the interest is in the partial derivatives of the exponent function $V(\boldsymbol{x})$ with respect to the first k components, i.e., $\boldsymbol{x}_{B_k}$, the correlation matrix σ is decomposed as

$$\Sigma = \begin{bmatrix}\Sigma_{B_kB_k} & \Sigma_{B_k\overline{B}_k}\\ \Sigma_{\overline{B}_kB_k} & \Sigma_{\overline{B}_k\overline{B}_k}\end{bmatrix}.$$

Plugging (7) into (15*), we have

$$\begin{aligned}
-V_{B_k}(\boldsymbol{x}) &= \int_0^\infty r^k f_{\boldsymbol{W}_{B_k}}(r\boldsymbol{x}_{B_k})\Pr(\boldsymbol{W}_{\overline{B}_k}\in[0,r\boldsymbol{x}_{\overline{B}_k}]\mid \boldsymbol{W}_{B_k}=r\boldsymbol{x}_{B_k})\mathrm{d}r\\
&= \int_0^\infty r^k \frac{\phi\left(\left(r\boldsymbol{x}^\circ_{B_k}\right)\right)^{1/\nu};\mathbf{0}_k,\Sigma_{B_kB_k}\right)}{\Phi^\circ(\boldsymbol{\infty};\mathbf{0},\Sigma)}\prod_{i=1}^k\left\{\frac{a_i}{\nu}(rx_i^\circ)^{1/\Delta-1}\right\}\\
&\quad\times\Phi^\circ\left(\left(r\boldsymbol{x}^\circ_{\overline{B}_k}\right)^{1/\nu};\Sigma_{\overline{B}_k}\Sigma^{-1}_{B_kB_k}\left(r\boldsymbol{x}^\circ_{B_k}\right)^{1/\nu},\Sigma_{\overline{B}_k\overline{B}_k}-\Sigma_{\overline{B}_kB_k}\Sigma^{-1}_{B_kB_k}\Sigma_{\overline{B}_kB_k}\right)\mathrm{d}r\\
&= \frac{\prod_{i\in B_k}\left\{x_i^{1/\nu-1}a_i^{1/\nu}\right\}}{\Phi^\circ(\boldsymbol{\infty};\mathbf{0},\Sigma)}\nu^{-k}\int_0^\infty r^{k/\nu}\phi\left(\left(r\boldsymbol{x}^\circ_{B_k}\right)^{1/\nu};\mathbf{0}_k,\Sigma_{B_kB_k}\right)\\
&\quad\times\Phi^\circ\left(\left(r\boldsymbol{x}^\circ_{\overline{B}_k}\right)^{1/\nu};\Sigma_{\overline{B}_kB_k}\Sigma^{-1}_{B_kB_k}\left(r\boldsymbol{x}^\circ_{B_k}\right)^{1/\nu},\Sigma_{\overline{B}_k\overline{B}_k}-\Sigma_{\overline{B}_kB_k}\Sigma^{-1}_{B_kB_k}\Sigma_{\overline{B}_kB_k}\right)\mathrm{d}r
\end{aligned}$$

Using the notation $Q_\Sigma(\boldsymbol{x}) = \left\{\boldsymbol{x}^\top\Sigma^{-1}\boldsymbol{x}\right\}^{1/2}$ and applying the change of variable $t = r^{1/\nu}Q_{\Sigma_{B_kB_k}}(\boldsymbol{x}^\circ_{B_k})$, leads to

$$
\begin{aligned}
-V_{B_k}(\boldsymbol{x}) &= \frac{\prod_{i\in B_k}\left\{x_i^{1/\nu-1}a_i^{1/\nu}\right\}}{\Phi^\circ(\infty;\mathbf{0},\Sigma)}\nu^{1-k}(2\pi)^{-k/2}|\Sigma_{B_kB_k}|^{-1/2}\int_0^\infty t^{k+\nu-1}Q_{\Sigma_{B_kB_k}}\left(\boldsymbol{x}^\circ_{B_k}\right)^{-k-\nu}e^{-t^2/2}\\
&\quad\times\Phi^\circ\left(\frac{\left(\boldsymbol{x}^\circ_{\overline{B}_k}\right)^{1/\nu}t}{Q_{\Sigma_{B_kB_k}}\left(\boldsymbol{x}^\circ_{B_k}\right)};\frac{\Sigma_{\overline{B}_kB_k}\Sigma^{-1}_{B_kB_k}\left(\boldsymbol{x}^\circ_{B_k}\right)^{1/\nu}t}{Q_{\Sigma_{B_kB_k}}\left(\boldsymbol{x}^\circ_{B_k}\right)},\Sigma_{\overline{B}_k\overline{B}_k}-\Sigma_{\overline{B}_kB_k}\Sigma^{-1}_{B_kB_k}\Sigma_{\overline{B}_kB_k}\right)\mathrm{d}t\\
&= \frac{\prod_{i\in B_k}\left\{x_i^{1/\nu-1}a_i^{1/\nu}\right\}}{\Phi^\circ(\infty;\mathbf{0},\Sigma)}2^{(\nu-2)/2}\nu^{1-k}\pi^{-k/2}|\Sigma_{B_kB_k}|^{-1/2}Q_{\Sigma_{B_kB_k}}\left(\boldsymbol{x}^\circ_{B_k}\right)^{-k-\nu}\Gamma\left(\tfrac{k+\nu}{2}\right)\\
&\quad\times T^\circ_{\nu+k}\left(\frac{\left(\boldsymbol{x}^\circ_{\overline{B}_k}\right)^{1/\nu}}{Q_{\Sigma_{B_kB_k}}\left(\boldsymbol{x}^\circ_{B_k}\right)};\frac{\Sigma_{\overline{B}_kB_k}\Sigma^{-1}_{B_kB_k}\left(\boldsymbol{x}^\circ_{B_k}\right)^{1/\nu}}{Q_{\Sigma_{B_kB_k}}\left(\boldsymbol{x}^\circ_{B_k}\right)},\frac{\Sigma_{\overline{B}_k\overline{B}_k}-\Sigma_{\overline{B}_kB_k}\Sigma^{-1}_{B_kB_k}\Sigma_{\overline{B}_kB_k}}{k+\nu}\right),
\end{aligned}
$$

which concludes the proof. □

From Lemma 4, we can demonstrate that, for $k = 1, \ldots, D-1$ we have

$$
\begin{aligned}
\lim_{\boldsymbol{x}_{\overline{B}_k}\downarrow\mathbf{0}_{D_k}} V_{B_k}(\boldsymbol{x}) &= \frac{\prod_{i\in B_k}\left\{a_ix_i^{1-\nu}\right\}^{1/\nu}2^{(\nu-2)/2}\Gamma\left(\frac{k+\nu}{2}\right)\nu^{1-k}}{\Phi^\circ(\infty;\mathbf{0},\Sigma)\pi^{k/2}|\Sigma_{B_kB_k}|^{1/2}Q_{\Sigma_{B_kB_k}}\left(\boldsymbol{x}^\circ_{B_k}\right)^{k+\nu}}\\
&\quad\times\lim_{\boldsymbol{x}_{\overline{B}_k}\downarrow\mathbf{0}_{D_k}} T^\circ_{\nu+k}\left(\frac{\left(\boldsymbol{x}^\circ_{\overline{B}_k}\right)^{1/\nu}}{Q_{\Sigma_{B_kB_k}}\left(\boldsymbol{x}^\circ_{B_k}\right)};\frac{\Sigma_{\overline{B}_kB_k}\Sigma^{-1}_{B_kB_k}\left(\boldsymbol{x}^\circ_{B_k}\right)^{1/\nu}}{Q_{\Sigma_{B_kB_k}}\left(\boldsymbol{x}^\circ_{B_k}\right)},\frac{\Sigma_{\overline{B}_k\overline{B}_k}-\Sigma_{\overline{B}_kB_k}\Sigma^{-1}_{B_kB_k}\Sigma_{\overline{B}_kB_k}}{k+\nu}\right)\\
&= \frac{\prod_{i\in B_k}\left\{a_ix_i^{1-\nu}\right\}^{1/\nu}2^{(\nu-2)/2}\Gamma\left(\frac{k+\nu}{2}\right)\nu^{1-k}}{\Phi^\circ(\infty;\mathbf{0},\Sigma)\pi^{k/2}|\Sigma_{B_k,B_k}|^{1/2}Q_{\Sigma_{B_kB_k}}\left(\boldsymbol{x}^\circ_{B_k}\right)^{k+\nu}}\\
&\quad\times T^\circ_{\nu+k}\left(\mathbf{0}_{D_k};\frac{\Sigma_{\overline{B}_kB_k}\Sigma^{-1}_{B_kB_k}\left(\boldsymbol{x}^\circ_{B_k}\right)^{1/\nu}}{Q_{\Sigma_{B_kB_k}}\left(\boldsymbol{x}^\circ_{B_k}\right)},\frac{\Sigma_{\overline{B}_k\overline{B}_k}-\Sigma_{\overline{B}_kB_k}\Sigma^{-1}_{B_kB_k}\Sigma_{\overline{B}_kB_k}}{k+\nu}\right)\\
&= \frac{\prod_{i\in B_k}\left\{a_ix_i^{1-\nu}\right\}^{1/\nu}2^{(\nu-2)/2}\Gamma\left(\frac{k+\nu}{2}\right)\nu^{1-k}}{\Phi^\circ(\infty;\mathbf{0},\Sigma)\pi^{k/2}|\Sigma_{B_k,B_k}|^{1/2}Q_{\Sigma_{B_kB_k}}\left(\boldsymbol{x}^\circ_{B_k}\right)^{k+\nu}}\times 0\\
&= 0,
\end{aligned}
$$

which implies that all densities on $\partial\Omega$ take the value zero.

D.4 Exact simulation algorithm

Similarly to Section C.4, the exact simulation algorithm for truncated extremal-t process arises from Algorithm 2 in Dombry *et al.* (2016), which requires simulating a process $\boldsymbol{W}/W(s_0)$ with distribution $P_{s_0}(A) = \mathrm{E}_{\boldsymbol{W}}[\mathbb{1}_{\{\boldsymbol{W}/W(s_0)\in A\}}W(s_0)]$ and where $\boldsymbol{W} = (W_0, W_1, \ldots, W_D) \equiv$

$(W(s_0), W(s_1), \ldots, W(s_D))$ is defined as in Theorem 3 to the exception of being $D+1$ dimensional. The distribution P_{s_0} can be written as,

$$\begin{aligned} P_{s_0}(A) &= \int_{\mathbb{R}_+^{D+1}} \mathbb{1}\left\{w_i/w_0 \in A_i, i = 1, \ldots, D\right\} w_0 \mathrm{d}F_{\boldsymbol{W}}(\boldsymbol{w}) \\ &= \int_{\mathbb{R}_+^{D+1}} \mathbb{1}\left\{w_i/w_0 \in A_i, i = 1, \ldots, D\right\} w_0 \frac{\phi\left((\boldsymbol{a}\boldsymbol{w})^{1/\nu}; \mathbf{0}_{D+1}, \Sigma\right)}{\Phi^\circ(\boldsymbol{\infty}; \mathbf{0}_{D+1}, \Sigma)} \prod_{i=0}^{D} (a_i w_i)^{1/\nu - 1} \frac{a_i}{\nu} \mathrm{d}\boldsymbol{w}, \end{aligned}$$

where $\boldsymbol{a} = (a_o, \ldots, a_D)$. Let $\boldsymbol{z} = (z_0, z_1, \ldots, z_D)$, with $z_0 = 1$ and $z_i = w_i/w_0, i = 1, \ldots, D$. The above equation can be written as

$$\begin{aligned} P_{s_0}(A) &= \int_{\mathbb{R}_+^{D+1}} \mathbb{1}\left\{z_i \in A_i, i = 1, \ldots, D\right\} w_0^{D+1} \frac{\phi\left((\boldsymbol{a}\boldsymbol{z})^{1/\nu}; \mathbf{0}_{D+1}, \Sigma\right)}{\Phi^\circ(\boldsymbol{\infty}; \mathbf{0}_{D+1}, \Sigma)} \prod_{i=0}^{D} (a_i w_0 z_i)^{1/\nu - 1} \frac{a_i}{\nu} \mathrm{d}w_0 d\boldsymbol{z}_{-1} \\ &= \int_{\mathbb{R}_+^{D+1}} \mathbb{1}\left\{z_i \in A_i, i = 1, \ldots, D\right\} w_0^{(D+1)/\nu} \frac{\phi\left((\boldsymbol{a}\boldsymbol{z})^{1/\nu}; \mathbf{0}_{D+1}, \Sigma\right)}{\Phi^\circ(\boldsymbol{\infty}; \mathbf{0}_{D+1}, \Sigma)} \prod_{i=0}^{D} (a_i z_i)^{1/\nu - 1} \frac{a_i}{\nu} \mathrm{d}w_0 d\boldsymbol{z}_{-1} \end{aligned}$$

Therefore, letting $\tilde{\boldsymbol{z}} = (\boldsymbol{a}\boldsymbol{z})^{1/\nu}$, the density of the distribution $P_{s_0}(\boldsymbol{z})$ is proportional to

$$\begin{aligned} f(\boldsymbol{z}) &\propto \prod_{i=1}^{D} \left\{z_i^{1/\nu - 1}\right\} \int_0^\infty w_0^{(D+1)/\nu} \exp\left\{-w_0^{2/\nu} \tilde{\boldsymbol{z}}^\top \Sigma^{-1} \tilde{\boldsymbol{z}}\right\} \mathrm{d}w_0 \\ &= \prod_{i=1}^{D} \left\{z_i^{1/\nu - 1}\right\} \int_0^\infty (2t)^{(D+1)/2} (\tilde{\boldsymbol{z}}^\top \Sigma^{-1} \tilde{\boldsymbol{z}})^{-(D+\nu+1)/2} e^{-t} \nu (2t)^{\nu/2 - 1} \mathrm{d}t \\ &\propto \prod_{i=1}^{D} \left\{z_i^{1/\nu - 1}\right\} \left(\tilde{\boldsymbol{z}}^\top \Sigma^{-1} \tilde{\boldsymbol{z}}\right)^{-(D+\nu+1)/2} \int_0^\infty t^{(D+\nu-1)/2} e^{-t} \mathrm{d}t \\ &\propto \prod_{i=1}^{D} \left\{z_i^{1/\nu - 1}\right\} \left(\tilde{\boldsymbol{z}}^\top \Sigma^{-1} \tilde{\boldsymbol{z}}\right)^{-(D+\nu+1)/2} \end{aligned}$$

Recall that B_D represents the set $\{1, \ldots, D\}$, given the block decomposition of the correlation matrix, $\Sigma = \begin{bmatrix} 1 & \Sigma_{0;B_D} \\ \Sigma_{0;B_D}^\top & \Sigma_{B_D;B_D} \end{bmatrix}$, the above equation can be written as

$$f(\boldsymbol{z}) \propto \prod_{i=1}^{D} \left\{z_i^{1/\nu - 1}\right\} \left[\frac{\tilde{\boldsymbol{z}}_{B_D}/\tilde{z}_0 - \Sigma_{B_D;0}}{\Sigma_{B_D;B_D} - \Sigma_{B_D;0}\Sigma_{B_D;0}} \left(\tilde{\boldsymbol{z}}_{B_D}/\tilde{z}_0 - \Sigma_{B_D;0}\right) + 1\right]^{-(D+\nu+1)/2},$$

which can be identified as the joint density function of $a_0 Z_i^\nu / a_i, i = 1, \ldots, D$, where $(Z_1, \ldots, Z_D)$ follows a multivariate truncated student-t distribution from below zero with scale parameter

$(\Sigma_{B_D;B_D} - \Sigma_{B_D;0}\Sigma_{B_D;0}^\top)/(\nu+1)$, location parameter $\Sigma_{B_D;0}$ and $\nu+1$ degrees of freedom. Algorithm 2 gives the pseudo-code of the exact simulation of the truncated extremal-t processes at finite location $\boldsymbol{s}_1, \ldots, \boldsymbol{s}_D$.

Algorithm 2: Exact simulation of truncated extremal-t processes at locations $\boldsymbol{s} = (\boldsymbol{s}_1, \ldots, \boldsymbol{s}_D)$.

Input: degree of freedom $\nu > 0$, $D \times D$ correlation matrix Σ
Output: $Z(\boldsymbol{s}_j)$, $j = 1, \ldots, D$.

Compute $\boldsymbol{a}$ from (D.1).
Simulate $E_0 \sim \text{Exp}(1)$ and set $z = 1/E_0$.
Set $Y(\boldsymbol{s}_1) = 1$ and simulate $Y(\boldsymbol{s})$ from a truncated-t distribution with location parameter $\Sigma_{-1;1}$, scale matrix $(\Sigma - \Sigma_{-1;1}\Sigma_{-1;1}^\top)/(\nu+1)$, and $\nu+1$ degrees of freedom at location $\boldsymbol{s}_i \neq \boldsymbol{s}_1$.
Set $W(\boldsymbol{s}_j) = a_0 Y^\nu(\boldsymbol{s}_j)/a_j$ for $j = 1, \ldots, D$.
Set $Z(\boldsymbol{s}_j) = W(\boldsymbol{s}_j)z$ for $j = 1, \ldots, D$.
for $i = 2, \ldots, D$ **do**
    Simulate $E_0 \sim \text{Exp}(1)$ and set $z = 1/E_0$.
    while $z > Z(\boldsymbol{s}_i)$ **do**
        Set $Y(\boldsymbol{s}_j) = 1$ and simulate $Y(\boldsymbol{s}_j)$, $j \neq i$ from a truncated-t distribution with location $\Sigma_{-j;j}$ and scale matrix $(\Sigma - \Sigma_{-j;j}\Sigma_{-j;j}^\top)/(\nu+1)$, and $\nu+1$ degrees of freedom.
        Let $W(\boldsymbol{s}_j) = a_0 Y^\nu(\boldsymbol{s}_j)/a_j$, $j = 1, \ldots, D$.
        if $W(\boldsymbol{s}_j) \times z < Z(\boldsymbol{s}_j), \forall j = 1, \ldots, i-1$ **then**
            Set $Z(s_j) = \max(Z(s_j), zW(s_j))$, $j = 1, \ldots, D$.
        Simulate $E_1 \sim \text{Exp}(1)$ and set $E_0 = E_0 + E_1$ and $z = 1/E_0$.

E Parameter estimation

E.1 Proof of Theorem 4

The distribution of the r-Pareto process with risk functional r is

$$\kappa_D(\cdot \cap \mathcal{A}_r)/\kappa_D(\mathcal{A}_r).$$

Suppose we have $\tilde{\boldsymbol{Y}} \in \Omega$ follow the angular distribution $\mathcal{H}/D$ and $\tilde{R}$ follows a unit Pareto distribution, i.e., $\Pr[\tilde{R} > x] = 1/x, x > 1$, we have

$$\Pr[\tilde{R}\tilde{\boldsymbol{Y}} \in \tilde{A}] = \tfrac{1}{D}\int_{\Omega}\int_{1}^{\infty} x^{-2}\mathbf{1}_{\{x\boldsymbol{w}\in\tilde{A}\}}\mathrm{d}x\mathcal{H}(\mathrm{d}\boldsymbol{w}).$$

If we have $\tilde{A} \subset \mathcal{A}_{r_0}$, then $x\boldsymbol{w} \in \tilde{A}$ leads to $r_0(x\boldsymbol{w}) = x\|w\|_1 = x > 1$. Therefore, we have

$$\Pr[\tilde{R}\tilde{\boldsymbol{Y}} \in \tilde{A}] = \tfrac{1}{D}\int_{\Omega}\int_{1}^{\infty} x^{-2}\mathbf{1}_{\{x\boldsymbol{w}\in\tilde{A}\}}\mathrm{d}x\mathcal{H}(\mathrm{d}\boldsymbol{w}) = \tfrac{1}{D}\int_{\Omega}\int_{0}^{\infty} x^{-2}\mathbf{1}_{\{x\boldsymbol{w}\in\tilde{A}\}}\mathrm{d}x\mathcal{H}(\mathrm{d}\boldsymbol{w}) = \kappa_D(\tilde{A})/D.$$

As we have $\Pr(\tilde{R}\tilde{\boldsymbol{Y}} \in \mathcal{A}_{r_0}) = \kappa_D(\mathcal{A}_{r_0})/D = 1/D$, it immediately follows that $\tilde{R}\tilde{\boldsymbol{Y}}$ is distributed as the r-Pareto process with risk functional r_0. Let $\tilde{r}(\cdot) = c_0 r(\cdot)$, then $\tilde{r}$ will be also a risk functional that is convex and homogeneous of order 1. Thus, we have $\tilde{r}(\boldsymbol{e}_i) = c_0/c_i \leq 1$. The complement set of $\mathcal{A}_{\tilde{r}}$ in $\mathbb{R}_+^D$, $\mathcal{A}_{\tilde{r}}^c = \{\boldsymbol{x} \in \mathbb{R}_+^D : \tilde{r}(\boldsymbol{x}) \leq 1\}$, is a convex set and contains points $\boldsymbol{e}_i, i = 1, \ldots, D$ on its boundary as well as the origin $\mathbf{0}$ since any homogenous function of order 1 maps the origin to the origin. For r_0, we know that $\mathcal{A}_{r_0}^c$ is a convex set and contains $\mathbf{0}$ and $\boldsymbol{e}_i, i = 1, \ldots D$ on its boundary, and $\mathcal{A}_{r_0}^c$ is the smallest convex set that contains points $\boldsymbol{e}_i, i = 1, \ldots, D$ and $\mathbf{0}$. Therefore, we have $\mathcal{A}_{r_0}^c \subset \mathcal{A}_{\tilde{r}}^c$, which implies that $\mathcal{A}_{\tilde{r}} \subset \mathcal{A}_{r_0}$. If we have $\tilde{A} \subset \mathcal{A}_{\tilde{r}}$, then we also have $\tilde{A} \subset \mathcal{A}_{r_0}$. Moreover, we can get

$$\Pr[\tilde{\boldsymbol{Z}}^{(r_0)} \in \tilde{A} | \tilde{\boldsymbol{Z}}^{(r_0)} \in \mathcal{A}_{\tilde{r}}] = \frac{1/D\kappa_D(\tilde{A}\cap\mathcal{A}_{\tilde{r}})}{1/D\kappa_D(\mathcal{A}_{\tilde{r}})} = \frac{\kappa_D(\tilde{A}\cap\mathcal{A}_{\tilde{r}})}{\kappa_D(\mathcal{A}_{\tilde{r}})} \tag{8}$$

Since κ_D is homogeneous of order -1 and $\mathcal{A}_{\tilde{r}} = 1/c_0\mathcal{A}_r$, it follows that

$$\begin{aligned}
&\Pr[\tilde{\boldsymbol{Z}}^{(r_0)} \in \tilde{A} | \tilde{\boldsymbol{Z}}^{(r_0)} \in \mathcal{A}_{\tilde{r}}] \\
&= \Pr[c_0\tilde{\boldsymbol{Z}}^{(r_0)} \in c_0\tilde{A} | c_0\tilde{\boldsymbol{Z}}^{(r_0)} \in \mathcal{A}_r] \\
&= \frac{\kappa_D(\tilde{A}\cap\mathcal{A}_r/c_0)}{\kappa_D(\mathcal{A}_r/c_0)} = \frac{c_0\kappa_D(c_0\tilde{A}\cap\mathcal{A}_r)}{c_0\kappa_D(\mathcal{A}_r)} = \frac{\kappa_D(c_0\tilde{A}\cap\mathcal{A}_r)}{\kappa_D(\mathcal{A}_r)}.
\end{aligned}$$

where $c_0\tilde{A} \subset \mathcal{A}_r$ since $\tilde{A} \subset \mathcal{A}_{\tilde{r}}$. Therefore, we have $\tilde{\boldsymbol{Z}}^{(r)} \overset{\mathrm{d}}{=} c_0\tilde{\boldsymbol{Z}}^{(r_0)} \mid r\left(c_0\tilde{\boldsymbol{Z}}^{(r_0)}\right) > 1$.

E.2 Comparing the angular and pairwise composite likelihoods

Consider samples of size $n = 500$ from the Brown-Resnick model defined on a 15×15 grid (i.e., $D = 225$) with 6 parameter combinations taking $\lambda = 3, 6$ and $\vartheta = 0.5, 1, 1.5$. For the angular likelihood ($10^\star$), the threshold is set as $r_0 = 30 \times D$, resulting in about 16 exceedances while the weights ζ_{B_k} in the composite likelihood ($9^\star$) take value 1 when the distance between two locations is less than $2\sqrt{2}$ and 0 otherwise, yielding 2268 pairs in the likelihood evaluation. After replicating the estimation procedure 300 times, violin plots of the estimators $\widehat{\lambda}$ and $\widehat{\vartheta}$ are displayed in Figure 1, using the angular (red) and pairwise composite likelihoods (blue), where the true parameter values are indicated by black dots. Results indicate that when in presence of relatively weak dependence (i.e., small λ and large ϑ), the bias exhibited by the angular likelihood estimates seems to increase, overestimating the range parameter λ and underestimating the smoothness parameter ϑ. The presence of bias is not surprising since data with weak dependence possess more effective independent samples compared to cases with stronger dependence, and has been observed in previous studies (see e.g. Engelke *et al.*, 2015; Huser *et al.*, 2016). As expected, the pairwise likelihood estimator produces unbiased estimates but uses 500 observations against approximately 16 for the angular likelihood. In terms of computation speed, the pairwise composite likelihood is about 476 times slower than the spectral likelihood in terms of core hours (average of 13.7 minutes with 16 CPU cores versus 9.2 seconds with 3 CPU cores). Overall, the angular likelihood approach does produce estimates that are biased although those aren't very large, in particular for stronger

dependence. As mentioned in Section 3.1 of the main manuscript, a censored likelihood approach could be taken at the expense of reducing the great computational gains. Furthermore, as dimension increases, the relative performance between the two approaches is expected to improve due to more available data (higher r_0, less bias).




Figure 1: Violin plots for 300 replicate estimates of $\boldsymbol{\theta} = (\lambda, \vartheta)$ for the Brown-Resnick process with $\lambda = 3, 6$ and $\vartheta = 0.5, 1, 1.5$ using the angular (red) and pairwise (blue) likelihoods. Black dots indicate the true values.

E.3 Comparing the angular likelihood and score matching approach

After considering the skewed Brown-Resnick r-Pareto process, we now generate 2000 samples from the truncated extremal-t r-Pareto process on a 8×8 grid with 8 parameter combinations using $\lambda = (3, 5)$, $\vartheta = (1, 1.5)$, and $\nu = 2, 3$ and L_1 and L_3 norms as risk functionals. The model is fitted using both the score matching and angular likelihood approaches with the outputs illustrated in Figure 4 when the L_3 and L_1-norm risk functional are used (respectively top and bottom two rows). We first note that there is barely any change in the performance of the estimators depending on the risk functional. No matter the norm used (L_3 or L_1), both estimation methods yield almost unbiased estimates for the smoothness parameter ϑ. Some bias is however present for the estimation of the range parameter λ, with the angular likelihood producing about 20% relative bias, while the score




Figure 2: Violin plots for the score matching (red) and angular likelihood (blue) estimates of λ (top) and b_2 (bottom) for the skewed Brown-Resnick r-Pareto process with L_3 norm risk functional and the same parameter combinations as in Figure 4 of the main manuscript. Black dots indicate the parameter true values.

matching method again shows numerical instability. Again, the angular likelihood shows greater computational efficiency, being about 10 times faster that the score matching method when using the L_3-norm risk functional (126 seconds on 3 CPU cores).

F Real-world applications

References

Azzalini, A. (2013) *The skew-normal and related families*. Volume 3. Cambridge University Press.

Dombry, C., Engelke, S. and Oesting, M. (2016) Exact simulation of max-stable processes. *Biometrika* **103**, 303–317.

Engelke, S., Malinowski, A., Kabluchko, Z. and Schlather, M. (2015) Estimation of Huesler–Reiss distributions and Brown–Resnick processes. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* **77**, 239–265.

Horrace, W. C. (2005) Some results on the multivariate truncated normal distribution. *Journal of multivariate analysis* **94**, 209–221.

Huser, R., Davison, A. C. and Genton, M. G. (2016) Likelihood estimators for multivariate extremes. *Extremes* **19**, 79–103.

Wadsworth, J. L. and Tawn, J. A. (2013) A new representation for multivariate tail probabilities. *Bernoulli* **19**, 2689–2714.

Wadsworth, J. L. and Tawn, J. A. (2014) Efficient inference for spatial extreme value processes associated to log-Gaussian random functions. *Biometrika* **101**, 1–15.




Figure 3: Violin plots for the score matching (red) and angular likelihood (blue) estimates of $\boldsymbol{\theta} = (\lambda, \vartheta, b_1, b_2)$ (top to bottom) for the skewed Brown-Resnick r-Pareto process with L_1 norm risk functional and the same parameter combinations as in Figure 4 of the main manuscript. Black dots indicate the parameter true values.




Figure 4: Violin plots for the score matching (red) and angular likelihood (blue) estimates of $\boldsymbol{\theta} = (\log(\lambda), \vartheta)$ for the truncated extremal-t r-Pareto processes with L_3 norm (top rows) and L_1 norm (bottom rows) risk functional, where $\lambda = c(3, 5), \vartheta = (1, 1.5)$ and ν is fixed to be either 2 (case 1–4) or 3 (case 5–8). Black dots indicate the true values.




Figure 5: Maps of bivariate empirical extremal coefficients with respect to two different reference point, and contours of the extremal coefficient of the Brown-Resnick r-Pareto model fitted using the angular likelihood (solid line) and score matching approach (dashed line) on the Florida rainfall data. The top and bottom rows respectively consider the L_1 and L_∞ norm risk functional.