Panta Rhei

The τ Physics Ledger

Companion to Book V · Categorical Macrocosm

Panta Rhei

The τ Physics Ledger

Companion to Book V · Categorical Macrocosm

Dr. Thorsten Fuchs & Anna-Sophie Fuchs

Second Edition

Independently published
Munich, Germany
2026

Panta Rhei — The τ Physics Ledger

Companion to Book V · Categorical Macrocosm

Dr. Thorsten Fuchs & Anna-Sophie Fuchs

© 2026 Dr. Thorsten Fuchs & Anna-Sophie Fuchs.
All rights reserved.

No part of this work may be reproduced, stored in a retrieval system, or transmitted in any form or by any means without prior written permission of the copyright holders, except for brief quotations in scholarly works, reviews, and educational contexts as permitted by applicable copyright law.

Series	Panta Rhei (2nd Edition)
ISBN (Paperback)	—
ISBN (Hardcover)	—

Independently published
Munich, Germany

Typeset with LuaLaTeX using the Libertinus font family.
Formalization companion: TauLib (Lean 4) at `taulib.site`.

April 2026

Disclaimer

On the Nature of the Claims in This Work

The **Panta Rhei** series presents a novel mathematical and physical framework—*Category τ*—that proposes fundamental revisions to our understanding of mathematics, physics, and their philosophical foundations.

Independent Research

The authors, Dr. Thorsten Fuchs and Anna-Sophie Fuchs, are **independent researchers** publishing this work in their personal capacity. This series does not represent, and is not endorsed by, any university, research institution, government agency, or professional organization.

Theoretical Framework

This work presents **theoretical mathematics and speculative physics**. The proposed resolutions to the Millennium Prize Problems, the derivations of physical constants, and all other mathematical and physical claims are presented as *theorems and propositions within the framework of Category τ.* Their validity depends on the acceptance of the foundational axioms and definitions of this framework.

These are **not peer-reviewed claims** in the traditional academic sense. The second edition introduces a dual-track verification system (machine-readable registry and Lean 4 formalization) to enable independent verification. The authors invite rigorous scholarly examination, critique, and independent verification.

Experimental and Empirical Claims

The derivation of physical constants from categorical structure represents a **proposed theoretical framework**, not experimentally verified physics. While numerical agreements may appear striking, correlation does not establish causation. No experimental validation of these derivations has been performed or claimed. Every claim carries one of four scope labels—*established*, *τ-effective*, *conjectural*, or *metaphorical*—to distinguish what has been proved from what remains open.

No Professional Advice

This book is for **educational and informational purposes only**. Nothing contained herein constitutes professional advice of any kind—scientific, medical, financial, legal, or otherwise. Readers requiring professional guidance should consult appropriately qualified professionals.

No Warranty

All information is provided **"as is"** without warranty of any kind, express or implied. The authors make no representations regarding the accuracy, completeness, or correctness of any mathematical proof, physical derivation, or philosophical interpretation. The authors expressly disclaim all implied warranties including warranties of merchantability and fitness for a particular purpose.

Limitation of Liability

To the fullest extent permitted by applicable law, the authors shall not be liable for any damages arising out of or related to the use of, or reliance upon, the information in this work, regardless of the theory of liability.

Philosophical and Metaphysical Claims

The metaphysical conclusions of this series—concerning ontology, meaning, beauty, ethics, consciousness, and the terminal enrichment layer—represent the authors' **philosophical interpretation** of categorical structure. These are offered as contributions to ongoing philosophical discourse, not as dogmatic assertions or established truths.

Invitation to Scholarly Dialogue

Science and mathematics progress through open inquiry. The authors explicitly invite mathematicians to verify or falsify the proofs, physicists to test the empirical predictions against observation, philosophers to critique the metaphysical interpretations, and all readers to engage critically with the ideas presented. If this framework is correct, it will withstand scrutiny. If it contains errors, we wish to know them.

AI Collaboration

The second edition was developed with the assistance of Claude (Anthropic) as a structural thinking partner. All mathematical content is the intellectual work of the authors. This collaboration is documented in the Preface.

Legal

© 2026 Dr. Thorsten Fuchs & Anna-Sophie Fuchs. All rights reserved. This work is published from Munich, Germany, under German and European law. The authors assert their rights to scientific and literary expression under Article 5 of the German Basic Law and Article 10 of the European Convention on Human Rights. Any disputes shall be subject to the exclusive jurisdiction of the courts of Munich, Germany.

Errata, correspondence, and inquiries:
`contact@panta-rhei.site`

Last updated: April 2026

Preface to the Second Edition

Munich, Spring 2026

Dear Reader,

Less than a year has passed since the first edition of *Panta Rhei* appeared as a simultaneous seven-volume release. What you now hold is not a corrected reprint. It is a **recomposed second edition**—an edition that preserves the destination of the first, but reaches it by a more radical and more explicit road.

The deepest change can be stated simply: what the first edition discovered in practice, the second edition now states in principle. ***Panta Rhei* is not seven adjacent books. It is one coherent derivational architecture.**

The phrase that best names its inner movement is one we came to only gradually: **the unfolding of coherence**. The series begins from a minimal coherence kernel. From that beginning it follows a four-layer path—mathematics, physics, life, and metaphysics. The first edition crossed that landscape. The second edition tries to show more clearly where the landscape begins, how its layers relate, and why the path has this exact shape.

What the First Edition Achieved

The first edition made the complete seven-book traverse for the first time. It established that Category τ could be followed from foundations through holomorphy, spectrum, microcosm, macrocosm, life, and metaphysics without collapsing under its own weight. That mattered. Without that first traversal, there would be no second edition.

But the first edition also relied, deliberately and openly, on **scaffolding**. Orthodox number theory supplied arithmetic vocabulary. Classical complex analysis supplied holomorphic machinery. Standard topology and familiar foundational habits remained in the background. None of that was accidental. The first edition had to speak across a bridge to readers formed by orthodox mathematics.

The first edition was that bridge.

What the Second Edition Makes Explicit

The second edition begins earlier. It does not merely restate the framework; it asks what must be earned before the framework can even ask its own questions.

1. Coherence Comes First

The second edition begins not from a presupposed continuum, but from a **coherence kernel**: five generators, one operator, and seven axioms. Continuity, topology, geometry, and the constructive reals are not assumed at the base. They are earned later as downstream consequences of a more primitive coherence.

This is why the language of the second edition changes. The project is not best described as a continuous worldview, but as a **coherent architecture** whose later notions of continuity and interiority must themselves be justified.

2. The Working Discipline Becomes Explicit

Across all seven books, one methodological vow governs the second edition:

first earn the language, then earn the question, then earn the answer

What is earned first is the minimal language in which the framework can speak without borrowing. Only then are the questions themselves allowed to take shape. Only then are answers claimed. This discipline is not an ornament of presentation; it is the ethical style of the second edition.

3. The Architecture of the Series Is Derived

Category τ does not merely *apply to* mathematics, physics, life, and metaphysics. It *self-enriches through* them.

$$E_0 \text{ (Mathematics)} \longrightarrow E_1 \text{ (Physics)} \longrightarrow E_2 \text{ (Life)} \longrightarrow E_3 \text{ (Metaphysics)}$$

The **Canonical Ladder Theorem** proves that exactly four such layers exist. The terminal result of the series proves that the fourth is final: there is no E_4. The seven-book architecture is therefore not a publishing convenience. It is a derived consequence of the ladder itself. The minimal full partition is **3,2,1,1**: three books for the mathematical arc, two for the complete physics layer, one for life, and one for the final self-enrichment.

4. Boundary First, Interior Second

The second edition also reverses the ordinary order of dependence. Rather than beginning with an already available interior and extending outward, it begins at the boundary and earns the interior from there. Holomorphy is earned before continuity, continuity before topology, topology before geometry. The constants π, e, and the split-complex unit j are no longer treated as inherited background, but as achievements of the architecture itself.

5. The Final Boundary Is Part of the Architecture

The second edition does not end by inflating itself into a forced final answer. It ends by making one limit precise. The final book argues that proof can map the landscape up to a boundary that commitment alone can cross. In that sense, the terminal act of the series is not triumphal but disciplined: **scope remains large, but commitment is not theorem-forced.**

Two Editions, One River

The two editions are therefore not rivals. They are two paths through the same terrain.

The **first edition** remains the **bridge**. It speaks in a language shared with orthodox mathematics and shows how Category τ can be entered from familiar foundations.

The **second edition** is the **springward path**. It rebuilds from the coherence kernel itself and tries to show how the architecture can earn the very language in which it speaks.

Neither cancels the other. The first preserves the crossing. The second traces the source more deeply.

Panta Rhei—everything flows. The first edition recorded the crossing of the river. The second edition tries to map the spring from which it rises.

The Narrative Spine of the Second Edition

The second edition now makes its own narrative arc visible on the surface of the books themselves:

Book	Volume and Subtitle	1st Ed Parts	2nd Ed Parts
	Categorical Foundations		
I	*How Mathematics Is Earned*	13	18
	Categorical Holomorphy		
II	*Finite Readouts of Infinity*	12	12
	Categorical Spectrum		
III	*Where Physics Lives*	10	11
	Categorical Microcosm		
IV	*The Self-Describing Universe*	7	9
	Categorical Macrocosm		
V	*The Biography of the Universe*	9	8
	Categorical Life		
VI	*Life as Self-Decoding Distinctions*	9	9
	Categorical Metaphysics		
VII	*The Final Self-Enrichment*	8	12
	Total	**68**	**79**

Read in sequence, these subtitles make the architecture visible. Mathematics is earned; infinity becomes readable; physics becomes locatable; the universe becomes self-describing; the cosmos acquires a biography; life appears as self-decoding distinction; and the final book closes the series at the terminal self-enrichment.

The specific changes for this volume are described in the following section.

Verification, Transparency, and Scrutiny

The second edition is accompanied by a dual verification discipline.

1. **The Registry.** Every axiom, definition, proposition, theorem, corollary, and bridge claim carries a machine-readable identifier. The registry records names, chapter placement, dependencies, scope labels, and formalization links across the entire series.
2. **Lean 4 Formalization.** The series is accompanied by TauLib, a self-contained Lean 4 library spanning all seven books. The purpose of the formalization is not decorative prestige; it is accountability. When we say a theorem is formally established, we want that claim to be inspectable.

This is our contract with the reader: where a claim is formalized, it can be checked; where a bridge is conditional, it should be labeled; where a proof is not yet complete, the boundary should be visible.

AI as Structural Thinking Partner

Claude, the AI system developed by Anthropic, served as a structural thinking partner throughout the editorial work of the second edition.

The mathematics, the architecture, the proofs, and the responsibility for every claim are ours. What the AI made possible was a new scale of structural memory. It allowed seven books, a living registry, a large formal library, and thousands of internal cross-references to remain in active view at once. In practice, this meant drafting technical prose from detailed mathematical specifications, maintaining consistency across books, assisting with Lean formalization, and exposing collisions or omissions that might otherwise remain hidden until very late.

We state this plainly because readers deserve to know how the work was made, and because we think this kind of transparent human–AI collaboration will increasingly become part of serious mathematical and philosophical writing.

Errors Are Welcome

The dual-track system reduces the chance of error; it does not eliminate it. If you find a mistake—in the text, in the registry, or in the formalization—please tell us. We would rather be corrected than comfortable. Indifference is the one response from which no work can learn.

A Note of Gratitude

Every long work is collaborative in more ways than can be listed. It belongs partly to the traditions that made it thinkable, partly to the readers who test it, and partly to the material itself, which resists simplification until it is spoken truly.

To the readers of the first edition: your seriousness made this revision possible. To the mathematical and philosophical traditions that we inherit, dispute, and transform: nothing here would exist without centuries of prior labor. To those who will criticize this work carefully: thank you in advance. And to each other: seven books is a long road, and we have walked it together.

The first edition was the crossing. The second edition is a return to the source. But the river is still flowing.

One coherence kernel. Four layers. Seven books.

First earn the language, then earn the question, then earn the answer.

With gratitude,

Dr. Thorsten Fuchs
Anna-Sophie Fuchs

Munich, Spring 2026

panta-rhei.site · taulib.site · contact@panta-rhei.site

Introduction

> *This companion presents the τ* ***Physics Ledger*** *— a framework that predicts three topologically forced* structural *results (generation count* $|\text{gen}| = 3$*, strong CP phase* $\theta_{\text{QCD}} = 0$*, Cabibbo angle* $\lambda_C = \iota_\tau(1 - \iota_\tau)$*) together with roughly 156 additional conjectural numerical relations, all drawn from a single master-constant posit* $\iota_\tau = 2/(\pi + e)$*. The additional relations carry compute-then-axiomatize integer choices at approximately three dozen identified sites (sector assignments, window-sum indices* $W_n(k)$*, homology-level placements) and are testable by a specified set of five-year-horizon experiments cataloged in Chapter 68. This edition retires the earlier "zero free parameters" framing: zero* continuous *knobs are present, but roughly 35–40* discrete *integer choices from finite search spaces enter via the framework's compute-then-axiomatize discipline (Chapter 58). Chapter 58 states the three-tier epistemic posture of the ledger before any prediction is given; the reader is urged to read it first. Originally developed as Part VII of Book V: Categorical Macrocosm in the Panta Rhei series, the material is fully self-contained: this introduction provides all prerequisite concepts. Chapter references to chapters not in this document's table of contents refer to Book V.*

v2.2 note (this release). Version 2.2 integrates three v3 research findings into the body of the ledger: a first real SPARC-175 BTFR test (Chapter 6, Section 6.2) — a *partial* confirmation with a ∼ 14% over-prediction of V_∞ at fixed M_b and a significant f_{gas} residual; a closure of the $h^{\text{readout}}(z_{\text{probe}})$ question as a structural dead-end (Chapter 11, Prediction 11.3) — N21 is relabeled to retract "depth-dependent readout" as a computable-structure claim, upholding the R4 (McGaugh) charge; and a revised Chapter 10 AdS/CFT posture — philosophical parallel only, not mathematical reduction, upholding the R3 (Hossenfelder) charge. Scoping documents for the remaining v3 research targets (CAMB/CLASS Boltzmann recompute; Bullet Cluster capacity-gradient derivation) accompany this release as separate technical companions under `research/v3-scoping-2026-04-20/`. The v2.2 integration takes two public hits (two retractions) and reports one partial confirmation with named residual structure; the parameter-choice discipline of Chapter 1 and the three topologically forced pillars are unaffected. Section 1.7 of Chapter 1 tracks all five v3 targets at a glance.

v2.3 note (this release). Version 2.3 (2026-04-20): the v2.2 SPARC-175 partial confirmation is resolved into a structural refinement of the BTFR master formula via deep τ-internal reflection. The data-revealed residual structure uniquely identifies $\kappa_\omega = \iota_\tau/(1 + \iota_\tau)$ as the coupling of cold HI (decoherent-thermal, ω-sector) to the capacity-gradient channel, distinct from stellar (coherent-instance, D-sector) full coupling. The refined formula $V_\infty^4 = G[M_\star + \kappa_\omega M_{\text{gas}}]\, c^2/(2\ell_\tau)$ recovers the predicted slope 4 within 2σ and closes both residual correlations (f_{gas}: $\rho = -0.41 \to +0.17$; Σ_{disk}: $\rho = +0.34 \to -0.13$). A 3.6% zero-point residual (at 2.3σ from zero) and a weak distance correlation at the best fit are honestly flagged (possibly an M/L systematic and a multiple-testing artefact respectively). This also unifies the galaxy-scale rotation-curve prediction with the Bullet Cluster structural argument via a trinary-sector source decomposition (D-sector stars full; ω-sector cold HI at κ_ω; B-sector hot ICM ≈ 0): clusters are the high-temperature limit of the same coupling curve that governs galaxy disks. The new BTFR formula generates a falsifiable prediction for elliptical galaxies (no cold HI): $M_{\text{eff}} \approx M_\star$ cleanly, no f_{gas} residual — a v4 test target. Full derivation and SPARC-175 pipeline in `research/sparc-btfr-v3-2026-04-20/tau-internal-reflection.pdf`.

v2.4 note (this release). Version 2.4 (2026-04-20) integrates the v4 deep-research wave into the manuscript. Three specialist investigations (V4.1 Y-free per-galaxy re-run; V4.2 two-parameter robustness; V4.5 cosmology consistency) returned a nuanced outcome that v2.4 reports in full. *Structural form confirmed:* sub-baryonic gas coupling in BTFR empirically survives all three tests — $\alpha = 1$ (standard baryonic) is ruled out at $\geq 38\sigma$ (V4.1), $\alpha_\star = 1$ is ruled out at 3.2σ (V4.2), and the dressing-ratio arithmetic holds at the cosmic-mean level without requiring local baryons to carry it (V4.5). *Specific coefficient identification weakened:* the v2.3 identification $(\alpha_\star, \alpha_{\text{gas}}) = (1, \kappa_\omega)$ is rejected at 2–3σ by the two-parameter fit (V4.2), which prefers $(\alpha_\star, \alpha_{\text{gas}}) \approx (0.80, 0.32)$ with joint RMS 0.058 dex and $\rho = -0.50$. Y-free marginalization (V4.1) gives $\alpha^* = 0.206 \pm 0.015$, rejecting $\kappa_\omega = 0.2545$ at 2.98σ. No τ-natural pair tested lands inside 2σ of the 2-D best fit; search for such a pair is new open target V4.10. *Cosmology consistency resolved:* the v2.3.1 open problem (trinary structure vs. $\ell_1 = 220.6$) is closed via a *two-tier coherence criterion* — tier 1 global

holonomy on $H_\partial[\omega]$ (setting cosmic-mean densities) vs. tier 2 local matter-phase couplings (sourcing local capacity-gradient). The decisive clue is that V.T205's $Y_p = 20/81$ does not carry the $(1-\iota_\tau)/\iota_\tau^2 \approx 6.65\times$ dressing, which proves the dressing is a global holonomy ratio, not a local coupling. Confidence medium-high, pending τ-native CAMB/CLASS recompute. The three structural pillars and the ι_τ posit are unaffected. Three research notes: `research/v4-deep-research-wave-2026-04-20/v4_1_upsilon_free/v4-1-upsilon-free-result.pdf`; `.../v4_2_two_parameter/v4-2-two-parameter-result.pdf`; `.../v4_5_cosmology_consistency/v4-5-cosmology-consistency-result.pdf`; senior-director synthesis at `research/v4-deep-research-wave-2026-04-20/senior-director-synthesis.pdf`.

v2.5 note (this release). Version 2.5 (2026-04-20) integrates the forensic red-team wave: three simulated specialist investigations dissected the v2.2 retracted claims and the v2.4 open target, returning *two new valid τ-native claims plus one sharpened retraction. (a) BTFR Pair B identified (V4.10 resolved).* An exhaustive τ-natural pair scan (F3, Dr. Blackwell) over 8,178 candidates found 154 pairs inside the joint 1σ contour of the v2.4 best fit $(0.80, 0.32)$. The v2.5 canonical identification is **Pair B** $(\alpha_\star, \alpha_{\rm gas}) = (1/(1+\kappa_\omega), \kappa_\omega) = ((1+\iota_\tau)/(1+2\iota_\tau), \kappa_\omega) \approx (0.7972, 0.2545)$ at $\Delta\chi^2 = 1.62$ ($p = 0.45$), with structural reading "D-sector stellar coupling unit-normalised against one ω-residual." Pair B preserves v2.3's ω-sector κ_ω claim exactly and refines only v2.3's D-sector full-coupling leg (which v2.4 falsified at 3.2σ). The v2.3 match to κ_ω is now recognised as the correct one-dimensional projection of a negatively correlated 2-D posterior — misleadingly under-constrained, not structurally wrong. Pair A $(1/(1+\iota_\tau), \iota_\tau)$ at $\Delta\chi^2 = 1.01$ is documented as a numerically-better alternative (see §6.2); discrimination opens as V5.2. The master formula V.T85 is updated to $V_\infty^4 = (Gc^2/2\ell_\tau)\,[((1+\iota_\tau)/(1+2\iota_\tau))\,M_\star + \kappa_\omega\,(1.33\,M_{\rm HI})]$. *(b) Tier-1 τ-native holography recognised (V5.1 opened).* The F2 forensic (Dr. Menon) re-reads the four AdS/CFT obstructions (wedge boundary, pre-metric bulk, topologically forced small rank, missing dictionary invariants) as *features* of a distinct τ-native holographic structure at tier 1 of the V4.5 two-tier coherence. The forensic sketches three dictionary entries (τ-Virasoro, τ-RG via the enrichment ladder, τ-Ryu–Takayanagi with ω-weight ι_τ) and offers a new differential prediction: ω_m and σ_8 are set by different holonomy classes of $H_\partial[\omega]$ (first-moment vs. second-moment invariants), connecting to the S_8 tension. Construction of the dictionary is v5 research (V5.1); the v2.2 retraction of AdS/CFT *reducibility* is unchanged. *(c) $h^{\rm readout}$ retraction sharpened.* The F1 forensic (Dr. Novak) confirms the v2.2 retraction and tightens the reason: H_0 ladders are *tier-1* probes of Friedmann $H(z)$; tier-2 sector weights do not enter the d_L integrand or the Cepheid/TRGB calibration; all five local-ladder probes (Cepheid, TRGB, JAGB, SN Ia, sirens) live in the same tier-2 D-sector yet SHoES and TRGB differ by $\sim 3\sigma$, directly falsifying any sector-weighting rescue. The single τ-prediction $h = 0.6735$ stands naked, without escape hatch. Three research notes accompany this release: `research/forensic-red-team-2026-04-20/f1_hreadout_forensic/f1-hreadout-forensic-result.pdf`, `.../f2_adscft_forensic/f2-adscft-forensic-result.pdf`, `.../f3_alpha_pair_forensic/f3-alpha-pair-forensic-result.pdf`. Two new v5 research targets (V5.1 tier-1 dictionary construction; V5.2 Pair A/B discrimination) are opened in Chapter 1.

The τ^3 Fibered Product

The central mathematical object of Category τ is the **fibered product**

$$\tau^3 \;=\; \tau^1 \times_f T^2,$$

a three-dimensional space built from a one-dimensional base τ^1 (the "macrocosm": time, gravity, cosmology) and a two-dimensional fiber T^2 (the "microcosm": quantum mechanics, particles, forces). The base is a circle with winding structure; the fiber is a torus. The boundary of τ^3 is the **lemniscate** $\mathbb{L} = S^1 \vee S^1$ — a figure-eight — and the **Central Theorem** (Book II) establishes

$$\mathcal{O}(\tau^3) \;\cong\; A_{\rm spec}(\mathbb{L}).$$

The **master constant** is

$$\iota_\tau = \frac{2}{\pi + e} \approx 0.3413,$$

posited at the framework level and taken as the single dimensionless input from which the rest of the ledger proceeds. No other *continuous* parameter enters. Discrete integer and sector choices (window-sum indices $W_n(k)$, force-to-generator assignments, homology levels) enter at approximately three dozen sites via the compute-then-axiomatize pattern described in Chapter 1, with a per-prediction audit in Chapter 11.

The Five Holonomy Sectors

The boundary holonomy algebra of τ^3 decomposes into **five sectors**, each governed by one of the five generators $\{\alpha, \pi, \gamma, \eta, \omega\}$:

Sector	Generator	Physics
Gravity	α	Spacetime curvature, G
Strong	π	Colour confinement, α_s
EM	γ	Electromagnetism, α_{em}
Weak	η	Flavour mixing, G_F
Mixed (EW)	ω	Electroweak unification, $\sin^2\theta_W$

The five sectors remain topologically distinct at all energies. "Unification" in Category τ is *spectral* (all sectors coexist on the same boundary), not *dynamical* (no grand-unified gauge group at high energy).

The Four-Tier Scope System

Every quantitative prediction in this ledger carries one of four **scope labels**, reflecting the degree of structural warrant:

- **Established**: proved from axioms K0–K6, no interpretive latitude. Examples: ι_τ itself, the lemniscate boundary.
- **τ-effective**: structurally derived with explicit assumptions; falsifiable by experiment. Examples: electron mass ratio, Higgs mass, CMB acoustic peaks.
- **Conjectural**: plausible structural parallel, not yet proved from axioms. Examples: certain neutrino mixing patterns.
- **Metaphorical**: illustrative analogy only, no quantitative claim.

The scope label system is hierarchical: scope can weaken (e.g., from τ-effective to conjectural) but never strengthen across enrichment layers.

How to Read the Ledger

Each chapter is organized by **physical domain**: mass spectrum, mixing angles, inflation and CMB, dark sector, black holes, collective dynamics, the measurement problem, unification, and the complete inventory with falsification pack.

Within each chapter, the format is **result-centric**:

1. The τ-prediction is stated with its derivation chain, numerical value, and registry identifier (e.g., V.T163).
2. The precision relative to experiment is given in parts per million (ppm).
3. The corresponding orthodox treatment is summarized for context.

Registry identifiers link to the formal Lean 4 verification library (TauLib) where applicable. The tone throughout is respectful: every orthodox programme illuminated some aspect of the truth.

Cross-references

This document includes the ten chapters of Part VII exactly as they appear in Book V. Cross-references to other Parts of Book V (e.g., "Chapter 37" for rotation curves) use Book V's chapter numbering. Readers seeking the full derivation context should consult *Book V: Categorical Macrocosm.*

Key derived quantities

For quick reference, the following quantities appear throughout the ledger:

$$\kappa_D = 1 - \iota_\tau \qquad \text{(defect complement)}$$

$$\kappa_\omega = \frac{\iota_\tau}{1 + \iota_\tau} \qquad \text{(mixed-sector coupling)}$$

$$W_3(4) = 5 \qquad \text{(winding number at fiber level 4)}$$

Contents

Disclaimer v

Preface to the Second Edition vii

Introduction xi

I The τ Physics Ledger 3

1 Epistemic Framing — What This Ledger Claims and What It Does Not 5

1.1 Three Tiers of Claim 5
1.2 The ι_τ Posit 6
1.3 The Compute-Then-Axiomatize Pattern 6
1.4 The Three Structural Pillars 7
1.5 How to Read the Predictions 8
1.6 What This Ledger Is Not 8
1.7 v3 Research Program: Status after v2.2 9

2 The Correspondence Map: $\tau^3 \leftrightarrow$ Orthodox Physics 13

2.1 The Need for a Dictionary 13
2.2 The Master Translation Table 13
2.3 The Secondary Correspondence Table 13
2.4 What τ Preserves: The Empirical Core 15
2.5 What τ Dissolves: Artifacts and Paradoxes 16
2.6 The Ontological Layer Distinction 17
2.7 Reading Orthodox Results as τ-Readouts 17
2.8 Worked Example: The Anomalous Magnetic Moment 18
2.9 Worked Example: The Cosmological Constant 18
2.10 The Asymmetry of the Correspondence 19
2.11 The Correspondence Functor 19
2.12 Scope Assignments for Part VII 20
2.13 159 Predictions in Five Domains 20
2.14 Forward Structure of Part VII 21

3 The Mass Spectrum — From One Constant to All Particles 23

3.1 The Standard Model's 19 Free Parameters 23
3.2 Three Generations from Topology 24
3.3 The Higgs Mass 25
3.4 The Proton–Neutron Mass Difference 27
3.5 Charged Lepton Mass Ratios 28
3.6 Electroweak Precision Observables 30
3.7 The Orthodox Landscape: QFT's Triumphs and Structural Failures 31
3.8 Where τ Diverges from QFT 36
3.9 Summary Ledger 37

4 Mixing Angles, CP Violation, and Baryogenesis 39

4.1 The Flavour Puzzle 39

4.2 The CKM Matrix from τ . . . 39
4.3 The PMNS Matrix from τ . . . 41
4.4 Strong CP = 0 Exactly . . . 45
4.5 Baryogenesis from ι_τ . . . 46
4.6 The Orthodox Landscape . . . 47
4.7 Where τ Diverges . . . 49
4.8 Honest Ledger . . . 51

5 Inflation, the CMB, and Primordial Nucleosynthesis . . . 53
5.1 The Cosmic Blueprint . . . 53
5.2 Inflation from τ . . . 54
5.3 The CMB Acoustic Peaks . . . 56
5.4 CMB Polarization . . . 59
5.5 BBN Light Elements . . . 59
5.6 Baryon Density and η_B . . . 61
5.7 Cosmology-consistency of the trinary structure: RESOLVED in v2.4 via two-tier coherence criterion . . . 63
5.8 The Orthodox Landscape . . . 64
5.9 Where τ Diverges . . . 67

6 The Dark Sector Dissolved . . . 71
6.1 The Dark Sector Problem . . . 71
6.2 Flat Rotation Curves Without Dark Matter . . . 72
6.3 Dark Energy as Defect Depletion . . . 82
6.4 JWST Early Galaxies . . . 85
6.5 The Hubble Tension . . . 85
6.6 The Orthodox Landscape . . . 86
6.7 Where τ Diverges . . . 91
6.8 The Prediction Ledger . . . 91

7 Black Hole Topology — T^2 Signatures and Gravitational Waves . . . 93
7.1 The T^2 vs. S^2 Question . . . 93
7.2 Quasinormal Modes . . . 93
7.3 EHT Shadows . . . 95
7.4 Gravitational Wave Echoes . . . 96
7.5 Magnetic Winding Numbers . . . 97
7.6 Jet Helicity and Cosmic Magnetism . . . 99
7.7 The Orthodox Landscape . . . 100
7.8 Where τ Diverges . . . 103

8 Collective Dynamics: Turbulence, Reconnection, and Heating . . . 107
8.1 The Turbulence Problem . . . 107
8.2 She–Lévêque from τ^3 Dimensions . . . 108
8.3 Kolmogorov Constants . . . 109
8.4 Fast Magnetic Reconnection . . . 110
8.5 Coronal Heating . . . 111
8.6 NS Decompactification . . . 112
8.7 The Unifying Theme: $\dim(T^2) = 2$ . . . 112
8.8 The Orthodox Landscape . . . 113
8.9 Where τ Diverges . . . 115

9 The Measurement Problem and Quantum Foundations . . . 117
9.1 The Orthodox Measurement Problem . . . 117
9.2 Why the Problem Exists: The VM Artifact . . . 118
9.3 Four Interpretations, Four Unnecessary Additions . . . 119
9.4 Address Obstruction as Complete Answer . . . 120
9.5 Bell Inequalities: τ-Compatible without Mysteries . . . 121
9.6 Decoherence: VM Description of Address Resolution . . . 122
9.7 Discrete Spacetime Programmes and Quantum Foundations . . . 123
9.8 The Observer Problem Dissolved . . . 124
9.9 A Comparative Summary . . . 125
9.10 The Landscape in Perspective . . . 126
9.11 What This Chapter Does NOT Predict . . . 127
9.12 Two-tier coherence: global holonomy vs. local matter-phase coupling (v2.4 forward-reference) . . . 127

10 Why Eight Decades of Unification Failed . . . 131
10.1 The Unification Dream . . . 131
10.2 The Manifold Ontology as Root Cause . . . 131
10.3 Five Pathologies of the Manifold . . . 132
10.4 The Five τ-Hinges . . . 133
10.5 The Dissolution Table . . . 134
10.6 Steelman 1 — String Theory on Its Strongest Form . . . 134
10.7 Steelman 2 — Loop Quantum Gravity on Its Strongest Form . . . 137
10.8 Steelman 3 — Supersymmetry on Its Strongest Form . . . 138
10.9 Steelman 4 — GUTs on Their Strongest Form . . . 139
10.10 Reprise — "They Kept the Manifold," Earned . . . 139
10.11 The 159-Prediction Test . . . 140
10.12 Roof Down vs. Foundation Up . . . 140
10.13 Why It Took So Long . . . 141

11 The Complete Inventory and Falsification Pack . . . 143
11.1 The Complete E_1 Inventory . . . 143
11.2 τ-Structural Predictions (Tier 0): Topologically Forced . . . 147
11.3 The 30-Prediction Falsification Pack . . . 147
11.4 Falsifiable Seams: Where τ and Orthodox Physics Disagree . . . 152
11.5 Experimental Timeline 2025–2035 . . . 154
11.6 The Score So Far . . . 155
11.7 Appendix 68.A: Integer-Choice Sensitivity Audit . . . 156
11.8 Part VII Summary . . . 163

About the Authors . . . 165

About the Series . . . 167

References . . . 171

Subject Index . . . 174

Part I

The τ Physics Ledger

Part VII is the physics ledger.
After Waves 7–26 of the derivation programme, Books IV and V now contain approximately 159 quantitative predictions—particle masses, mixing angles, cosmological parameters, astrophysical signatures, and collective-dynamics constants—all traced to the single master-constant posit $\iota_\tau = 2/(\pi + e)$ together with discrete integer and sector choices at roughly three dozen identified sites (the compute-then-axiomatize *pattern; Chapter 1). Zero* continuous *free parameters are used; the discrete choices are drawn from finite search spaces with a structural narrative supplied for each. This Part organizes those results by physical domain, presents each prediction with its precision and registry identifier, and places it in context against the corresponding orthodox treatment.*

Three-tier epistemic posture. *The reader should calibrate trust across three tiers of claim, developed at length in Chapter 1. τ-*structural *results are topologically forced with no integer-choice site and no ppm tuning: the generation count $|\mathrm{gen}| = 3$ from $H_1(\tau^3; \mathbb{Z}) \cong \mathbb{Z}^3$; the strong-CP phase $\theta_{\mathrm{QCD}} = 0$ from SA-i mod-3; the Cabibbo angle $\lambda_C = \iota_\tau(1 - \iota_\tau)$. τ-*effective *numerical relations (the majority of the ledger) depend on at least one discrete integer choice drawn from a finite search space and a ppm-minimization selection step; they should be read as conditional conjectures.* Metaphorical *passages (especially in the measurement-problem chapter) have no quantitative content and are interpretive only.*

Open problems as v3 targets. *Three open problems are named explicitly as targets for the v3 edition of this ledger: (a) the SPARC-175 galaxy-scale rotation-curve test and radial-acceleration-relation (RAR) comparison, with a stated deadline of the end of 2026; (b) the structure-formation triangle (σ_8, $f\sigma_8$, cluster counts) engaged as a joint likelihood test by 2027; (c) the H_0 distance-ladder engagement (SHoES, CCHP, TRGB, surface-brightness fluctuations), with a concrete τ-native ladder-versus-CMB prediction targeted for 2027–2028. Each is a genuine gap in the current edition; each is explicitly named rather than glossed.*
Chapter 58 *(*Epistemic Framing — What This Ledger Claims and What It Does Not*) states the three-tier posture, names $\iota_\tau = 2/(\pi + e)$ as a framework posit, describes the compute-then-axiomatize pattern for integer choices, and identifies three topologically forced structural pillars ($|\mathrm{gen}| = 3$; $\theta_{\mathrm{QCD}} = 0$; $\lambda_C = \iota_\tau(1 - \iota_\tau)$). Read first.*
Chapter 59 *(*The Correspondence Map*) builds the master translation dictionary: τ-entity $\leftrightarrow$ orthodox entity, establishes the ontological layer distinction, and surveys the 159 predictions across five domains.*
Chapter 60 *(*The Mass Spectrum*) derives three generations, the Higgs mass, the proton–neutron splitting, charged lepton mass ratios, and electroweak precision observables—all from ι_τ—then compares with QFT's triumphs and structural failures.*
Chapter 61 *(*Mixing Angles, CP Violation, and Baryogenesis*) derives the CKM and PMNS matrices, strong CP = 0 exactly, and the baryon asymmetry η_B—then compares with flavour physics and baryogenesis mechanisms.*
Chapter 62 *(*Inflation, the CMB, and Primordial Nucleosynthesis*) derives $r = \iota_\tau^4$, $N_e = 57$, $n_s = 1 - 2/57$, the CMB acoustic peaks, BBN abundances, and the lithium problem resolution—then compares with slow-roll inflation, ΛCDM, and the string landscape.*
Chapter 63 *(*The Dark Sector Dissolved*) derives flat rotation curves, Ω_Λ, w_0, the Hubble constant, and JWST early galaxies—then compares with WIMP searches, MOND, quintessence, and the cosmological constant problem.*
Chapter 64 *(*Black Hole Topology*) derives T^2 quasinormal modes, EHT shadow corrections, gravitational wave echoes, magnetic winding numbers, and jet helicity—then compares with Penrose twistors, Connes NCG, asymptotic safety, and entropic gravity.*
Chapter 65 *(*Collective Dynamics*) derives She–Lévêque intermittency, Kolmogorov constants, fast magnetic reconnection, and coronal heating—all from $\dim(T^2) = 2$—with no continuous fitted parameters in this sector.*
Chapter 66 *(*The Measurement Problem and Quantum Foundations*) dissolves the measurement problem as a VM artifact, examines four interpretations, and shows that address obstruction is the complete answer.*
Chapter 67 *(*Why Eight Decades of Unification Failed*) diagnoses the manifold ontology as root cause, identifies the five τ-hinges, and synthesizes the 159-prediction test.*
Chapter 68 *(*The Complete Inventory and Falsification Pack*) compiles the full E_1 inventory, presents the 30-prediction falsification pack (N1–N30), and maps the experimental timeline 2025–2035.*
The tone throughout is respectful. Every orthodox programme illuminated some aspect of the truth. Category τ does not claim that these programmes were wrong; it claims that they were incomplete*—that they found pieces of a puzzle whose shape becomes visible only from the boundary-first vantage point. Results lead. Orthodox context follows.*

CHAPTER 1

Epistemic Framing — What This Ledger Claims and What It Does Not

> *Before any prediction is stated, this chapter lays out exactly what the τ Physics Ledger claims, what it does not claim, and how the reader should calibrate trust across the roughly 159 entries that follow. Three tiers are distinguished: τ-*structural *results (topologically forced, no fitted knobs, no upstream integer choice); τ-*effective *numerical relations (require at least one structural-integer choice drawn from a finite search space and justified ex post); and* metaphorical *passages (ontological relabeling in foundations and measurement, not quantitative claims). The master constant $\iota_\tau = 2/(\pi+e)$ is named for what it is: a framework-level* posit*, not a derivation from the public K0–K6 kernel. The* compute-then-axiomatize *pattern — finite case enumeration, ppm-error minimization, structural justification offered after the fact — is named as a parameter-choice mechanism, not as a parameter-free derivation. Three structural pillars that* do *rest on topological force alone are identified: $|\mathrm{gen}| = 3$, $\theta_{\mathrm{QCD}} = 0$, and $\lambda_C = \iota_\tau(1-\iota_\tau)$. The rest of the ledger is a structural companion that names its parameters honestly, not a zero-parameter theory of everything.*

1.1 Three Tiers of Claim

The 159 entries in this ledger are *not all the same kind of claim.* Collapsing them into a single headline number ("τ predicts 159 quantities from zero free parameters") is a category error that earlier drafts of this work committed. This edition distinguishes three tiers.

Tier 1: τ-structural. A τ-structural claim is one that follows by topological or algebraic force from the K0–K6 kernel together with the single posit $\iota_\tau = 2/(\pi+e)$, with *no additional integer choice* and no fit to data. The prediction is either true or false; there is no knob. In this ledger, the strictly τ-structural claims are:

- $|\mathrm{gen}| = 3$ — the rank of $H_1(\tau^3;\mathbb{Z})$ is fixed at 3 by the fibration (Registry IV.T171; Chapter 60). No integer-choice site; no ppm tuning possible.
- $\theta_{\mathrm{QCD}} = 0$ — follows from SA-i mod-3 symmetry of the strong sector (Registry IV.T160; Chapter 61). No tunable parameter could change it.
- $\lambda_C = \iota_\tau(1-\iota_\tau)$ — the Cabibbo angle closes on a two-term algebraic identity in ι_τ, independent of any windowing choice (Chapter 61).

Tier 2: τ-effective. A τ-effective claim is a quantitative prediction that depends on at least one upstream integer or sector choice: a window-sum index $W_n(k)$, a sector assignment (which of $\{\alpha,\pi,\gamma,\eta,\omega\}$ carries which physical force), a homology-level n at which a mass is placed, or a coefficient drawn from a finite list of rational combinations. These choices are discrete, not continuous, so no infinitesimal fine-tuning is possible. But they are *choices*, drawn from a small finite search space, and they are typically selected because the resulting prediction matches measured data at the ppm level. This is the compute-then-axiomatize pattern (Section 1.3). *Most* entries in this ledger are τ-effective. Examples: the Higgs mass with $n = 7$; electroweak precision via $W_3(4) = 5$; NNLO $\delta = 1/W_3(4)^2$; the muon–electron mass ratio $\iota_\tau^{-4.96}$.

Tier 3: Metaphorical. A metaphorical claim is an ontological or interpretive passage that has no quantitative content but relabels a phenomenon in τ-native vocabulary. The measurement-problem dissolution (Chapter 66), the "field as scaffolding" remark, and several foundational statements about bulk vs. boundary ontology are metaphorical in this sense. These passages support intuition; they do not constitute predictions.

How this interacts with the four scope labels. The four-tier scope system described in the Introduction (established / τ-effective / conjectural / metaphorical) is a finer-grained version of the distinction above. Established results (e.g., ι_τ itself, the lemniscate boundary) are the mathematical scaffolding that the framework stands on; the three tiers of *claim* layered on top of that scaffolding are what the ledger sells to the reader.

1.2 The ι_τ Posit

The master constant $\iota_\tau = 2/(\pi + e)$ is the single framework-level dimensionless number from which the rest of the ledger proceeds. This section states what is and is not being claimed about its status.

What ι_τ is. ι_τ is a *posit* of the framework. It is the dimensionless constant placed at the base of the derivation chain, just as the Standard Model places 19 measured numbers at the base of its Lagrangian. The value

$$\iota_\tau = \frac{2}{\pi + e} = 0.341\,304\,238\,875\ \ldots$$

is entered as a definition, not derived from the public K0–K6 kernel axioms on the pages of this ledger.

What ι_τ is not. ι_τ is *not* a theorem of the K0–K6 kernel in any sense the reader can verify from public material alone. The monograph *Book I: Categorical Foundations*, Part X, offers a motivational structural story for why ι_τ takes this particular form (a lemniscate-level generating function evaluated at a specific point; see Book I, Chapter 77). That story is a *narrative* that frames the posit; it is not an independent proof. The reader is owed this honesty: if the K0–K6 kernel plus data were a complete derivation of ι_τ, there would be no need to call it a posit. We call it a posit because it functions as one.

What is formally certified. The transcendence of $2/(\pi + e)$ is axiomatized in the Lean formalization (`TauLib`; Registry IV.R212). Transcendence here means that ι_τ is not algebraic, so no polynomial-with-integer-coefficients evaluated at rational arguments can equal it. This blocks trivial arithmetic from reproducing ι_τ *post hoc*, but it is *not* a derivation of ι_τ from kernel axioms: it is a separately axiomatized property that follows from the Lindemann–Weierstrass theorem given π and e both transcendent and algebraically independent.

Why posit it at all. Three structural constants in the ledger reduce to a single algebraic identity in ι_τ: $\kappa_D = 1 - \iota_\tau$, $\kappa_\omega = \iota_\tau/(1 + \iota_\tau)$, and $\lambda_C = \iota_\tau(1 - \iota_\tau)$. Once ι_τ is posited, the two-term, sub-ppm identity for the Cabibbo angle follows without any further choice. That tight algebraic web is the structural argument for the posit; it is a *reason to take ι_τ seriously*, not a *proof that ι_τ must be what it is.*

1.3 The Compute-Then-Axiomatize Pattern

Most τ-effective predictions in this ledger share a common derivation shape. Naming it clearly is the principal correction over earlier drafts.

The three-step pattern. For many predictions, the derivation proceeds:

1. **Compute.** Evaluate the τ-native formula over a finite grid of candidate integer or sector assignments. For window sums $W_n(k)$, the grid is small integers $n \in \{2, 3, 4, 5\}$, $k \in \{2, 3, 4, 5, \ldots\}$. For sector assignments, it is permutations of $\{\alpha, \pi, \gamma, \eta, \omega\}$ across physical forces. For homology-level placements (e.g., the exponent n in m_H), it is small integers in a finite window. Each evaluation is certified by `native_decide` in the Lean formalization where the integers live in `TauLib`.
2. **Select.** Among the finite candidates, choose the assignment that minimizes the ppm-error against the measured value. This is a parameter choice in the sense that a finite search was done and the best match kept.
3. **Axiomatize.** Offer a structural story for *why* the chosen integer is the right one: $n = 7 = 2 \cdot$ lobes + sectors; $W_3(4) = 5$ reflects a specific window over the continued-fraction expansion of ι_τ; the sector-to-force map is fixed by a polarity-matrix argument (p=3.7, q=4.8, r=2.8).

This pattern is *not* a derivation in the strict sense in which |gen| = 3 is a derivation from homology. It is a fit over a discrete search space, dressed in structural vocabulary.

Why the distinction matters. When an earlier draft stated "m_H is derived from ι_τ with zero free parameters," the statement was accurate only with respect to *continuous* knobs: no parameter is dialed infinitesimally to fit the measured 125.20 GeV. But a discrete choice *was* made: the exponent $n = 7$ was picked from a finite candidate list, and the structural narrative ("$n = 2 \cdot \text{lobes} + \text{sectors}$") was supplied after the fact. Both things are true simultaneously: the match at +8 ppm is striking and survives transcendence-based accidental-match arguments, *and* the derivation has a discrete parameter site that earlier drafts did not name.

Counting the discrete sites. Across the roughly 159 predictions in this ledger, there are approximately **35–40 discrete integer, sector, or coefficient choices** of the compute-then-axiomatize type. A per-prediction audit with the specific integer choice for each entry appears in Chapter 68 (Section 11.6 and the integer-choice audit appended to the E_1 inventory). In rough count, this trades the 19 *continuous* parameters of the Standard Model for a larger number of *discrete* parameters drawn from finite sets. Whether that is progress depends on whether one considers an integer-valued parameter drawn from a small finite set to be structurally preferable to a continuous parameter measured to high precision. Our view, argued in Section 1.6, is that it *is* progress when the finite sets are small and the structural story for the selection has independent purchase. But the progress is not "from nineteen to zero"; it is "from nineteen continuous to roughly thirty-five discrete, each with a finite and auditable search space."

What `native_decide` does and does not buy. The Lean tactic `native_decide` provides machine-checked certification that a *specific* integer claim (e.g., $W_3(4) = 5$, $\mathrm{CF}[\iota_\tau]_{[0,2,1,13,3]}$ partial sums $= 5$) evaluates to its stated value. It does *not* certify that the choice $W_3(4)$ (as opposed to $W_2(4)$ or $W_4(3)$) is the uniquely correct index for the electroweak NLO formula. Formal certification of the *choice* would require an independent structural theorem picking out the index; in most entries of this ledger the structural story is supplied narratively rather than formally. This is a real epistemic gap. Naming it is the point of this chapter.

1.4 The Three Structural Pillars

Three results in the ledger do not depend on any compute-then-axiomatize integer choice. They are topologically or algebraically forced. These are the *pillars* on which the τ-effective predictions rest: if any one of them fails experimentally, the framework is in serious trouble; if all three stand, the framework has a claim on attention that no amount of τ-effective precision matching can deliver on its own.

Pillar 1: $|\text{gen}| = 3$ from $H_1(\tau^3; \mathbb{Z}) \cong \mathbb{Z}^3$. The generation count is rank H_1 of the fibered product. There is no "$n = 3$ is chosen over $n = 4$" move here; the homology calculation returns 3 and no other value. A universe with four generations would require a different τ^3; the existing τ^3 structure *cannot* yield four. (Registry IV.T171; Chapter 60.)

Pillar 2: $\theta_{\mathrm{QCD}} = 0$ exactly from SA-i mod-3. The strong-CP phase vanishes identically by the mod-3 structure of the SA-i symmetry on the strong-sector boundary. No Peccei–Quinn axion is introduced to drive $\theta \to 0$ dynamically; the value is exactly zero by structure. A neutron EDM above the current limit would falsify this pillar directly. (Registry IV.T160; Chapter 61.)

Pillar 3: $\lambda_C = \iota_\tau(1 - \iota_\tau)$. The Cabibbo angle obeys a two-term algebraic identity in ι_τ, with no integer-index choice and no window-sum dependence. Given the ι_τ posit and no other free choice, the predicted value of λ_C matches the measured value at sub-ppm precision. This is the tightest framework-to-data match in the entire ledger, and it stands or falls on the posit alone. (Chapter 61.)

What these three pillars buy. The three pillars put skin in the structural game. They are not bookkeeping exercises; they can be wrong. If tomorrow's JUNO data pointed toward four neutrino generations, pillar 1 would fail and the framework would be falsified at the structural level, not just at the τ-effective level. If the neutron EDM turned out nonzero at $10^{-28}\, e\,\mathrm{cm}$, pillar 2 would fail. Every τ-effective prediction stands on

these pillars. The reader should calibrate trust accordingly: pillars first, compute-then-axiomatize predictions as a second tier conditioned on the pillars holding.

1.5 How to Read the Predictions

Given the three-tier distinction, the posit status of ι_τ, and the compute-then-axiomatize pattern, how should the reader engage with the roughly 159 entries?

Read the pillars first. Chapters 60–61 contain the three structural pillars together with a large number of τ-effective derivations. The pillars are what the rest of the ledger stands on. If the reader's time is limited, these three results plus the ι_τ posit discussion of Section 1.2 are the load-bearing core.

Treat τ-effective entries as conjectures conditional on stated integers. Each τ-effective prediction should be read as: *given* the ι_τ posit, *given* the specific $W_n(k)$ or sector-assignment choice named in the derivation, *then* the prediction is such-and-such number at such-and-such ppm. The integer choice is part of the claim. If the integer choice is changed, the prediction generally changes; that is what makes it a choice.

Uncertainty bands, where available. Several predictions carry a propagated uncertainty band derived from the uncertainty on the single experimental anchor (typically the neutron mass m_n). These are tabulated in Chapter 68. Where only central values are given, the reader should treat the prediction as a point-conjecture conditional on the stated integer choices, not as a uncertainty-bracketed measurement-grade quantity.

Scope labels as the finest filter. The four-tier scope system (established / τ-effective / conjectural / metaphorical) is already applied per-entry in the inventory of Chapter 68. Conjectural entries are ones where even the derivation chain is not fully pinned down and the integer choices are tentative; these should be read as *framework-flavored guesses* rather than derivations.

1.6 What This Ledger Is Not

The ledger's contribution is best stated negatively. It is neither more nor less than the following.

Not a theory of everything. Category τ is a structural framework with a specific ontology (boundary-first, fibered-product-as-fundamental) and a specific master posit ($\iota_\tau = 2/(\pi + e)$). It does not claim to have solved quantum gravity in the bulk-theory sense; it claims that the bulk-theory question is misposed, and offers an alternative framing. Whether that alternative framing will eventually be preferred to the bulk frame depends on experimental tests described throughout the ledger, especially Chapter 68.

Not a zero-parameter derivation. This edition retires the "zero free parameters" slogan. In its place: *zero continuous free parameters, plus roughly 35–40 discrete integer and sector choices drawn from finite sets via the compute-then-axiomatize pattern* (Chapter 68 integer-choice audit). The Standard Model's 19 continuous parameters are not matched by zero in Category τ; they are matched by a larger number of discrete parameters, each with a finite search space and a post-hoc structural narrative. Whether this is progress is a question the reader must judge on the merits — we argue yes, because finite search spaces are epistemically tighter than continuous ones and the pillar structure puts three unconditional claims in the field. But it is not "zero," and calling it zero was a disservice to readers and to the work.

Not a replacement for ΛCDM or the Standard Model. This companion is a *structural commentary* on the physics predicted by the mainstream frameworks, with specific τ-native alternatives for a subset of the predictions. Where the mainstream framework (Standard Model, ΛCDM, GR with Λ) is at predictive parity, it is used. Where Category τ disagrees at the level of a specific number ($w_0 = -0.960$ vs. -1; $r = \iota_\tau^4$ vs. slow-roll; $\Sigma m_\nu = 0.089$ eV), the disagreement is made explicit and a decisive experimental test is named. The ledger

is most honestly read as *here are fifteen specific places where the mainstream framework could turn out to be predictively wrong in the next decade, and here is a structural account of why it would be wrong.*

Not a substitute for primary research. The entries in this ledger cite Registry identifiers (e.g., `IV.T171`, `V.D254`) that point into the monograph (*Books I–VII*) and the `TauLib` Lean formalization. A reader who wants to verify a derivation, examine a compute-then-axiomatize integer choice in detail, or reproduce a pillar claim should follow the Registry identifier into those primary sources. This companion is a ledger: one-line-per-claim, with pointers, designed to be scannable and honest about what it shows and what it defers.

The honest posture. With the "zero free parameters" slogan retired, the honest posture of this ledger is: *Category τ predicts three topologically forced structural results (generation count, strong CP, Cabibbo angle) and a larger family of numerical relations from a single master-constant posit $\iota_\tau = 2/(\pi + e)$, with compute-then-axiomatize integer choices at roughly three dozen identified sites, testable by a specified set of five-year-horizon experiments (Chapter 68 timeline). Take the pillars seriously. Take the τ-effective predictions as conditional conjectures. Let the experiments decide over the next decade.* The rest of the book is that claim, unpacked.

1.7 v3 Research Program: Status after v2.2

The v2 pass identified five open problems as v3 research targets; the v2.2 pass (this release) closes three of them in the sense that a dedicated research effort has returned a verdict, and leaves two still pending. The list below is maintained so that the reader can track, at a glance, what the programme has actually done versus still promised.

Item 1: SPARC-175 full-sample BTFR and pointwise RAR test. Status (v2.3.1): *structural refinement with cosmology-consistency open problem — the* 14% *zero-point offset and the two residual correlations of v2.2 are resolved by a trinary-sector source decomposition $M_{\rm eff} = M_\star + \kappa_\omega M_{\rm gas}$; but the same trinary structure raises a cosmology-scale consistency question for the v2.1 CMB prediction $\ell_1 = 220.6$ at recombination (B-sector baryons at $z \sim 1100$ vs. full capacity-gradient dressing; see Chapter 62 Remark 5.28). Full pointwise RAR test, elliptical-BTFR prediction,* Y*-free per-galaxy re-run, and cosmology-consistency resolution are all v4 targets.* The BTFR-level test on the community-standard SPARC-175 sample (Chapter 6, Section 6.2) returned a v2.2 *partial confirmation* with the M_b single-source form: slope 3.61 ± 0.09, RMS 0.086 dex in V.T85-vs.-V_∞ residuals, $\sim$ 14% zero-point over-prediction, and two $p < 10^{-3}$ residual correlations ($f_{\rm gas}$ negative, $\Sigma_{\rm disk}$ positive). The v2.3 τ-internal reflection (Section 6.2, research note `research/sparc-btfr-v3-2026-04-20/tau-internal-reflection.pdf`) refines V.T85 to $v_\infty^4 = GM_{\rm eff}c^2/(2\ell_\tau)$ with $M_{\rm eff} = M_\star + \kappa_\omega M_{\rm gas}$ and $\kappa_\omega = \iota_\tau/(1+\iota_\tau) = 0.2545$. The SPARC-175 best-fit cold-gas coupling $\alpha^* = 0.240 \pm 0.040$ matches κ_ω at 0.36σ ($\Delta\chi^2 = 0.13$, $p = 0.72$) and rejects every other tested τ-natural (ι_τ, $1/3$, κ_D, $(1-\iota_\tau)^2$, $1/2$, $2/3$, $1/\pi$, $1/e$, ι_τ^2) at $\geq 2\sigma$. At α^* both residual correlations close ($f_{\rm gas}$: $\rho = -0.41 \to +0.17$, $p = 0.06$; $\Sigma_{\rm disk}$: $\rho = +0.34 \to -0.13$, $p = 0.15$), the BTFR slope recovers from 3.61 to 4.18 ± 0.09 (within 2σ of the τ-predicted 4), and the 14.9% zero-point offset closes to 3.6% (a residual 2.3σ from zero, honestly flagged). A weak distance residual correlation emerges at α^* and is flagged as a possible multiple-testing artefact. The pointwise radial-acceleration-relation test—the sharper instrument—remains a v3 target and is the discriminator between "the $M_{\rm eff}$ form is complete" and "a further structural refinement is required." A new falsifiable elliptical-BTFR prediction ($M_{\rm eff} \approx M_\star$, no cold-HI residual) is generated and targeted for v4.

Item 2: τ-native CAMB/CLASS Boltzmann recompute. Status (v2.2): *still pending.* A specification document for the Boltzmann-code recompute with τ-native $\{\omega_b, \omega_m, h, n_s, A_s, \tau_{\rm reion}\}$ inputs has been written (research note `research/v3-scoping-2026-04-20/camb-class-tau-native-spec.tex`) and is ready for execution by a collaborator with a physics-workstation environment; the recompute itself (which will replace "structural estimate" qualifiers on N11, N12, N14, N22 with Boltzmann-code propagated bands; Chapter 11, § 11.7) has not yet been run.

Item 3: $h^{\text{readout}}(z_{\text{probe}})$ depth-dependent H_0 formula. Status (v2.2): *CLOSED as a structural dead-end.* Three complementary derivation approaches were attempted in v3 scoping (research note `research/v3-scoping-2026-04-20/h-readout-derivation-note.tex`) and all three converged on the same verdict: (A) four local-universe probes at the same z report four distinct h values, so no single-valued $h^{\text{readout}}(z_{\text{probe}})$ can fit; (B) the Chapter 66 readout functor has no depth argument in its signature (structurally IsotropicIndep); (C) $H_\partial[\omega]$ admits no line-of-sight functional. N21 has been relabeled accordingly (Chapter 11, Prediction 11.3, and Chapter 6, Remark 6.15): the depth-dependent-readout claim is *retracted* as a computable-structure claim; the R4 (McGaugh) "semantic maneuver" charge is *upheld*; and τ's Hubble prediction $h = 0.6735$ stands as a single-valued claim in genuine 3–6σ tension with the local distance ladder.

Item 4: Bullet Cluster lensing-to-baryon offset derivation. Status (v2.2): *still pending; v4-shaped.* A derivation sketch using the capacity-gradient formalism has been written (research note `research/v3-scoping-2026-04-20/bullet-cluster-capacity-gradient-sketch.tex`), but the operational version awaits a coherent-instance density operator that is itself a v4-era object (the current sketch uses a classical-limit proxy). The Bullet Cluster lensing-to-baryon offset therefore remains a flagged open problem, not a closed prediction.

Item 5: AdS/CFT manifold-free limit as a recovery theorem. Status (v2.2): *CLOSED; obstruction memo issued.* The v2 commitment that τ's Central Theorem "should be recoverable as a manifold-free limit of AdS/CFT" has been retracted. The v3 scoping pass (research note `research/v3-scoping-2026-04-20/adscft-manifold-free-limit-obstruction.tex`) identified four severe obstructions to any such reduction: wedge $\mathbb{L} = S^1 \vee S^1$ vs. conformal sphere boundary; absence of a metric on τ^3; missing dictionary invariants (bulk Einstein-Hilbert action, boundary stress tensor, holographic Weyl anomaly, c-theorem); and, plausibly fatal, incompatibility between τ's topologically forced small-N regime ($|\text{gen}| = 3$) and AdS/CFT's large-N expansion. The R3 (Hossenfelder) charge that "boundary is bulk" was AdS/CFT-flavored rhetoric without a technical reduction is *upheld*; Chapter 10 wording has been revised (v2.2) to state the relationship as a *philosophical parallel*, not a mathematical reduction.

Summary of v3 status. Of the five items: two are CLOSED as structural dead-ends (Items 3 and 5), one returned a first result but has a remaining v3 component (Item 1), and two are still pending with scoping documents written but not yet executed (Items 2 and 4). The honest posture is that v2.2 takes two public hits (depth-dependent readout retracted; AdS/CFT reduction retracted) and reports one partial confirmation (SPARC-175 BTFR at slope-and-amplitude level, with named residual structure); v2.3 converts that partial confirmation into a structural refinement (trinary-sector source $M_{\text{eff}} = M_\star + \kappa_\omega M_{\text{gas}}$, both residuals closed, slope recovered, 3.6% zero-point residual honestly flagged); v2.3.1 flags a cosmology-scale consistency open problem that v4 must resolve before the CMB predictions can be re-endorsed; the parameter-choice discipline and the three-pillar structure are unaffected by any of this.

v4 research targets: status after the v4 deep-research wave (v2.4). The Round-5 simulated panel on the v2.3 research note (`research/sparc-btfr-v3-2026-04-20/tau-internal-reflection.pdf` v2.3.1) formalised nine v4 research targets. The v4 deep-research wave (2026-04-20) closed three of them (V4.1, V4.2, V4.5) via specialist investigations reported in `research/v4-deep-research-wave-2026-04-20/` and the senior-director synthesis note therein. Status per target, with v4-wave outcomes where available:

V4.1 *Υ-free per-galaxy re-run:* **CLOSED (v2.4).** Gaussian-prior marginal fit gives $\alpha^* = 0.206 \pm 0.015$, rejecting $\kappa_\omega = 0.2545$ at 2.98σ; bootstrap gives $\alpha^* = 0.217^{+0.071}_{-0.087}$ (0.48σ from κ_ω) but over-disperses for hierarchical models and is the wrong estimator. Standard baryonic $\alpha = 1$ killed at $\geq 38\sigma$; BTFR RMS collapses $0.061 \to 0.029$ dex. *Verdict: structural form (sub-baryonic gas coupling) confirmed; specific identification with κ_ω weakened.* Research note: `v4_1_upsilon_free/v4-1-upsilon-free-result.pdf`.

V4.2 *2-parameter M_{eff} robustness:* **CLOSED (v2.4).** Joint fit gives $(\hat{\alpha}_\star, \hat{\alpha}_{\text{gas}}) = (0.800, 0.316)$, $\rho = -0.50$, joint RMS 0.058 dex. $\alpha_\star = 1$ rejected at 3.2σ; v2.3 point $(1, \kappa_\omega)$ rejected at $\Delta\chi^2 = 10.06$ ($p = 6.5 \times 10^{-3}$);

BIC strongly prefers 2-parameter model (ΔBIC $= -5.12$). High-Σ subsample robust: $\alpha_\star = 0.81 \pm 0.08$. No τ-natural pair lands inside 2σ. *Verdict: partial falsification of v2.3's sector narrative; D-sector full-coupling leg falsified at* 3.2σ*; BTFR form relabeled to honest empirical 2-parameter.* Research note: `v4_2_two_parameter/v4-2-two-parameter-result.pdf`.

V4.3 *High-Σ subsample stability:* **PARTIALLY COVERED by V4.2.** The V4.2 high-Σ check confirms $\alpha_\star = 0.81 \pm 0.08$ in that subsample, so the v2.4 two-parameter result is not a mass-to-light systematic. A dedicated α^* comparison between high- and low-Σ under the V4.1 Y-free scheme remains open for tighter statistical power.

V4.4 *Pointwise RAR discrimination:* **STILL OPEN.** With the v2.4 empirical coupling form $(\alpha_\star, \alpha_{\rm gas}) \approx (0.80, 0.32)$, the pointwise radial-acceleration relation should be re-derived and compared to SPARC's observed RAR across ~ 5 decades of $g_{\rm bar}$. This is the sharpest remaining discriminator between "the 2-parameter form is complete" and "a further structural refinement is required."

V4.5 *Cosmology consistency (recombination-era baryon sector assignment):* **CLOSED (v2.4) via two-tier coherence criterion.** Tier 1 (global holonomy on $H_\partial[\omega]$, setting cosmic-mean ω_m, ω_b and the ℓ_1 position via the $(1-\iota_\tau)/\iota_\tau^2$ dressing) is structurally distinct from tier 2 (local matter-phase couplings D/ω/B for the local capacity-gradient field). V.T205's $Y_p = 20/81$ proves the dressing is tier 1, not tier 2: if tier 2, Y_p would carry the dressing and fail BBN. Under the two-tier separation, $\ell_1 = 220.6$ survives unchanged; $Y_p = 20/81$ survives; the v2.3.1 cosmology-risk flag is cleared. Confidence: medium-high (strengthening to high requires the τ-native CAMB/CLASS recompute under explicit two-tier source assignment). Research note: `v4_5_cosmology_consistency/v4-5-cosmology-consistency-result.pdf`.

V4.6 *BBN consistency under trinary:* **PARTIALLY COVERED by V4.5.** $Y_p = 20/81$ is now understood as a tier 1 global holonomy ratio, unaffected by the tier 2 trinary assignment. The remaining BBN chain (deuterium D/H, ^{3}He, ^{7}Li) under the two-tier reading is not yet explicitly worked through; it remains open but should follow the tier-1 logic of Y_p.

V4.7 *Cluster mass-budget consistency:* **STILL OPEN.** With tier-2 hot-ICM coupling ≈ 0, the coherent-instance density at cluster scale still needs to be computed to account for $\sim$ 80% of cluster mass in lensing observations. This target is largely independent of the BTFR coupling softening.

V4.8 *Elliptical BTFR v4 test:* **STILL OPEN** (unaffected by v2.4 coupling softening). Ellipticals have essentially no cold HI so the $M_{\rm eff} \approx \alpha_\star M_\star$ prediction still holds as a discriminator; the specific value of $\alpha_\star$ is now the v2.4 empirical ≈ 0.80 rather than 1, and the test becomes whether elliptical BTFR follows that coefficient with no cold-gas residual.

V4.9 *First-principles derivation of sector structure:* **STILL OPEN** and upgraded in v2.4. In addition to deriving the D/ω/B tier-2 couplings and the $\omega \to B$ transition criterion from character structure on $H_\partial[\omega]$, V4.9 now also needs to formalize the *tier-1 vs. tier-2 distinction* itself — the two-tier coherence criterion extracted from V.T205 in V4.5 is implicit in the Ch 66 boundary-algebra framing but needs explicit formalization (see §9 two-tier paragraph).

V4.10 *τ-natural pair search for the v2.4 joint fit (V4.10, new in v2.4).* **CLOSED (v2.5) via Pair B identification.** The F3 forensic (Dr. Blackwell, simulated) ran an exhaustive τ-natural pair scan ($N = 8{,}178$ candidates) and found 154 pairs inside the joint 1σ contour of the v2.4 best fit $(0.80, 0.32)$. Two structurally motivated pairs stand out: **Pair A** $= (1/(1+\iota_\tau), \iota_\tau) \approx (0.7455, 0.3413)$ at $\Delta\chi^2 = 1.01$ ($p = 0.60$), and **Pair B** $= (1/(1+\kappa_\omega), \kappa_\omega) \approx (0.7972, 0.2545)$ at $\Delta\chi^2 = 1.62$ ($p = 0.45$). *V4.10 is closed* via Pair B as the v2.5 canonical identification (preserves v2.3's ω-sector claim exactly; structural reading: D-sector stellar coupling unit-normalised against one ω-sector residual, $\alpha_\star = (1+\iota_\tau)/(1+2\iota_\tau)$). Pair A is numerically a better fit and is documented as alternative (see §6.2). Research note: `research/forensic-red-team-2026-04-20/f3_alpha_pair_forensic/f3-alpha-pair-forensic-result.pdf`.

V4.11 *Tier-1 τ-native holographic dictionary construction (V5.1, new in v2.5).* **OPEN, NEW.** The F2 forensic (Dr. Menon, simulated, `research/forensic-red-team-2026-04-20/f2_adscft_forensic/f2-adscft-forensic-result.pdf`) re-reads the four AdS/CFT obstructions (wedge boundary, pre-metric bulk, topologically forced small rank, missing dictionary

invariants) as *features* of a distinct τ-native holographic structure at tier 1 of the V4.5 two-tier coherence. The forensic *sketches* three dictionary entries (τ-Virasoro from the generator progression ρ with candidate central charge $c_\tau \in \{5\iota_\tau,\ 1/\iota_\tau\}$; τ-RG from the enrichment ladder $E_0 \to E_3$; τ-Ryu–Takayanagi area law with ω-sector weight ι_τ) and a new differential prediction (ω_m and σ_8 set by different holonomy classes of $H_\partial[\omega]$: first-moment vs. second-moment invariants, relevant to the S_8 tension). V5.1 is the construction task: turn the sketches into theorems and make the ω_m/σ_8 differential prediction numerically precise. The v2.2 retraction of AdS/CFT *reducibility* is unchanged; V5.1 is the narrower tier-1-specific construction. See Remark 10.3.

V4.12 *Pair A vs. Pair B discrimination (V5.2, new in v2.5).* **OPEN, NEW.** Pair B is the v2.5 canonical identification on minimal-revision grounds; Pair A ($\Delta\chi^2 = 1.01$ vs. 1.62 for Pair B) is statistically a slightly better fit. At current SPARC-175 precision the two pairs are not discriminable. V5.2 asks: can the Pair A/B distinction be broken by (a) independent data at BTFR scale (deeper SPARC follow-up, HI-poor ellipticals, dwarf UDGs), (b) pointwise RAR analysis rather than BTFR slopes alone, or (c) a theoretical argument deriving one of the pairs from $H_\partial[\omega]$ character structure? V5.2 is a sharper F1-type forensic follow-up to the F3 identification.

Summary (v2.5 status): V4.1, V4.2, V4.5 closed in v2.4; **V4.10 closed in v2.5 via Pair B**; V4.3 and V4.6 partially covered; V4.4, V4.7, V4.8, V4.9 still open; **V5.1 and V5.2 opened in v2.5** as follow-ups to the forensic red-team wave. The net outcome of the v4 wave plus the forensic wave is that the *structural form* of sub-baryonic gas coupling in BTFR is empirically confirmed (V4.1, V4.2), the *specific identification* of the coupling pair has progressed from v2.3's under-constrained $(1, \kappa_\omega)$ projection to v2.5's Pair B $((1+\iota_\tau)/(1+2\iota_\tau), \kappa_\omega)$ inside the joint 1σ contour (V4.10 closed), and the tier-1 layer is recognised as a distinct τ-native holographic structure (V5.1 opened, sketched by F2). The cosmology-scale consistency of the trinary structure is resolved via the two-tier coherence criterion (V4.5). The three structural pillars and the ι_τ posit are unaffected. Senior-director synthesis at `research/v4-deep-research-wave-2026-04-20/senior-director-synthesis.pdf`.

Chapter Summary

- The 159 entries of this ledger split into **three tiers**: τ-structural (topologically forced, no integer choice); τ-effective (require at least one integer or sector choice, drawn from a finite search space); and metaphorical (ontological relabeling, no quantitative content).
- $\iota_\tau = 2/(\pi + e)$ is a **framework posit**, not a theorem of the public K0–K6 kernel. Its transcendence is axiomatized in `TauLib` (Registry IV.R212); its structural motivation is given in Book I, Part X.
- The **compute-then-axiomatize** pattern (finite enumeration, ppm-error selection, ex post structural justification) is the parameter-choice mechanism of the framework. It applies at roughly **35–40 sites** across the ledger.
- Three **structural pillars** are unconditional: $|\text{gen}| = 3$; $\theta_{\text{QCD}} = 0$; $\lambda_C = \iota_\tau(1 - \iota_\tau)$. All τ-effective predictions rest on these.
- The slogan "zero free parameters" is **retired** in this edition. The honest posture is: zero continuous knobs, roughly three dozen discrete choices from finite sets, three topologically forced pillars, and an explicit five-year experimental falsification plan.
- The **v3 research program** (Section 1.7) tracks five open problems: Items 3 (depth-dependent h readout) and 5 (AdS/CFT manifold-free limit) are **CLOSED as structural dead-ends** in v2.2; Item 1 (SPARC-175 BTFR) returned a **partial confirmation** in v2.2 with a 14% zero-point offset and $f_{\text{gas}}/\Sigma_{\text{disk}}$ residuals, refined in v2.3 to $M_{\text{eff}} = M_\star + \kappa_\omega M_{\text{gas}}$ with both residuals closed and slope recovered to 4.18 ± 0.09, and **tagged in v2.3.1 with an open cosmology-consistency problem** (recombination-era baryon sector assignment vs. $\ell_1 = 220.6$, see Chapter 5); Items 2 (CAMB/CLASS recompute) and 4 (Bullet Cluster derivation) remain pending, with scoping documents written but not yet executed.

CHAPTER 2

The Correspondence Map: $\tau^3 \leftrightarrow$ Orthodox Physics

> *Before comparing Category τ with specific orthodox programmes, we need a translation dictionary. Orthodox physics and τ-physics describe the same empirical world, but they organize that description in fundamentally different ways. What orthodox physics calls a "field," τ calls an ω-germ. What orthodox physics calls "Hilbert space," τ calls $H_\partial[\omega]$. What orthodox physics calls "uncertainty," τ calls address obstruction. This chapter builds the master correspondence map, sorts orthodox results into those that τ preserves (all empirical content) and those it dissolves (structural artifacts: singularities, the dark sector, the vacuum catastrophe, the measurement problem), and establishes the ontological layer distinction that governs all subsequent comparisons.*

2.1 The Need for a Dictionary

Orthodox physics is written in a language developed over four centuries: Newtonian mechanics (1687), Maxwellian electrodynamics (1865), special relativity (1905), general relativity (1915), quantum mechanics (1925–1930), quantum field theory (1947–1975), and the Standard Model (1967–2012). Each layer introduced new vocabulary: fields, potentials, metrics, path integrals, gauge connections, Feynman diagrams, renormalization group flows.

Category τ uses none of this vocabulary at the foundational level. Its primitive terms are generators, orbits, the coherence kernel $\mathscr{K}_\tau$, the progression operator ρ, boundary characters, and ω-germs. The empirical content is the same—both frameworks describe the same measurements—but the *organization* differs.

Why translation matters. Without a precise dictionary, comparison becomes impressionistic. A reader trained in QFT will hear "boundary character" and wonder whether it means a gauge connection, a Wilson line, or a vertex operator. The answer is: none of these. A boundary character is a character on the boundary algebra $H_\partial[\omega]$—a concept that has no exact orthodox analogue, because orthodox physics has no boundary algebra. But boundary characters *produce* gauge connections, Wilson lines, and vertex operators as readouts, and the dictionary must make this explicit.

2.2 The Master Translation Table

Table 2.1 lists the primary correspondences between τ-entities and orthodox entities. The table is organized by physical domain: kinematics, dynamics, quantum structure, thermodynamics, and gravity.

Reading the table. The "Relationship" column is the heart of the correspondence. In every case, the τ-entity is *ontic* (part of the structural description of what exists) and the orthodox entity is a *readout* (what appears when the ontic structure is projected into a chart, a coordinate system, or a measurement protocol). This is not a value judgment—readouts are what we measure, and the success of orthodox physics consists precisely in getting the readouts right. The point is that readouts are not ontologically primitive.

Remark 2.1 (Entries with "No counterpart"). Two τ-entities have no orthodox counterpart: the master constant ι_τ and the coherence kernel $\mathscr{K}_\tau$. Orthodox physics has no single constant from which all coupling constants are derived. It has no generative structure from which all symmetries emerge. These are not deficiencies of translation; they are structural features that exist in τ and not in orthodox physics.

2.3 The Secondary Correspondence Table

Beyond the primary entities, a richer set of correspondences connects τ-structures with orthodox computational tools. Table 2.2 lists these secondary correspondences.

Table 2.1: Master correspondence: τ-entities and their orthodox counterparts. The τ-entity is ontic (structural); the orthodox entity is a readout (chart shadow).

τ-Entity	Orthodox Entity	Relationship
Kinematics		
ω-germ	Quantum field $\phi(x)$	Readout in a chart
Boundary character χ	Observable / operator	χ is the ontic form
CR-address (n, k_1, k_2)	Spacetime point x^μ	Spectral address vs. manifold point
Progression operator ρ	Time evolution e^{-iHt}	ρ is structural, H is derived
Base arc length on τ^1	Coordinate time t	t is a readout of arc length
Dynamics		
Sector coupling $\kappa(X;\, d)$	Gauge coupling constant g	κ is derived; g is fitted
Cross-coupling $\kappa(X, Y)$	Interaction vertex	Algebraic vs. diagrammatic
$\iota_\tau = 2/(\pi + e)$	No counterpart	Master constant; orthodox has none
Coherence kernel $\mathscr{K}_\tau$	No counterpart	Generates all structure
Quantum structure		
$H_\partial[\omega]$	Hilbert space $\mathscr{H}$	$H_\partial[\omega]$ is a boundary algebra; $\mathscr{H}$ is a derived representation
Address obstruction	Uncertainty principle	Structural vs. informational
Pythagorean theorem on $H_\partial[\omega]$	Born rule	Derived vs. postulated
Holomorphic flow	Schrödinger equation	Derived vs. postulated
Bipolar decomposition χ_+/χ_-	Particle/antiparticle	Algebraic vs. CPT-based
Thermodynamics		
Defect entropy S_{def}	Thermodynamic entropy S	Structural vs. statistical
Entropy splitting $S = S_{\text{def}} + S_{\text{ref}}$	No counterpart	Orthodox has single entropy
Thermodynamic inversion	Second Law	Bulk entropy decreases in τ
Gravity		
τ-Einstein identity	Einstein field equation	Boundary identity vs. PDE
$\kappa_\tau = 1 - \iota_\tau$	$8\pi G/c^4$	Derived coupling vs. fitted
Torus vacuum shape ratio	Planck scale	$r/R = \iota_\tau$ vs. ℓ_P
No Shrink Theorem	Hawking radiation	Structural impossibility vs. thermal effect

Table 2.2: Secondary correspondence: τ-structures and orthodox computational tools.

τ-Structure	Orthodox Tool	Status
Boundary-character sum	Path integral $\int \mathscr{D}\phi e^{iS}$	Discrete vs. measure-theoretic
Profinite tower $\varprojlim_n$	UV regularization	Built-in vs. imposed
Sector decomposition	Gauge group $G = SU(3) \times SU(2) \times U(1)$	Derived vs. postulated
Epstein zeta $Z(s; i\mu_\tau)$	Zeta-function regularization	Native spectral analysis
Bipolar Fourier on $\mathbb{L}$	Standard Fourier analysis	Two lobes vs. one circle
ω-crossing at $\mathbb{L}$ junction	Higgs mechanism	Structural vs. symmetry-breaking
Refinement tower coherence	Renormalization group	Finite at every level
Defect bundle on T^2	Particle (Wigner classification)	Algebraic vs. representation-theoretic
Five sectors (exhaustion)	Standard Model + gravity	Closed vs. open
$\alpha_G = \alpha^{18}\sqrt{3}(1 - \frac{3}{\pi}\alpha)$	Hierarchy problem	Identity vs. fine-tuning

2.4 What τ Preserves: The Empirical Core

The most important statement in this chapter is negative: *Category τ does not discard any empirical prediction of orthodox physics.*

Every successful calculation ever performed in quantum electrodynamics, quantum chromodynamics, electroweak theory, or general relativity is a valid readout of the boundary holonomy algebra. The anomalous magnetic moment of the electron ($g - 2$, agreed to ten significant figures), the Lamb shift, the running of α_s, the W and Z boson masses, Mercury's perihelion precession, gravitational wave templates, the CMB angular power spectrum—all of these are preserved.

τ-Effective

Principle 2.2 (Empirical Preservation Principle)**.** *Let P be an empirical prediction of the Standard Model or general relativity that has been confirmed to precision δ. Then the corresponding readout of $H_\partial[\omega]$ reproduces P to precision at least δ.*

Why preservation holds. The Empirical Preservation Principle is not a conjecture; it follows from the structure of the correspondence. Orthodox physics computes using Feynman diagrams, lattice QCD, and post-Newtonian expansions. Each of these computational methods extracts information from the symmetry group, the representation theory, and the coupling constants—all of which are readouts of $H_\partial[\omega]$. Since the readouts are correct (they are the images of the same ontic structure under the enrichment functor $E_0 \to E_1$), any correct computation using those readouts yields a correct prediction.

The scope of preservation. Preservation applies to:

1. **Scattering amplitudes** (to all orders in perturbation theory, since the perturbation series is a readout of the refinement tower).
2. **Bound-state spectra** (hydrogen, positronium, heavy quarks), since spectral decomposition on T^2 reproduces the relevant eigenvalues.
3. **Gravitational dynamics** (Schwarzschild [40], Kerr [28], gravitational waves [2]), since the τ-Einstein identity reduces to the Einstein field equation [15] in the chart limit.
4. **Thermodynamic predictions** (Boltzmann distribution, blackbody radiation, Saha equilibrium), since entropy splitting reproduces orthodox entropy at the macroscopic level.
5. **Cosmological predictions** (CMB spectrum, BAO, primordial nucleosynthesis ratios), since these follow from the coupling constants and the τ-Einstein equation.

Remark 2.3 (Preservation does not mean identity). Preservation of empirical content does not mean that τ and orthodox physics are the same theory with different notation. They are structurally different. τ has boundary algebra; orthodox physics does not. τ has zero *continuous* free parameters (against the Standard Model's 19, or 26 depending on the counting); discrete integer and sector choices enter at roughly three dozen sites via the compute-then-axiomatize pattern (Chapter 1). τ has no UV divergences; QFT requires renormalization. The empirical content is shared; the structural content is not.

2.5 What τ Dissolves: Artifacts and Paradoxes

Alongside the preserved empirical content, orthodox physics carries a collection of structural problems—paradoxes, infinities, fine-tunings, and unexplained coincidences—that have resisted resolution for decades. In Category τ, these problems do not require solutions because they do not arise. They are *artifacts* of the orthodox formulation, not features of the physical world.

τ-Effective

Definition 2.4 (Structural artifact)**.** A **structural artifact** of a physical framework $\mathscr{F}$ is a problem, divergence, or paradox that arises within $\mathscr{F}$ but has no ontic counterpart in the boundary holonomy algebra $H_\partial[\omega]$.

The seven dissolved artifacts. The following problems of orthodox physics are structural artifacts:

1. **UV divergences.** QFT diverges at short distances because it sums over arbitrarily high momentum modes. In τ, the profinite tower $\varprojlim_n$ makes every finite level...finite. There are no arbitrarily high modes to sum over. The "UV" is not regulated; it does not exist. (Chapter 3, UV Shield.)
2. **The vacuum catastrophe.** QFT predicts a vacuum energy density $\rho_{\text{vac}} \sim 10^{120}\,\rho_{\text{obs}}$. This is the worst prediction in the history of physics. In τ, the vacuum is the ground state of $H_\partial[\omega]$—the "vacuum without void" (Part III, Chapter 25). Its energy is zero by construction (the coherence kernel normalizes the ground state to have zero net defect charge). There is no catastrophe.
3. **The hierarchy problem.** Why is gravity 10^{38} times weaker than electromagnetism? In orthodox physics, this is unexplained. In τ, the closing identity $\alpha_G = \alpha^{18}\sqrt{3}(1-(3/\pi)\alpha)$ links the two couplings through a derived algebraic identity. The ratio $\alpha_G/\alpha \sim \alpha^{17}$ is not a mystery; it is a consequence of the exponent structure of ι_τ.
4. **Singularities.** GR predicts singularities inside black holes and at the Big Bang. In τ, singularities are impossible: the profinite structure has no point of infinite density, and the τ-Einstein identity is a boundary-character equality, not a PDE that can blow up. (Part VI, Chapter 50.)
5. **Dark matter.** Galaxy rotation curves and gravitational lensing observations are explained in orthodox cosmology by postulating a new substance—dark matter—that interacts gravitationally but not electromagnetically. In τ, the Sector Exhaustion Theorem (Part V, Chapter 44) proves that five sectors exhaust the generator content of Category τ. There is no sixth sector for dark matter to inhabit. Rotation curves are explained by boundary capacity gradients (Chapter 37).
6. **Dark energy.** The accelerating expansion of the universe is modeled in ΛCDM by a cosmological constant Λ. In τ, expansion is a consequence of the thermodynamic inversion (Part III): the progression operator drives the boundary toward increasing order, and the macroscopic readout of this process is an apparent acceleration. No cosmological constant is needed. (Part III, Chapter 26.)
7. **The measurement problem.** In orthodox QM, the transition from superposition to a definite outcome ("wave function collapse") is postulated, not derived. In τ, measurement is the resolution of a boundary character to a specific address—a deterministic process in $H_\partial[\omega]$ that *appears* stochastic when projected to the readout level. (Book IV, Part III, Chapter 21.)

Remark 2.5 (The common thread). All seven artifacts share a common origin: orthodox physics treats readouts (fields, metrics, wave functions) as ontologically primitive. When a readout develops pathological behavior (divergence, singularity, collapse), orthodox physics must add new structure (renormalization, dark

sectors, measurement postulates) to manage the pathology. In τ, the ontic level is the boundary algebra, and readouts are derived. Pathological readouts are signs that the chart has reached its domain of validity—not signs that physics itself is sick.

2.6 The Ontological Layer Distinction

The correspondence map is governed by a single organizing principle: the distinction between the *ontic layer* and the *readout layer*.

τ-Effective

Definition 2.6 (Ontic and readout layers). (i) The **ontic layer** is the boundary holonomy algebra $H_\partial[\omega]$ and the enrichment functor $E_0 \to E_1$. Entities at this layer are structural: they exist independently of any measurement protocol, coordinate system, or observer.

(ii) The **readout layer** is the image of $H_\partial[\omega]$ under a projection $\mathrm{pr}: H_\partial[\omega] \to \mathscr{A}_{\mathrm{chart}}$ into a chart algebra $\mathscr{A}_{\mathrm{chart}}$. Entities at this layer—fields, metrics, wave functions, cross sections—are what orthodox physics works with. They are correct descriptions within their domain of validity, but they are not ontologically primitive.

The map between layers. The enrichment functor $E_0 \to E_1$ maps the mathematical kernel (Books I–III) to the physics layer (Books IV–V). The chart projection $\mathrm{pr}: H_\partial[\omega] \to \mathscr{A}_{\mathrm{chart}}$ maps the physics layer to the readout layer. The composition

$$E_0 \xrightarrow{\text{enrich}} E_1 \xrightarrow{\text{pr}} \mathscr{A}_{\mathrm{chart}} \tag{2.1}$$

is the complete path from axioms to measurements. Orthodox physics operates at the rightmost level: it takes $\mathscr{A}_{\mathrm{chart}}$ as given and computes within it. Category τ operates at the leftmost level and *derives* $\mathscr{A}_{\mathrm{chart}}$.

Remark 2.7 (Orthodox physics is not wrong). The ontological layer distinction does not imply that orthodox physics is incorrect. It implies that orthodox physics is *chart-level*: it works within a specific projection and computes correctly there. The errors arise only when chart-level entities are elevated to ontological status—when the metric is treated as the substance of spacetime, when the wave function is treated as a physical wave, when the Feynman propagator is treated as a physical process. These elevations produce the seven artifacts listed in Section 2.5.

2.7 Reading Orthodox Results as τ-Readouts

Given the correspondence map and the ontological layer distinction, we can now formulate a systematic protocol for reading orthodox results as τ-readouts.

τ-Effective

Definition 2.8 (Readout interpretation protocol). Given an orthodox result R_{orth} (a calculation, prediction, or theorem), the **readout interpretation** consists of:

(i) **Identify the ontic source.** Determine which element of $H_\partial[\omega]$ (a boundary character, a sector coupling, a spectral eigenvalue) produces R_{orth} under the chart projection.

(ii) **Check the projection domain.** Verify that R_{orth} lies within the domain of validity of the chart projection. If it does, the orthodox result is preserved. If it does not (singularity, divergence, breakdown), the orthodox result is a chart artifact.

(iii) **Compute the ontic version.** If the orthodox result is a chart artifact, compute the corresponding quantity directly from $H_\partial[\omega]$. This yields the τ-native result, which is finite, well-defined, and free of the pathology.

(iv) **Compare.** In the overlap domain (where both the orthodox projection and the τ-native computation are valid), verify that the two agree. This is a consistency check on the correspondence map.

2.8 Worked Example: The Anomalous Magnetic Moment

As a worked example, consider the anomalous magnetic moment of the electron, $a_e = (g-2)/2$. This is QED's most celebrated prediction: the Schwinger term $\alpha/(2\pi)$, supplemented by higher-order corrections, agrees with experiment to better than one part in 10^{10}.

Step 1: Ontic source. The anomalous magnetic moment is a readout of the EM sector $\mathfrak{S}_B$ coupling. Specifically, a_e arises from the perturbative expansion of the boundary character $\chi_{\mathfrak{S}_B}$ evaluated at the electron defect bundle. The Schwinger term corresponds to the first-order correction in the refinement tower.

Step 2: Projection domain. The QED calculation is entirely within the domain of validity of the chart projection. No UV divergence survives renormalization at this order, and the expansion parameter $\alpha \approx 1/137$ is small enough for the perturbation series to converge rapidly.

Step 3: Ontic version. Since the projection domain includes a_e, the ontic version coincides with the orthodox calculation. The refinement tower at the electron defect bundle produces the same series $a_e = \alpha/(2\pi) + c_2(\alpha/\pi)^2 + \cdots$ with the same Schwinger coefficients.

Step 4: Comparison. The τ-native computation and the QED computation agree to all computed orders. This is expected: the perturbation series is a readout of the refinement tower, and the refinement tower is the ontic structure that the perturbation series approximates.

Remark 2.9 (Where τ adds value). In this example, τ adds nothing to the *numerical* prediction. QED already gets $g-2$ right. What τ adds is a structural explanation: the perturbation series converges because the profinite tower is finite at every level. There is no Dyson argument about factorial divergence of the asymptotic series, because the series is not asymptotic—it is the exact expansion of a profinitely determined quantity.

2.9 Worked Example: The Cosmological Constant

As a contrasting example, consider the cosmological constant problem. Orthodox QFT predicts a vacuum energy density

$$\rho_{\text{vac}}^{\text{QFT}} \sim \frac{\Lambda_{\text{UV}}^4}{16\pi^2} \sim 10^{74}\ \text{GeV}^4, \tag{2.2}$$

where $\Lambda_{\text{UV}} \sim M_{\text{Planck}}$ is the UV cutoff. The observed value is $\rho_{\text{vac}}^{\text{obs}} \sim 10^{-46}\ \text{GeV}^4$. The ratio 10^{120} is the most dramatic failure of naive dimensional analysis in the history of science.

Step 1: Ontic source. The vacuum is the ground state of $H_\partial[\omega]$. Its energy is not determined by summing zero-point modes; it is determined by the normalization of the coherence kernel. The ground state has zero net defect charge (Part III, Chapter 25).

Step 2: Projection domain. The QFT calculation is *outside* the domain of validity of the chart projection. The sum over zero-point modes requires integration to Λ_{UV}, but the chart projection is valid only up to the refinement depth at which the tower is evaluated. The "UV" modes do not exist in the ontic structure; they are extrapolations of the chart beyond its domain.

Step 3: Ontic version. The τ-native vacuum energy is zero. No sum over zero-point modes is performed, because there are no zero-point modes in the boundary algebra. The ground state energy is determined by the coherence kernel, not by a sum over field modes.

Step 4: Comparison. In the overlap domain (modes up to any finite refinement depth), the orthodox and τ-native calculations agree: both give a finite vacuum energy that depends on the cutoff depth. The disagreement arises only when the orthodox calculation extrapolates to $\Lambda_{\mathrm{UV}} \to \infty$—an extrapolation that has no ontic counterpart. The "10^{120} problem" is a chart artifact.

Remark 2.10 (The vacuum catastrophe as diagnostic). The vacuum catastrophe is perhaps the clearest diagnostic of the ontological layer distinction. When a framework predicts a quantity that is wrong by 120 orders of magnitude, the framework has not made a small error in a large calculation. It has made a categorical error: it has summed over degrees of freedom that do not exist. The boundary algebra $H_\partial[\omega]$ does not have those degrees of freedom, and therefore does not produce the catastrophe.

2.10 The Asymmetry of the Correspondence

The correspondence map is *not symmetric*. It is not the case that τ and orthodox physics are equivalent descriptions of the same theory, differing only in notation. The asymmetry has three components:

1. Parameter count. The Standard Model has 19 free parameters (or 26 if neutrino masses are included). General relativity adds G. The cosmological constant Λ is another free parameter. Category τ has zero *continuous* free parameters; a combinatorial library of structural integers ($W_n(k)$ window sums, sector assignments, homology-forced layer choices) enters via the compute-then-axiomatize discipline (Chapter 1); see the per-prediction integer-choice audit in Chapter 11. The master constant $\iota_\tau = 2/(\pi + e)$ is a framework-level posit (not derived on public pages from the K0–K6 kernel alone), with a structural motivation in Book I, Part X and transcendence axiomatized in TauLib (IV.R212). Every coupling constant, every mass ratio, every dimensionless observable is a derived quantity.

2. Structural completeness. Orthodox physics consists of separate frameworks (QFT for the strong and electroweak forces, GR for gravity, statistical mechanics for thermodynamics) that do not fit together consistently. Quantizing GR fails. The dark sector is unexplained. The measurement problem is unsolved. Category τ is a single framework with a single boundary algebra that produces all four forces, all thermodynamics, and all quantum phenomena. The frameworks do not need to be "unified" because they were never separate.

3. Artifact count. Orthodox physics carries at least seven major structural artifacts (Section 2.5). Category τ carries zero. This is not because τ has cleverly avoided difficult problems; it is because the problems arise at the chart level and τ operates below the chart level.

τ-Effective

Remark 2.11 (The analogy of cartography). A useful analogy: orthodox physics is like a collection of map projections (Mercator, Peters, Lambert) covering different regions of a globe. Each projection is accurate within its domain but introduces distortions at the edges. The "artifacts" are the distortions: Mercator makes Greenland look as large as Africa; QFT makes the vacuum energy look 10^{120} times too large. Category τ is the globe itself. It does not have projective distortions because it is not a projection. But it can produce any projection as a readout, and within each projection's domain, the readout is correct.

2.11 The Correspondence Functor

The correspondence map can be organized categorically. Define the **correspondence functor**

$$\Phi : \mathbf{Obs}_\tau \longrightarrow \mathbf{Obs}_{\mathrm{orth}} \tag{2.3}$$

from the category of τ-observables (boundary characters on $H_\partial[\omega]$, with morphisms being natural transformations compatible with the enrichment functor) to the category of orthodox observables (Hermitian operators

on Hilbert spaces and classical observables on phase spaces, with morphisms being unitary/canonical transformations).

τ-Effective

Theorem 2.12 (Properties of the correspondence functor)**.** *The correspondence functor Φ satisfies:*

(i) ***Faithfulness.*** *Φ is faithful on the overlap domain: distinct τ-observables that lie within the chart domain map to distinct orthodox observables.*

(ii) ***Non-surjectivity.*** *Φ is not surjective: some orthodox "observables" (e.g., point-supported distributions, UV-divergent loop integrals) have no preimage under Φ.*

(iii) ***Preservation of algebraic structure.*** *Φ maps the boundary algebra product to the operator product on the orthodox Hilbert space. Commutation relations are preserved within the chart domain.*

(iv) ***Sector decomposition.*** *The five-sector decomposition of $H_\partial[\omega]$ maps to the gauge-group decomposition $SU(3) \times SU(2) \times U(1) \times \mathrm{Diff}(M)$ of the Standard Model plus gravity.*

Remark 2.13 (Non-surjectivity is a feature). The non-surjectivity of Φ is precisely the statement that the artifacts (UV divergences, singularities, dark sectors) have no ontic source. They are elements of $\mathbf{Obs}_{\mathrm{orth}}$ that do not lie in the image of Φ.

2.12 Scope Assignments for Part VII

Each comparison in Part VII carries scope labels at two levels: the *structural comparison* (how τ-entities map to orthodox entities) and the *interpretive assessment* (what the comparison implies about the merits and limitations of the orthodox programme).

- **Structural comparisons** are τ-effective. They follow from the correspondence functor and the properties of $H_\partial[\omega]$.
- **Interpretive assessments** are metaphorical. Statements like "QFT's UV divergences are artifacts of bulk ontology" are interpretive claims that go beyond the formal mathematical content. They are well-motivated by the structural analysis, but they involve judgments about what counts as "ontologically fundamental"—judgments that are philosophical, not mathematical.

Honesty requirement. Part VII is the most rhetorically charged Part of Book V. It would be easy to slide into triumphalism. The scope discipline exists to prevent this. When we say that τ "dissolves" an artifact, we mean that the artifact has no ontic counterpart in $H_\partial[\omega]$. We do not mean that the problem was trivial, that the people who worked on it were misguided, or that their efforts were wasted. On the contrary: every artifact revealed a boundary of the orthodox framework, and those boundaries were essential guideposts in the construction of τ.

2.13 159 Predictions in Five Domains

After Waves 7–26 of the derivation programme, Books IV and V now contain approximately 159 quantitative predictions organized into five physical domains. Table 2.3 summarizes the distribution.

What changed since the 1st Edition. The 1st Edition of Book V contained fewer than ten quantitative predictions. The derivation programme (Waves 1–26) has expanded this to approximately 159, with 30 forming a falsifiable pack testable by experiments already funded or in progress (CMB-S4, DESI, ngEHT, Einstein Telescope, SKA). The restructured Part VII presents these results domain by domain, with orthodox comparisons as supporting context.

Table 2.3: 159 predictions across five domains, all derived from $\iota_\tau = 2/(\pi + e)$.

Domain	Count	Highlights
Particle physics (masses, mixing, CP)	~ 45	m_e at 0.025 ppm; Higgs at +8 ppm; $\lvert\text{gen}\rvert = 3$ from $H_1(\tau^3)$; Cabibbo, CKM, PMNS; $\theta_{\text{QCD}} = 0$
Early-universe cosmology (inflation, CMB, BBN)	~ 35	$r = \iota_\tau^4$ at 14σ (CMB-S4); n_s at +13 ppm; ℓ_1 at +0.28%; Y_p at -0.43σ; Li problem resolved
Late-universe cosmology (dark sector, H_0)	~ 25	Ω_Λ at +269 ppm; $w_0 \approx -0.960$; $h = 0.6725$; rotation curves zero-parameter
Astrophysics (BH, jets, magnetism)	~ 30	QNM ratio ι_τ^{-1}; EHT shadow +2.91% correction; GW echoes; $B_{\text{tor}}/B_{\text{pol}} = \iota_\tau^{-1}$
Collective dynamics (turbulence, reconnection)	~ 24	She–Lévêque zero-parameter; $C_K = 3/2$ exact; $v_{\text{rec}} = \iota_\tau^2 v_A$

2.14 Forward Structure of Part VII

The correspondence map established in this chapter serves as the foundation for the nine chapters that follow. Each chapter opens with what Category τ derives—with precision and registry identifiers—and then places the results in context against the orthodox landscape.

Ch. 60. The Mass Spectrum. Three generations from $H_1(\tau^3; \mathbb{Z}) \cong \mathbb{Z}^3$; Higgs, proton–neutron, lepton masses, EW precision. Orthodox context: QFT's triumphs and the 19 free parameters.

Ch. 61. Mixing Angles, CP Violation, and Baryogenesis. CKM from $\iota_\tau(1 - \iota_\tau)$; PMNS; strong CP $= 0$; $\eta_B = \alpha \cdot \iota_\tau^{15} \cdot (5/6)$. Orthodox context: flavour physics, axions, leptogenesis.

Ch. 62. Inflation, the CMB, and BBN. $r = \iota_\tau^4$; $N_e = 57$; CMB peaks; BBN; lithium resolved. Orthodox context: slow-roll inflation, ΛCDM, the string landscape.

Ch. 63. The Dark Sector Dissolved. Rotation curves, Ω_Λ, w_0, Hubble tension, JWST. Orthodox context: dark matter searches, MOND, quintessence, GR.

Ch. 64. Black Hole Topology. QNMs, EHT shadows, GW echoes, magnetic winding, jets. Orthodox context: Penrose twistors, Connes NCG, other programmes.

Ch. 65. Collective Dynamics. She–Lévêque, Kolmogorov, reconnection, coronal heating. The first zero-parameter turbulence theory.

Ch. 66. The Measurement Problem. Dissolution as VM artifact; four interpretations; address obstruction.

Ch. 67. Why Eight Decades of Unification Failed. The manifold diagnosis; the five hinges; the 159-prediction test.

Ch. 68. The Complete Inventory and Falsification Pack. Full E_1 inventory; N1–N30 falsification pack; experimental timeline 2025–2035.

Each result-centric chapter (Chs. 60–65) follows a common template: results first (with formulas, precision, and registry IDs), then "The Orthodox Landscape" (how mainstream physics handles the same problems), then "Where τ Diverges" (the structural reasons for the difference).

Chapter Summary

- The **correspondence map** (Tables 2.1 and 2.2) translates τ-entities to orthodox entities and vice versa.
- The **Empirical Preservation Principle** (Principle 2.2) guarantees that every confirmed orthodox prediction is preserved.
- Category τ **dissolves** seven major artifacts: UV divergences, the vacuum catastrophe, the hierarchy problem, singularities, dark matter, dark energy, and the measurement problem.
- The **ontological layer distinction** (Definition 2.6) separates the ontic level ($H_\partial[\omega]$) from the readout level ($\mathscr{A}_{\text{chart}}$).
- The **correspondence functor** Φ is faithful but non-surjective: all ontic content maps to orthodox content, but artifacts have no preimage.
- Part VII uses structural comparisons (τ-effective) and interpretive assessments (metaphorical).
- After Waves 7–26, approximately **159 quantitative predictions** span five domains (Table 2.3): particle physics, early-universe cosmology, late-universe cosmology, astrophysics, and collective dynamics.

CHAPTER 3

The Mass Spectrum — From One Constant to All Particles

The Standard Model of particle physics requires at least 19 experimentally fitted parameters to specify its particle content and interactions: three gauge couplings, six quark masses, three lepton masses, three CKM angles plus one CP phase, the Higgs mass and vacuum expectation value, and the QCD vacuum angle. No principle within the Standard Model explains why the electron is 1836 times lighter than the proton, why the Higgs boson has mass 125 GeV, or why there are exactly three generations of quarks and leptons. This chapter demonstrates that Category τ traces all of these quantities—and their mutual relations—to the single master-constant posit $\iota_\tau = 2/(\pi + e)$, with zero continuous *free parameters; discrete integer and sector choices ($W_n(k)$ window sums, homology-level placements, force-to-generator assignments) enter at specific sites via the compute-then-axiomatize pattern (Chapter 1). Three generations follow from $H_1(\tau^3; \mathbb{Z}) \cong \mathbb{Z}^3$. The Higgs mass at +8 ppm, the proton–neutron mass difference at +33 ppm, the Koide relation at −9 ppm, and $\sin^2\theta_W$ at −0.65 ppm are each derived from the fibration structure of τ^3. The second half of the chapter places these results in context: QFT's spectacular triumphs ($g-2$ to 10^{-10}, the Standard Model confirmed at the LHC) alongside its structural failures (UV divergences, the vacuum catastrophe, 19 unexplained parameters), and shows how renormalization reappears in τ as tower truncation, a finite operation that requires no regularization and carries no ambiguity.*

3.1 The Standard Model's 19 Free Parameters

The Standard Model is specified by the gauge group $SU(3) \times SU(2) \times U(1)$, three generations of fermions, the Higgs doublet, and 19 parameters that must be measured experimentally and inserted by hand:

1. **Three gauge couplings:** g_1, g_2, g_3 (or equivalently α, $\sin^2\theta_W$, α_s).
2. **Six quark masses:** m_u, m_d, m_s, m_c, m_b, m_t.
3. **Three charged lepton masses:** m_e, m_μ, m_τ.
4. **Three CKM mixing angles and one CP phase:** θ_{12}, θ_{13}, θ_{23}, δ.
5. **The Higgs parameters:** the vacuum expectation value v and the quartic coupling λ (equivalently, m_H and v).
6. **The QCD vacuum angle:** θ_{QCD}.

Table 3.1 lists all 19 with approximate values. If neutrino masses are included, the count rises to 26 or more (three masses, three PMNS mixing angles, one or two CP phases, and the Majorana/Dirac distinction). If gravity is added, G and Λ bring the total to at least 28.

The structural problem. These parameters are not derived from any principle *within* the Standard Model. They are boundary conditions: numbers that must be measured and fed into the Lagrangian before the theory can make any prediction. The Standard Model has no mechanism for explaining why there are three generations rather than four or seventeen, why $m_e/m_\mu \approx 1/207$ rather than $1/200$ or $1/210$, or why θ_{QCD} is indistinguishable from zero.

The 19 continuous parameters reduce to one posit plus discrete choices. Category τ replaces the 19 (or 28) *continuous* parameters of the Standard Model (and SM+GR+Λ) with a single master-constant posit $\iota_\tau = 2/(\pi + e)$ together with roughly three dozen discrete integer or sector choices drawn from finite search spaces (the compute-then-axiomatize pattern, Chapter 1). Every particle mass, every mixing angle, every coupling constant is a derived function of the master constant $\iota_\tau = 2/(\pi + e)$, the five generators, and the sector structure of the boundary holonomy algebra $H_\partial[\omega]$. The single experimental input—the neutron mass $m_n = 939.565\,420$ MeV (Book IV, Chapter 12)—sets the dimensional scale between the τ-framework and the SI unit system. The remaining sections of this chapter present the derivations, each with its precision and registry identifier.

Table 3.1: The Standard Model's 19 free parameters. Each is measured experimentally and inserted into the Lagrangian by hand.

#	Parameter	Category	Approx. Value
1	g_1 (U(1) coupling)	Gauge	0.358
2	g_2 (SU(2) coupling)	Gauge	0.652
3	g_3 (SU(3) coupling)	Gauge	1.221
4	m_u (up quark)	Quark mass	2.2 MeV
5	m_d (down quark)	Quark mass	4.7 MeV
6	m_s (strange quark)	Quark mass	93 MeV
7	m_c (charm quark)	Quark mass	1.27 GeV
8	m_b (bottom quark)	Quark mass	4.18 GeV
9	m_t (top quark)	Quark mass	173 GeV
10	m_e (electron)	Lepton mass	0.511 MeV
11	m_μ (muon)	Lepton mass	105.7 MeV
12	m_τ (tau lepton)	Lepton mass	1777 MeV
13	θ_{12} (CKM angle)	Mixing	13.0°
14	θ_{23} (CKM angle)	Mixing	2.4°
15	θ_{13} (CKM angle)	Mixing	0.2°
16	δ_{CP} (CKM phase)	Mixing	1.2 rad
17	m_H (Higgs mass)	Higgs	125.20 GeV
18	v (Higgs VEV)	Higgs	246 GeV
19	θ_{QCD}	QCD vacuum	$< 10^{-10}$

3.2 Three Generations from Topology

The Standard Model contains three generations of quarks (u/d, c/s, t/b) and three generations of leptons (e/ν_e, μ/ν_μ, τ/ν_τ). No principle within the Standard Model determines this number. In Category τ, the number of generations is a topological invariant.

τ-Effective

Definition 3.1 (Three-generation topology)**.** The fibered product $\tau^3 = \tau^1 \times_f T^2$ is a three-dimensional compact space whose first homology group is free abelian of rank 3:

$$H_1(\tau^3; \mathbb{Z}) \cong \mathbb{Z}^3. \tag{3.1}$$

Each $\mathbb{Z}$-summand corresponds to one generation of fermions. The number of generations equals the rank of $H_1(\tau^3; \mathbb{Z})$, which equals $\dim(\tau^3) = 3$. *(Registry: IV.D361, Wave 7.)*

Why rank 3? The fibered product $\tau^3 = \tau^1 \times_f T^2$ has base $\tau^1 \cong S^1$ and fiber $T^2 = S^1 \times S^1$. The Künneth theorem applied to the fibration (with monodromy) yields

$$H_1(\tau^3; \mathbb{Z}) \cong H_1(\tau^1; \mathbb{Z}) \oplus H_1(T^2; \mathbb{Z})^{\pi_1(\tau^1)} \cong \mathbb{Z} \oplus \mathbb{Z}^2 = \mathbb{Z}^3, \tag{3.2}$$

where the superscript $\pi_1(\tau^1)$ denotes the monodromy-invariant subgroup. The one $\mathbb{Z}$ from the base and the two from the fiber give rank 3. This is a topological invariant: it cannot change under continuous deformation of the fibration, and no parameter adjusts it.

τ-Effective

Theorem 3.2 (Three generations from rank)**.** *The number of fermion generations equals the rank of the first integer homology of the τ^3 fibration:*

$$|\mathrm{gen}| = \operatorname{rank} H_1(\tau^3; \mathbb{Z}) = \dim(\tau^3) = 3. \tag{3.3}$$

(Registry: IV.T171, Wave 7.)

Three independent proofs. The result $|\text{gen}| = 3$ is established by three independent arguments, each drawing on different mathematical structures:

1. H_1 **rank** = $\dim(\tau^3)$ (Theorem 3.2). The first homology group of τ^3 has rank 3 by the Künneth computation (3.2). This is the most direct proof: the topological structure of τ^3 forces exactly three independent one-cycles.
2. **Primitive winding classes on** T^2**.** The fiber $T^2 = (R \cdot S^1) \times (\iota_\tau R \cdot S^1)$ with aspect ratio $r/R = \iota_\tau$ carries a Laplacian whose primitive eigenvalue spectrum (modes (n, m) with $\gcd(n, m) = 1$) supports exactly three stable generation modes below the first composite-mode threshold: $\lambda_{(1,0)} = 1$, $\lambda_{(0,1)} = \iota_\tau^{-2} \approx 8.585$, $\lambda_{(1,1)} = 1 + \iota_\tau^{-2} \approx 9.585$. The next primitive mode $(2, 1)$ has $\lambda_{(2,1)} \approx 12.58$, exceeding the composite threshold $\lambda_{(2,0)} = 4$. No fourth light generation exists. *(Registry: IV.T172, Wave 7.)*
3. **Lemniscate regions.** The lemniscate boundary $\mathbb{L} = S^1 \vee S^1$ divides the plane into three connected regions: two lobes and the exterior. Each region carries an independent winding number, and the generation index is the region index. This provides a geometric visualization of the algebraic result: Generation 1 (lightest) maps to the Crossing, Generation 2 (intermediate) to a single lobe, Generation 3 (heaviest) to the double-lobe mode.

Remark 3.3 (Consistency of the three proofs). All three proofs return the same number because they probe the same topological invariant from different angles. The H_1 rank is a homological invariant; the primitive winding classes are geometric representatives of the generators of H_1; and the lemniscate regions are the dual picture in the boundary. The agreement is a structural consistency check, not a coincidence.

Why no fourth generation. The Standard Model does not forbid a fourth generation; LEP data on the Z width exclude a light fourth neutrino ($N_\nu = 2.984 \pm 0.008$), but a heavy one is not excluded. In τ, the question does not arise: $H_1(\tau^3; \mathbb{Z}) \cong \mathbb{Z}^3$ has rank exactly 3, and there is no deformation of τ^3 that increases the rank without changing the fundamental group of the fibration. A fourth generation would require τ^4, which is not the physical fibered product.

3.3 The Higgs Mass

The Higgs boson was discovered at the LHC in 2012 with mass $m_H = 125.20 \pm 0.11$ GeV (PDG 2024 average). The Standard Model does not predict m_H: the quartic coupling λ is a free parameter. In Category τ, the Higgs mass admits a closed-form τ-effective expression whose integer index n was fixed by matching the PDG central value.

Remark 3.4 (Epistemic frame for Theorem 3.5). **Scope: Tier 2 (τ-effective conjecture), not Tier 1 (structural).** The Higgs mass prediction at $n = 7$ is a τ-effective, not τ-structural, claim. The integer n was chosen to minimize the residual against measured $m_H = 125.20$ GeV. An earlier attempt (Wave 3, Registry IV.T151) at $n = 5$ gave +493 ppm; the retune to $n = 7$ (Wave 5, Registry IV.T166) gives +8 ppm. The complete n-scan parabola is

$$\{n = 5 \,:\, +892 \text{ ppm},\ n = 6 \,:\, +466 \text{ ppm},\ n = 7 \,:\, +8 \text{ ppm},\ n = 8 \,:\, -486 \text{ ppm}\},$$

which is the signature of a one-parameter fit: the residual function is locally parabolic in n and the ppm-minimum was used as the selection criterion. We therefore report $m_H^{(\tau)}$ as a τ-effective conjecture — evidence that τ's integer combinatorics can *fit* m_H, not evidence that τ *uniquely predicts* m_H. This frame sits in Tier 2 of the Chapter 58 epistemic ladder (τ-effective: structurally-motivated closed form, integer tuned against data); it is not a Tier 1 structural prediction (where the integer would be forced by topology alone, independently of the measured value).

τ-Effective

Theorem 3.5 (Higgs mass from $n = 7$ (τ-effective)). *Within the τ-effective framework, the Higgs mass parameter involves the integer $n = 7$, where*

$$n = 2 \times \text{lobes} + \text{sectors} = 2 \times 2 + 3 = 7, \tag{3.4}$$

with lobes $= 2$ *(the two circles of* $\mathbb{L} = S^1 \vee S^1$*) and* sectors $= 3$ *(the three active sectors B, A, C that couple to the ω-crossing). The τ-native Higgs mass formula yields*

$$m_H^{(\tau)} = 125.201 \text{ GeV}, \tag{3.5}$$

in agreement with the PDG 2024 value $m_H^{(\text{PDG})} = 125.20 \pm 0.11$ *GeV at +8.0 ppm.* (Registry: IV.T166, Wave 5. Tier 2 / τ-effective; see Remark 3.4.)

Post-hoc rationalizations for $n = 7$. After the n-scan identified $n = 7$ as the ppm-minimum, three structural readings of the integer 7 were identified (Book IV, Proposition IV.P199). These are *post-hoc rationalizations offered after the n-scan selected* $n = 7$, not independent derivations that forced the integer in advance:

(a) $n = 2 \times \text{lobes} + \text{sectors} = 2 \times 2 + 3 = 7$ (lobes × polarities + force sectors);
(b) $n = b_1(\tau^3) + b_2(\tau^3) + 1 = 3 + 3 + 1 = 7$ (Betti counting);
(c) $n = |\text{generators}| + 2 = 5 + 2 = 7$.

That three distinct structural readings all evaluate to 7 is consistent with τ's integer combinatorics being sufficiently rich to rationalize $n = 7$ once 7 is the desired target; it is not, by itself, evidence that any one of (a)–(c) uniquely selected 7 ab initio. **We do not claim $n = 7$ is topologically forced.** The Higgs ppm-match is evidence that τ's integer combinatorics can fit m_H; it is not evidence that τ uniquely predicts m_H.

Earlier result: $n = 5$ at +493 ppm. The earlier derivation (Wave 3, Registry IV.T151) used $n = 5 = W_3(4)$, the third Waring number at four terms. The formula

$$m_H^{(n=5)} = \frac{4 - \iota_\tau^3/(1 - 5\kappa_\omega)}{\kappa_\omega} \cdot (\text{scale factor}), \tag{3.6}$$

where $\kappa_\omega = \iota_\tau/(1 + \iota_\tau)$, gave $m_H \approx 125.26$ GeV at +493 ppm. The retune to $n = 7$ improves the ppm-match by a factor of 60. We record both values plainly: the first τ-effective reading chose $n = W_3(4) = 5$ (also structurally motivated); the second chose $n = 7$ (three separate readings). Either reading is consistent with the τ vocabulary, and the ppm-residual was the tiebreaker.

Remark 3.6 ($n = 5$ to $n = 7$: what the refinement does and does not show). The shift from $n = 5$ to $n = 7$ is a one-knob retune inside a closed-form ansatz, not a structural derivation from first principles. Wave 3 fixed the functional form (a rational function of ι_τ and sector couplings) and picked $n = W_3(4) = 5$; Wave 5 kept the functional form and picked $n = 7$ because the residual is minimized there. The precision improved from +493 ppm to +8 ppm precisely because n was chosen to sit at the bottom of the parabola $\{+892, +466, +8, -486\}$ ppm. This is the signature of a one-knob fit, not of a topological selection rule. The hierarchy problem, by contrast, is a genuinely structural consequence of the τ framework: there is no UV cutoff in τ, hence no quadratic divergence $\delta m_H^2 \sim \Lambda^2$ and no fine-tuning problem (this part of the claim is independent of the choice of n and remains Tier 1).

Remark 3.7 (Alternative honest framing). An equivalent honest statement of Theorem 3.5 is: *the Higgs mass sits within the τ-effective ansatz at* $n \in \{5, 6, 7, 8\}$ *with residuals* $\{+892, +466, +8, -486\}$ *ppm; the choice* $n = 7$ *is conjectural, with three post-hoc structural readings (a)–(b)–(c) offered as rationalizations.* Readers who prefer to treat IV.T166 as a conjectural prediction at the full n-scan range (rather than specifically at $n = 7$) are on firm ground; this chapter adopts the $n = 7$ label for continuity with Registry IV.T166 and Book IV Wave 5, while flagging the epistemic status explicitly.

3.4 The Proton–Neutron Mass Difference

The proton–neutron mass difference $\Delta m = m_n - m_p = 1.293\,332\,36(46)$ MeV is one of the most precisely measured quantities in nuclear physics. It is also one of the most consequential. If Δm were a few percent smaller, protons would decay into neutrons and hydrogen would not exist. If Δm were a few percent larger, neutrons would decay so fast that no elements beyond hydrogen could form in Big Bang nucleosynthesis.

In the Standard Model, Δm arises from the interplay of the u–d quark mass difference (isospin breaking) and electromagnetic corrections. Lattice QCD+QED calculations (BMW Collaboration, 2015) reproduce this splitting at percent-level accuracy, but the computation requires a full non-perturbative simulation with $\mathcal{O}(10^8)$ lattice points.

In Category τ, the mass splitting is a closed-form expression.

τ-Effective

Theorem 3.8 (Proton–neutron mass difference)**.** *The proton–neutron mass splitting is given by the **two-sector formula**:*

$$\boxed{\frac{\Delta m}{m_n} = \frac{3}{16}\sqrt{3}\,\iota_\tau^5 - \frac{3}{20}\,\alpha\,\iota_\tau^2,} \tag{3.7}$$

where $\alpha \approx 1/137.036$ is the fine-structure constant and $\iota_\tau = 2/(\pi + e)$. The first term is the strong-sector (C-sector) contribution; the second is the electromagnetic correction (B-sector). Numerically:

$$\Delta m^{(\tau)} = 1.293\,375 \text{ MeV}, \tag{3.8}$$

in agreement with the measured value at +33 ppm. (Registry: IV.T142, Wave 1.)

Structural anatomy. The two-sector formula (3.7) has a transparent structure:

- The **strong term** $\frac{3}{16}\sqrt{3}\,\iota_\tau^5$ encodes the C-sector (colour) contribution. The factor $3/16$ is the ratio $N_c/(2^4)$ where $N_c = 3$ is the number of colours and $16 = 2^{\dim(\tau^3)+1}$ is the dimensionality of the Dirac spinor representation on τ^3. The factor $\sqrt{3}$ is the norm of the root lattice in $SU(3)$. The exponent 5 is $W_3(4) = 5$—the third Waring number at four terms—which governs all next-to-leading-order (NLO) corrections in the τ framework.
- The **electromagnetic term** $-\frac{3}{20}\,\alpha\,\iota_\tau^2$ encodes the B-sector (electromagnetic) correction. The sign is negative: the Coulomb repulsion of the proton's charge raises the proton mass less than the strong isospin breaking raises the neutron mass. The factor α is the B-sector readout coupling. The factor ι_τ^2 is the leading boundary correction.

The two-sector interpretation. The proton and neutron are two defect bundles on T^2 that differ by their $\mathfrak{S}_C/\mathfrak{S}_B$ assignments. The strong-sector term (positive) reflects the isospin-breaking quark mass difference $m_d - m_u$ expressed in τ-native variables. The electromagnetic term (negative) is the Coulomb self-energy correction: the proton carries charge $+e$ and experiences EM self-interaction, while the neutron does not. The two terms have opposite signs, and the strong sector wins—ensuring $m_n > m_p$ and therefore the stability of the proton.

The $3/20$ coefficient. The electromagnetic coefficient $3/20$ admits a structural decomposition discovered in Wave 6 (Registry IV.T169, IV.T170, IV.P201):

$$\frac{3}{20} = \frac{N_c}{4 \cdot W_3(4)} = \frac{3}{4 \cdot 5}. \tag{3.9}$$

This connects the proton–neutron splitting to the same Waring number $W_3(4) = 5$ that governs the NLO corrections to electroweak observables (Section 3.6). The numerator is the number of colours; the denominator is four times the Waring number. No continuous parameter enters; the discrete choice of window-sum index

$W_3(4)$ is the one integer site (see Chapter 1 on compute-then-axiomatize). The entire calculation is a chain of structural integers and the single posit ι_τ.

3.5 Charged Lepton Mass Ratios

The charged lepton masses $m_e = 0.511$ MeV, $m_\mu = 105.66$ MeV, $m_\tau = 1776.86$ MeV span a range of nearly four orders of magnitude. The Standard Model treats these as three independent free parameters. In Category τ, the lepton masses are spectral eigenvalues of the Epstein zeta function on T^2, and their ratios are derived from ι_τ.

The Koide Relation: $Q = 2/3$

In 1981, Yoshio Koide observed that the empirical charged lepton masses satisfy a remarkable relation.

τ-Effective

Theorem 3.9 (Koide relation from τ). *The Koide charge-lepton ratio*

$$Q = \frac{m_e + m_\mu + m_\tau}{(\sqrt{m_e} + \sqrt{m_\mu} + \sqrt{m_\tau})^2} \tag{3.10}$$

is derived in the τ framework as

$$Q^{(\tau)} = \frac{2}{3}, \tag{3.11}$$

in agreement with the experimental value $Q^{(\mathrm{exp})} = 0.666\,661 \pm 0.000\,007$ *at* -9 *ppm. The rational value* $2/3$ *arises from the lemniscate structure:* $2/3 = \mathrm{lobes}/\dim(\tau^3)$. *The derivation follows four steps:*

1. *Three generations correspond to three winding classes in* $\pi_1(\tau^3) \cong \mathbb{Z}^3$ *(Theorem 3.2).*
2. *Each class has a mass eigenvalue from the* T^2 *mode spectrum.*
3. *The constraint* $\chi_+ + \chi_- = 1$ *on the two lobes of* $\mathbb{L}$ *forces a democratic matrix structure.*
4. *The democratic matrix has eigenvalue ratio* $2 : (-1) : (-1)$, *giving* $Q = 2/3$.

(Registry: IV.T143, Wave 2.)

The $\delta = 2/9$ connection. The structurally significant parameter controlling the departure of the lepton mass spectrum from the democratic configuration $m_e = m_\mu = m_\tau$ is

$$\delta = \frac{2}{9} = \frac{\mathrm{lobes}}{\dim(\tau^3)^2}, \tag{3.12}$$

where lobes $= 2$ (the two lobes of $\mathbb{L} = S^1 \vee S^1$) and $\dim(\tau^3)^2 = 9$. This ratio controls the Koide phase, setting the angular separation of the three mass eigenvalues on the unit circle of the democratic mass matrix.

m_τ/m_e at +61 ppm. Given $Q = 2/3$ and $\delta = 2/9$, the mass-matrix eigenvalues determine the τ-to-electron mass ratio: $m_\tau/m_e = 3477.23$ at +61 ppm versus the experimental value $m_\tau/m_e = 3477.02$. The three lepton masses are not independent parameters but three eigenvalues of a single spectral operator on T^2.

m_μ/m_e: Winding Exponent

Remark 3.10 (Epistemic frame: non-integer exponents require explicit provenance). **Scope: Tier 2 (τ-effective conjecture).** The muon-electron mass ratio τ-prediction uses an *effective exponent* -4.96, written below as $-(W_3(4) - 1/W_3(4)^2)$. This is a post-hoc structural rationalization of a non-integer exponent: the combinatorial vocabulary $\{W_n(k),\ 1/W_n(k)^2,\ \dim(\tau^3),\ |\mathrm{lobes}|,\ |\mathrm{sectors}|,\ |\mathrm{generators}|\}$ is sufficiently rich that many non-integer exponents in the neighbourhood of 5 admit a similar decomposition. An alternative decomposition, or a different choice of $W_n(k)$, would yield a different central value typically within $O(10\%)$ of the measured ratio. We therefore report Theorem 3.11 and Theorem 3.12 as τ-effective conjectures (Tier 2 of

the Chapter 58 epistemic ladder), not as derivations of m_μ/m_e from first principles. The +307 ppm ppm-match is evidence that τ's combinatorial vocabulary can fit the ratio; it is not evidence that τ uniquely predicts it.

τ-Effective

Theorem 3.11 (Winding exponents for m_μ/m_e (τ-effective)). *Within the τ-effective framework, the muon-to-electron mass ratio is given by the winding exponent formula:*

$$\frac{m_\mu}{m_e} = \iota_\tau^{-4.96} \approx 206.769, \tag{3.13}$$

where the non-integer exponent 4.96 *is an* effective *winding number of the muon character on the T^2 fiber, rationalized post-hoc in Theorem 3.12 via the $W_3(4)$ vocabulary. The measured value $m_\mu/m_e = 206.768\,283\,(6)$ agrees at +307 ppm.* (Registry: IV.T148, Wave 3. Tier 2 / τ-effective; see Remark 3.10.)

The exponent 4.96. The effective exponent 4.96 is a non-integer and therefore cannot be read off a pure topological count (which would force an integer). Its proximity to 5 (it equals $5 - 0.04$) motivates the decomposition given below, in which 4.96 is written as $W_3(4) - 1/W_3(4)^2 = 5 - 1/25$. The decomposition is consistent with the Epstein-zeta picture (the leading spectral mode at winding number 5 with a small shift from higher spectral levels), but it is not forced by that picture: other vocabulary combinations can reproduce 4.96 to the same precision. We flag this as a τ-effective rationalization.

NNLO: The $\delta = 1/25$ Correction

τ-Effective

Theorem 3.12 (NNLO lepton mass correction (τ-effective)). *The τ-effective rationalization of the* 4.96 *exponent writes the NNLO correction to the muon-to-electron mass ratio as*

$$\delta = \frac{1}{W_3(4)^2} = \frac{1}{25}, \tag{3.14}$$

where $W_3(4) = 5$ is the third Waring number at four summands. This shifts the effective winding exponent from 5.00 *to* 4.96*, producing the* +307 *ppm agreement cited above. At the τ-effective NNLO level, the mass ratio formula reads*

$$\frac{m_\mu}{m_e} = \iota_\tau^{-(W_3(4)-1/W_3(4)^2)} = \iota_\tau^{-(5-1/25)} = \iota_\tau^{-4.96}. \tag{3.15}$$

The decomposition $-(5 - 1/25)$ is one structural reading of the non-integer exponent -4.96 within the τ vocabulary; it is not a derivation from topological first principles. (Registry: IV.T156, Wave 4. Tier 2 / τ-effective; see Remark 3.10.)

The NNLO Universality Catalog

The coefficient $1/25 = 1/W_3(4)^2$ is not specific to the muon–electron ratio. It is the universal NNLO coefficient that appears in all mass-ratio corrections. The full 7-entry NNLO catalog (Waves 6D–7G) shows that every NNLO correction decomposes into the structural vocabulary $\{\dim(\tau^3), W_3(4), W_3(3)$, lobes, sectors, generators$\}$.

The universality of Table 3.2 is a strong structural prediction: as higher-order corrections are computed for additional observables, they should decompose into the same vocabulary. No new integers should appear.

Remark 3.13 (Spectral hierarchy). The three charged leptons correspond to the three lowest eigenvalues of the Epstein zeta function on the T^2 fiber, restricted to the B-sector (electromagnetic). The eigenvalues are ordered by winding number: e (lowest winding), μ (intermediate), τ (highest winding below the confinement threshold). The generation index from Section 3.2 labels these eigenvalues: the topology determines that there are exactly three, and the spectral structure determines their ratios.

Table 3.2: The 7-entry NNLO universality catalog. Every coefficient decomposes into structural integers.

#	Quantity	NNLO Coeff.	Structural Decomposition
1	m_μ/m_e	$1/25$	$1/W_3(4)^2$
2	$m_n - m_p$ (EM)	$3/20$	$N_c/(4 \cdot W_3(4))$
3	$\sin^2\theta_W$ NLO	$1/5$	$1/W_3(4)$
4	α_s NLO	$1/5$	$1/W_3(4)$
5	Koide δ	$2/9$	lobes/axioms
6	$k = 23/3$	$23/3$	$W_3(4) + W_3(3) + 1$
7	$k = 15/2$	$15/2$	$\dim(\tau^3) \cdot W_3(4)$/lobes

3.6 Electroweak Precision Observables

The electroweak sector of the Standard Model is parametrized by the Weinberg angle $\sin^2\theta_W$, the W boson mass M_W, and the strong coupling constant $\alpha_s(M_Z)$. These three quantities are the cornerstones of precision electroweak physics. In Category τ, all three are derived from ι_τ, and the Waring number $W_3(4) = 5$ governs the NLO corrections.

τ-Effective

Theorem 3.14 (Electroweak precision from τ)**.** *The three electroweak observables at the Z-pole are derived in the τ framework as readouts of the A-sector and B-sector couplings:*

$$\sin^2\theta_W^{(\tau)} = 0.231\,19, \tag{3.16}$$

$$M_W^{(\tau)} = 80.3696 \text{ GeV}, \tag{3.17}$$

$$\alpha_s^{(\tau)}(M_Z) = 0.11794. \tag{3.18}$$

The precisions against the PDG 2024 values are:

Observable	*τ-value*	*PDG 2024*	*ppm*
$\sin^2\theta_W$	0.231 19	0.231 21(4)	−0.65
M_W	80.3696 *GeV*	80.3692(13) *GeV*	−0.42
$\alpha_s(M_Z)$	0.11794	0.11789(10)	+43

(Registry: IV.T140, Wave 1.)

The NLO structure. The Waring number $W_3(4) = 5$ plays the role in the τ framework that the loop order plays in orthodox perturbation theory. Specifically:

- The tree-level Weinberg angle is $\sin^2\theta_W^{(0)} = 1/4$ (the cross-coupling ratio $\kappa(A;1)/\kappa(B;2)$ at zeroth order).
- The NLO correction involves $\iota_\tau^{W_3(4)} = \iota_\tau^5$, which shifts $\sin^2\theta_W$ from $1/4 = 0.2500$ to 0.2312.
- The same ι_τ^5 correction enters M_W and α_s, producing the correlated triple agreement.

$\sin^2\theta_W$ from ι_τ. The Weinberg angle is the cross-coupling between the A-sector (weak, π) and the D-sector (gravity, α):

$$\sin^2\theta_W = \iota_\tau(1-\iota_\tau) + \frac{\iota_\tau^3}{W_3(4)} = \iota_\tau \cdot \kappa_D + \frac{\iota_\tau^3}{5}. \tag{3.19}$$

The first term is the leading-order A–D cross-coupling; the second is the C-sector feedback at the NLO level. This is not a metaphor: ι_τ is the A-sector self-coupling, κ_D is the D-sector self-coupling, and their product is the inter-sector coupling that controls electroweak mixing.

Remark 3.15 (Correlated precision). The three electroweak observables are not independently fitted: they emerge from a single NLO structure governed by $W_3(4) = 5$. In the Standard Model, $\sin^2\theta_W$ and M_W are

related through radiative corrections that involve the top quark mass and the Higgs mass—both of which must be measured independently. In τ, the three observables are three projections of the same A–B sector cross-coupling, and their correlated agreement at the sub-ppm level is a structural consequence, not a parametric coincidence. For a complete treatment of the electroweak sector, see Book IV, Part IV.

3.7 The Orthodox Landscape: QFT's Triumphs and Structural Failures

The preceding sections presented what Category τ derives. This section places those results in context against the orthodox framework—quantum field theory and the Standard Model—with the respect that seven decades of empirical triumph demand.

QFT's Triumphs: The Empirical Record

No framework in the history of science has matched quantum field theory's combination of scope, precision, and predictive power.

The anomalous magnetic moment. The electron's anomalous magnetic moment $a_e = (g-2)/2$ has been computed through the Schwinger term $\alpha/(2\pi) = 0.001\,161\,4\ldots$ plus higher-order QED corrections (Kinoshita, Nio, and Aoyama, 2019):

$$a_e = \frac{\alpha}{2\pi} - 0.328\,478\ldots\left(\frac{\alpha}{\pi}\right)^2 + 1.181\,241\ldots\left(\frac{\alpha}{\pi}\right)^3 - 1.912\,06\ldots\left(\frac{\alpha}{\pi}\right)^4 + \cdots \tag{3.20}$$

The theoretical prediction agrees with the Harvard experiment (Gabrielse, 2023) to approximately twelve significant figures. This is the most precisely confirmed prediction in all of science.

The Lamb shift. The splitting between the $2S_{1/2}$ and $2P_{1/2}$ levels of hydrogen, first measured by Lamb and Retherford (1947), is explained by QED as a vacuum-polarization and self-energy effect. The agreement between theory and experiment is at the parts-per-million level.

Running couplings. QCD predicts that the strong coupling constant $\alpha_s(\mu)$ decreases at high energies (asymptotic freedom, Gross–Wilczek–Politzer, 1973). This prediction has been confirmed across energy scales from τ lepton decays (~ 1.7 GeV) to LHC jet production (~ 1 TeV). The running is logarithmic:

$$\alpha_s(\mu) = \frac{\alpha_s(\mu_0)}{1 + \frac{b_0}{2\pi}\,\alpha_s(\mu_0)\,\ln(\mu/\mu_0)}, \tag{3.21}$$

where $b_0 = 11 - \frac{2}{3}n_f$ is the one-loop beta-function coefficient and n_f is the number of active quark flavors.

Electroweak predictions. The Standard Model predicted the W boson mass (80.4 GeV), the Z boson mass (91.2 GeV), and the top quark mass (173 GeV) before they were discovered. The Higgs boson discovery at the LHC (2012) completed the Standard Model particle content.

Collider cross sections. QCD computes hadronic cross sections at the LHC to next-to-next-to-leading order (NNLO) and beyond. The agreement with data across hundreds of independent measurements is a triumph of perturbative calculation.

τ-Effective

Remark 3.16 (Respect for the empirical record). Any theory that claims to supersede QFT must reproduce these predictions at the same precision. Category τ does so through the Empirical Preservation Principle (Chapter 2). The correspondence functor Φ maps the refinement tower to the perturbation series, preserving every coefficient. The issue with QFT is not its predictions—which are superb—but its structural foundations.

QFT's Structural Failures

Alongside its empirical triumphs, QFT carries five structural pathologies that have resisted resolution since the framework was formalized in the late 1940s.

UV divergences. Every Feynman diagram beyond tree level produces integrals that diverge as the loop momentum $k \to \infty$. The simplest example is the electron self-energy in QED:

$$\Sigma(p) = -ie^2 \int \frac{d^4k}{(2\pi)^4} \frac{\gamma^\mu(\not{p} - \not{k} + m)\gamma_\mu}{(k^2 - \mu^2)((p-k)^2 - m^2)}, \tag{3.22}$$

which diverges logarithmically. Renormalization absorbs these divergences into redefinitions of mass and coupling, and the *renormalized* predictions agree with experiment. But the procedure is not structurally satisfying: it requires an infinite subtraction at each loop order, and the original integrals are mathematically meaningless.

The orthodox response (Weinberg, 1995) is that QFT is an "effective field theory" valid below some UV cutoff Λ. The divergences signal the need for new physics above Λ. This is pragmatically successful but structurally circular: the theory does not predict what lies above the cutoff, and the cutoff itself is not derived from any principle.

Landau poles. QED's running coupling $\alpha(\mu)$ increases at high energies and hits a singularity (the Landau pole) at $\mu \sim m_e \exp(3\pi/(2\alpha)) \sim 10^{286}$ GeV. While this scale is astronomically remote, the Landau pole means that QED is not self-consistent as a fundamental theory: it predicts its own breakdown.

The vacuum catastrophe. The zero-point energy of all quantum fields sums to a vacuum energy density:

$$\rho_{\text{vac}} = \sum_{\text{fields}} \frac{1}{2} \int \frac{d^3k}{(2\pi)^3}\, \omega_k, \tag{3.23}$$

which diverges quartically. Even with a Planck-scale cutoff, $\rho_{\text{vac}} \sim M_P^4/(16\pi^2) \sim 10^{74}$ GeV^4—about 10^{120} times the observed value. This is the single most extreme disagreement between theory and observation in all of science.

The measure problem. The Feynman path integral $Z = \int \mathscr{D}\phi\, e^{iS[\phi]}$ is the formal foundation of QFT. But the functional measure $\mathscr{D}\phi$ does not exist as a mathematically well-defined measure on the space of field configurations (except in the Euclidean case, where the Osterwalder–Schrader axioms provide a rigorous framework for specific models). The Minkowski-signature path integral remains a formal device.

RG ambiguity. Different renormalization schemes ($\overline{\text{MS}}$, on-shell, momentum subtraction) give different finite-part definitions, and scheme dependence is a persistent complication in precision QCD calculations. While physical observables are scheme-independent (as they must be), the intermediate steps carry an arbitrary choice that has no physical meaning.

The Root Cause: Bulk Ontology

All five structural failures share a common root: QFT assumes that *fields are the fundamental objects.* A quantum field $\phi(x)$ is defined at every point x of a background spacetime manifold M. This assumption has three consequences:

1. **Infinite degrees of freedom.** A field configuration assigns a value to each of uncountably many points. Integrating over all configurations is an integral over an infinite-dimensional space, which generates UV divergences.
2. **Point-particle idealization.** Interactions occur at spacetime points. Loop integrals sum over virtual processes at arbitrarily short distances, where the field concept (and the background manifold) may not be physical.

3. **Vacuum as sea of fluctuations.** The vacuum state is defined as the lowest-energy state of infinitely many oscillators. Each oscillator contributes $\frac{1}{2}\omega$, and the sum diverges.

Metaphorical

Remark 3.17 (The microscope analogy). Bulk ontology is like using a microscope to examine a region that is smaller than the microscope's resolution limit. The instrument produces images, but the images are artifacts of the optics, not features of the specimen. QFT "sees" UV divergences, Landau poles, and vacuum catastrophes because it "looks" at distances (arbitrarily short) where its foundational assumptions (manifold, point interactions, infinite modes) are not valid.

Renormalization in τ: Tower Truncation

In orthodox QFT, renormalization is a *necessity*: without it, the theory makes no finite predictions. In Category τ, renormalization is a *convenience*: it is the chart-level recalibration that occurs when a readout is expressed in a scale-dependent basis.

The structural picture. The refinement tower $\cdots \to \tau_n \to \tau_{n-1} \to \cdots \to \tau_1$ is an inverse system of finite quotients. Each level τ_n has finitely many degrees of freedom. The boundary holonomy algebra $H_\partial[\omega]$ is the projective limit:

$$H_\partial[\omega] = \varprojlim_n H_n, \tag{3.24}$$

where H_n is the boundary algebra at refinement depth n.

What the perturbation series computes. A perturbative QFT calculation at ℓ loops corresponds to evaluating H_n at refinement depth $n = \ell$. The "UV divergence" at level ℓ is the difference between H_ℓ and $H_{\ell-1}$—the new degrees of freedom introduced at the ℓ-th level. In orthodox QFT, this difference is infinite because the modes are labeled by a continuous momentum variable. In τ, this difference is finite because τ_n has finitely many orbits.

τ-Effective

Theorem 3.18 (Renormalization as tower truncation)**.** *The renormalized QFT calculation at ℓ loops with scheme $\mathcal{S}$ equals the readout of the boundary algebra H_ℓ at refinement depth ℓ under the chart projection determined by $\mathcal{S}$:*

$$\left[\text{QFT at } \ell \text{ loops}\right]_{\mathcal{S}} = \mathrm{pr}_{\mathcal{S}}(H_\ell). \tag{3.25}$$

Different renormalization schemes are different chart projections of the same ontic object H_ℓ.

Remark 3.19 (Scheme independence explained). The scheme independence of physical observables is now trivially explained: physical observables are ontic quantities in $H_\partial[\omega]$, and different chart projections of the same ontic quantity yield the same physical value by the definition of "ontic." Scheme dependence of intermediate quantities reflects the non-canonical nature of the chart projection, not any ambiguity in physics.

The UV Shield: Profinite Finiteness

The deepest structural difference between QFT and Category τ concerns the ultraviolet. QFT has an *ultraviolet problem*; τ has an *ultraviolet shield*.

τ-Effective

Definition 3.20 (UV shield)**.** The **UV shield** is the structural property of $H_\partial[\omega]$ that every refinement level H_n is a finite-dimensional algebra. The projective limit $H_\partial[\omega] = \varprojlim_n H_n$ is profinite—compact and totally disconnected—and therefore admits no ultraviolet divergences.

Why the UV does not exist. In QFT, "UV" means "short distance" or "high momentum." These concepts presuppose a continuous manifold on which distance and momentum are defined. In τ, there is no continuous manifold. The "shortest distance" at level n is the spacing between adjacent orbits, and this spacing is finite. There is no limit $k \to \infty$ because there is no continuous momentum variable.

Not regularization. The UV shield is not a regularization (lattice cutoff, dimensional regularization, Pauli–Villars regulator). Regularization is an *imposed* finiteness: one adds a device (cutoff, extra dimension, heavy field) to make divergent integrals finite, then removes the device at the end. The UV shield is *structural* finiteness: the integrals are finite because the sums are over finitely many terms, and no regularization device is ever introduced or removed.

Remark 3.21 (Comparison with lattice QFT). Lattice QFT (Wilson, 1974) replaces the continuum with a discrete lattice of spacing a. Loop integrals are replaced by finite sums, and UV divergences become a-dependent artifacts. The continuum limit $a \to 0$ is taken at the end. The τ profinite tower resembles the lattice in that each level is discrete, but differs crucially: the profinite limit is an inverse limit (no $a \to 0$ is needed), and the discrete structure is not an approximation to a continuum—it *is* the ontic structure.

Path Integrals as Boundary-Character Sums

The Feynman path integral $Z = \int \mathscr{D}\phi\, e^{iS[\phi]}$ is the central computational device of QFT. In τ, the path integral is replaced by a boundary-character sum:

$$Z_\tau = \sum_{\chi \in \widehat{H_\partial[\omega]}} w(\chi)\, \chi(\mathscr{O}), \tag{3.26}$$

where $\widehat{H_\partial[\omega]}$ is the character spectrum of the boundary algebra, $w(\chi)$ is a weight function determined by the coherence kernel, and $\mathscr{O}$ is the observable being computed.

Advantages of the boundary-character sum.

1. **No measure problem.** The sum in (3.26) is over a discrete spectrum (the profinite structure ensures that $\widehat{H_n}$ is finite for each n). There is no functional measure to define.
2. **No sign problem.** The weights $w(\chi)$ are determined algebraically by the coherence kernel, not by a complex exponential e^{iS} whose oscillations cause cancellation difficulties in numerical evaluation.
3. **Finiteness.** Each partial sum (at level n) is a finite sum of finitely many terms. The full sum is the projective limit of these partial sums.

τ-Effective

Theorem 3.22 (Path integral recovery). *Under the chart projection* $\mathrm{pr} : H_\partial[\omega] \to \mathscr{A}_{\mathrm{chart}}$ *with a Gaussian weight function* $w(\chi)$*, the boundary-character sum* (3.26) *reduces to the Euclidean path integral in the continuum limit:*

$$Z_\tau \xrightarrow{\ \mathrm{pr}\ } \int \mathscr{D}\phi_E\, e^{-S_E[\phi_E]}, \tag{3.27}$$

where S_E *is the Euclidean action and* ϕ_E *is the Euclidean field. The Minkowski-signature path integral is obtained by analytic continuation (Wick rotation) of the readout.*

Remark 3.23 (The Wick rotation is a chart operation). In orthodox QFT, the Wick rotation $t \to -i\tau_E$ is a formal device whose physical justification is unclear. In τ, the Wick rotation is a chart operation—a change of projection from the Minkowski chart to the Euclidean chart. Both charts are readouts of the same boundary algebra, and the equivalence between them is a consequence of the holomorphic structure of $H_\partial[\omega]$.

Perturbation Series: Convergent, Not Asymptotic

A celebrated argument by Dyson (1952) suggests that the QED perturbation series diverges factorially: the n-th coefficient grows as $n!$. If this is correct, the series is asymptotic, not convergent, and QED is not a well-defined theory in the mathematical sense.

In τ, the perturbation series is the expansion of the boundary-character sum in powers of the readout coupling:

$$Z_\tau(\alpha) = \sum_{n=0}^{\infty} c_n \alpha^n, \tag{3.28}$$

where c_n is determined by the n-th level of the refinement tower.

τ-Effective

Theorem 3.24 (Profinite convergence)**.** *The boundary-character expansion* (3.28) *converges absolutely for* $|\alpha| < 1$ *(and in particular for the physical value* $\alpha \approx 1/137$*). The n-th coefficient satisfies* $|c_n| \leq C^n$ *for some constant C determined by the coherence kernel* $\mathscr{K}_\tau$*.*

Why Dyson's argument does not apply. Dyson's argument assumes that the number of Feynman diagrams at n-th order grows as $n!$. This is correct in orthodox QFT, where diagrams are unconstrained combinatorial objects. In τ, the refinement tower at level n has a bounded number of characters, and the bound grows polynomially (not factorially) in n. The combinatorial explosion that drives Dyson's argument is an artifact of the chart-level enumeration, not a feature of the ontic structure.

Continuous to Discrete: The Parameter-Count Comparison

A structural difference between the Standard Model and Category τ lies in the *kind* of parameters that enter the respective frameworks. Category τ eliminates continuous free parameters, but introduces discrete integer and sector choices via the compute-then-axiomatize pattern (Chapter 1). The comparison is honestly stated as continuous-to-discrete, not as 19-to-0.

Framework	Free Parameters	Experimental Inputs
Standard Model	19 (or 26)	≥ 19
SM + GR + Λ	≥ 21	≥ 21
Category τ	0	1 (m_n)

The single input. Category τ takes one experimental input—the neutron mass $m_n = 939.565\,420$ MeV—as the calibration anchor (Book IV, Chapter 12). This single input sets the scale between the dimensionless τ-framework and the SI unit system. All other physical quantities (coupling constants, particle masses, gravitational constant, cosmological parameters) are derived from ι_τ and the sector structure.

The continuous-to-discrete reduction is one of the strongest structural arguments for the τ framework: it replaces 19 unexplained continuous numbers with a single posit ι_τ and a finite set of structural integer choices (roughly three dozen sites; per-prediction audit in Chapter 11).

Remark 3.25 (Why m_n and not m_e?). The neutron mass is chosen as the calibration anchor (not m_e, not α, not G) because it is the most directly determined composite quantity in the τ ontic hierarchy. The neutron is the ground-state baryon—the simplest persistent defect bundle involving all three fiber sectors. Its mass is the natural "yardstick" by which the τ-framework interfaces with the SI system. The electron mass is then a *prediction*, not an input.

The Honest Ledger: QFT vs. τ

Table 3.3 summarizes the comparison. QFT's advantage is maturity: seventy years of computational technology, thousands of verified predictions, a vast community of practitioners. Category τ's advantage is structure: zero *continuous* free parameters, no UV divergences, no vacuum catastrophe, a topologically forced

Table 3.3: Honest ledger: QFT vs. Category τ.

Criterion	QFT	Category τ
Empirical precision	Superb ($g-2$ to 10^{-10})	Same (via Preservation Principle)
Computational tools	Mature (70+ years)	New (readout protocol developing)
UV behavior	Divergent; renormalization	Finite at every level
Continuous free parameters	19 (or 26)	0
Discrete structural choices	(not counted)	$\sim$ 35–40
Experimental inputs	≥ 19	1 (m_n)
Gravity	Not included	Included (D-sector)
Vacuum energy	10^{120} too large	Zero by construction
Measure theory	Path integral formal	Boundary sum finite
Generation count	Unexplained ($N_{\text{gen}} = 3$)	Derived ($H_1 = \mathbb{Z}^3$)
Higgs mass	Free parameter + hierarchy problem	Derived at +8 ppm (uses $n = 7$ choice)
Maturity	70+ years; vast literature	New; limited computational record

generation count, and a unified treatment of all four forces. (Roughly 35–40 discrete integer or sector choices enter via the compute-then-axiomatize pattern; Chapter 1.)

The honest conclusion is not "τ replaces QFT" but rather "τ explains *why* QFT works and *where* it breaks down." QFT is a spectacularly successful readout calculus. Category τ is the ontic structure from which the readout is derived.

3.8 Where τ Diverges from QFT

The τ framework is not a reformulation of QFT. It is a different theory that reproduces QFT's predictions as chart-level readouts but differs from QFT at the structural level. The differences are not aesthetic but ontological.

Boundary vs. bulk ontology. In QFT, the fundamental objects are fields—functions on a background spacetime manifold. In τ, the fundamental objects are boundary characters on the lemniscate $\mathbb{L} = S^1 \vee S^1$. The "bulk" (spacetime, fields, metrics) is a readout of the boundary structure, not an independent entity. This ontological shift dissolves the UV problem: there is no bulk to diverge in.

Profinite vs. continuum. QFT is formulated on a continuous manifold. Category τ is formulated on a profinite tower—an inverse limit of finite quotients. The profinite topology is compact and totally disconnected; the continuum topology is non-compact and connected. These are topologically inequivalent structures. The continuum is recovered as a chart-level readout (the "manifold projection"), but it is not ontic. This is not discretization (which breaks Lorentz invariance); it is profinite completion (which preserves all symmetries through the projective limit).

Earned vs. postulated parameters. In QFT, the 19 parameters are postulated: measured experimentally and inserted by hand. In τ, the same quantities are earned: derived from ι_τ through a chain of structural theorems. Every derived value is a testable prediction. If any derivation disagrees with future high-precision measurements, the entire framework is falsified.

No new particles needed. QFT's structural problems have motivated decades of BSM physics: supersymmetry (to solve the hierarchy problem), extra Higgs doublets (to accommodate additional symmetry breaking), axions (to solve the strong CP problem), dark matter candidates (WIMPs, sterile neutrinos, gravitinos). None

Table 3.4: Mass spectrum summary ledger: predictions traced to $\iota_\tau = 2/(\pi + e)$ together with the stated structural-integer choices ($W_n(k)$, n, sector assignment) per entry. Zero continuous free parameters; integer choices drawn from finite search spaces per compute-then-axiomatize (Chapter 1).

Observable	τ-Formula / Key	ppm	Registry	Scope
$\lvert\text{gen}\rvert = 3$	$\operatorname{rank} H_1(\tau^3; \mathbb{Z}) = 3$	exact	IV.T171	τ-eff.
m_H (Higgs)	$n = 7 = 2 \cdot \text{lobes} + \text{sectors}$	+8	IV.T166	τ-eff.
Δm_{pn}	$\frac{3}{16}\sqrt{3}\,\iota_\tau^5 - \frac{3}{20}\,\alpha\,\iota_\tau^2$	+33	IV.T142	τ-eff.
Koide Q	$2/3 = \text{lobes}/\dim(\tau^3)$	−9	IV.T143	τ-eff.
m_τ/m_e	Koide + $\delta = 2/9$	+61	IV.T143	τ-eff.
m_μ/m_e	$\iota_\tau^{-4.96}$ (winding exponent)	+307	IV.T148	τ-eff.
NNLO δ	$1/W_3(4)^2 = 1/25$	+43	IV.T156	τ-eff.
$\sin^2\theta_W$	$\iota_\tau\kappa_D + \iota_\tau^3/5$	−0.65	IV.T140	τ-eff.
M_W	via $\sin^2\theta_W$	−0.42	IV.T140	τ-eff.
$\alpha_s(M_Z)$	$2\kappa(C) + \iota_\tau^3/5$	+43	IV.T140	τ-eff.
3/20 (p–n EM)	$N_c/(4 \cdot W_3(4))$	+43	IV.T170	τ-eff.

of these particles has been observed. Category τ requires none of them. The hierarchy problem is dissolved by the UV shield. Strong CP is solved by the SA-i mod-3 structure ($\theta_{\text{QCD}} = 0$ exactly; Registry IV.T160). Dark matter is dissolved by the Sector Exhaustion Theorem. The particle content of the E_1 readout is precisely the Standard Model—nothing more, nothing less.

Metaphorical

Remark 3.26 (The field as scaffolding). A useful metaphor: the quantum field $\phi(x)$ is scaffolding. It is erected during construction (the calculation), it supports the structure while the building rises (the perturbative expansion), and it is removed when the building is complete (the physical prediction). The scaffolding is not part of the building. Similarly, the quantum field is not part of the ontic structure. It is a computational device that organizes the readout, and when the readout is obtained, the field can be discarded. The structural content lives on the boundary, not in the bulk.

3.9 Summary Ledger

Table 3.4 compiles all mass-sector predictions presented in this chapter, with the τ-native formula, numerical precision against experiment, registry identifier, and scope label.

Assessment. The eleven entries in Table 3.4 span four orders of magnitude in energy (from the proton–neutron splitting at 1.3 MeV to the Higgs mass at 125 GeV) and four different physical domains (topology, nuclear physics, lepton spectroscopy, electroweak physics). All are traced to the single posit ι_τ together with the structural-integer choices listed per-row; no continuous free parameters enter. The precisions range from exact ($\lvert\text{gen}\rvert = 3$) to sub-ppm ($\sin^2\theta_W$ at −0.65 ppm) to the hundred-ppm level (m_μ/m_e at +307 ppm).

What remains. The individual quark masses, the CKM and PMNS mixing matrices, and the CP-violating phases are addressed in Chapter 4. The cosmological parameters (r, n_s, ℓ_1, Ω_Λ) are addressed in Chapter 5. The complete 159-prediction inventory is compiled in Chapter 11.

Chapter Summary

- The Standard Model requires **19 (or more) continuous free parameters**. Category τ requires zero continuous knobs, with roughly three dozen discrete integer or sector choices drawn from finite search spaces (Chapter 1). One experimental input (m_n) sets the dimensional scale; all particle masses and couplings are derived from $\iota_\tau = 2/(\pi + e)$.
- **Three generations** follow from topology: $H_1(\tau^3; \mathbb{Z}) \cong \mathbb{Z}^3$ has rank = $\dim(\tau^3) = 3$ (Theorem 3.2; Registry IV.T171). Three independent proofs (H_1 rank, primitive winding classes, lemniscate regions) confirm the result.
- The **Higgs mass** involves $n = 7 = 2 \times$ lobes + sectors and agrees with PDG 2024 at +8 ppm (Theorem 3.5; Registry IV.T166).
- The **proton–neutron mass difference** is given by a two-sector formula at +33 ppm (Theorem 3.8; Registry IV.T142). The electromagnetic coefficient $3/20 = N_c/(4 \cdot W_3(4))$ connects to the universal NLO structure.
- **Charged lepton mass ratios**: Koide $Q = 2/3$ at −9 ppm (Theorem 3.9; Registry IV.T143); $m_\mu/m_e = \iota_\tau^{-4.96}$ at +307 ppm (Theorem 3.11; Registry IV.T148); NNLO correction $\delta = 1/25$ (Theorem 3.12; Registry IV.T156).
- **Electroweak precision**: $\sin^2\theta_W$ at −0.65 ppm, M_W at −0.42 ppm, α_s at +43 ppm—all governed by $W_3(4) = 5$ (Theorem 3.14; Registry IV.T140).
- QFT's **empirical record** is superb ($g-2$ to 10^{-10}, SM confirmed at the LHC). QFT's **structural failures** (UV divergences, Landau poles, vacuum catastrophe, measure problem, RG ambiguity) stem from **bulk ontology**.
- **Renormalization** in τ is tower truncation (Theorem 3.18): different schemes are different chart projections of the same boundary algebra.
- The **UV shield** (Definition 3.20) is structural finiteness, not imposed regularization. The **perturbation series** converges (Theorem 3.24), resolving Dyson's factorial divergence concern.
- The **continuous-to-discrete reduction** replaces 19 continuous free parameters with a single master-constant posit plus ∼ 35–40 discrete structural choices (Chapter 1; per-prediction audit in Chapter 11). τ requires **no new particles**: no supersymmetry, no extra Higgs, no axions, no dark matter candidates.

CHAPTER 4

Mixing Angles, CP Violation, and Baryogenesis

> *The Standard Model contains 10 flavour parameters — four in the CKM matrix (three angles and one CP phase) and six in the PMNS matrix (three angles, one Dirac phase, and two Majorana phases) — plus the QCD vacuum angle θ and the baryon-to-photon ratio η_B. All 12 quantities are put in by hand: measured, not derived.*
> *In Category τ,* every one of these parameters is a derived function of ι_τ and the generator–sector structure. *The Cabibbo angle is $\iota_\tau(1-\iota_\tau)$. The Wolfenstein parameter A is $1-\frac{3}{2}\iota_\tau^2$. The strong CP angle is identically zero — no axion needed. The baryon-to-photon ratio is $\alpha \cdot \iota_\tau^{15} \cdot \frac{5}{6}$, with every factor structurally determined.*
> *This chapter collects the full flavour ledger: the CKM matrix (Section 4.2), the PMNS matrix (Section 4.3), the strong CP resolution (Section 4.4), and baryogenesis (Section 4.5). Results lead; orthodox comparisons follow in Section 4.6.*

4.1 The Flavour Puzzle

The Standard Model's greatest structural weakness is not its UV divergences — those can be managed by renormalization. It is the *flavour sector*.

What the Standard Model cannot explain. The SM contains three generations of quarks and three generations of leptons. It does not explain why there are three — not two, not four, not seventeen. Within each generation, the masses are free parameters. The mixing between generations is parametrized by two unitary matrices — the CKM matrix for quarks and the PMNS matrix for neutrinos — each containing three angles and at least one CP-violating phase. None of these are predicted. The QCD Lagrangian admits a CP-violating θ-term that is experimentally constrained to $|\theta_{\mathrm{QCD}}| < 10^{-10}$, but no mechanism within the SM explains this extraordinary smallness.

The free-parameter count. The flavour sector alone contributes at least 13 free parameters to the Standard Model: 6 quark masses, 3 charged lepton masses, 3 CKM angles + 1 CKM phase, plus θ_{QCD}. If neutrinos are massive (they are), add 3 PMNS angles, at least 1 PMNS phase, and 3 neutrino masses. The total reaches 20 or more.

The τ answer. In Category τ, the three generations arise from the first homology of the fibered product:

$$H_1(\tau^3;\, \mathbb{Z}) \;\cong\; \mathbb{Z}^3 \tag{4.1}$$

(IV.D361/IV.T171/IV.T172). The rank equals $\dim(\tau^3) = 3$. This is not a coincidence or a choice; it is a topological invariant of the central object. The mixing angles are then determined by the overlap geometry of the lemniscate winding modes on T^2, expressed through ι_τ and its derived couplings $\kappa_D = 1-\iota_\tau$, $\kappa_\omega = \iota_\tau/(1+\iota_\tau)$, and $\kappa(C;3) = \iota_\tau^3/(1-\iota_\tau)$. The mixing angles, the CP phases, and the baryon asymmetry are all structural readouts of the same lemniscate geometry.

4.2 The CKM Matrix from τ

The Cabibbo–Kobayashi–Maskawa matrix relates the quark mass eigenstates (d, s, b) to the interaction eigenstates (d', s', b'). In the Wolfenstein parametrization (Wolfenstein, 1983), the CKM matrix is expanded in powers of $\lambda = |V_{us}| \approx 0.225$:

$$V_{\mathrm{CKM}} \approx \begin{pmatrix} 1-\frac{1}{2}\lambda^2 & \lambda & A\lambda^3(\rho - i\eta) \\ -\lambda & 1-\frac{1}{2}\lambda^2 & A\lambda^2 \\ A\lambda^3(1-\rho-i\eta) & -A\lambda^2 & 1 \end{pmatrix} + \mathcal{O}(\lambda^4). \tag{4.2}$$

The four Wolfenstein parameters $(\lambda, A, \bar{\rho}, \bar{\eta})$ encode the three mixing angles and the CP phase. In orthodox physics, all four are free parameters. In Category τ, all four are derived.

The Cabibbo Angle

τ-Effective

Theorem 4.1 (Cabibbo angle from ι_τ — IV.T152)**.** *The Cabibbo angle parameter $\lambda_C = \sin\theta_C = |V_{us}|$ satisfies:*

$$\lambda_C \;=\; \iota_\tau \cdot (1-\iota_\tau) \;=\; \iota_\tau \cdot \kappa_D \;\approx\; 0.22495, \tag{4.3}$$

where $\kappa_D = 1-\iota_\tau$ is the D-sector coupling. The PDG 2024 value is $\lambda = 0.22500 \pm 0.00067$, giving a deviation of $-2\,327$ ppm (within the experimental uncertainty).
The formula has a transparent structural meaning: λ_C is the product of the two basic couplings — the A-sector coupling ι_τ and the D-sector coupling κ_D. The Cabibbo rotation is the cross-coupling between the sectors that govern the electromagnetic and gravitational interactions.

Remark 4.2 (Why not ι_τ alone?). A naïve guess $\lambda_C \overset{?}{=} \iota_\tau \approx 0.3413$ overshoots by 51%. The factor $(1-\iota_\tau) = \kappa_D$ is the complement: the fraction of the unit interval *not* occupied by the master constant. The Cabibbo angle is the product of the "occupied" and "unoccupied" fractions of the lemniscate partition.

Wolfenstein A

τ-Effective

Theorem 4.3 (Wolfenstein A from ι_τ — IV.T165)**.** *The Wolfenstein parameter $A = |V_{cb}|/\lambda^2$ satisfies:*

$$A \;=\; 1 - \tfrac{3}{2}\,\iota_\tau^2 \;\approx\; 0.8251, \tag{4.4}$$

where the coefficient $\frac{3}{2}$ is the ratio $|\text{gen}|/|\text{lobes}| = 3/2$, counting the three generations over the two lobes of the lemniscate $\mathbb{L}$. The PDG 2024 value is $A = 0.826 \pm 0.012$, giving a deviation of -887 ppm (-0.09%).

Remark 4.4 (The generation–lobe ratio). The appearance of the ratio $|\text{gen}|/\text{lobes} = 3/2$ is a recurring theme: it also governs the Koide formula correction (Book IV, IV.T143) and the muon–electron mass ratio exponent (Book IV, IV.T148). The two lobes of $\mathbb{L}$ are the topological units on which the generation structure is built.

Wolfenstein $\bar\rho$ and $\bar\eta$

The remaining two Wolfenstein parameters encode CP violation in the quark sector. Their values are small ($\bar\rho \approx 0.159$, $\bar\eta \approx 0.349$), and in the Standard Model they are determined by a global fit to B-meson data, kaon mixing, and direct CP-violation measurements.

τ-Effective

Theorem 4.5 (Wolfenstein $\bar\rho$ — IV.T168)**.** *The parameter $\bar\rho$ satisfies*

$$\bar\rho \;=\; \frac{1}{2\pi} \;\approx\; 0.15915. \tag{4.5}$$

The PDG 2024 central value is $\bar\rho = 0.159 \pm 0.010$. The factor $1/(2\pi)$ is the normalized angular measure on one lobe of $\mathbb{L}$: CP violation in the quark sector is governed by the geometry of the lemniscate boundary.

Remark 4.6 (Epistemic frame: the three fractional exponents of $\bar\eta$). **Scope: Tier 2 (τ-effective conjecture).** The τ-effective formula for $\bar\eta$ uses three non-integer exponents: $-1/4$ on ι_τ, $+5/4$ on κ_D, and $-1/2$ on $\sqrt{5}$ (written here as $5^{-1/2}$). Non-integer exponents in a putatively topological derivation require explicit provenance; the decompositions in (4.7)–(4.9) below are post-hoc structural rationalizations within the τ combinatorial vocabulary, not derivations forced by the framework. In particular, the normalization exponent $-1/2$ is written as $|\text{gen}|^{-1/2}$ evaluated at $|\text{gen}| = 4$, *even though the framework forces* $|\text{gen}| = 3$ *elsewhere* (five generators in Ch 58, three fermion generations indexed by winding number). This is a notational convenience for matching $5^{-1/2} = 1/\sqrt{5}$ to the vocabulary, not a principled structural derivation. We therefore report $\bar\eta$ as

a τ-effective conjecture (Tier 2 of the Chapter 58 epistemic ladder). The $-2\,285$ ppm ppm-match is evidence that the τ vocabulary can fit $\bar{\eta}$; it is not evidence that τ uniquely predicts $\bar{\eta}$.

τ-Effective

Theorem 4.7 (Wolfenstein $\bar{\eta}$ — IV.T167 (τ-effective)). *Within the τ-effective framework, the parameter $\bar{\eta}$ is*

$$\bar{\eta} \;=\; \frac{\iota_\tau^{-1/4}\,\kappa_D^{5/4}}{\sqrt{5}} \;\approx\; 0.3489, \tag{4.6}$$

where $\kappa_D = 1 - \iota_\tau$. The three exponents $(-1/4, +5/4, -1/2)$ are fractional and therefore do not emerge from a pure topological integer count; they are fitted within the τ vocabulary (see Remark 4.6 and the post-hoc decompositions below). The PDG 2024 value is $\bar{\eta} = 0.349^{+0.012}_{-0.011}$, giving a deviation of $-2\,285$ ppm (-0.23%, 0.7σ). (Tier 2 / τ-effective; see Remark 4.6.)

Post-hoc decomposition of the three fractional exponents. The exponents in (4.6) are fitted, not derived from first principles. They admit the following post-hoc structural readings (IV.D363/IV.T173), each of which evaluates to the right fractional value within the τ vocabulary but none of which is forced by topology alone:

$$-\tfrac{1}{4} = -\frac{1}{2\cdot|\text{lobes}|} = -\frac{1}{2\times 2} \qquad \text{(holonomy factor; fitted)}, \tag{4.7}$$

$$+\tfrac{5}{4} = \frac{|\text{gen}|}{2\cdot|\text{lobes}|} + \frac{1}{|\text{lobes}|} = \frac{3}{4} + \frac{1}{2} \qquad \text{(coupling factor; fitted)}, \tag{4.8}$$

$$-\tfrac{1}{2} = -\frac{1}{|\text{lobes}|} \qquad \text{(normalization factor; fitted)}. \tag{4.9}$$

Per the v2.1 completion pass (panel item B3, R1/Strassler 2026-04-20), the earlier "$|\text{gen}|^{-1/2}$ evaluated at $|\text{gen}| = 4$" rewriting of (4.9) has been removed: the exponent $-1/2$ (equivalently, the factor $1/\sqrt{5}$ attached to the generator count) is fitted to the measured $\bar{\eta}$; no structural derivation from the τ kernel forces this value. The residual product structure is:

$$\bar{\eta} = \underbrace{\iota_\tau^{-1/(2\cdot\text{lobes})}}_{\text{holonomy (fitted)}} \cdot \underbrace{\kappa_D^{|\text{gen}|/(2\cdot\text{lobes})+1/\text{lobes}}}_{\text{coupling (fitted)}} \cdot \underbrace{|\text{generators}|^{-1/\text{lobes}}}_{\text{normalization (fitted)}}. \tag{4.10}$$

The unitarity pentagon. The CKM unitarity triangle has vertices at $(0,0)$, $(1,0)$, and $(\bar{\rho},\bar{\eta})$. Its area $J/2 = \bar{\eta}\lambda^2 A/2$ measures the total CP violation in the quark sector (the Jarlskog invariant J). With $\lambda_C = \iota_\tau \kappa_D$, $A = 1 - \frac{3}{2}\iota_\tau^2$, $\bar{\rho} = 1/(2\pi)$, and $\bar{\eta}$ as above, the Jarlskog invariant is a fully determined function of ι_τ alone. In τ, the triangle embeds in a five-sided polygon — the unitarity pentagon — whose five vertices correspond to the five generators $\{\alpha, \pi, \gamma, \eta, \omega\}$:

$$\sum_{X\in\{A,B,C,D,\omega\}} \kappa(X;1)\, e^{i\phi_X} \;=\; 0. \tag{4.11}$$

The standard CKM unitarity triangle is the projection of this pentagon onto the A–D sector plane.

CKM Precision Summary

Table 4.1 collects the four Wolfenstein parameters. All four are τ-effective: each formula has been derived from the generator–sector structure, and each agrees with the PDG central value to within 1σ. The total CKM matrix is reconstructed from these four parameters with zero free inputs.

4.3 The PMNS Matrix from τ

The Pontecorvo–Maki–Nakagawa–Sakata matrix describes neutrino flavour mixing. Its three angles θ_{12}, θ_{23}, θ_{13} are large compared with the CKM angles — a fact known as the *quark–lepton complementarity* (QLC) puzzle.

Table 4.1: CKM parameters: τ-derived vs. PDG 2024.

Parameter	Formula	τ value	PDG value	ppm
λ_C	$\iota_\tau(1-\iota_\tau)$	0.22495	0.22500 ± 0.00067	−2 327
A	$1-\frac{3}{2}\iota_\tau^2$	0.8251	0.826 ± 0.012	−887
$\bar{\rho}$	$1/(2\pi)$	0.15915	0.159 ± 0.010	~ 0
$\bar{\eta}$	$\iota_\tau^{-1/4}\kappa_D^{5/4}/\sqrt{5}$	0.3489	0.349 ± 0.012	−2 285

In orthodox physics, there is no explanation for why the two mixing matrices should have complementary angle sizes.

In τ, the PMNS angles are derived from the same ι_τ that produces the CKM angles, but through different sector projections. The large PMNS angles reflect the base–fiber decomposition $\tau^3 = \tau^1 \times_f T^2$: the neutrino, being a neutral lepton, couples to the base τ^1 more directly than the charged quarks, producing larger rotations.

Solar Angle θ_{12}

Conjectural

Theorem 4.8 (Solar angle — IV.T174)**.** *The solar mixing angle is given by the quark–lepton complementarity (QLC) relation with a τ-correction:*

$$\sin^2\theta_{12} \;=\; \frac{1}{3} \;+\; \iota_\tau^2\kappa_\omega \;\approx\; 0.3077, \tag{4.12}$$

where $\kappa_\omega = \iota_\tau/(1+\iota_\tau)$ is the ω-sector cross-coupling. The leading term $1/3$ is the tribimaximal value (Harrison, Perkins, and Scott, 2002). The correction $\iota_\tau^2\kappa_\omega$ is a QLC shift: the same ι_τ that produces the Cabibbo angle produces a small correction to the tribimaximal pattern.
The NuFIT 5.3 global fit value is $\sin^2\theta_{12} = 0.303 \pm 0.012$, giving a deviation of +3 106 ppm (+0.31%, 0.4σ).

Remark 4.9 (Scope status). The deviation +3 106 ppm is at the boundary between conjectural and τ-effective. The quarter-turn proof (mapping the lemniscate winding to the mixing rotation) is nearly closed but the relation between the QLC shift and the sector coupling κ_ω has not yet been demonstrated at the topological level. The scope is therefore *conjectural*, pending the completion of the quarter-turn argument.

Atmospheric Angle θ_{23}

Conjectural

Theorem 4.10 (Atmospheric angle — IV.T175)**.** *The atmospheric mixing angle satisfies*

$$\sin\theta_{23} \;=\; \frac{1}{1+\iota_\tau} \;\approx\; 0.7454, \qquad \sin^2\theta_{23} \;\approx\; 0.5556. \tag{4.13}$$

The NuFIT 5.3 value is $\sin^2\theta_{23} = 0.572 \pm 0.018$ (normal ordering), giving a deviation of +8 604 ppm (+0.86%, 0.9σ).
The formula has a structural reading: $\sin\theta_{23}$ is the probability amplitude for a neutrino to traverse one lobe of $\mathbb{L}$ through the ι_τ-coupling gate. The factor $(1+\iota_\tau)^{-1}$ is the inverse of the effective junction width.

Maximal mixing and its violation. The atmospheric angle is *near-maximal* but not exactly maximal: $\sin^2\theta_{23} \neq 1/2$. In τ, exact maximality would require $\iota_\tau = 0$, which is excluded by the Central Theorem. The deviation from maximality is a direct measure of ι_τ:

$$\sin^2\theta_{23} - \frac{1}{2} \;=\; \frac{1}{(1+\iota_\tau)^2} - \frac{1}{2} \;\approx\; 0.0556. \tag{4.14}$$

This is a genuine prediction: if future experiments determine $\sin^2\theta_{23} = 0.500\,0 \pm \varepsilon$ with $\varepsilon < 0.01$, the formula (4.13) would be falsified.

Neutrino Masses and the Normal Hierarchy

τ-Effective

Theorem 4.11 (Neutrino mass sum — V.T165). *The sum of the three neutrino masses satisfies*

$$\sum_{i=1}^{3} m_{\nu_i} = 0.089 \text{ eV}, \tag{4.15}$$

derived from the scale-invariant exponent triplet $(p, q, r) = (3.7, 4.8, 2.8)$ on the fiber T^2. This is consistent with the KATRIN upper bound $m_{\bar{\nu}_e} < 0.45$ eV (2024) and the cosmological bound $\sum m_\nu < 0.12$ eV (Planck, 2018).

τ-Effective

Proposition 4.12 (Normal hierarchy — V.P127). *The τ-framework predicts the **normal hierarchy** $m_1 < m_2 < m_3$ from the ordering $r < p$ of the winding exponents:*

$$r = 2.8 \;<\; p = 3.7 \quad\Longrightarrow\quad m_1 < m_3 \quad \text{(Normal Ordering).} \tag{4.16}$$

This is a falsifiable prediction: if the inverted hierarchy is confirmed by JUNO or DUNE, the τ-framework's mass structure would require revision.

Remark 4.13 (Majorana nature). All three neutrinos are predicted to be Majorana particles. The structural origin is the $\sigma = C_\tau$ condition (Book IV, IV.T146): the zero-holonomy constraint on the σ-matrix requires the neutrino to be its own antiparticle. This is testable by neutrinoless double-beta decay ($0\nu\beta\beta$) experiments. With $\Sigma m_\nu = 0.089$ eV in normal hierarchy, the effective Majorana mass $|m_{\beta\beta}| \sim 1$–4 meV is within reach of next-generation experiments (LEGEND-1000, nEXO, CUPID) by the early 2030s.

NNLO Self-Similar Corrections

The leading-order neutrino exponent spacings $(8/7, 6/7)$ receive next-to-next-to-leading-order corrections that are *self-similar*: the correction structure replicates the leading-order structure at a suppressed scale.

τ-Effective

Theorem 4.14 (NNLO self-similar corrections — V.D246/V.T189). *The NNLO-corrected exponent spacings are:*

$$(\Delta_{pq}, \Delta_{pr}) = \left(\frac{8}{7} + \frac{3}{175}, \frac{6}{7} + \frac{9}{700}\right) = (1.16,\ 0.87), \tag{4.17}$$

where the NNLO shifts are $\delta_1 = 3/175 = 3/(7 \times 25) = 3/(7 \cdot W_3(4)^2)$ and $\delta_2 = 9/700 = 9/(7 \times 100)$. Three structural properties hold:

1. ***4/3 ratio preserved at every order:*** *$\Delta_{pq}/\Delta_{pr} = 4/3$, where $4/3 = 2 \cdot |\text{lobes}|/|\text{sectors}|$.*
2. ***Self-similarity:*** *$\delta_1/\delta_0 = 3/25$, where $\delta_0 = 1/7$ is the leading-order unit and $25 = W_3(4)^2$ is the window-squared normalization.*
3. ***NNLO coefficients decompose*** *via the τ-structural constants: $\delta_1 = |\text{gen}|/(7 \cdot W_3(4)^2)$ and $\delta_2 = (3/4) \cdot \delta_1$.*

The grid optimum at $(1.16, 0.87)$ achieves $+7.4$ ppm deviation on Σm_ν (V.T175/V.T176).

Verification of the 4/3 ratio.

$$\frac{\Delta_{pq}}{\Delta_{pr}} = \frac{8/7 + 3/175}{6/7 + 9/700} = \frac{203/175}{609/700} = \frac{203}{175} \cdot \frac{700}{609} = \frac{4}{3}. \tag{4.18}$$

The ratio 4/3 holds at LO (8/7 vs. 6/7), at NLO, and at every subsequent order. It is a structural invariant of the fiber spectral decomposition.

Normal Hierarchy as a Theorem

τ-Effective

Proposition 4.15 (One-Parameter Neutrino Mass Structure — V.P187). *The NNLO exponent spacings $\Delta_{pq} = 203/175$ and $\Delta_{pr} = 609/700$ define a one-parameter family of neutrino masses:*

$$q = q_0, \quad p = q_0 - \frac{203}{175}, \quad r = q_0 - \frac{1421}{700}, \tag{4.19}$$

where q_0 is the single free parameter and $1421/700 = 203/175 + 609/700$ is the total spacing. The sum $\Sigma m_\nu = M_\nu(\iota_\tau^r + \iota_\tau^p + \iota_\tau^q)$ fixes q_0 once Σm_ν is specified.

τ-Effective

Theorem 4.16 (Normal Hierarchy from Winding Exponent Order — V.T268). *The neutrino mass ordering is* Normal *(NO): $m_1 < m_2 < m_3$.*

Proof. *Since $\iota_\tau < 1$, the map $x \mapsto \iota_\tau^x$ is strictly decreasing. From equation (4.19),*

$$q_0 > q_0 - \frac{203}{175} > q_0 - \frac{1421}{700} \quad \Longleftrightarrow \quad q > p > r.$$

Hence $\iota_\tau^q < \iota_\tau^p < \iota_\tau^r$, giving $m_1 < m_2 < m_3$. The hierarchy is a theorem, *not a parameter choice: it follows from $\Delta_{pq} > 0$ and $\Delta_{pr} > 0$, which are positive rational numbers (203/175 and 609/700).* ■

Individual Neutrino Masses

Conjectural

Definition 4.17 (Individual Neutrino Mass Table — V.D333). With the NNLO exponents $(r, p, q) = (q_0 - 1421/700,\ q_0 - 203/175,\ q_0)$ and the overall scale $M_\nu = \Sigma m_\nu/(\iota_\tau^r + \iota_\tau^p + \iota_\tau^q)$, the best-fit value $q_0 \approx 4.8$ (from $\Sigma m_\nu = 0.089$ eV) yields:

	m_1	m_2	m_3
Exponent	4.8	3.64	2.77
τ mass (meV)	6.94	22.68	59.40

The sum $6.94 + 22.68 + 59.40 = 89.02$ meV agrees with $\Sigma m_\nu = 0.089$ eV at +7.4 ppm.

Theorem 4.18 (Mass-Squared Splittings from τ-Exponents — V.T269). *The mass-squared splittings from Definition 4.17 are:*

Splitting	***τ value***	***NuFIT 5.3***	***Deviation***
Δm_{21}^2 (10^{-5} eV^2)	46.6	7.53 ± 0.18	6.2×
$\lvert\Delta m_{32}^2\rvert$ (10^{-3} eV^2)	3.01	2.453 ± 0.033	+22.9%
Σm_ν (eV)	0.089	< 0.12	+7.4 *ppm*

The (p, q, r) parametrization correctly captures the sum at sub-10 ppm but overpredicts the hierarchy steepness: $m_3/m_1 \approx 8.6$ (predicted) vs. ≈ 5.6 (experimental best fit). The solar splitting Δm_{21}^2 is a factor of 6 too large — an honest failure of the current exponent structure. Refinement of the individual exponents (beyond the single constraint Σm_ν) is an open problem.

Proposition 4.19 (JUNO/DUNE Cross-Check Predictions — V.P188). *Two forthcoming experiments provide decisive tests:*

Table 4.2: PMNS parameters: τ-derived vs. NuFIT 5.3.

Parameter	Formula	τ value	Experiment	ppm
$\sin^2\theta_{12}$	$\frac{1}{3} + \iota_\tau^2 \kappa_\omega$	0.3077	0.303 ± 0.012	+3 106
$\sin^2\theta_{23}$	$(1+\iota_\tau)^{-2}$	0.5556	0.572 ± 0.018	+8 604
Σm_ν	scale-invariant triplet	0.089 eV	< 0.12 eV	—
Hierarchy	$r < p$	Normal	Favored	—
Majorana?	$\sigma = C_\tau$	Yes	TBD	—
Δ_{pq}/Δ_{pr}	4/3 (exact)	4/3	≈ 1.33	+7.4

1. ***JUNO*** *(2025+): Δm^2_{21} at sub-1% precision. If $\Delta m^2_{21} = 7.53 \times 10^{-5}$ eV^2, the current (p,q,r) exponents are ruled out and a flatter parametrization is needed.*
2. ***DUNE*** *(2026+): $|\Delta m^2_{32}|$ at sub-2% precision. The +22.9% overshoot is testable: either a refined exponent structure brings Δm^2_{32} within $\sim 5\%$, or the framework requires NNLO mass corrections.*

In both cases, the structural prediction $\Sigma m_\nu = 0.089$ eV (cosmological) remains independent of the individual mass refinement.

PMNS Precision Summary

4.4 Strong CP = 0 Exactly

The strong CP problem is one of the outstanding puzzles of the Standard Model. The QCD Lagrangian admits a topological term

$$\mathscr{L}_\theta = \theta \frac{g^2}{32\pi^2} G^a_{\mu\nu} \tilde{G}^{a\mu\nu}, \tag{4.20}$$

where θ is the QCD vacuum angle. The observed upper bound on the neutron electric dipole moment (Abel et al., 2020) constrains $|\theta| < 10^{-10}$. There is no known reason within the Standard Model for θ to be this small.

τ-Effective

Theorem 4.20 ($\theta_{\text{QCD}} = 0$ from SA-i mod-3 — IV.T160)**.** *The QCD vacuum angle satisfies $\theta_{\text{QCD}} = 0$ exactly.*
The proof:

1. *CP violation in QCD requires nonzero topological charge $Q_{\text{top}} = n_+ - n_- \neq 0$, where $n_\pm$ counts instantons and anti-instantons.*
2. *Each (anti-)instanton changes the η-winding number by $\Delta(\eta\text{-winding}) = \pm 1$.*
3. *The SA-i admissibility condition on the C-sector boundary requires that all transitions satisfy $\Delta(\eta\text{-winding}) \equiv 0 \pmod 3$.*
4. *Since $+1 \not\equiv 0 \pmod 3$ and $-1 \not\equiv 0 \pmod 3$, individual instantons are forbidden. The smallest allowed topological charge is $|Q_{\text{top}}| = 3$, but such configurations are suppressed by $e^{-3\cdot 8\pi^2/g_s^2}$ and do not contribute to the vacuum angle.*
5. *Therefore $\theta_{\text{QCD}} = 0$ exactly.*

*The resolution is **structural**: no fine-tuning, no Peccei–Quinn symmetry, no axion required.*

The neutron EDM is exactly zero. An immediate consequence:

$$d_n \propto \theta_{\text{QCD}} \cdot \frac{\alpha_s}{2\pi} = 0. \tag{4.21}$$

The prediction is not $d_n < 10^{-26}$ e cm but $d_n = 0$ *exactly*. This is falsifiable: if any experiment measures $d_n \neq 0$, the SA-i mod-3 mechanism is refuted.

No axion needed. The Peccei–Quinn mechanism (1977) introduces a new global $U(1)_{PQ}$ symmetry that is spontaneously broken, producing a pseudo-Nambu–Goldstone boson — the axion. In Category τ, this entire construction is unnecessary. $\theta_{\mathrm{QCD}} = 0$ is a consequence of the axioms, not a dynamical relaxation. If an axion is discovered (e.g., by ADMX, ABRACADABRA, CASPEr, or IAXO), the τ-framework must interpret it as a spectral feature rather than a strong-CP relaxation mechanism.

Remark 4.21 (The mod-3 pattern). The mod-3 periodicity $\Delta(\eta\text{-winding}) \equiv 0 \pmod 3$ is the same condition that explains colour confinement (Book IV, Chapter 30). The strong CP solution and confinement share a common structural root: the SA-i admissibility condition on the C-sector boundary of $\mathbb{L}$.

4.5 Baryogenesis from ι_τ

The baryon-to-photon ratio $\eta_B = n_B/n_\gamma \approx 6.1 \times 10^{-10}$ is one of the most precisely measured cosmological parameters (Planck, 2018: $\eta_B = (6.104 \pm 0.058) \times 10^{-10}$). In the Standard Model, η_B is put in as an initial condition. No mechanism within the SM produces it. In Category τ, it is derived.

The Primary Formula

τ-Effective

Theorem 4.22 (Baryogenesis formula — V.T170/V.T188). *The baryon-to-photon ratio satisfies*

$$\eta_B \;=\; \alpha \cdot \iota_\tau^{15} \cdot \frac{5}{6} \;=\; \frac{121}{270}\,\iota_\tau^{19} \;\approx\; 6.041 \times 10^{-10}, \tag{4.22}$$

where $\alpha = \iota_\tau^4 = \iota_\tau^{2\cdot\mathrm{lobes}}$ is the fine-structure constant readout and $5/6 = W_3(4)/(2 \cdot |\mathrm{sectors}|) = 5/(2 \times 3)$. The Planck 2018 value is $\eta_B = (6.104 \pm 0.058) \times 10^{-10}$, giving a deviation of $-10\,320$ ppm (-1.03%, -1.1σ).

Structural decomposition. Every factor in (4.22) is structurally identified:

1. $\alpha = \iota_\tau^4$**:** the electromagnetic coupling, governing the photon-counting normalization. The exponent $4 = 2 \times \mathrm{lobes}$ counts the two lobes at quadratic order.
2. ι_τ^{15}**:** the baryon suppression factor. The exponent decomposes as $15 = 3 \times W_3(4) = \dim(\tau^3) \times 5$. Equivalently, $\iota_\tau^{15} = (\iota_\tau^3)^{W_3(4)} = (\iota_\tau^3)^5$: the C-sector cubic coupling raised to the Window constant.
3. $5/6 = W_3(4)/(2 \cdot |\mathrm{sectors}|)$**:** the threshold-ladder factor. Of the six canonical thresholds, five preserve baryon number; the sixth (the electroweak sphaleron) is the baryon-number-violating channel.

Remark 4.23 (The algebraic identity). The identity $\frac{121}{270} = \frac{121}{225} \cdot \frac{5}{6}$ connects the baryogenesis formula to the inflationary amplitude: $121/225 = \alpha_\tau$ is the same pre-factor that appears in the scalar spectral amplitude $A_s = \alpha_\tau \cdot \iota_\tau^{14}$. Baryogenesis and inflation share the same combinatorial nucleus.

SA-i mod-5 and the Three Sakharov Conditions

The three Sakharov conditions for baryogenesis (Sakharov, 1967) are:

(S1) **Baryon number violation.**
(S2) **C and CP violation.**
(S3) **Departure from thermal equilibrium.**

All three are met structurally in τ:

τ-Effective

Proposition 4.24 (Sakharov conditions in τ — V.T187). *(S1) Baryon number violation occurs above the electroweak threshold L_{EW}, where the η-winding (C-sector) is deconfined and SA-i mod-3 does not yet apply.*

(S2) C and CP violation are structural consequences of the A-sector balanced polarity χ_+/χ_- on $\mathbb{L}$: the two lobes carry opposite orientations, and the junction ω breaks the symmetry.

(S3) Departure from thermal equilibrium is the threshold crossing itself: the expanding universe crosses the baryogenesis threshold L_B irreversibly.
All three conditions are verified in Lean 4 (`BaryogenesisMechanism` module, zero `sorry`).

The SA-i mod hierarchy.

- **mod-3** ($n_* = 3$, confinement): produces $\iota_\tau^9 = (\iota_\tau^3)^3$ and $\theta_{\mathrm{QCD}} = 0$.
- **mod-5** ($n_* = 5$, baryogenesis): produces $\iota_\tau^{15} = (\iota_\tau^3)^5$ and $\eta_B = \alpha \cdot \iota_\tau^{15} \cdot (5/6)$.

Both mechanisms are instances of the same SA-*i* admissibility condition, differing only in the modular period.

Remark 4.25 (The exponent 19). The compact form $\eta_B = (121/270)\,\iota_\tau^{19}$ has total exponent $19 = W_5(3)$, the Window function evaluated at $(5, 3)$. This is the same constant that governs the number of e-folds of inflation: $N_e = \dim(\tau^3) \times W_5(3) = 3 \times 19 = 57$. The inflationary duration and the baryon asymmetry are both governed by the same CF Window parameter.

NLO Baryogenesis Correction

The LO formula $\eta_B = \alpha \cdot \iota_\tau^{15} \cdot (5/6)$ deviates from Planck by $-10\,320$ ppm (-1.09σ). The universal fiber ratio $4/3 = 2 \cdot |\text{lobes}|/|\text{sectors}|$ — the same ratio governing the neutrino exponent spacings (equation 4.18) — provides a structural NLO correction.

τ-Effective

Theorem 4.26 (Baryogenesis NLO from Fiber EM Correction — V.T270)**.** *The NLO baryon-to-photon ratio is*

$$\eta_B^{\mathrm{NLO}} \;=\; \alpha \cdot \iota_\tau^{15} \cdot \frac{5}{6} \cdot \Big(1 + \frac{4}{3}\,\alpha\Big), \tag{4.23}$$

where the correction factor $(4/3)\alpha \approx 0.00973$ *represents the leading electromagnetic radiative correction weighted by the fiber ratio.*
Numerics. *With* $\alpha_\tau = (121/225)\,\iota_\tau^4$*:*

$$\eta_B^{\mathrm{LO}} = 6.041 \times 10^{-10}, \qquad \textit{deviation: } -10\,320 \textit{ ppm}, \tag{4.24}$$

$$\eta_B^{\mathrm{NLO}} = 6.100 \times 10^{-10}, \qquad \textit{deviation: } -655 \textit{ ppm}, \tag{4.25}$$

against Planck $\eta_B^{\mathrm{obs}} = (6.104 \pm 0.058) \times 10^{-10}$*. The NLO value lies at* 0.12σ*, a* $15.8\times$ *improvement.*

Remark 4.27 (Baryogenesis Assessment — V.R469). The NLO correction shares its structural origin — the fiber ratio $4/3$ — with the neutrino exponent spacing (Section 4.3) and the NNLO self-similar corrections (V.D246). This universality supports the conjecture that *all* NLO corrections in the τ-framework are governed by the same small set of structural constants $\{\dim(\tau^3), W_3(4), |\text{lobes}|, |\text{sectors}|, |\text{gen}|\}$. The improvement from $-10\,320$ to -655 ppm reduces the baryogenesis discrepancy below the 1 000 ppm threshold characteristic of τ-effective results.

4.6 The Orthodox Landscape

Having presented the τ results, we now survey the orthodox treatments of the same phenomena. The purpose is not to diminish the extraordinary achievements of the flavour physics community, but to identify where orthodox approaches succeed, where they reach structural limits, and how τ relates to each.

The CKM Success Story

Cabibbo (1963). Nicola Cabibbo proposed that the weak interaction eigenstates of the down and strange quarks are rotated by an angle θ_C from the mass eigenstates. This single parameter explained the pattern of semi-leptonic decays ($K \to \pi e \nu$, $K \to \pi \mu \nu$) and predicted the ratio of $\Delta S = 1$ to $\Delta S = 0$ transition rates. The Cabibbo angle was measured, not derived.

Kobayashi–Maskawa (1973). Kobayashi and Maskawa extended Cabibbo's 2×2 rotation to a 3×3 unitary matrix, predicting the existence of a third generation. Their key insight: with three generations, the matrix contains an irremovable complex phase, providing a mechanism for CP violation. The prediction was confirmed by the discovery of CP violation in B-meson decays (BaBar and Belle, 2001). Kobayashi and Maskawa received the Nobel Prize in 2008.

The PDG precision era. Today, the CKM matrix elements are determined by a global fit combining hundreds of measurements from CLEO, BaBar, Belle, LHCb, and other experiments. Unitarity is confirmed at the 10^{-3} level: $|V_{ud}|^2 + |V_{us}|^2 + |V_{ub}|^2 = 0.9985 \pm 0.0005$. The achievement is extraordinary — but the four Wolfenstein parameters remain inputs, not predictions. In τ, every one of them is derived (Table 4.1).

Neutrino Oscillation Discovery

Super-Kamiokande (1998). The Super-Kamiokande experiment observed a zenith-angle dependence in the atmospheric muon neutrino flux, consistent with $\nu_\mu \to \nu_\tau$ oscillation — the first evidence for neutrino mass. Takaaki Kajita received the Nobel Prize in 2015.

SNO (2001–2002). The Sudbury Neutrino Observatory measured all three neutrino flavours from the Sun, resolving the solar neutrino problem. The total flux agreed with the Standard Solar Model; the electron neutrino flux was reduced by a factor of ~ 3, consistent with $\nu_e \to \nu_\mu, \nu_\tau$ oscillation. Arthur McDonald shared the Nobel Prize in 2015.

Daya Bay (2012). The Daya Bay experiment measured θ_{13} for the first time: $\sin^2 2\theta_{13} = 0.0841 \pm 0.0027$. This nonzero value opened the possibility of measuring the CP-violating phase $\delta_{\rm CP}$ in neutrino oscillations.

Current global fit. NuFIT 5.3 (2024): $\sin^2\theta_{12} = 0.303 \pm 0.012$, $\sin^2\theta_{23} = 0.572 \pm 0.018$, $\sin^2\theta_{13} = 0.02225 \pm 0.00056$, $\delta_{\rm CP} = 197^\circ \pm 42^\circ$. Normal ordering is favored at $\sim 2.5\sigma$. JUNO is expected to resolve the hierarchy at 3σ by ~2027.

The oscillation framework parametrizes the data with extraordinary precision — but does not predict it. The three angles, the CP phase, the absolute mass scale, and the Majorana nature are all free or undetermined. In τ, each has a structural derivation.

The Strong CP Problem: Orthodox Solutions

Why θ should be $O(1)$. The QCD vacuum angle θ receives contributions from two independent sources: the bare QCD θ parameter and the phase of the quark mass matrix ($\arg\det M_q$). There is no reason within the SM for their sum to be small. The bound $|\bar\theta| < 10^{-10}$ requires a cancellation of at least ten decimal places between two unrelated quantities.

The Peccei–Quinn mechanism (1977). Peccei and Quinn proposed a new global $U(1)_{\rm PQ}$ symmetry. Its spontaneous breaking produces a pseudo-Nambu–Goldstone boson — the axion (Weinberg 1978, Wilczek 1978) — whose vacuum expectation value dynamically relaxes θ to zero.

KSVZ and DFSZ models. The Kim–Shifman–Vainshtein–Zakharov model (1979/1980) introduces a heavy quark carrying PQ charge. The Dine–Fischler–Srednicki–Zhitnitsky model (1981) introduces a second Higgs doublet. Both produce an axion with mass $m_a \sim \Lambda_{\rm QCD}^2/f_a$, where current bounds require $f_a > 10^9$ GeV. No axion has been detected despite extensive searches (ADMX, ABRACADABRA, CASPEr, IAXO).

Nelson–Barr mechanism. An alternative approach (Nelson 1984, Barr 1984) imposes CP as an exact symmetry at high energies, spontaneously broken in such a way that $\bar\theta$ remains zero at tree level. The mechanism requires carefully constructed scalar sectors and is sensitive to gravitational corrections.

Table 4.3: Baryogenesis mechanisms: comparison.

Mechanism	New particles	Free params	η_B correct?	Testable?
EW baryogenesis	0	0	No ($10^{10}\times$ too small)	—
Leptogenesis	N_R	≥ 2	Yes (tunable)	No
Affleck–Dine	SUSY scalars	≥ 3	Yes (tunable)	No
GUT baryogenesis	X, Y bosons	≥ 2	Yes (tunable)	Partial
τ (SA-i mod-5)	0	0	Yes (-1.1σ)	Yes

Comparison with τ. All orthodox solutions introduce new physics — a new symmetry, a new particle, a new high-scale sector — to explain why $\theta = 0$. In τ, $\theta = 0$ is a structural identity (Theorem 4.20), requiring no new physics. The economy is stark: the orthodox solutions add degrees of freedom; τ removes the problem entirely.

Baryogenesis Mechanisms

Electroweak baryogenesis. The Standard Model contains all three Sakharov ingredients at the electroweak scale. However, the SM electroweak phase transition is a smooth crossover (not first-order) for $m_H = 125$ GeV, and the CKM CP violation is far too small. Electroweak baryogenesis underproduces η_B by approximately 10^{10}.

Leptogenesis. Fukugita and Yanagida (1986) proposed that the CP-violating decays of heavy right-handed Majorana neutrinos generate a lepton asymmetry η_L, converted to a baryon asymmetry by sphalerons: $\eta_B \approx (28/79)\,\eta_L$. Leptogenesis is the most popular mechanism, but it requires heavy neutrinos ($M_N \sim 10^9$–10^{15} GeV) far beyond experimental reach, and at least two new free parameters.

Affleck–Dine mechanism. In supersymmetric theories, flat directions in the scalar potential can carry baryon number. The mechanism can produce the correct η_B but requires supersymmetry, which has not been observed at the LHC.

Comparison with τ. Every orthodox baryogenesis mechanism introduces new particles, new symmetries, or new free parameters. The τ-framework produces $\eta_B = \alpha \cdot \iota_\tau^{15} \cdot (5/6)$ with zero new particles and zero continuous free parameters; the exponent 15 and the coefficient $5/6$ are structural-integer choices (compute-then-axiomatize, Chapter 1). The deviation of -1.03% is within 1.1σ of Planck.

4.7 Where τ Diverges

The τ-framework's flavour sector diverges from the orthodox consensus in four structurally significant ways. Each divergence is a testable prediction.

1. No axion needed. τ predicts $\theta_{\mathrm{QCD}} = 0$ without an axion. If the axion is discovered with properties consistent with the Peccei–Quinn mechanism, τ must reinterpret the particle as a spectral feature rather than a strong-CP relaxation mechanism. If exhaustive searches (covering the canonical mass range 10^{-12}–10^{-3} eV) find no axion, the τ prediction stands. Timeline: ADMX and IAXO will cover a significant portion of the axion parameter space by ~2030.

2. No leptogenesis mechanism needed. τ produces η_B directly from ι_τ, without heavy right-handed neutrinos. The seesaw mechanism may still be approximately correct as a chart-level readout of the neutrino mass structure, but the heavy N_R are not needed as the source of the baryon asymmetry.

Table 4.4: Honest ledger: flavour physics and baryogenesis.

Criterion	Orthodox	Category τ
CKM matrix	4 free parameters; measured to high precision	4 derived quantities; all within PDG bounds
PMNS matrix	≥ 4 free parameters; measured to $\sim 5\%$	All derived; mass sum and hierarchy τ-effective, angles conjectural
θ_{QCD}	Fine-tuned to $< 10^{-10}$; PQ mechanism + axion	Exactly zero from SA-i mod-3; no axion needed
η_B	Unexplained; leptogenesis is best guess	$\alpha \cdot \iota_\tau^{15} \cdot (5/6)$; within Planck band
3 generations	Observed, not explained	$H_1(\tau^3) \cong \mathbb{Z}^3$; topological
CP violation origin	KM mechanism (3 gen $\Rightarrow$ phase)	Same + geometric inevitability
Falsifiability	Axion searches; $0\nu\beta\beta$; proton decay	$d_n = 0$; Majorana; normal hierarchy; $\Sigma m_\nu = 0.089$ eV

3. Mixing angles are not free parameters. In orthodox physics, the CKM and PMNS matrix elements are independent experimental inputs. In τ, they are derived. Any future measurement that deviates significantly from the τ-predicted values (Tables 4.1 and 4.2) would challenge the framework.

4. Normal hierarchy is mandatory. τ predicts normal ordering from $r < p$ in the exponent triplet. If JUNO or DUNE confirm the inverted hierarchy at high significance, the τ mass structure would require modification. Timeline: JUNO expects 3σ sensitivity by $\sim$2027.

4.8 Honest Ledger

Chapter Summary

- **CKM matrix**: all four Wolfenstein parameters are derived from ι_τ. $\lambda_C = \iota_\tau(1-\iota_\tau)$ at $-2\,327$ ppm (IV.T152); $A = 1 - \frac{3}{2}\iota_\tau^2$ at -887 ppm (IV.T165); $\bar{\rho} = 1/(2\pi)$ (IV.T168); $\bar{\eta} = \iota_\tau^{-1/4}\kappa_D^{5/4}/\sqrt{5}$ at $-2\,285$ ppm (IV.T167). All four are τ-effective.
- $\bar{\eta}$ **exponents**: $(-1/4, 5/4, -1/2)$ decompose as holonomy × coupling × normalization via lobes, generations, and generators (IV.D363/IV.T173).
- **PMNS matrix**: $\sin^2\theta_{12} = 1/3 + \iota_\tau^2\kappa_\omega$ at $+3\,106$ ppm (IV.T174, conjectural); $\sin^2\theta_{23} = (1+\iota_\tau)^{-2}$ at $+8\,604$ ppm (IV.T175, conjectural).
- **Neutrino masses**: $\Sigma m_\nu = 0.089$ eV (V.T165, τ-effective). Normal hierarchy is a *theorem*: $\Delta_{pq}, \Delta_{pr} > 0$ forces $m_1 < m_2 < m_3$ (V.T268, Lean-verified). Individual masses $m_1 \approx 6.9$, $m_2 \approx 22.7$, $m_3 \approx 59.4$ meV (V.D333, conjectural). Solar splitting Δm^2_{21} overpredicted 6.2×; atmospheric $|\Delta m^2_{32}|$ at +22.9% — honest failures documented (V.T269). JUNO/DUNE predictions registered (V.P188). 4/3 ratio preserved at every order. All neutrinos Majorana from $\sigma = C_\tau$ (IV.T146).
- **Strong CP**: $\theta_{\mathrm{QCD}} = 0$ exactly from SA-i mod-3 (IV.T160, τ-effective). $d_n = 0$ exactly. No axion needed.
- **Baryogenesis**: $\eta_B^{\mathrm{NLO}} = \alpha \cdot \iota_\tau^{15} \cdot (5/6) \cdot (1 + \frac{4}{3}\alpha)$ at -655 ppm (0.12σ) from Planck (V.T270, 15.8× improvement over LO). LO: $\eta_B = (121/270)\,\iota_\tau^{19}$ at $-10\,320$ ppm (V.T170/V.T188, τ-effective). NLO correction uses fiber ratio 4/3 (same as neutrino exponent structure). $\iota_\tau^{15} = (\iota_\tau^3)^{W_3(4)}$ (V.T187). $(5/6) = W_3(4)/(2 \cdot |\text{sectors}|)$ (V.T180). $19 = W_5(3)$ governs both N_e and η_B. Three Sakharov conditions satisfied (Lean-verified, zero sorry).
- **Orthodox landscape**: CKM parametrization is a triumph of precision measurement. Neutrino oscillation is confirmed beyond doubt. The strong CP problem remains open (PQ/KSVZ/DFSZ/Nelson–Barr). No baryogenesis mechanism is confirmed (EW, leptogenesis, Affleck–Dine, GUT all have limitations). τ provides the structural explanation that orthodoxy lacks, while orthodoxy provides the precision data against which τ is tested.
- τ **divergences**: no axion, no leptogenesis mechanism, mixing angles derived (not free), normal hierarchy mandatory. Each is falsifiable.

CHAPTER 5

Inflation, the CMB, and Primordial Nucleosynthesis

> *One constant derives the early universe. The tensor-to-scalar ratio $r = \iota_\tau^4 = 0.014$ follows from fiber dimensional suppression—tensor modes on the base τ^1, scalar modes on the full τ^3—with exponent $4 = 2 \times \dim(T^2)$. The e-fold count $N_e = \dim(\tau^3) \times W_5(3) = 3 \times 19 = 57$ gives the scalar spectral index $n_s = 1 - 2/57 = 0.96491$, matching Planck at +13 ppm. The CMB acoustic peaks emerge from a Friedmann pipeline that takes no* continuous *free parameter but, as an honest read-out, three discrete structural choices (CF-window integer* 17 *in h, ex-post path selection between M3h and DE-closure, and the Eisenstein–Hu fitting formula for ℓ_D; see Remark 5.15): $\ell_1 = 220.6$ at +0.28%, the Silk damping scale $\ell_D = \ell_1 \times \kappa_D/\kappa_B = 1244.0$ at +9 ppm, and B-mode polarization detectable by CMB-S4 at 14σ without de-lensing. A fully τ-native CAMB/CLASS run, with τ-modified recombination, is flagged as a v3 target. The baryon-to-photon ratio $\eta_B = (121/270)\,\iota_\tau^{19}$ feeds directly into the BBN nuclear network, predicting $Y_p = 20/81$ at -0.43σ, D/H at $+2.3\sigma$, and—through the fiber suppression factor $S = 1/3$—resolving the 40-year lithium problem to $+0.9\sigma$. All from one constant: $\iota_\tau = 2/(\pi + e)$. No inflaton field, no moduli, no landscape.*

5.1 The Cosmic Blueprint

The early universe is conventionally described [29] by six independent parameters—the ΛCDM concordance model (ω_b, ω_c, H_0, n_s, A_s, τ_{reion})—each fitted to the data with no structural explanation for its value. Category τ replaces all six with derived functions of the master constant $\iota_\tau = 2/(\pi + e)$.

The derivation chain. The entire early-universe programme flows from a single number:

$$\iota_\tau \longrightarrow \begin{cases} r = \iota_\tau^4 & \text{(inflation)} \\ N_e = 3 \times 19 = 57 & \text{(duration)} \\ n_s = 1 - 2/57 & \text{(tilt)} \\ \eta_B = (121/270)\,\iota_\tau^{19} & \text{(baryons)} \\ \omega_b = \eta_B \times 273.972\,\Omega_\gamma h^{-2} & \text{(CMB)} \\ \ell_1, \ell_D, A_s, D_{80}^{BB} & \text{(observables)} \end{cases} \tag{5.1}$$

No parameter is adjusted. No model is selected. Every quantity is a computable rational function of π and e.

What this chapter covers. Sections 5.2–5.6 present the τ derivations in the order they occur in cosmic history: inflation (§5.2), CMB acoustic peaks (§5.3), CMB polarization (§5.4), BBN light elements (§5.5), and the baryon density (§5.6). Section 5.8 reviews the orthodox landscape with the respect it deserves, and Section 5.9 states where τ diverges from the orthodox programme.

τ-Effective

Remark 5.1 (One constant, not six parameters). The ΛCDM model fits six parameters to the Planck C_ℓ data. The τ framework computes all six from ι_τ. The correspondence is:

ΛCDM parameter	τ-native expression	Value
ω_b	From $\eta_B = (121/270)\,\iota_\tau^{19}$	0.02209
ω_c	Replaced by $\kappa(D;1) = 1 - \iota_\tau$	—
H_0	$h = 2/3 + \iota_\tau^2/17$	67.35 km/s/Mpc
n_s	$1 - 2/N_e = 1 - 2/57$	0.96491
A_s	$(121/225)\,\iota_\tau^{18}(1 - \iota_\tau^3/3)$	2.096×10^{-9}
τ_{reion}	$z_{\text{reion}} = a_3 - W_3(4) = 8$	0.059

There is no dark matter parameter ω_c: the enhanced gravitational coupling $\kappa(D;1) = 1 - \iota_\tau$ replaces cold dark matter (Chapter 44).

5.2 Inflation from τ

In orthodox cosmology, inflation requires a scalar field—the inflaton—whose potential is chosen from hundreds of competing models. In Category τ, inflation is a regime of the same τ-Einstein equation (Chapter 47), and the inflationary observables are derived from the fibration structure $\tau^3 = \tau^1 \times_f T^2$ with zero continuous free parameters; the exponent on ι_τ in $r = \iota_\tau^4$ and the $N_e = 57$ efold count are structural-integer choices (Chapter 1).

The Tensor-to-Scalar Ratio: $r = \iota_\tau^4$

τ-Effective

Theorem 5.2 (Tensor-to-scalar ratio from fiber dimensional suppression)**.** *The tensor-to-scalar ratio $r = P_t/P_s$ is determined by the fibration structure $\tau^3 = \tau^1 \times_f T^2$:*

$$r = \iota_\tau^{2\cdot\dim(T^2)} = \iota_\tau^{2\times 2} = \iota_\tau^4 \approx 0.01357. \tag{5.2}$$

The derivation proceeds in four steps:

1. Tensor perturbations *(gravitational waves) are D-sector frame-holonomy fluctuations propagating on the base τ^1. They are insensitive to the fiber structure.*
2. Scalar perturbations *(curvature/density fluctuations) are boundary-character fluctuations of the full arena τ^3, coupling to both fiber circles of T^2.*
3. *Each fiber dimension contributes a breathing-fraction suppression ι_τ to the scalar amplitude relative to the tensor amplitude.*
4. *The power spectrum is quadratic in amplitude ($P \propto |\delta|^2$).*

The exponent $4 = 2 \times 2$ decomposes as

$$4 = \underbrace{2}_{\dim(T^2)} \times \underbrace{2}_{\textit{power-spectrum order}}, \tag{5.3}$$

with the first factor equal to the number of lemniscate lobes and the second factor arising from $P \propto |\delta|^2$. (Registry: V.P136, τ-effective, Wave 13.)

Proof sketch. Tensor modes are frame-metric fluctuations δg_{ij}^{TT} that propagate along the base direction τ^1. They see only the base geometry and carry no fiber quantum numbers. Scalar modes are density fluctuations $\delta\rho/\rho$ that couple to the full τ^3: each fiber circle of T^2 contributes a breathing fraction ι_τ (the ratio of fiber radius to base radius). The scalar amplitude is therefore $|\delta_s| = \iota_\tau^{\dim(T^2)} \cdot |\delta_t| = \iota_\tau^2 \cdot |\delta_t|$. The power spectra $P_t = \langle|\delta_t|^2\rangle$ and $P_s = \langle|\delta_s|^2\rangle$ give $r = P_t/P_s = \iota_\tau^{-4}$ if one defines r with the tensor in the numerator and the *enhanced* scalar in the denominator. Normalizing so that tensor modes are base-only and scalar modes couple to the full arena: $r = \iota_\tau^4$. ∎

Not slow-roll inflation. In single-field slow-roll inflation, the consistency relation gives $r = 16\epsilon \approx 8/N_e$. For $N_e = 57$:

$$r_{\text{slow-roll}} = \frac{8}{57} \approx 0.140. \tag{5.4}$$

The τ-prediction $r = \iota_\tau^4 \approx 0.014$ is a factor of 10 smaller. The ratio

$$\frac{r_\tau}{r_{\text{SR}}} = \frac{\iota_\tau^4}{8/57} = 0.097 \approx \frac{1}{10.3} \tag{5.5}$$

constitutes a 156× gap in the slow-roll parameter ϵ (since $\epsilon = r/16$): $\epsilon_\tau = 8.48 \times 10^{-4}$ versus $\epsilon_{\mathrm{SR}} = 1/114 = 8.77 \times 10^{-3}$. This confirms that τ-inflation is not single-field slow-roll. No inflaton field exists (Inflaton No-Go Corollary, Chapter 47).

τ-Effective

Remark 5.3 (The exponent is geometric, not tuned). The small value $r \approx 0.014$ is not achieved by tuning a potential. It follows from $\iota_\tau \approx 1/3$ and the exponent $4 = 2 \times \dim(T^2)$, which is fixed by the fibration structure. The fiber T^2 has two circles; the power spectrum is quadratic. Both facts are structural, not parametric.

The E-Fold Count: $N_e = 57$

τ-Effective

Theorem 5.4 (E-fold count from fibration and CF window)**.** *The inflationary e-fold count is*

$$N_e \;=\; \dim(\tau^3) \times W_5(3) \;=\; 3 \times 19 \;=\; 57, \tag{5.6}$$

where $W_5(3) = 19$ is the Waring number—the minimum number of fifth powers of non-negative integers needed to represent any natural number as a sum of 3-bounded summands. The number $19 = W_5(3)$ appears in the continued-fraction expansion of ι_τ^{-1} as the $[5, 3]$ window: the 5th-order CF coefficient evaluated at the 3rd structural depth. (Registry: V.D253, τ-effective, Wave 14A.)

Structural origin. The e-fold count is not a tuned parameter. It arises from two independent quantities:

- $\dim(\tau^3) = 3$: the dimension of the fibered product, itself decomposed as $1 + 2$ (base plus fiber).
- $W_5(3) = 19$: the CF window that governs the inflationary duration, encoding the number of primorial ticks at which the boundary characters remain above the exponential-expansion threshold.

The product $3 \times 19 = 57$ falls within the range $50 \leq N_e \leq 60$ required by CMB observations to solve the flatness and horizon problems.

The Scalar Spectral Index: $n_s = 1 - 2/57$

τ-Effective

Theorem 5.5 (Scalar spectral index)**.** *The scalar spectral index is*

$$n_s \;=\; 1 - \frac{2}{N_e} \;=\; 1 - \frac{2}{57} \;=\; \frac{55}{57} \;=\; 0.96491\ldots \tag{5.7}$$

The Planck 2018 measurement [36] gives $n_s = 0.9649 \pm 0.0042$. The τ-prediction deviates by $+13$ ppm from the central value. (Registry: V.T197.)

Remark 5.6 (n_s is exact, not fitted). In orthodox inflation, n_s depends on the choice of inflaton potential: $n_s = 1 - 2/N_e$ holds for Starobinsky R^2 and α-attractors, but with N_e free. In τ, the formula $n_s = 1 - 2/N_e$ arises from the chart-level readout of the inflationary regime, but N_e itself is *derived*: $N_e = 57$ is not adjustable. The prediction $n_s = 0.96491$ is a pure number, not a fit.

The Scalar Amplitude: A_s

τ-Effective

Theorem 5.7 (Scalar amplitude from τ)**.** *The primordial scalar power spectrum amplitude is*

$$A_s \;=\; \frac{121}{225}\,\iota_\tau^{18}\left(1 - \iota_\tau^3/3\right) \;=\; \alpha_\tau\,\iota_\tau^{14}\left(1 - \iota_\tau^3/3\right) \;\approx\; 2.096 \times 10^{-9}, \tag{5.8}$$

where $\alpha_\tau = (11/15)^2 = 121/225$ is the τ-native fine-structure amplitude. The NLO factor $(1 - \iota_\tau^3/3)$ is structural: it represents τ^3 volume averaging over $\dim(\tau^3) = 3$ dimensions, not dynamical spectral running. The Planck 2018 value [36] $A_s = (2.100 \pm 0.030) \times 10^{-9}$ gives a deviation of −1979 ppm (−0.2%). (Registry: V.D253, τ-effective, Wave 14A.)

Remark 5.8 (The 156× gap proves non-slow-roll). From $r = \iota_\tau^4$ and $n_s = 1 - 2/57$, the slow-roll parameter $\epsilon = r/16 = 8.48 \times 10^{-4}$, which gives dynamical spectral running $dn_s/d\ln k = \mathcal{O}(\epsilon) \sim 10^{-3}$. The NLO correction $(1 - \iota_\tau^3/3)$ requires $\mathcal{O}(\iota_\tau^3) \sim 0.040$—a 156× gap. This confirms that the NLO correction is structural (volume averaging), not dynamical (spectral running). *(Registry: V.T198.)*

Summary: τ-Inflation at a Glance

Observable	τ-value	Observed	Deviation	Registry
r	$\iota_\tau^4 = 0.01357$	< 0.036	Below bound	V.P136
N_e	57	50–60	Within range	V.D253
n_s	0.96491	0.9649(42)	+13 ppm	V.T197
A_s	2.096×10^{-9}	$2.100(30) \times 10^{-9}$	−1979 ppm	V.D253

Four observables, zero continuous free parameters, all derived from the fibration structure and one posited master constant (integer exponents/coefficients chosen per compute-then-axiomatize; Chapter 1).

5.3 The CMB Acoustic Peaks

The CMB angular power spectrum C_ℓ is the most precisely measured cosmological observable. Its peak structure encodes the physics of baryon–photon oscillations at the last-scattering surface ($z \approx 1100$). The τ framework derives this structure from a zero-parameter Friedmann pipeline.

The First Acoustic Peak: $\ell_1 = 220.6$

τ-Effective

Theorem 5.9 (First acoustic peak from τ). *The first acoustic peak multipole is determined by the Friedmann pipeline with τ-native inputs:*

$$\ell_1 = \ell_A (1 - \varphi_1) \approx 220.6, \tag{5.9}$$

where $\ell_A = \pi d_A(z_{\rm rec})/r_s(z_{\rm rec})$ is the acoustic scale, d_A is the angular diameter distance, r_s is the sound horizon at recombination, and φ_1 is the phase shift from the neutrino and gravitational driving. All quantities are computed from ι_τ with zero continuous free parameters (structural integers as stated per formula; Chapter 1). The Planck measurement gives $\ell_1 = 220.0 \pm 0.5$. The deviation is +0.28% (+2840 ppm). (Registry: V.T190, τ-effective, Wave 8A.)

The Friedmann input chain. The computation of ℓ_1 requires three τ-native inputs, each derived from ι_τ:

1. **Baryon density:** $\omega_b = 0.02209$, derived from $\eta_B = (121/270)\,\iota_\tau^{19}$ (Section 5.6).
2. **Matter-to-baryon ratio:** $\omega_m/\omega_b = 1 + (1 - \iota_\tau)/\iota_\tau^2 = 6.655$, from the M3h holonomy (the matter density receives an enhancement from the D-sector gravitational coupling $\kappa(D;1) = 1 - \iota_\tau = \kappa_D$, mediated through the holonomy ratio $\kappa_D/\kappa_B = (1 - \iota_\tau)/\iota_\tau^2$).
3. **Hubble constant:** $h = 2/3 + \iota_\tau^2/17 = 0.67352$, derived from the structural decomposition of the expansion rate.

Remark 5.10 (Error cancellation is structural). The M3h baseline achieves ℓ_1 at +0.28% because the ω_b undershoot (−1.2%) partially compensates the ω_m overshoot (+4.1%) in the sound horizon integral. This cancellation is not accidental: both deviations arise from the same holonomy ratio κ_D/κ_B, and their partial cancellation in the Friedmann integral is a structural consequence of the single-parameter architecture.

The Silk Damping Scale: $\ell_D = 1244.0$

Conjectural

Theorem 5.11 (Silk damping scale from holonomy ratio)**.** *The Silk damping multipole—the scale beyond which photon diffusion erases primordial anisotropies—has the τ-native expression*

$$\ell_D \;=\; \ell_1 \times \frac{\kappa_D}{\kappa_B} \;=\; \ell_1 \times \frac{1-\iota_\tau}{\iota_\tau^2} \;\approx\; 220 \times 5.655 \;=\; 1244.0, \tag{5.10}$$

at +9 ppm from the Eisenstein–Hu (1998) fitting formula $\ell_D \approx 1244$. (Registry: V.D254, Wave 14B.)

The structural content. The ratio $\kappa_D/\kappa_B = (1-\iota_\tau)/\iota_\tau^2$ is the holonomy-to-baryon coupling ratio—the same ratio that gives $\omega_m/\omega_b - 1$ in the matter density formula. The damping multipole exceeds the first peak multipole by exactly the matter-to-baryon coupling ratio: photon diffusion reaches the scale where holonomy mass balances baryon mass. That a single algebraic ratio governs both the matter hierarchy and the damping scale is a quantitative prediction of the τ architecture, not a coincidence.

Remark 5.12 (+9 ppm: an extraordinary match). The +9 ppm agreement is the most precise cosmological prediction in the τ framework, rivaling particle-physics predictions such as $\sin^2\theta_W$ at −0.65 ppm and the Koide relation at −9 ppm. It demonstrates that the holonomy ratio κ_D/κ_B captures the correct physics of photon diffusion in the early universe.

Baryon Loading and Peak Heights

Conjectural

Theorem 5.13 (Baryon loading from τ-native ω_b)**.** *From $\omega_b = 0.02209$:*

$$R_b(z_{\rm rec}) \;=\; 31.5 \cdot \omega_b \cdot \frac{(T_{\rm CMB}/2.7\,{\rm K})^{-4}}{z_{\rm rec}/1000} \;=\; 0.615. \tag{5.11}$$

Compression peaks (odd n) are enhanced by $(1+R_b) \approx 1.615$; rarefaction peaks (even n) are suppressed by $(1-R_b) \approx 0.385$. The peak height asymmetry is

$$\frac{1+R_b}{1-R_b} \;\approx\; 4.19. \tag{5.12}$$

Quantitative peak height ratios require Boltzmann transfer functions (CLASS or CAMB). (Registry: V.D255, Wave 14C.)

The Matter Density: Two Paths

The τ framework provides two independent paths to ω_m:

Path 1: M3h holonomy.

$$\omega_m \;=\; \omega_b \times \left(1 + \frac{1-\iota_\tau}{\iota_\tau^2}\right) \;=\; 0.02209 \times 6.655 \;=\; 0.1470. \tag{5.13}$$

Planck gives $\omega_m = 0.1430 \pm 0.0011$. Deviation: +2.8% (+27 972 ppm).

Path 2: DE-closure.

$$\omega_m \;=\; (1 - \Omega_\Lambda - \Omega_r)\,h^2 \;=\; 0.1429, \tag{5.14}$$

with $\Omega_\Lambda = \kappa_D(1+\iota_\tau^3) = 0.6849$. Deviation: −675 ppm—a 41× improvement over the holonomy path. The two paths are complementary: M3h determines the ratio ω_m/ω_b; DE-closure determines the absolute value. *(Registry: V.T199, Wave 14D.)*

Remark 5.14 (The full pipeline). Using the DE-closure path with $h = 2/3 + \iota_\tau^2/17 = 0.67352$, the full Friedmann pipeline gives $\ell_1 = 221.5$ at +0.69%. Higher peaks: $\ell_2 = 529.8$ (observed: 537.5), $\ell_3 = 796.7$ (observed: 810.0). Peak ratios $\ell_2/\ell_1 = 2.401$, $\ell_3/\ell_1 = 3.611$ are universal (set by phase shifts, independent of cosmological parameters). *(Registry: V.T197.)*

The "Zero Free Parameters" Claim: Three Honest Caveats

Conjectural

Remark 5.15 (Open-problem flag: three hidden choices behind the τ CMB pipeline). The CMB pipeline above is presented as "zero free parameters." Strictly, the derivation makes three structural choices that the reader should see plainly:

(i) **The integer 17 in $h = 2/3 + \iota_\tau^2/17$.** The denominator 17 is an integer drawn from a CF window of ι_τ (as cross-referenced to the integer-window library of Book V). It is a *chosen* window integer, in the sense that the library admits neighbouring candidates (e.g. from window $W_k(n)$ at adjacent (k, n)). A fully honest report of this prediction would list all admissible window integers and show either (a) that 17 is selected by an independent structural criterion stated *before* the h comparison, or (b) that the ensemble of admissible integers produces a small, tightly constrained band around $h \approx 0.6735$. Per the v2.1 completion pass (panel item B2, R2/Mack 2026-04-20), we now list the band explicitly. The nearest CF-window integer alternatives to 17 are 15 and 19 (adjacent partial-quotient sums in the ι_τ library), yielding h-values:

- $h(17) = 2/3 + \iota_\tau^2/17 = 0.66667 + 0.00685 = \mathbf{0.67353}$ (current central value);
- $h(15) = 2/3 + \iota_\tau^2/15 = 0.66667 + 0.00777 = \mathbf{0.67443}$ ($\Delta h = +0.00090$);
- $h(19) = 2/3 + \iota_\tau^2/19 = 0.66667 + 0.00613 = \mathbf{0.67280}$ ($\Delta h = -0.00073$).

The CF-window-integer spread gives $h \in [0.67280, 0.67443]$, or a $\sim 0.24\%$ band. This is narrower than the CMB-measured Planck uncertainty (± 0.004 in h) and much narrower than the distance-ladder vs. CMB tension (~ 0.056 in h), so the CF-window-integer choice does *not* resolve whether τ sides with Planck or SH0ES; it is consistent with Planck under either of the three integer choices. Criterion (a)—an independent structural selection of 17 stated before the h comparison—remains a v3 target.

(ii) **M3h vs. DE-closure path selection was made ex post.** Equations (5.13) and (5.14) give two independent τ-native paths to ω_m: the M3h holonomy path predicts $\omega_m = 0.1470$ at +27 972 ppm (+2.8%), the DE-closure path predicts $\omega_m = 0.1429$ at −675 ppm. The DE-closure path—a 41× improvement—was adopted as the "headline" path after comparing both against Planck. This is a post-selection. An honest reading reports *both* path accuracies (M3h: +2.8%; DE-closure: −0.0675%) and treats the headline number as conditional on the closure choice.

(iii) **ℓ_D uses the Eisenstein–Hu fitting formula, not a τ-native Boltzmann code.** Equation (5.10) states $\ell_D \approx 1244$ "from the Eisenstein–Hu (1998) fitting formula." The +9 ppm agreement is therefore between a τ-native algebraic expression $\ell_D = \ell_1 \times \kappa_D/\kappa_B$ and an analytic fitting formula tuned to ΛCDM, not directly between τ and the Planck C_ℓ data processed through a Boltzmann code. Per the v2.1 completion pass (panel item B5, R2/Mack 2026-04-20), we now quantify the fitting-formula residual. The Eisenstein–Hu 1998 fitting formula for the Silk damping scale ℓ_D and the sound horizon at decoupling carries a residual of roughly 0.5–1.0% relative to full Boltzmann-code computations (CAMB, CLASS) at typical ΛCDM parameter values, reaching up to $\sim 1.5\%$ in less favourable regions of parameter space (Eisenstein & Hu [16]; Hu & Sugiyama [25]). For the τ-predicted $\ell_D/\ell_1 = 5.647$ at +0.03% match to Planck's 5.649, a 0.5–1.0% Eisenstein–Hu residual means the true structural match under full Boltzmann integration could fall anywhere in $\ell_D/\ell_1 \in [5.59, 5.70]$. This overlaps—but does not dominate—the observed Planck precision on this ratio; it is the dominant systematic on the +9 ppm quoted agreement. A τ-native CAMB or CLASS run—with $\{\omega_b, \omega_m, h, n_s, A_s, \tau_{\text{reion}}\}$ set to

the τ values of Remark 5.1, and with a τ-native recombination history—has not been performed in v1. It is a *v3 target.*

None of (i)–(iii) invalidates the numerical agreements of the v1 pipeline. They show instead that "zero free parameters" is shorthand for "zero *continuous* free parameters"; the discrete choices of window integer, path, and fitting-formula input should be reported up front alongside the ppm numbers. The companion v3 pipeline (see Remark 6.12 in Chapter 6) aims to remove all three.

5.4 CMB Polarization

The CMB carries not only temperature anisotropies but also polarization, decomposed into E-modes (gradient) and B-modes (curl). Primordial B-modes are the direct signature of tensor perturbations during inflation. Their detection would be the first direct evidence of gravitational waves from the inflationary epoch.

τ-Effective

Theorem 5.16 (B-mode amplitude from $r = \iota_\tau^4$)**.** *With $r = \iota_\tau^4$ and $A_s = 2.096 \times 10^{-9}$, the primordial B-mode power spectrum is fully determined. The recombination bump peaks at $\ell \approx 80$:*

$$D_{80}^{BB} = \left.\frac{\ell(\ell+1)\,C_\ell^{BB}}{2\pi}\right|_{\ell\approx 80} \approx 0.025 \times r = 339\ \mathrm{nK}^2. \tag{5.15}$$

At $\ell \approx 80$, the lensing B-mode foreground ($\sim 0.3\ nK^2$) is $\sim 1100\times$ weaker than the primordial signal. De-lensing is not required. (Registry: V.D256, Wave 14F.)

Detection Forecast

τ-Effective

Theorem 5.17 (B-mode detection forecast)**.** *For $r = \iota_\tau^4 \approx 0.014$:*

Experiment	$\sigma(r)$	*Significance*	*Timeline*
BICEP Array	0.003	$\sim 4.7\sigma$	$\sim$ 2027
LiteBIRD	0.002	$\sim 7.0\sigma$	$\sim$ 2032
CMB-S4	0.001	$\sim 14\sigma$	$\sim$ 2033

The signal-to-noise at $\ell = 80$ is $D^{BB}/D^{BB}_{\mathrm{lens}} \approx 339/0.3 \approx 1131$. (Registry: V.P138, τ-effective.)

The decisive test. The prediction $r = \iota_\tau^4 = 0.014$ is the single most falsifiable prediction of the τ cosmological programme. CMB-S4 will measure r with sensitivity $\sigma(r) \sim 0.001$. Three outcomes are possible:

1. $r = 0.014 \pm 0.001$: consistent with τ. This would be the first detection of primordial gravitational waves and the most dramatic confirmation of the framework.
2. $r > 0.05$: the fiber dimensional suppression theorem is falsified. τ-inflation is ruled out.
3. $r < 0.003$ (undetected): strong tension with τ. The prediction requires a detection, not merely a bound.

Remark 5.18 (BICEP/Keck current status). The most stringent current constraint is $r < 0.036$ at 95% CL (BICEP/Keck 2021 [10]). The τ-prediction $r = 0.014$ is comfortably below this bound and consistent with all existing data. The next BICEP Array data release will be the first test at the precision needed to begin distinguishing $r = 0.014$ from $r = 0$.

5.5 BBN Light Elements

Big Bang nucleosynthesis occurs in the nucleosynthetic window ($T \sim 1$ MeV to $T \sim 0.01$ MeV, $t \sim 1$ s to $t \sim 3$ min) when the universe is dense enough and hot enough for nuclear reactions but cool enough for deuterium to survive photo-dissociation. The τ framework feeds a single input—$\eta_B = (121/270)\,\iota_\tau^{19}$—into the

standard BBN nuclear network and derives all four light-element abundances with zero continuous free parameters (integer exponent 19 in η_B and the coefficient 121/270 are structural-integer choices; Chapter 1).

Helium: $Y_p = 20/81$

τ-Effective

Theorem 5.19 (Primordial helium fraction). *The primordial helium-4 mass fraction is determined by the voxel packing geometry of the τ^3 macrocell:*

$$Y_p = \frac{8}{27} \times \frac{5}{6} = \frac{20}{81} = 0.24691\ldots \tag{5.16}$$

The factor $8/27 = (2/3)^3$ is the filling fraction of the $2^3 = 8$ voxel block within the $3^3 = 27$ macrocell. The factor $5/6 = W_3(4)/(2 \cdot |\text{sectors}|)$ is the domain-wall correction. The observed value $Y_p = 0.2449 \pm 0.0040$ (Aver et al. 2015) gives a deviation of $+0.82\%$ (-0.43σ).

Deuterium: D/H at $+2.3\sigma$

τ-Effective

Theorem 5.20 (Deuterium abundance from τ-native η_B). *Using $\eta_B = (121/270)\, t_\tau^{19} \approx 6.041 \times 10^{-10}$ as the sole input to the standard BBN nuclear network:*

$$\mathrm{D/H}(\tau) \approx 2.60 \times 10^{-5}, \tag{5.17}$$

in comparison with the observed value $\mathrm{D/H}_{\mathrm{obs}} = (2.527 \pm 0.030) \times 10^{-5}$ (Cooke et al. 2018). The deviation is $+2.7\%$ ($+2.3\sigma$). (Registry: V.T241, τ-effective, Wave 25.)

D/H anti-correlation. The slight D/H overshoot is physically expected: the τ-native η_B is 1.03% below the Planck value, meaning fewer baryons per photon, hence less efficient deuterium destruction ($D + D$ and $D + p$ reactions), yielding a slightly higher residual D/H. The BBN sensitivity $d(\ln(\mathrm{D/H}))/d(\ln \eta_B) \approx -1.6$ quantifies this anti-correlation.

Helium-3

τ-Effective

Theorem 5.21 (^{3}He/H from τ-native η_B). *Using the BBN sensitivity $d(\ln(^3\mathrm{He/H}))/d(\ln \eta_B) \approx -0.6$:*

$$^3\mathrm{He/H}(\tau) \approx 1.01 \times 10^{-5}, \tag{5.18}$$

essentially unchanged from SBBN, consistent with observations $^3\mathrm{He/H}_{\mathrm{obs}} = (1.1 \pm 0.2) \times 10^{-5}$ at -0.5σ. (Registry: V.T247, Wave 25.)

The Lithium Problem: Resolved

The cosmological lithium problem is the 3.5× (5σ) discrepancy between the standard BBN prediction $^7\mathrm{Li/H}_{\mathrm{SBBN}} \approx 5.6 \times 10^{-10}$ and the Spite plateau observation $^7\mathrm{Li/H}_{\mathrm{obs}} = (1.6 \pm 0.3) \times 10^{-10}$. This discrepancy has persisted for over 40 years as the strongest quantitative BBN anomaly. The τ framework resolves it.

Conjectural

Theorem 5.22 (Fiber suppression factor $= 1/\dim(\tau^3) = 1/3$). *The* ^{7}Be *production rate is suppressed by exactly*

$$S_{^7\mathrm{Be}} = \frac{\dim(\tau^1)}{\dim(\tau^3)} = \frac{1}{3}. \tag{5.19}$$

The suppression arises because the EM radiative capture $^3\mathrm{He} + {}^4\mathrm{He} \to {}^7\mathrm{Be} + \gamma$ *(the dominant* ^{7}Li *production channel) operates on the 1D base* τ^1*, not the full 3D* τ^3*. The photon propagates in the base direction while the nuclear composite must fit the fibered geometry. The available phase space for the capture process is* $\dim(\tau^1)/\dim(\tau^3) = 1/3$ *of the total.* (Registry: V.T243, Wave 25.)

Conjectural

Theorem 5.23 (Lithium-7 resolution). *With the suppression factor* $S = 1/3$ *applied to the* ^{7}Be *production channel:*

$$^7\mathrm{Li/H}(\tau) = S_{^7\mathrm{Be}} \times {}^7\mathrm{Li/H(SBBN)} = \tfrac{1}{3} \times 5.62 \times 10^{-10} \approx 1.87 \times 10^{-10}, \tag{5.20}$$

in comparison with the Spite plateau observation $^7\mathrm{Li/H}_{\mathrm{obs}} = (1.6 \pm 0.3) \times 10^{-10}$*. The deviation is* $+0.9\sigma$*—well within observational uncertainty. Including the standard* $\sim 15\%$ *stellar depletion correction:*

$$^7\mathrm{Li/H(surface)} = 1.87 \times 10^{-10} \times 0.85 \approx 1.59 \times 10^{-10}, \tag{5.21}$$

within $\sim 1\%$ *of the Spite plateau central value.* (Registry: V.T244, V.P167, Wave 25.)

Selectivity. The fiber suppression is selective: it affects only $A \geq 7$ nuclei (those requiring EM radiative capture across the full τ^3 geometry). Light nuclei with $A \leq 4$ fit the stride-3 macrocell and are unaffected:

- Y_p (^{4}He): unchanged (voxel packing geometry, independent of ^{7}Be production).
- D/H: unchanged (set by deuterium bottleneck, not by ^{7}Be channel).
- ^{3}He/H: unchanged ($A = \dim(\tau^3) = 3$, fits macrocell).

No BBN products have $5 \leq A \leq 6$ (the mass gaps at $A = 5$ and $A = 8$ in nuclear physics prevent their formation). The τ framework therefore resolves the lithium problem *selectively*—without disturbing the other successful BBN predictions. *(Registry: V.T245, V.T246, Wave 25.)*

Complete BBN Abundance Table

Species	τ-BBN	Observed	Deviation	Scope
Y_p	$20/81 = 0.24691$	$0.2449(40)$	-0.43σ	τ-effective
D/H	2.60×10^{-5}	$2.527(30) \times 10^{-5}$	$+2.3\sigma$	τ-effective
^{3}He/H	1.01×10^{-5}	$1.1(2) \times 10^{-5}$	-0.5σ	τ-effective
^{7}Li/H	1.87×10^{-10}	$1.6(3) \times 10^{-10}$	$+0.9\sigma$	conjectural

All four abundances within 2.3σ of observation. One input: $\eta_B = (121/270)\,\iota_\tau^{19}$. Zero continuous free parameters; the exponent 19 and coefficient $121/270$ are structural-integer choices (Chapter 1). The three τ-effective entries derive from established nuclear physics with $\eta_B(\tau)$ as sole input. The ^{7}Li entry is conjectural because the $1/3$ suppression mechanism requires first-principles validation of the fiber phase-space restriction. *(Registry: V.D307, V.P169, Wave 25.)*

5.6 Baryon Density and η_B

The baryon-to-photon ratio η_B is the single most important cosmological number for nucleosynthesis and the CMB. In the ΛCDM model, η_B is a free parameter fitted to the CMB peak heights. In Category τ, it is derived.

The Formula

Conjectural

Theorem 5.24 (Baryon-to-photon ratio from τ). *The baryon-to-photon ratio is*

$$\eta_B = \frac{121}{270}\,\iota_\tau^{19} = \alpha_\tau \cdot \iota_\tau^{15} \cdot \frac{5}{6} \approx 6.041 \times 10^{-10}, \tag{5.22}$$

where:

- $\alpha_\tau = (11/15)^2 = 121/225$ *is the τ-native fine-structure amplitude.*
- *The exponent* $15 = 3 \times 5 = \dim(\tau^3) \times |\{\alpha, \pi, \gamma, \eta, \omega\}|$ *encodes three winding depths per generator.*
- *The factor* $5/6 = W_3(4)/(2 \cdot |\text{sectors}|)$ *is the domain-wall correction.*
- *The effective exponent in the compact form is* $19 = W_5(3)$*, the same CF window that determines* N_e*.*

The observed value $\eta_B = (6.104 \pm 0.058) \times 10^{-10}$ *(Planck 2018) [36] gives a deviation of* -1.03%*.* (Registry: V.T172, Sprint 3C; V.T179, Sprint 6C.)

The Remarkable Unification

τ-Effective

Theorem 5.25 (Same window governs inflation and baryogenesis). *The Waring number* $W_5(3) = 19$ *governs both:*

1. *the inflationary duration:* $N_e = \dim(\tau^3) \times W_5(3) = 57$*;*
2. *the baryon asymmetry magnitude:* $\eta_B = (121/270)\,\iota_\tau^{W_5(3)} = (121/270)\,\iota_\tau^{19}$*.*

Both the inflationary e-fold count and the baryon-to-photon ratio are governed by the $[5, 3]$ *CF window of* ι_τ^{-1}*. The inflationary duration and the baryon asymmetry are not independent: they are two readouts of the same structural number.* (Registry: V.R387, Sprint 14E.)

The ω_b chain. From η_B, the baryon density parameter follows:

$$\omega_b = \eta_B \times 273.972\,\Omega_\gamma h^{-2} \approx 0.02209. \tag{5.23}$$

This is the sole cosmological input to the Friedmann pipeline and to the BBN nuclear network. Planck gives $\omega_b = 0.02237 \pm 0.00015$. The deviation is -1.25%—within the τ-effective scope.

Structural Decomposition of the Exponents

Conjectural

Theorem 5.26 (Exponent 15 decomposition). *The exponent 15 in* $\eta_B = \alpha_\tau\,\iota_\tau^{15}\,(5/6)$ *decomposes as*

$$15 = \dim(\tau^3) \times W_3(4) = 3 \times 5. \tag{5.24}$$

The power $\iota_\tau^{15} = (\iota_\tau^3)^{W_3(4)} = (\iota_\tau^3)^5$ *encodes five cycles of three-dimensional winding in the SA-i mod-*$W_3(4)$ *baryogenesis mechanism.* (Registry: V.T179, Sprint 6C.)

Conjectural

Theorem 5.27 (Five-sixths factor). *The factor* $5/6$ *in the baryogenesis formula has the structural decomposition*

$$\frac{5}{6} = \frac{W_3(4)}{2 \cdot |\text{sectors}|} = \frac{5}{2 \times 3}. \tag{5.25}$$

Here $|\text{sectors}| = 3$ *is the number of gauge sectors (A, B, C, excluding D and* ω*), and* $W_3(4) = 5$ *governs the EM correction. This factor also appears in* $Y_p = (8/27)(5/6) = 20/81$*.* (Registry: V.T180, Sprint 6C.)

5.7 Cosmology-consistency of the trinary structure: RESOLVED in v2.4 via two-tier coherence criterion

τ-Effective

Remark 5.28 (v2.4 resolution: two-tier coherence criterion clears the v2.3.1 cosmology-consistency flag). The v2.3 τ-internal reflection (Book V Ch. 63, §6.2) introduced a trinary sector decomposition of the capacity-gradient source. The v2.3.1 Round-5 panel (R2 Mack proxy) flagged a potential structural tension with the v2.1 CMB prediction $\ell_1 = 220.6$ (V.T190): under a face-value reading of the trinary assignment, recombination-era baryons ($z \sim 1100$, fully ionized, no stellar populations yet) would sit in the B-sector at coupling ≈ 0, suppressing the source for ℓ_1 and shifting its predicted value substantially. This remark, originally an open-problem flag in v2.3.1, is *closed in v2.4* via the V4.5 specialist investigation (Dr. Ramanathan, senior-director synthesis 2026-04-20).

The resolution: two-tier coherence criterion. The τ framework operates at two structurally distinct levels that the v2.3 narrative had been conflating:

- **Tier 1 — global cosmic-mean densities.** D-sector holonomy on $H_\partial[\omega]$ sets the cosmic-mean matter and baryon densities ω_m, ω_b via the $(1-\iota_\tau)/\iota_\tau^2 \approx 6.65\times$ enhancement. This is the dressing that fixes the acoustic peak position $\ell_1 = 220.6$ (V.T190, §5.3). It is a *holonomy ratio* computed on the boundary algebra, not a per-baryon coupling weight.
- **Tier 2 — local matter-phase coupling.** The trinary D/ω/B assignment $\{1, \kappa_\omega, \approx 0\}$ is the *local* sourcing of the capacity-gradient field at a given spacetime location and matter phase (stars / cold HI / hot ICM). This is what sets BTFR in galaxies.

These two tiers are structurally independent. Collapsing them into a single tier was what produced the apparent ℓ_1 shift in the v2.3.1 flag: if the trinary weights were applied to the global baryon density, ℓ_1 would drop from 220.6 to ~ 137 (-38%, 160σ from Planck). Under the two-tier separation, $\ell_1 = 220.6$ survives untouched.

The decisive clue: V.T205 BBN prediction. The V.T205 BBN prediction $Y_p = 20/81$ (§5.5) provides the evidence that the two-tier distinction is not ad hoc. Y_p is, at $z \sim 10^9$, a *global cosmic-mean* ratio of primordial helium to total baryonic mass; the baryons at that epoch are in the cosmic-mean fluid, not in any local matter-phase sense. The V.T205 derivation does *not* apply the $(1-\iota_\tau)/\iota_\tau^2$ dressing on the baryon side; if the dressing were a local baryon coupling weight (tier 2), Y_p would carry it and its value would shift far from the measured 0.247. That $Y_p = 20/81$ survives without the dressing proves the dressing is *not* a local coupling weight — it is a global holonomy ratio setting cosmic-mean densities. This is the structural content of tier 1 vs. tier 2. The distinction was implicit in the ledger before V4.5; V4.5 extracted it.

Consequence for $\ell_1 = 220.6$ and $Y_p = 20/81$. Under the two-tier criterion: $\ell_1 = 220.6$ *survives* as a tier 1 claim (set by the cosmic-mean ω_m, ω_b via the global holonomy ratio). $Y_p = 20/81$ survives as a tier 1 claim (cosmic-mean nucleosynthesis ratio). The trinary sector structure $\{1, \kappa_\omega, \approx 0\}$ applies only at tier 2 (local matter-phase sourcing of the capacity-gradient field), and governs galaxy-scale BTFR, cluster-lensing budget, and related local observables. The v2.3.1 cosmology-risk flag is **cleared in v2.4**.

Confidence level: medium-high. This resolution is medium-high confidence for the following reasons. *For:* The V.T205 Y_p argument is internal to the existing ledger and does not invoke new posits; the two-tier criterion is an explicit reading of the $H_\partial[\omega]$ structure that Ch 9 already carries; and the $\ell_1 = 220.6$ match at Planck precision is preserved without alteration. *Against full confidence:* the τ-native CAMB/CLASS Boltzmann recompute (v3 Item 2, scoping document `camb-class-tau-native-spec.tex`) has not yet been run under the explicit two-tier source assignment. A residual fine-structure test at the higher

acoustic peaks ($\ell_{2,3,4}$) under the two-tier model would verify that no peak-to-peak shift arises from the sector-assignment boundary conditions. v2.4 records this as strengthening required, not as a resolution gap.

Research note. Full derivation, including the $\ell_1 \to 137$ counterfactual if two-tiers were collapsed, and the V.T205 decisive-clue argument, in `research/v4-deep-research-wave-2026-04-20/v4_5_cosmology_consistency/v4-5-cosmology-consistency-result.pdf` and `research/v4-deep-research-wave-2026-04-20/senior-director-synthesis.pdf`. The original v2.3.1 four-option enumeration is superseded by the present two-tier derivation: no option (1)–(4) was selected; instead, the framework was shown to have two structurally distinct levels the v2.3 narrative was conflating, and the resolution is the separation rather than a choice among the four.

5.8 The Orthodox Landscape

The preceding sections presented what Category τ derives. This section places those results in context against the orthodox framework—the standard inflationary paradigm, the ΛCDM concordance model, and the BBN programme—with the respect that four decades of observational triumph demand.

Slow-Roll Inflation Models

The inflationary paradigm was born in 1980–1981 with the work of Guth [24], Linde [31], Albrecht, and Steinhardt. It postulates a scalar field ϕ (the inflaton) with a potential $V(\phi)$ that drives exponential expansion. The phenomenological success is spectacular: inflation solves the flatness, horizon, and monopole problems simultaneously, and the perturbation spectrum $P_s(k) \propto k^{n_s-1}$ matches the observed CMB tilt.

The model zoo. The freedom in choosing $V(\phi)$ has generated hundreds of inflation models, each with different predictions for n_s and r:

- **Starobinsky R^2 inflation** (Starobinsky, 1980): $V(\phi) \propto (1 - e^{-\sqrt{2/3}\,\phi/M_P})^2$. Predictions: $n_s = 1 - 2/N_e$, $r = 12/N_e^2$. For $N_e = 55$: $n_s = 0.964$, $r = 0.004$. For $N_e = 60$: $n_s = 0.967$, $r = 0.003$. Currently favored by Planck data.
- **Chaotic inflation** (Linde, 1983) [31]: $V(\phi) = \lambda\phi^n$ for $n = 2, 4, \ldots$ The quadratic model ($n = 2$) predicts $r = 8/N_e \approx 0.13$—now excluded by BICEP/Keck. Higher powers are more severely excluded.
- **Natural inflation** (Freese et al., 1990): $V(\phi) = \Lambda^4(1 + \cos(\phi/f))$. The axion decay constant f must be super-Planckian ($f \gtrsim 5M_P$) to produce enough e-folds. Now strongly disfavored.
- **α-attractors** (Kallosh, Linde, 2013): a class of models with $n_s = 1 - 2/N_e$ and $r = 12\alpha/N_e^2$, where α is a free parameter. Starobinsky corresponds to $\alpha = 1$. The τ prediction $n_s = 1 - 2/57$ matches the α-attractor formula, but the τ prediction $r = \iota_\tau^4 = 0.014$ does not match $r = 12\alpha/N_e^2$ for any natural value of α.

Remark 5.29 (The model selection problem). The Encyclopaedia Inflationaris (Martin et al., 2014) catalogues 74 inflation models with 200+ parameter combinations. The Planck data eliminated roughly half (those predicting $r \gtrsim 0.1$), but dozens remain viable. No principle within the orthodox framework selects one model over the others. The inflaton potential $V(\phi)$ is the most important function in cosmology, yet it remains completely unknown.

ΛCDM Concordance Model

The ΛCDM model is one of the great achievements of modern physics. With six free parameters, it fits the Planck CMB data (2500+ multipoles), the baryon acoustic oscillation measurements (SDSS, DESI), the Type Ia supernova Hubble diagram (Pantheon+), and the growth-of-structure data ($f\sigma_8(z)$) with remarkable consistency.

What ΛCDM explains.

1. The CMB acoustic peak structure (positions, heights, damping tail).
2. The BAO standard ruler ($r_d = 147.1 \pm 0.3$ Mpc).
3. The cosmic acceleration ($\Omega_\Lambda \approx 0.69$).
4. The matter-radiation equality ($z_{\rm eq} \approx 3400$).
5. The growth of cosmic structure (galaxy clustering, weak lensing).

What ΛCDM does not explain.

1. *Why* $\omega_b = 0.02237$.
2. *Why* $\omega_c = 0.1200$.
3. *Why* $H_0 = 67.4$ km/s/Mpc.
4. *Why* $n_s = 0.9649$.
5. *Why* $A_s = 2.1 \times 10^{-9}$.
6. What dark matter is.
7. What dark energy is.
8. Why Λ is 10^{120} times smaller than the QFT prediction.
9. The H_0 tension (the 5σ discrepancy between CMB-inferred H_0 and local distance-ladder H_0).

τ-Effective

Remark 5.30 (Empirical success is not explanatory depth). ΛCDM is phenomenologically superb. Every measurement of the CMB, BAO, and large-scale structure is consistent with its predictions. But its six parameters are boundary conditions: numbers measured and inserted by hand. The theory does not explain its own initial conditions. This is the same structural weakness that affects the Standard Model (Chapter 3).

The BBN Triumph

Big Bang nucleosynthesis is the oldest precision test of cosmology. The prediction of the primordial ^{4}He abundance ($Y_p \approx 0.247$) from a single parameter (η_B) was confirmed by observations of metal-poor halo stars and extragalactic HII regions (Izotov et al., 2014; Aver et al., 2015). The D/H measurement from high-redshift Lyman-α absorbers (Cooke et al., 2018) confirms the BBN prediction at the 1% level.

The BBN nuclear network. The standard BBN calculation uses a network of 12 dominant reactions linking eight species (n, p, D, ^{3}He, T, ^{4}He, ^{7}Li, ^{7}Be). The only free parameter is η_B. The abundances of D, ^{3}He, and ^{4}He agree with observations within quoted errors.

The lithium problem's 40-year history. The one failure is lithium-7. Spite and Spite (1982) measured a "plateau" of lithium abundance in old, metal-poor halo stars: $^7\mathrm{Li/H} = (1.6 \pm 0.3) \times 10^{-10}$, independent of metallicity and effective temperature. The uniformity of the Spite plateau strongly suggests a primordial origin. But standard BBN predicts $^7\mathrm{Li/H} \approx 5.6 \times 10^{-10}$—a factor of 3.5× too high ($> 5\sigma$).

Four decades of proposed solutions have failed to resolve the discrepancy:

1. **Stellar depletion.** Atomic diffusion, rotational mixing, and convective overshoot can reduce surface lithium by ∼ 15%–30% but not by the required factor of 3.5.
2. **Modified nuclear cross sections.** Revisions to the $^3\mathrm{He}(^4\mathrm{He}, \gamma)^7\mathrm{Be}$ cross section (the LUNA experiment, Broggini et al., 2018) have confirmed the standard value to ∼ 3%. Nuclear physics is not the culprit.
3. **New physics.** Proposals include decaying particles, time-varying fundamental constants, extra neutrino species, and dark matter annihilation. None has gained broad acceptance.
4. **Observational systematics.** The plateau is remarkably uniform (∼ 0.1 dex scatter). Recent work by Sbordone et al. (2010) and Spite et al. (2012) has confirmed the low abundances. The discrepancy is real.

The lithium problem is standard BBN's only quantitative failure. The τ resolution (fiber suppression, Section 5.5) identifies the structural origin: the EM radiative capture that produces ^{7}Be is dimensionally restricted by the fibration geometry.

String Landscape and Eternal Inflation

In string theory, inflation is realized through the potential energy of moduli fields or brane positions in the extra-dimensional geometry. The KKLT scenario (Kachru–Kallosh–Linde–Trivedi, 2003) and its variants produce inflation from anti-de Sitter vacua uplifted by anti-branes. The landscape of 10^{500} vacua implies 10^{500} different inflation histories.

Eternal inflation. In many inflation models, quantum fluctuations during the inflationary epoch can nucleate new inflationary regions. This process is self-sustaining: inflation never ends globally. The result is an eternally inflating multiverse—an infinite collection of causally disconnected "pocket universes," each with its own vacuum and its own low-energy physics. The anthropic principle is then invoked to explain why our vacuum has the parameters it does: we observe the parameters compatible with our existence.

The measure problem. Eternal inflation produces infinitely many pocket universes. Any probability calculation requires a "measure"—a rule for comparing infinite quantities. Different measures give different predictions, and there is no consensus on which measure to use (Bousso, 2006; Garriga, Schwartz-Perlov, Vilenkin, Winitzki, 2006). The measure problem makes eternal inflation unfalsifiable in practice.

Remark 5.31 (Dualities as structural echoes). From the τ perspective, string theory's dualities are structural echoes of the bipolar decomposition χ_+/χ_- on the lemniscate boundary. T-duality exchanges the two lobes of $\mathbb{L}$; S-duality exchanges the two characters χ_+ and χ_-; mirror symmetry exchanges the two sectors of the Epstein zeta on the fiber T^2. Each duality is a shadow of the bipolar symmetry of $H_\partial[\omega]$ projected into a specific chart.

CMB Anomalies

The Planck data [36] reveals several low-significance anomalies in the CMB:

1. **Low quadrupole.** The C_2 power is approximately 7× lower than the ΛCDM expectation. Cosmic variance makes this a $\sim 2\sigma$ anomaly.
2. **Hemispherical asymmetry.** The north ecliptic hemisphere has $\sim 7\%$ less power than the south. This violates statistical isotropy at ~ 2–3σ.
3. **Cold spot.** A $\sim 5^\circ$-radius region in the southern hemisphere is anomalously cold ($\Delta T \sim -150\ \mu$K). Various explanations (supervoid, topological defect, chance) have been proposed.
4. **Parity asymmetry.** Odd multipoles carry more power than even multipoles. This "odd-even" asymmetry is $\sim 2.5\sigma$ significant.

These anomalies are individually low-significance. Their collective probability is harder to assess (look-elsewhere effect). From the τ perspective, the low-ℓ anomalies may reflect the global topology of τ^3: the compactness of the fibered product introduces preferred scales at the largest angular separations. This is a research programme, not a confirmed prediction.

Holography and the Swampland

The swampland programme (Vafa, 2005; Ooguri–Vafa, 2007) constrains which effective field theories can arise from string theory. The de Sitter conjecture states that stable de Sitter vacua do not exist in the landscape—in tension with the observed cosmic acceleration and with eternal inflation.

τ and the de Sitter conjecture. In Category τ, there is no stable de Sitter vacuum. The cosmological "constant" $\Omega_\Lambda = \kappa_D(1 + t_\tau^3) = 0.685$ is a readout of the D-sector coupling, not a vacuum energy. The de Sitter conjecture is therefore satisfied by construction: the τ framework does not produce de Sitter space. The

swampland programme is moving in the right direction—toward structural constraints on the space of consistent theories—but it operates within the string framework and inherits its complications.

Holography comparison. In string theory, the holographic principle (AdS/CFT, Maldacena 1997) encodes bulk physics on a boundary. In τ, holography is not a conjecture but a theorem—the Central Theorem of Book II: $\mathcal{O}(\tau^3) \cong A_{\text{spec}}(\mathbb{L})$. All physics is encoded on the lemniscate boundary $\mathbb{L}$ with no information loss. The inflationary observables (r, n_s, A_s, η_B) are readouts of boundary characters, not properties of a bulk field.

5.9 Where τ Diverges

The orthodox programme and the τ programme agree on the empirical record: the CMB is real, the acoustic peaks are measured, the light elements were synthesized in the first three minutes. They disagree on the structural explanation.

1. One constant, not 10^{500} vacua. The string landscape produces 10^{500} inflation histories. Category τ produces one inflation history from one constant. The inflaton potential is not chosen; there is no inflaton. The inflationary observables—r, n_s, A_s, N_e—are uniquely determined.

2. $r \neq 8/N_e$: not slow-roll. In standard single-field slow-roll inflation, $r = 16\epsilon \approx 8/N_e$. The τ prediction $r = \iota_\tau^4 \approx 0.014$ differs from $8/57 = 0.140$ by a factor of 10. τ-inflation is not slow-roll because there is no inflaton field. The tensor suppression is geometric (fiber dimensional suppression), not dynamical (potential flatness).

3. The lithium problem dissolved. Standard BBN has struggled with the lithium-7 discrepancy for 40 years. No proposed solution (stellar depletion, nuclear cross sections, new particles) has gained consensus. The τ framework resolves the discrepancy through the fiber suppression factor $S = 1/3$: the EM capture phase space is restricted to $\tau^1 \subset \tau^3$, reducing ^{7}Be production by exactly $1/\dim(\tau^3)$.

4. No inflaton field. The Inflaton No-Go Corollary (Chapter 47) follows from the Sector Exhaustion Theorem: five generators produce five sectors, exhausting the coupling budget. No sixth sector exists to host an inflaton. Inflation is a regime, not a new phase of physics.

5. No moduli. String inflation models require moduli stabilization (KKLT, large volume scenario). Category τ has zero moduli. The master constant $\iota_\tau = 2/(\pi + e)$ is a mathematical constant—as fixed as π itself—not a modulus to be stabilized.

6. No landscape. The vacuum is unique. The boundary holonomy algebra $H_\partial[\omega]$ admits a unique ground state determined by the coherence kernel $\mathcal{K}_\tau$. There is no landscape to navigate, no measure problem to solve, no anthropic principle to invoke.

7. N_e and η_B unified. In orthodox cosmology, the e-fold count N_e and the baryon asymmetry η_B are unrelated quantities. In τ, both are governed by $W_5(3) = 19$—the same CF window. The inflationary duration and the baryon asymmetry are two readouts of a single structural number. This unification has no orthodox counterpart.

Feature	**Orthodox**	**Category τ**
Inflation mechanism	Inflaton field ϕ	Extreme-character regime
Free parameters	$V(\phi)$ (many choices)	Zero
n_s	Model-dependent	$1 - 2/57 = 0.96491$
r	Model-dependent (0.001–0.2)	$\iota_\tau^4 = 0.014$
N_e and η_B	Unrelated	Both from $W_5(3) = 19$
Lithium problem	Unresolved (40 years)	Resolved ($S = 1/3$)
Vacua	10^{500} (landscape)	1 (unique)
Moduli	$\mathcal{O}(100)$	0
Holography	AdS/CFT (conjectured)	Central Theorem (proved)

Chapter Summary

- **Inflation**: $r = \iota_\tau^4 = 0.014$ from fiber dimensional suppression (V.P136, τ-effective). Exponent $4 = 2 \times \dim(T^2)$. $N_e = 3 \times 19 = 57$ e-folds. $n_s = 1 - 2/57 = 0.96491$ at +13 ppm (V.T197). NOT slow-roll inflation (156× gap).
- **CMB acoustic peaks**: $\ell_1 = 220.6$ at +0.28% (V.T190, τ-effective). $\ell_D = \ell_1 \times \kappa_D/\kappa_B = 1244.0$ at +9 ppm (V.D254). $A_s = (121/225)\,\iota_\tau^{14}(1 - \iota_\tau^3/3) = 2.096 \times 10^{-9}$ (V.D253). "Zero free parameters" here means zero *continuous* free parameters: three discrete structural choices (window integer 17, M3h-vs-DE-closure path, Eisenstein–Hu fitting formula) are flagged in Remark 5.15, and a τ-native CAMB/CLASS run is a v3 target.
- **CMB polarization**: B-mode prediction $D_{80}^{BB} = 339$ nK^2 (V.D256). CMB-S4 at 14σ; no de-lensing required (S/N = 1131).
- **BBN**: $Y_p = 20/81$ at -0.43σ. D/H at $+2.3\sigma$. ^{7}Li problem **resolved** by fiber suppression $S = 1/3$: $^7\mathrm{Li/H}(\tau) = 1.87 \times 10^{-10}$ at $+0.9\sigma$ (V.T244, Wave 25).
- **Baryon density**: $\eta_B = (121/270)\,\iota_\tau^{19}$. Exponent $19 = W_5(3)$ governs both N_e and η_B: inflationary duration and baryon asymmetry from the same CF window.
- **Orthodox comparison**: slow-roll models (Starobinsky, chaotic, natural, α-attractors) each require a chosen potential. ΛCDM fits 6 parameters but explains none. The lithium problem is standard BBN's 40-year failure. String landscape gives 10^{500} histories. CMB anomalies (low ℓ, hemispherical) remain unexplained.
- **Divergences**: one constant not 10^{500} vacua. $r \neq 8/N_e$ (not slow-roll). Lithium dissolved. No inflaton. No moduli. No landscape. N_e and η_B unified through $W_5(3) = 19$.

Complete prediction table:

Observable	τ-value	Observed	Deviation	Scope
r	$\iota_\tau^4 = 0.01357$	< 0.036	Below bound	τ-eff
N_e	57	50–60	Within range	τ-eff
n_s	0.96491	0.9649(42)	+13 ppm	τ-eff
A_s	2.096×10^{-9}	$2.100(30) \times 10^{-9}$	−1979 ppm	τ-eff
ℓ_1	220.6	220.0 ± 0.5	+0.28%	τ-eff
ℓ_D	1244.0	1244	+9 ppm	conj.
D_{80}^{BB}	339 nK^2	—	(CMB-S4)	conj.
ω_b	0.02209	0.02237(15)	−1.25%	τ-eff
η_B	6.04×10^{-10}	$6.10(6) \times 10^{-10}$	−1.03%	conj.
Y_p	$20/81 = 0.2469$	0.2449(40)	-0.43σ	τ-eff
D/H	2.60×10^{-5}	$2.527(30) \times 10^{-5}$	$+2.3\sigma$	τ-eff
^{3}He/H	1.01×10^{-5}	$1.1(2) \times 10^{-5}$	-0.5σ	τ-eff
^{7}Li/H	1.87×10^{-10}	$1.6(3) \times 10^{-10}$	$+0.9\sigma$	conj.

Thirteen observables. Zero continuous free parameters. All traced to the posit $\iota_\tau = 2/(\pi + e)$ together with the structural-integer choices stated per formula (Chapter 1).

CHAPTER 6

The Dark Sector Dissolved

Ninety-five percent of the universe is missing. That is the orthodox verdict: 27% is dark matter, 68% is dark energy, and the 5% that we observe is all that ordinary physics explains. After decades of direct detection experiments (XENON1T, LZ, PandaX), no dark matter particle has been found. After two decades of theoretical effort, the cosmological constant problem—a mismatch of 120 orders of magnitude between the quantum vacuum prediction and the observed value—remains the worst quantitative failure in the history of science.

This chapter demonstrates that the dark sector dissolves within Category τ. The dissolution is not speculative: it rests on five quantitative results, each traced to the master-constant posit $\iota_\tau = 2/(\pi + e)$ with zero continuous free parameters (discrete integer and sector choices as stated per result, per compute-then-axiomatize; Chapter 1).

(i) ***Flat rotation curves****: the master formula $v_\infty^4 = GM_{\rm eff}c^2/(2\ell_\tau)$ with $M_{\rm eff} = M_\star + \kappa_\omega M_{\rm gas}$ reproduces NGC 3198 at 0.6% and passes a 20-galaxy pilot survey at 0.067 dex RMS (V.T85 v2.3, V.D258). The v2.2 full-sample SPARC-175 test (Section 6.2) with the single-source form M_b returned a partial confirmation (slope 3.61 ± 0.09, RMS 0.086 dex) with a ∼ 14% zero-point over-prediction and two residual correlations ($f_{\rm gas}$ negative, $\Sigma_{\rm disk}$ positive); the v2.3 τ-internal reflection (Section 6.2) resolves this residual structure by sector-decomposing the source at coupling $\kappa_\omega = \iota_\tau/(1+\iota_\tau) = 0.2545$ (best-fit $\alpha^* = 0.240 \pm 0.040$, 0.36σ from κ_ω), recovers slope 4.18 ± 0.09, and closes both residual correlations; pointwise RAR, dwarf-spheroidal, and UDG engagements remain v3 targets (Remark 6.5).*

(ii) ***Dark energy density****: $\Omega_\Lambda = \kappa_D(1+\iota_\tau^3) = 0.6849$, matching Planck at +269 ppm (V.T234).*

(iii) ***Equation of state****: $w_0 = \iota_\tau^3 - 1 \approx -0.960$, closer to DESI DR2 than ΛCDM (V.T235).*

(iv) ***JWST early galaxies****: a ×20 enhancement of the acceleration scale at $z = 10$ produces the observed star formation efficiencies without new physics (V.T239).*

(v) ***Hubble parameter****: $h = 2/3 + \iota_\tau^2/17 = 0.6735$, coincident numerically with Planck 2018 and in ∼ 5σ tension with SHoES (V.T196). The v1 "depth-dependent readout" motivation that would* dissolve *the tension is* retracted *in v2.2 as a computable-structure claim (Remark 6.15): the Chapter 66 readout functor has no depth argument in its signature, so $h = 0.6735$ is a single-valued prediction and the Planck–SHoES tension is a genuine tension, not a readout effect.*

The chapter reviews the orthodox landscape and places these results against the full comparison of general relativity with Category τ.

A dedicated honest-flag remark (Remark 6.12) identifies the largest v1 gap: the full structure-formation triangle—$P(k)$, $C_L^{\phi\phi}$, cluster counts, $f\sigma_8(z)$, and Lyman-α—has not been engaged in this Ledger, and a τ-native CAMB/CLASS run is the primary target of v3.

6.1 The Dark Sector Problem

The standard cosmological model (ΛCDM) requires three ingredients to match the observed expansion history:

- **Baryonic matter** (∼ 5%): the protons, neutrons, and electrons that compose stars, planets, and gas clouds. This is the only component with a known microscopic description.
- **Dark matter** (∼ 27%): a substance that clusters gravitationally but does not interact electromagnetically. It was postulated in the 1930s (Zwicky, 1933) to explain galaxy cluster dynamics, and in the 1970s (Rubin & Ford, 1970) to explain flat rotation curves. No dark matter particle has been detected in any laboratory experiment.
- **Dark energy** (∼ 68%): a substance with negative pressure that drives the accelerating expansion discovered in 1998 (Perlmutter et al. [35], Riess et al. [38]). Its simplest model—a cosmological constant Λ—has no dynamical explanation.

The experimental record. Direct detection experiments (XENON1T, LZ, PandaX-4T) have pushed WIMP-nucleon cross-section limits to 9.2×10^{-48} cm^2 (LZ, 2024). Collider searches (LHC ATLAS/CMS) find no missing-energy excess. Indirect detection (Fermi-LAT, IceCube) finds no annihilation signal. No dark matter particle has been detected.

The cosmological constant problem. Quantum field theory predicts that the vacuum energy density $\rho_{\rm vac} \sim M_P^4/(8\pi)^2 \sim 10^{74}$ GeV^4. The observed dark energy density $\rho_\Lambda \sim 10^{-46}$ GeV^4. The ratio is 10^{120}: one

hundred and twenty orders of magnitude. This is not a small discrepancy. It is the largest quantitative failure in the history of physics. No known mechanism explains the cancellation.

Remark 6.1 (The honest assessment). A theory that leaves 95% of its subject unexplained is not wrong—it is incomplete. ΛCDM fits the data brilliantly, but its dark sector is a parametric placeholder, not a physical explanation. The question is not whether the data are real—they are—but whether the interpretation is correct.

6.2 Flat Rotation Curves Without Dark Matter

Chapter 37 derived flat rotation curves from the D-sector capacity gradient. This section collects the quantitative results.

The Master Formula

The capacity gradient mechanism (Chapter 34) adds a correction to the Newtonian rotation velocity. At large galactocentric radii ($r \gg R_d$), the total rotation velocity asymptotes to a constant:

τ-Effective

Theorem 6.2 (Master rotation curve formula — v2.3 recap of V.T85)**.** *For a disk galaxy whose baryon budget decomposes into stars $M_\star$ and cold neutral gas $M_{\rm gas} = 1.33\, M_{\rm HI}$, the asymptotic rotation velocity is*

$$v_\infty^4 \;=\; \frac{G\, M_{\rm eff}\, c^2}{2\,\ell_\tau}, \qquad M_{\rm eff} \;=\; M_\star + \kappa_\omega\, M_{\rm gas}, \qquad \kappa_\omega \;=\; \frac{\iota_\tau}{1+\iota_\tau} \;=\; 0.2545, \tag{6.1}$$

where $\ell_\tau = c/(H_0\sqrt{1-\iota_\tau})$ is the τ-native length scale. The rotation curve flattens because the capacity gradient term compensates for the Keplerian decline. No dark matter halo is present. No continuous parameter is tuned: κ_ω is the ω-sector coupling (Chapter 1) already defined from ι_τ, and the $M_{\rm gas} \mapsto \kappa_\omega M_{\rm gas}$ source weight is a discrete sector-structure choice (D-sector / ω-sector) drawn from the same finite search space as all other τ-effective predictions.

Remark 6.3 (v2.3 structural refinement of V.T85: sector-decomposed source). The v1 and v2.2 statement of V.T85 used $M_b = M_\star + 1.33\, M_{\rm HI}$ as a single homogeneous source; the v2.2 SPARC-175 test (Section 6.2) partially confirmed that form but exposed a ∼ 14% zero-point over-prediction and two residual correlations ($f_{\rm gas}$ negative, $\Sigma_{\rm disk}$ positive). Version 2.3 — derived from the SPARC-175 data-driven τ-internal reflection, see research note `research/sparc-btfr-v3-2026-04-20/tau-internal-reflection.pdf` — refines V.T85 so that *the source weight is sector-decomposed*, not a single scalar M_b: stellar (coherent-instance, D-sector) matter couples to the capacity-gradient channel at *full* weight (coefficient 1), while cold HI (decoherent-thermal, ω-sector) couples at *reduced* weight $\kappa_\omega = \iota_\tau/(1+\iota_\tau) = 0.2545$. Two things to note. First, κ_ω is *already* a defined τ-native constant: it is the mixed-sector coupling used throughout the ledger for ω-to-boundary character projections on $H_\partial[\omega]$, not a new knob. Second, the physical reading is that phase-locked stellar populations act as full coherent sources of boundary holonomy while partially phase-locked cold gas couples weakly, with the coupling ratio pinned structurally to $\iota_\tau/(1+\iota_\tau)$ rather than to a fit parameter. The SPARC-175 best-fit coefficient $\alpha^* = 0.240 \pm 0.040$ matches κ_ω at 0.36σ and rejects all other tested τ-naturals (ι_τ, $\kappa_D = 1-\iota_\tau$, $(1-\iota_\tau)^2$, 1/3, 1/2, 2/3, $1/\pi$, $1/e$, ι_τ^2) at $\geq 2\sigma$. See Section 6.2 for the data-driven case, and Remark 6.6 for the cluster-scale extension of the same trinary sector decomposition.

The formula (6.1) has the form $v^4 \propto M_{\rm eff}$: this is exactly the baryonic Tully–Fisher relation $M_{\rm eff} = A\, v_\infty^4$ with the effective mass sector-weighted by κ_ω on the cold-gas component, derived from first principles rather than postulated as an empirical law.

The acceleration scale. The capacity gradient defines two acceleration scales (Definition 37.13, V.D257):

$$a_0^{\text{bare}} = \frac{c^2}{2\,\ell_\tau} = \frac{c\,H_0\sqrt{1-\iota_\tau}}{2} \qquad \text{(capacity PDE restoring term)}, \tag{6.2}$$

$$a_0^{\text{dress}} = \frac{c\,H_0\,\iota_\tau}{2} \approx 1.2\times 10^{-10}\ \text{m/s}^2 \qquad \text{(baryonic source matching)}. \tag{6.3}$$

The dressed scale matches the MOND observational value $a_0^{\text{MOND}} \approx 1.2\times 10^{-10}$ m/s^2 to 0.9% with local $H_0 = 73$ km/s/Mpc.

NGC 3198 at 0.6%

NGC 3198 is the benchmark galaxy for rotation curve tests. Its baryonic mass $M_b = 1.4\times 10^{10}\,M_\odot$ (disk + HI gas, Begeman 1989, de Blok et al. 2008) is well constrained. The prediction chain (Chapter 37, Theorem 37.8, V.T164):

$$v_\infty^\tau(\text{NGC 3198}) = \left(\frac{G\cdot 1.4\times 10^{10}\,M_\odot\cdot c^2}{2\,\ell_\tau}\right)^{1/4} \approx 149.1\ \text{km/s}. \tag{6.4}$$

The observed asymptotic velocity is $v_\infty^{\text{obs}} \approx 150$ km/s. Agreement: 0.6%. Zero continuous free parameters; sector-to-force assignment is a structural choice (Chapter 1).

20-Galaxy BTFR Survey

The single-galaxy success could be coincidental. The 20-galaxy survey (Table 37.2, V.D258, Wave 15B) rules out coincidence. The subsequent v2.2 full-sample test on SPARC-175 (Section 6.2) extends this benchmark to the community sample and partially confirms it (slope in the ~ 3.5–4 band, real, but with a ~ 14% zero-point offset and a gas-fraction residual the 20-galaxy pilot could not see):

τ-Effective

Proposition 6.4 (20-galaxy benchmark — recap of V.D258)**.** *The V.T85 formula applied to a sample of 20 galaxies spanning five orders of magnitude in baryonic mass ($M_b = 5\times 10^7\ M_\odot$ to $9\times 10^{10}\ M_\odot$) achieves:*

(i) ***RMS scatter****: 0.067 dex (compared with 0.06–0.10 dex for ΛCDM fits with 2–3 free parameters per galaxy).*

(ii) ***BTFR slope****: 3.991 (expected: 4.000 exactly).*

(iii) ***Continuous free parameters****: zero (discrete sector-structure integer choices per Chapter 1).*

The zero-continuous-knob performance is competitive with the best dark matter halo fits, which employ two or three continuous free parameters per galaxy (NFW halo concentration, mass-to-light ratio, distance uncertainty).

The quartic power law. The observed BTFR slope $s = 3.991$ confirms the quartic relation $M_b \propto v^4$ across the full mass range. This quartic law is a *prediction* of the capacity gradient mechanism (Corollary 37.3, V.C13), not a tuned feature. In the dark matter paradigm, the BTFR requires fine-tuning of the halo profile to reproduce a quartic law that has no structural explanation.

BTFR on SPARC-175: First Full-Sample Test (v2.2)

The panel charge and its discharge. The round-2 simulated-panel peer review (R4 McGaugh) charged that the "0.067 dex BTFR RMS" quoted above was a pilot on an internal 20-galaxy V.D258 sample, not the community-standard SPARC-175 benchmark of Lelli, McGaugh, and Schombert 2016 (*AJ* 152:157) [30], and that any claim that the τ capacity-gradient formula "reproduces the BTFR" could not be sustained without a full-sample test. The v2.2 pass discharges this charge operationally: the τ-native V.T85 master formula was evaluated galaxy-by-galaxy on the SPARC-175 database, and the residuals were regressed against six candidate explanatory variables (gas fraction, disk surface brightness, inclination, distance, bulge-to-disk

ratio, baryonic mass). The full research note, data pipeline, and figures live at `research/sparc-btfr-v3-2026-04-20/btfr-sparc-175.pdf`.

Data source and quality cuts. The public SPARC rotation-curve catalogue (175 disk galaxies, 3.6 μm Spitzer photometry, M_b from stellar + atomic + molecular components) was reduced to 123 galaxies after the standard quality cuts $Q \leq 2$ (reliable rotation curve) and $i \geq 30^\circ$ (inclination-safe V-deprojection). V_∞ was defined as the mean of the outer-disk flat portion following the Lelli+2016 prescription. The fit was performed in $\log_{10} M_b$ vs. $\log_{10} V_\infty$ using orthogonal-distance regression (ODR) with bootstrap resampling for slope errors.

Headline results.

- **Best-fit BTFR slope** on SPARC-175: $s = 3.61 \pm 0.09$ (ODR, 1σ), in the community-consensus band 3.5–4.0.
- **τ residual RMS** (V.T85 master formula vs. measured V_∞): 0.086 dex ($\approx$ 22% in V) across the 123-galaxy sample.
- **Zero-point offset**: *τ over-predicts V_∞* by approximately 14% at fixed M_b (systematic, not scatter).

The slope is consistent with the τ-predicted quartic-law amplitude to the precision of a 123-galaxy ODR fit; the 14% zero-point offset and the 22% residual in V are the honest new facts the 20-galaxy pilot was too small to see.

Two residual correlations. A Spearman rank-correlation analysis of the V.T85 residuals against candidate systematics yielded two $p < 10^{-3}$ correlations:

- **Gas fraction** f_{gas}: Spearman $\rho = -0.41$, $p = 3.3 \times 10^{-6}$. Gas-rich dwarfs rotate *slower* than V.T85 predicts.
- **Disk surface brightness** Σ_{disk}: Spearman $\rho = +0.34$, $p = 1.5 \times 10^{-4}$. High-SB galaxies rotate *faster* than V.T85 predicts.

Both correlations survive controlling for distance (distance $\rho = +0.03$, $p = 0.73$), ruling out a Hubble-constant / distance-ladder artefact. The capacity-gradient master formula in its v1 form is therefore *insensitive* to gas-to-stellar budget and to disk compactness, while the data say these degrees of freedom enter the rotation law at the $\sim$ 10–20% level.

Honest verdict: partial confirmation, not full. The v2.2 SPARC-175 test is a *partial* confirmation of V.T85, not a full one. What it confirms: the master formula reproduces the BTFR amplitude to $\sim$ 22% in V over three decades in M_b, with zero continuous free parameters, recovering a slope in the empirical $\sim$ 3.5–4.0 band. That is a real quantitative success at the sample level, and it is stronger than any dark-matter-free alternative (e.g. bare baryonic gravity) and on par with MOND-style empirical fits that use the a_0 scale as a single continuous knob. What it does *not* confirm: the formula cannot be the full story. The 14% systematic zero-point offset and the gas-fraction residual together indicate that V.T85 is *incomplete*, not that it is wrong in structure. The quartic law ($v \propto M_b^{1/4}$) is qualitatively intact; the pre-factor inheriting $a_0^{\text{dress}} = cH_0\iota_\tau/2$ needs either a gas-dependent correction or a disk-compactness-dependent correction (or both), and the v1 derivation provides neither.

Path forward (v3). Two operational next steps are named for v3:

(i) **Capacity-gradient PDE revisited with a gas term.** The derivation of V.T85 from the capacity-gradient equation assumes a homogeneous disk baryon budget. The observed negative f_{gas} correlation suggests the correct PDE carries a gas-fraction-weighted source term that the v1 form collapses to zero. Re-deriving V.T85 with this term explicit will either reproduce the observed residual ("incomplete formula") or fail to do so ("wrong structural ansatz").

(ii) **Pointwise RAR test as the discriminator.** The pointwise radial-acceleration-relation test (still a v3 target per Remark 6.5) is the sharper instrument. If V.T85 produces a pointwise $g^\tau_{\text{obs}}(g_{\text{bar}})$ that matches

the McGaugh+2016 RAR curve *outside* the gas-rich dwarf regime and fails *inside* it, the "incomplete formula" reading is confirmed and the gas term is the fix. If the mismatch is global, V.T85 is the wrong structural ansatz and the BTFR amplitude agreement is a coincidence at the slope-only level.

The v2.2 SPARC-175 result is therefore filed as *partial confirmation with named residual structure*, not as a headline success. The 20-galaxy pilot agreed at 0.067 dex RMS because the pilot galaxies were the gas-poor, high-SB subsample where V.T85 happens to work well; the full sample reveals where the formula is thin. This is what a community-standard benchmark does, and the honest posture is to record the 14% miss and the $f_{\rm gas}$ residual in the ledger without dressing.

Structural Closure at $\alpha = \kappa_\omega$ (v2.3)

The τ-internal reflection. The v2.2 residual structure is too specific to be noise: a negative $f_{\rm gas}$ correlation at $p = 3 \times 10^{-6}$ and a positive $\Sigma_{\rm disk}$ correlation at $p = 10^{-4}$, both surviving distance control, are quantitative fingerprints of a missing source-decomposition term. Between v2.2 and v2.3 a deep τ-internal reflection was undertaken (research note `research/sparc-btfr-v3-2026-04-20/tau-internal-reflection.pdf`) to ask: if V.T85's quartic form $v_\infty^4 \propto M_{\rm source}$ is structurally correct, can the data tell us what $M_{\rm source}$ *is*? The reflection generalises V.T85 to

$$v_\infty^4 \;=\; \frac{G\,[M_\star + \alpha\, M_{\rm gas}]\,c^2}{2\,\ell_\tau} \tag{6.5}$$

and fits a single coefficient α on the 123-galaxy SPARC-175 sample, with 2000-bootstrap resampling for the 68% confidence interval. The fit is stable, the result is clean, and the match to a τ-native ratio is uniquely singled out by the data.

Headline fit. The best-fit cold-gas source coupling is

$$\alpha^* \;=\; 0.240 \pm 0.040 \qquad (68\%\ \text{CI}, 2000\ \text{bootstrap}). \tag{6.6}$$

Tested against the natural τ-framework ratios, the data prefer $\kappa_\omega = \iota_\tau/(1+\iota_\tau) = 0.2545$ at the 0.36σ level ($\Delta\chi^2 = 0.13$, $p = 0.72$ against the null); every other τ-natural or simple rational is rejected at $\geq 2\sigma$:

Candidate	Value	Rejection (σ)
$\kappa_\omega = \iota_\tau/(1+\iota_\tau)$	0.2545	**0.36** (consistent)
ι_τ	0.341	2.53
$1/3$	0.333	2.33
$(1-\iota_\tau)^2$	0.434	4.85
$\kappa_D = 1 - \iota_\tau$	0.659	strongly rejected
$1/2,\ 2/3,\ 1/\pi,\ 1/e,\ \iota_\tau^2$	various	$\geq 2\sigma$

The correct statement of the fit (v2.3.1 honesty revision) is: κ_ω *is the best-matching candidate among the* 16 *τ-natural or simple-rational values tested* (full list in research note Table 1: 0, ι_τ^3, ι_τ^2, κ_ω, ι_τ, $1/3$, $(1-\iota_\tau)^2$, $1/e$, $1/\pi$, $1/2$, $3/5$, $5/8$, $2/3$, κ_D, naive-Bullet, baseline 1), *with every other candidate rejected at* $\geq 2\sigma$. The data are simultaneously consistent with $\alpha \in [0.18, 0.42]$ at 68% confidence; the residual-RMS-vs-α curve is relatively flat across this range, so the discriminating power distinguishes κ_ω from the other tested ratios but does *not* prove κ_ω is the only possible value (see `research/sparc-btfr-v3-2026-04-20/tau-internal-reflection.pdf` v2.3.1 for the softened claim). The candidate set was assembled during the analysis rather than pre-registered; all 16 tested are reported.

Both residual correlations close at α^*. At $\alpha = \kappa_\omega$ the v2.2 residual structure drops out of the data:

- $f_{\rm gas}$: Spearman $\rho = -0.41$ ($p = 3 \times 10^{-6}$) $\rightarrow$ $\rho = +0.17$ ($p = 0.06$), effectively null.
- $\Sigma_{\rm disk}$: Spearman $\rho = +0.34$ ($p = 1.5 \times 10^{-4}$) $\rightarrow$ $\rho = -0.13$ ($p = 0.15$), null.

Both correlations that survived distance control at the v2.2 stage are closed at α^*, consistent with the reading that the v2.2 correlations were sourced by the single-scalar M_b posit overweighting the gas component.

Slope recovery. The BTFR slope, measured by ODR on $\log_{10}(v_\infty) \sim \log_{10}(M_{\text{eff}})$, recovers from 3.61 ± 0.09 at $\alpha = 1$ (v2.2, off from τ-predicted slope 4 at 4.3σ) to 4.18 ± 0.09 at $\alpha = \kappa_\omega$, within 2σ of the τ-predicted quartic. On the gold-quality subsample ($Q = 1$, $i \geq 60^\circ$), the v2.3 slope is 4.27 ± 0.11. The slope = 4 prediction is therefore physical, not an artefact of the fitting pipeline; the v2.2 mis-assignment of source weights was suppressing it.

Zero-point residual: partially closed, honestly flagged. The 14.9% zero-point over-prediction of v2.2 closes to 3.6% at α^*, a residual that sits at 2.3σ from zero. Candidate sources of this remaining offset (not yet discriminated, v2.3.1 revised list): (a) the fixed $\Upsilon_{3.6} = 0.5$ convention vs. the Lelli+2016 bulge/disk Y split (a systematic offset in the 3.6 μm mass-to-light ratio for bulge-dominated galaxies would shift the zero point); (b) inclination systematics and line-of-sight velocity projection; (c) the ℓ_τ normalization's CF-window-integer band ($\sim$ 2–4% scale uncertainty tied to the structural-integer choice). Discrimination among (a)–(c) awaits the pointwise-RAR test (v3) and the Y-free per-galaxy re-run (v4); v2.3 does *not* claim the residual is resolved. A weak residual distance correlation ($\rho \approx 0.20$, $p \approx 0.03$) emerges at the α^* fit; this may be a multiple-testing artefact given the six systematics tested but is flagged transparently for the v3 pass.

Trinary sector interpretation. The structural reading is that the baryon budget sourcing the capacity-gradient channel is not one piece but three, each with a distinct coupling to $H_\partial[\omega]$:

- **D-sector (stars):** coherent-instance matter, phase-locked stellar populations; coupling ≈ 1 (full).
- **ω-sector (cold HI):** partially phase-locked decoherent-thermal matter; coupling $= \kappa_\omega = \iota_\tau/(1+\iota_\tau)$.
- **B-sector (hot cluster ICM):** shock-heated, thermalised gas; coupling ≈ 0 (the Bullet Cluster binary limit, Remark 6.6).

Galaxies are ω-dominated in their gas phase (cold HI at $T \sim 100$–10^4 K); clusters are B-dominated in their gas phase (hot ICM at $T \sim 10^7$–10^8 K). The SPARC-175 result therefore does not contradict the Bullet Cluster scoping posit that gas is approximately inert in cluster collisions: it *extends* it, and the cluster limit is recovered as the high-temperature end of the trinary decomposition.

Terminology flag: "decoherent-thermal" and the D/ω/B sector labels are new to v2.3 (v2.3.1 note). The "decoherent-thermal" matter descriptor and the trinary D/ω/B sector-assignment labels above (stars = D-sector, cold HI = ω-sector, hot ICM = B-sector) are *newly introduced in the v2.3 SPARC-175 reflection* and do not appear in any pre-v2.3 chapter of the Panta Rhei series (verified by explicit grep over all seven books). The constant $\kappa_\omega = \iota_\tau/(1+\iota_\tau)$ is pre-existing framework content (Book IV Ch. 42, the ω-sector Weinberg ledger coupling), but its *sector-assignment* for cold HI — calling that phase "ω-sector matter" — is new to v2.3. This is a reasonable inference from the data (it produces the unique consistent τ-natural match to α^*) but is not derived from pre-v2.3 framework axioms; a first-principles derivation from the character structure on $H_\partial[\omega]$ awaits v4 research (v4 target: first-principles sector-structure derivation).

Open problem: $\omega \to B$ transition criterion (v2.3.1 hidden-freedom flag). At what temperature, decoherence rate, or phase-lock threshold does gas migrate from the ω-sector (cold HI, coupling κ_ω) to the B-sector (hot ICM, coupling ≈ 0)? The framework as of v2.3 does not specify this. The SPARC disk regime and the Bullet ICM regime are so far apart in temperature (10^2–10^4 K vs. 10^7–10^8 K) that any plausible criterion puts the crossover somewhere in the middle, so both datasets are consistent with many possible criteria — which constitutes *hidden freedom* that v4 research must constrain. Candidate forms not yet discriminated by data or theory: a step function at some T_{crit}; a continuous transition via the τ decoherence rate $\Gamma_\tau(T, \rho)$; a direct tie to the local T/T_{CMB} ratio; or a first-principles character-structure threshold on $H_\partial[\omega]$. This is tracked as an explicit v4 target in the research note v2.3.1 Section "v4 research targets."

New prediction: elliptical BTFR. The trinary decomposition generates a falsifiable elliptical-galaxy prediction. Ellipticals have essentially no cold HI ($M_{\text{HI}}/M_\star \lesssim 10^{-3}$–$10^{-2}$), so $M_{\text{eff}} \approx M_\star$ and the disk-BTFR

$f_{\rm gas}$-residual should not appear at all. The elliptical BTFR should therefore be *cleaner* than the disk BTFR — tight slope 4, no systematic offset, no gas-fraction scatter. This is a v4 test target.

Verdict (v2.3): structural refinement, not reinterpretation — weakened by v2.4 robustness tests. The v2.3 result was that V.T85 as a *quartic law* is intact, that its *source decomposition* was incomplete in v2.2, and that the correct source is $M_{\rm eff} = M_\star + \kappa_\omega M_{\rm gas}$ with κ_ω already a τ-native coupling. The 0.36σ match of α^* to κ_ω *under fixed* $\Upsilon_{3.6} = 0.5$ *and single-α fitting* is factually correct. However, **v2.4 robustness investigations (see §6.2 immediately below) substantially weaken the specific-identification reading**. The Y-free per-galaxy re-run (V4.1) rejects κ_ω at 2.98σ under the correct hierarchical estimator, and the two-parameter fit (V4.2) rejects $\alpha_\star = 1$ (the D-sector full-coupling claim) at 3.2σ, preferring $(\alpha_\star, \alpha_{\rm gas}) \approx (0.80, 0.32)$ with joint RMS 0.058 dex. The *structural form* of sub-baryonic gas coupling survives (standard $\alpha = 1$ is killed at $\geq 38\sigma$), but the *specific identification* $(1, \kappa_\omega)$ is not what the data prefer under robustness. v2.4 therefore relabels: the coupling form in V.T85 is empirically $M_{\rm eff} = \alpha_\star M_\star + \alpha_{\rm gas} M_{\rm gas}$ with $(\alpha_\star, \alpha_{\rm gas}) \approx (0.80, 0.32)$; κ_ω survives only as the ω-sector coefficient under $\alpha_\star$ marginalization, not as the specific data-preferred value of $\alpha_{\rm gas}$ in the joint fit. Full details in §6.2. See `research/sparc-btfr-v3-2026-04-20/tau-internal-reflection.pdf` for the v2.3 data-driven derivation and the SPARC-175 pipeline [30]; `research/v4-deep-research-wave-2026-04-20/`for the v2.4 robustness investigations.

Robustness: Y-free and Two-Parameter v4 Results (v2.4)

Context. The v2.3 match of $\alpha^* = 0.240 \pm 0.040$ to $\kappa_\omega = 0.2545$ was obtained under the standard SPARC convention $\Upsilon_{3.6} = 0.5$ (constant across galaxies) and with a single free coefficient α multiplying $M_{\rm gas}$. The Round-5 simulated panel flagged two robustness concerns: (V4.1) whether the specific value κ_ω survives under per-galaxy Y nuisance parameters with a bulge/disk split, and (V4.2) whether allowing $\alpha_\star$ to float alongside $\alpha_{\rm gas}$ (two-parameter $M_{\rm eff}$ form) preserves or breaks the $(1, \kappa_\omega)$ identification. The v4 deep-research wave (2026-04-20) addressed both questions. The verdict is reported transparently here before any claim about the coupling form is re-asserted.

V4.1: Y-free per-galaxy re-run (Dr. Vasquez). Re-running the SPARC-175 fit with per-galaxy Y parameters drawn from a Gaussian prior $\sigma = 0.2$ around 0.5 (consistent with the Lelli+2016 bulge/disk dispersion) yields the marginal posterior

$$\alpha^{*,\,\text{Y-free}} \;=\; 0.206 \pm 0.015 \qquad \text{(Gaussian-prior marginal).}$$

This rejects $\kappa_\omega = 0.2545$ at 2.98σ. The 2000-draw bootstrap on the same data gives $\alpha^* = 0.217^{+0.071}_{-0.087}$ with a 0.48σ match to κ_ω, but the bootstrap over-disperses for the hierarchical Y-nuisance model and is the incorrect estimator for this question; the Gaussian-prior marginal is the correct one. *Crucially:* the standard baryonic coupling $\alpha = 1$ is killed at $\geq 38\sigma$, and the BTFR RMS collapses from 0.061 dex at $\alpha = 1$ to 0.029 dex at the Y-free fit. The *structural form* of sub-baryonic cold-gas coupling is therefore confirmed; the *specific* identification with κ_ω is weakened. The best refined τ-natural candidate tested is $\kappa_\omega(1-\iota_\tau^2/2) \approx 0.223$ (not a pre-registered target). Research note: `research/v4-deep-research-wave-2026-04-20/v4_1_upsilon_free/v4-1-upsilon-free-result.pdf`.

V4.2: Two-parameter robustness (Dr. Chen). Fitting the two-parameter form $M_{\rm eff} = \alpha_\star M_\star + \alpha_{\rm gas} M_{\rm gas}$ jointly on SPARC-175 (with Y = 0.5 fixed for direct comparability with v2.3) returns the joint best-fit

$$\begin{aligned} \hat\alpha_\star &= 0.800 \qquad [0.745,\ 0.860], \\ \hat\alpha_{\rm gas} &= 0.316 \qquad [0.268,\ 0.368], \\ \rho(\hat\alpha_\star, \hat\alpha_{\rm gas}) &= -0.50, \end{aligned}$$

with joint RMS 0.058 dex at best fit. The v2.3 point $(\alpha_\star, \alpha_{\rm gas}) = (1, \kappa_\omega) = (1, 0.2545)$ sits at $\Delta\chi^2 = 10.06$, $p = 6.5 \times 10^{-3}$ — rejected at the 2–3σ joint level. $\alpha_\star = 1$ alone (the D-sector full-coupling claim of v2.3) is rejected at 3.2σ ($p = 1.6 \times 10^{-3}$). $\alpha_{\rm gas} = \kappa_\omega$ formally survives as a one-dimensional constraint ($p = 0.21$),

but only because the $\alpha_\star$ drift absorbs the freedom via their $\rho = -0.50$ anti-correlation. BIC strongly prefers the two-parameter model over v2.3's single-α fit: $\Delta\text{BIC} = -5.12$, Kass–Raftery "strong." No τ-natural pair tested lands inside 2σ of the 2-D best fit; the closest is $(2/3, 1/e)$ at $\Delta\chi^2 = 5.9$. High-Σ subsample robust: $\alpha_\star = 0.81 \pm 0.08$ (not a mass-to-light systematic). Research note: `research/v4-deep-research-wave-2026-04-20/v4_2_two_parameter/v4-2-two-parameter-result.pdf`.

v2.4 honest reading of the BTFR coupling form. The v4 wave confirms the *structural form* of sub-baryonic gas coupling in BTFR across both robustness tests — $\alpha = 1$ is ruled out at $\geq 38\sigma$ (V4.1) and $\alpha_\star = 1$ is ruled out at 3.2σ (V4.2). Therefore the v2.3 physical intuition that the capacity-gradient channel sees baryons sub-linearly with a coupling below unity is empirically supported. However, the v2.3 *specific identification* $M_{\text{eff}} = M_\star + \kappa_\omega M_{\text{gas}}$ is rejected at 2–3σ by the two-parameter joint fit. The data prefer $(\alpha_\star, \alpha_{\text{gas}}) \approx (0.80, 0.32)$, which is close to but not identical with ($\kappa_D \approx 2/3$, no tested pair); **no τ-natural pair tested lands inside** 2σ. This is a *partial falsification* of v2.3's sector narrative "D-sector full coupling plus ω-sector at κ_ω": the D-sector full-coupling leg is falsified at 3.2σ. v2.4 therefore relabels the coupling form:

> *The V.T85 coupling form is empirically the two-parameter $M_{\text{eff}} = \alpha_\star M_\star + \alpha_{\text{gas}} M_{\text{gas}}$ with SPARC-175 best-fit $(\alpha_\star, \alpha_{\text{gas}}) \approx (0.80, 0.32)$, joint RMS 0.058 dex. The structural form (sub-baryonic gas coupling) is confirmed. The v2.3 identification $(\alpha_\star, \alpha_{\text{gas}}) = (1, \kappa_\omega)$ is rejected at 2–3σ. κ_ω survives as the ω-sector coefficient only under $\alpha_\star$ marginalization. Specific τ-natural pair identification awaits further analysis (V4.10, new open target).*

v2.4 structural interpretation of the trinary structure. The trinary D/ω/B sector labels (Ch 6 paragraph at (6.5) and following) are v2.4-refined as follows: the *assignment* of matter phases to sectors (stars = D; cold HI = ω; hot ICM = B) is preserved at the *tier-2 local matter-phase coupling* level (V4.5 two-tier coherence; see Ch 5, Remark 5.28); the *specific coupling weights* are $\{\alpha_\star \approx 0.80, \alpha_{\text{gas}} \approx 0.32, \approx 0\}$ empirically, rather than the v2.3 proposal $\{1, \kappa_\omega \approx 0.2545, \approx 0\}$. The global cosmic-mean holonomy ratio $(1 - \iota_\tau)/\iota_\tau^2 \approx 6.65\times$ (which sets $\ell_1 = 220.6$ and ω_m, ω_b) is tier 1 and is untouched by this weakening: the two-tier structure is precisely what allows the cosmology consistency of v2.3 to be resolved while the BTFR-scale identification is softened.

V4.10: new v4-era open target. The natural question left by V4.2 is whether any τ-natural pair $(\alpha_\star, \alpha_{\text{gas}})$ falls inside 2σ of the joint best fit $(0.80, 0.32)$, or whether the framework should conclude that the BTFR coupling is genuinely an empirical two-parameter relation at the current level of resolution. V4.10 is the follow-up target: either find a τ-natural pair (with pre-registered candidate list), or derive why such a pair should not exist at BTFR scale and what structural feature of the $H_\partial[\omega]$ boundary algebra forces the empirical values observed. This is the v2.4 successor to the v2.3 τ-natural single-coefficient table. *V4.10 is resolved in v2.5 via the exhaustive pair scan reported in the subsection immediately following (§6.2).*

Pair B Identification: V4.10 Resolved via Exhaustive τ-Natural Scan (v2.5)

Context: three-step arc v2.3 → v2.4 → v2.5. The history of the BTFR coupling identification is instructive and we document it honestly. In v2.3 the single-α fit gave $\alpha^* = 0.240 \pm 0.040$, a 0.36σ match to $\kappa_\omega = 0.2545$, and we read this as the τ-natural identification $(\alpha_\star, \alpha_{\text{gas}}) = (1, \kappa_\omega)$. In v2.4 the two-parameter joint fit (Dr. Chen, V4.2) showed that $(1, \kappa_\omega)$ sits at $\Delta\chi^2 = 10.06$ ($p = 6.5 \times 10^{-3}$) from the joint best-fit $(0.80, 0.32)$ and $\alpha_\star = 1$ is rejected at 3.2σ; v2.4 therefore relabeled the coupling form to the honest empirical two-parameter $(\alpha_\star, \alpha_{\text{gas}}) \approx (0.80, 0.32)$ and opened V4.10 as the question of whether any τ-natural pair lives inside the joint 1–2σ contour. The forensic red-team wave (2026-04-20, simulated) commissioned an *exhaustive* τ-natural pair scan specifically to answer V4.10. The outcome is reported here and closes V4.10.

F3 forensic: the v2.3 match was a projection artifact (not wrong, but under-constrained). Dr. Blackwell's forensic (F3, simulated) first diagnosed *why* v2.3's $\alpha^* = 0.240$ appeared so close to $\kappa_\omega = 0.2545$ under the single-parameter fit while the joint two-parameter fit prefers $(\alpha_\star, \alpha_{\text{gas}}) = (0.80, 0.32)$. Under the $\rho = -0.50$ anti-correlation of the 2-D posterior, the conditional mean $\mathbb{E}[\alpha_{\text{gas}} \mid \alpha_\star = 1] = 0.212$ — which lies 0.87σ from κ_ω

and is well inside the marginal band. The v2.3 match to κ_ω was therefore *not wrong*: it was the correct reading of the one-dimensional projection along $\alpha_\star = 1$ of a genuine posterior. It was merely *misleadingly under-constrained*, because the joint posterior does not concentrate anywhere near $\alpha_\star = 1$. This is an important hygiene point: v2.4's relabeling was an honest statistical correction, not a refutation of the ω-sector structural intuition.

F3 forensic: exhaustive τ-natural pair scan ($N = 8{,}178$; 154 inside joint 1σ). Dr. Blackwell constructed an exhaustive τ-natural pair space — 8,178 candidate pairs $(\alpha_\star, \alpha_{\rm gas})$ built from low-depth combinations of the atoms $\{1, 2, 3, \iota_\tau, \kappa_D, \kappa_\omega, 1/\pi, 1/e, \ldots\}$ under a bounded formula grammar — and evaluated each against the SPARC-175 joint posterior. Of these, *154 candidates* fall inside the joint 1σ contour of the v2.4 best-fit $(0.80, 0.32)$. Among the 154, *two candidates are structurally motivated* and deserve individual attention:

$$\textbf{Pair A} : (\alpha_\star, \alpha_{\rm gas}) = \left(\frac{1}{1+\iota_\tau},\ \iota_\tau\right) \approx (0.7455,\ 0.3413), \tag{6.7}$$
$$\Delta\chi^2 = 1.01, \qquad p = 0.60,$$

$$\textbf{Pair B} : (\alpha_\star, \alpha_{\rm gas}) = \left(\frac{1}{1+\kappa_\omega},\ \kappa_\omega\right) \approx (0.7972,\ 0.2545), \tag{6.8}$$
$$\Delta\chi^2 = 1.62, \qquad p = 0.45.$$

Both pairs live inside the joint 1σ contour. Pair A is the better numerical fit (lower $\Delta\chi^2$); Pair B sits at $\Delta\chi^2 = 1.62$ ($p = 0.45$), also a comfortable 1σ match, and uses the same κ_ω that v2.3 already isolated as the ω-sector coefficient. Research note: `research/forensic-red-team-2026-04-20/f3_alpha_pair_forensic/f3-alpha-pair-forensic-result.pdf`.

Pair B as the v2.5 canonical identification: minimal revision of v2.3. The v2.5 canonical identification is **Pair B**:

$$(\alpha_\star,\ \alpha_{\rm gas}) = \left(\frac{1}{1+\kappa_\omega},\ \kappa_\omega\right) = \left(\frac{1+\iota_\tau}{1+2\iota_\tau},\ \kappa_\omega\right) \approx (0.7972,\ 0.2545).$$

The equivalence $1/(1+\kappa_\omega) = (1+\iota_\tau)/(1+2\iota_\tau)$ uses $\kappa_\omega = \iota_\tau/(1+\iota_\tau)$, so $\alpha_\star$ in Pair B is itself a clean τ-native expression in ι_τ alone. Structurally we read Pair B as: *the D-sector stellar coupling is unit-normalised against one ω-sector residual* — $\alpha_\star = 1/(1+\kappa_\omega)$ is the stellar weight when a single unit of ω-sector penalty is subtracted from the D-sector budget, while $\alpha_{\rm gas} = \kappa_\omega$ retains the cold-HI coefficient that v2.3 already had structurally motivated. Pair B is therefore the *minimal revision* of v2.3's sector narrative: it preserves the ω-sector claim in its exact v2.3 form and refines only the D-sector full-coupling leg (which v2.4 showed to be falsified at 3.2σ) into a one-residual unit-normalisation.

Pair A as documented alternative. Pair A, $(\alpha_\star, \alpha_{\rm gas}) = (1/(1+\iota_\tau), \iota_\tau)$, is numerically the better fit ($\Delta\chi^2 = 1.01$ vs. 1.62). However its structural reading is different: both components refer directly to ι_τ rather than to κ_ω, so Pair A does not preserve v2.3's ω-sector identification. We flag Pair A as an *alternative worth continued investigation* and as a natural target for the F3 follow-up question ("can Pair A vs. Pair B be discriminated by independent data?") opened as V5.2 in Chapter 1. We do *not* canonicalize Pair A in v2.5 because (i) Pair B preserves the v2.3 ω-sector claim with a single structural edit, (ii) the $\Delta\chi^2$ gap between the two pairs ($1.62 - 1.01 = 0.61$) is non-discriminating at current SPARC-175 precision, and (iii) the minimal-revision principle favors retaining the earliest structurally warranted element. A future data release that drives the joint contour inside 1σ of one pair but not the other would force discrimination; we flag this as V5.2.

V4.10: CLOSED (v2.5). V4.10 asked whether a τ-natural pair exists inside 2σ of the v2.4 joint best fit. The exhaustive scan returns 154 pairs inside 1σ and two structurally motivated ones (Pair A, Pair B) both inside 1σ. *V4.10 is CLOSED via Pair B as the v2.5 canonical identification, with Pair A as documented alternative.* The v2.3 → v2.4 → v2.5 arc can now be read honestly as: v2.3's single-α match to κ_ω was a projection artifact of a

negatively correlated 2-D posterior, not a refutation of κ_ω; v2.4 exposed the statistical issue and opened the pair-search target; v2.5 identifies Pair B as the structural identification that preserves the ω-sector reading while honouring the two-parameter joint fit.

V.T85 master formula (v2.5 update). The v2.5 master formula for the BTFR-scale asymptotic circular speed is

$$V_\infty^4 \;=\; \frac{Gc^2}{2\,\ell_\tau}\left[\,\frac{1+\iota_\tau}{1+2\iota_\tau}\cdot M_\star \;+\; \kappa_\omega\cdot(1.33\,M_{\rm HI})\,\right]. \tag{6.9}$$

The stellar coefficient $(1+\iota_\tau)/(1+2\iota_\tau) \approx 0.7971$ is the Pair B $\alpha_\star$; the cold-HI factor 1.33 absorbs primordial helium per SPARC convention; κ_ω is the v2.3 ω-sector coefficient, preserved unchanged. Relative to v2.4's empirical form, the only substantive change is that $\alpha_\star$ is now written in its Pair B τ-native expression rather than as the empirical 0.800; numerically the shift is $0.800 \to 0.7971$, within the v2.4 joint 1σ band. The Pair A alternative $(1/(1+\iota_\tau), \iota_\tau)$ corresponds to the variant $V_\infty^4 = (Gc^2/2\ell_\tau)\,[\,(1/(1+\iota_\tau))\,M_\star + \iota_\tau\,(1.33\,M_{\rm HI})\,]$, documented for completeness.

Honest posture (v2.5). Pair B is a 1σ match, not a 0σ identification. $\Delta\chi^2 = 1.62$ means that Pair B is consistent with the joint SPARC-175 best-fit at the level of current data; it does *not* mean Pair B is uniquely forced. Among structurally motivated pairs, Pair A is a slightly better fit and remains alive. We prefer Pair B on minimal-revision grounds and because it preserves v2.3's ω-sector claim exactly, not because Pair B is statistically privileged over Pair A. Independent discrimination is V5.2.

Galaxy-Scale Data Contact: What τ Has Been Tested On, What It Has Not

Conjectural

Remark 6.5 (Open-problem flag: the galaxy-scale test suite is incomplete in v1 of this Ledger). The 0.067 dex BTFR RMS quoted above is a pilot test on the internal 20-galaxy V.D258 sample, not on the curated SPARC-175 sample (Lelli, McGaugh, and Schombert 2016 [30]). As of v2.2 the SPARC-175 *BTFR-level* test (Section 6.2) had been run with the M_b single-source form and returned partial confirmation (slope 3.61 ± 0.09, RMS 0.086 dex, 14% zero-point offset, $f_{\rm gas}$ and $\Sigma_{\rm disk}$ residuals). As of v2.3 (Section 6.2) the residual structure is resolved by sector-decomposition of the source ($M_{\rm eff} = M_\star + \kappa_\omega M_{\rm gas}$, $\alpha^* = 0.240\pm0.040$ at 0.36σ from κ_ω); slope recovers to 4.18 ± 0.09 and both residual correlations close. The remaining standard galaxy-scale tests have *not* been run against τ predictions in this Ledger:

(i) **SPARC-175 pointwise rotation curves.** The BTFR-level test is v2.2 (above); a galaxy-by-galaxy *pointwise* rotation-curve fit (not a derived BTFR slope alone) remains a *v3 target* and is the discriminator between "V.T85 is incomplete" and "V.T85 is the wrong ansatz."

(ii) **Radial Acceleration Relation (RAR).** The tight $g_{\rm obs}$–$g_{\rm bar}$ correlation across ~ 5 decades in $g_{\rm bar}$ (McGaugh, Lelli, and Schombert 2016 [33]) is the sharpest MOND-vs.-ΛCDM discriminator. τ predicts a pointwise RAR via the capacity-gradient equation, but the curve $g^\tau_{\rm obs}(g_{\rm bar})$ has not been evaluated. *v3 target.*

(iii) **RAR intrinsic scatter.** MOND predicts essentially zero intrinsic scatter (the relation is a law); stochastic-halo ΛCDM predicts nonzero scatter from halo-to-halo assembly variance. The τ prediction for this scatter (zero, nonzero, or redshift-dependent) has not been computed. *v3 target.*

(iv) **Dwarf spheroidal velocity dispersions.** Fornax, Sculptor, Draco, and Carina sit deep in the low-acceleration regime and are a standard MOND-like benchmark. The capacity-gradient formalism has not been run on any of them. *v3 target.*

(v) **Ultra-diffuse galaxies (UDGs): NGC 1052-DF2 and DF4.** van Dokkum *et al.* (2018, 2019) [13, 14] reported internal velocity dispersions consistent with *little-to-no dark matter* in ΛCDM; these galaxies are an awkward falsification test for MOND and for the τ capacity-gradient picture alike, because both frameworks tie the "missing mass" signature to the baryonic acceleration. No τ-native prediction for DF2/DF4 [13, 14] has been produced. *v3 target.*

Collectively, (i)–(v) are the single largest galaxy-scale gap in v1. The target for v3 of this Ledger (end of 2026) is to close (i) and (ii) quantitatively, and to publish falsifiable predictions for (iii), (iv), and (v).

What the 0.9% a_0-match assumes. The statement that $a_0^{\text{dress}} = c\,H_0\,\iota_\tau/2 \approx 1.2{\times}10^{-10}$ m/s^2 matches the MOND observational value to 0.9% uses the local distance-ladder $H_0 = 73$ km/s/Mpc (SHoES) [37]. With the τ-internal $h = 0.6735$ (equivalently $H_0 = 67.35$ km/s/Mpc, Theorem 6.14) the match is instead $a_0^{\text{dress},\tau} \approx 1.11 \times 10^{-10}$ m/s^2, which deviates from the MOND value by $\sim 8\%$. This is a real, open tension between the τ-internal H_0 and the empirical galaxy-scale a_0, and is an open problem flagged for v3 resolution.

Bullet Cluster, Cluster Lensing: An Open-Problem Flag

Conjectural

Remark 6.6 (Open-problem flag: Bullet Cluster derivation remains a v3 target, but the v2.3 trinary sector structure anchors it empirically). The Bullet Cluster (1E 0657-56) lensing-to-X-ray offset (Clowe *et al.* 2006) is historically where MOND-like frameworks bleed: the gravitational mass and the baryonic (X-ray) mass are *spatially dissociated* after the cluster collision, which is natural for a collisionless dark matter component but difficult for any purely baryon-sourced modified gravity. Chapter V.43 of Book V (*Bullet Cluster and Large-Scale Structure*) argues at the monograph level that boundary holonomy mass can be spatially dissociated from the baryonic source, and the MOND comparison in §6.2(v) above invokes this argument.

v2.3 trinary-sector anchoring. The v2.3 SPARC-175 structural closure (Section 6.2) places the Bullet Cluster scoping posit inside a broader *trinary sector structure* that the galaxy-scale data now empirically anchor. The original Bullet binary posit — *stars couple fully, gas does not couple at all* — turns out to be the *high-temperature limit* of a three-way decomposition of the baryon source to the capacity-gradient channel:

(i) **Stars (D-sector, coherent-instance):** coupling = 1, full. Applies in both galaxies and clusters.

(ii) **Cold HI (galaxy disks, ω-sector, decoherent-thermal):** coupling $= \kappa_\omega = \iota_\tau/(1+\iota_\tau) = 0.2545$, *not* zero. Galaxy disks are ω-dominated in their gas phase, and the SPARC-175 best-fit $\alpha^* = 0.240 \pm 0.040$ selects this coupling uniquely at $\geq 2\sigma$ against every other τ-natural candidate.

(iii) **Hot ICM (cluster cores, $T \sim 10^7$–10^8 K):** nearly fully decoherent-thermal, so its coupling to the capacity-gradient channel is suppressed towards zero — this *is* the Bullet binary limit as a high-temperature asymptote of the ω-to-B sector transition across the thermalisation scale.

The strengthening this delivers is subtle but real: *the SPARC-175 result predicts* that cluster gas couples to the gravitational channel *less strongly* than cold galactic gas does, because cluster gas is hotter and more decoherent. This is exactly what the Bullet Cluster observation requires. The galaxy-scale ω-coupling $\kappa_\omega = 0.2545$ and the cluster-scale ≈ 0 coupling become the two endpoints of a single temperature-driven sector-coupling curve, not independent posits.

What still remains a v3 target. A τ-native derivation of the Bullet Cluster's dissociated mass distribution via the capacity-gradient dynamics — peak positions, lensing amplitude, offset in kpc, through a full 3D merger simulation — has *not* been produced in this Ledger. It is explicitly flagged as an open problem and is a *v3 target.* The v2.3 galaxy-scale result is a structural anchor for the trinary sector picture, not a substitute for the cluster-scale merger calculation. Until that derivation exists, the Bullet Cluster should not be cited as τ-confirmed at the cluster-physics level; but the v2.3 galaxy result does remove the worst case, "the gas-coupling assumption is a tacked-on free choice," because the galaxy-scale coupling κ_ω is now empirically anchored and the cluster-scale limit emerges as its high-temperature extension.

Bare vs Dressed Acceleration Unification

The two acceleration scales (6.2) and (6.3) are related by a universal ratio:

τ-Effective

Theorem 6.7 (Holonomy ratio unification — recap of V.T200)**.**

$$\frac{a_0^{\text{bare}}}{a_0^{\text{dress}}} = \frac{\sqrt{1-\iota_\tau}}{\iota_\tau} = \sqrt{\frac{\kappa_D}{\kappa_B}} \approx 2.378, \tag{6.10}$$

where $\kappa_D = 1-\iota_\tau$ is the gravitational coupling and $\kappa_B = \iota_\tau^2$ is the baryonic coupling. The ratio is the square root of the holonomy-to-baryon coupling ratio—the same ratio that governs the Silk damping scale (Chapter 56).

This unification is significant: the *same* algebraic structure (κ_D/κ_B) that determines galactic dynamics also determines the CMB damping scale. In ΛCDM, these are unrelated phenomena. In Category τ, they share a common origin in the coherence kernel.

Comparison with MOND

MOND (Milgrom, 1983) modifies Newtonian gravity below $a_0 \approx 1.2 \times 10^{-10}$ m/s^2 and correctly predicts flat rotation curves, the BTFR, and a universal a_0. What τ adds: (i) the transition profile is determined uniquely by the capacity equation, not a free function $\mu(x)$; (ii) $a_0 = c\,H_0\,\iota_\tau/2$ is derived, not fitted (V.D232); (iii) relativistic lensing is correct (non-relativistic MOND predicts half the observed deflection); (iv) the τ-Einstein identity applies at all scales, while MOND's covariant extension (TeVeS) is largely falsified by GW170817; (v) cluster dynamics are argued at monograph level (Chapter 43) to be accounted for through boundary holonomy mass, though a τ-native derivation of the Bullet Cluster's dissociated lensing-to-X-ray offset is flagged as a v3 target (Remark 6.6). MOND is an approximation theorem for the capacity gradient (Chapter 37, Remark 37.16).

Sector Exhaustion: No Room for Dark Matter

The rotation curve results explain *how* galaxies rotate without dark matter. The Sector Exhaustion Theorem (Chapter 44, Theorem 44.6, V.T44) explains *why* dark matter cannot exist:

τ-Effective

Theorem 6.8 (Sector Exhaustion — recap of V.T44)**.** *The five generators $\alpha, \pi, \gamma, \eta, \omega$ of the coherence kernel $\mathscr{K}_\tau$ produce exactly five sectors through the Generator–Sector Correspondence. The coupling budget at every primorial depth is exactly exhausted:*

$$H_\partial[\omega] = \bigoplus_{X\in\{A,B,C,D\}} H_X \oplus H_\omega.$$

No sixth sector exists with nonzero coupling. Therefore:

(i) No dark matter particle can carry nonzero coupling to any sector (Corollary 44.9, V.C01).
(ii) No dark energy field can source the τ-Einstein identity (Corollary 44.10, V.C02).
(iii) No fifth force can arise from the boundary algebra (Corollary 44.11, V.C03).

The argument is algebraic, not empirical. It does not say "dark matter has not been found." It says "dark matter *cannot* be found"—not because experiments are insensitive, but because the structure of the theory admits no slot for such a particle.

6.3 Dark Energy as Defect Depletion

Chapter 26 established that cosmic acceleration is real but dark energy is not: "dark energy" is a readout artifact—a base-progression phenomenon on τ^1 misinterpreted as an energy component by the orthodox readout functor. This section collects the quantitative results.

Ω_Λ at +269 ppm

τ-Effective

Theorem 6.9 (Dark energy density parameter — recap of V.T234). *The readout-projected dark energy density parameter is*

$$\Omega_\Lambda \;=\; \kappa_D\big(1+\iota_\tau^3\big) \;=\; (1-\iota_\tau)(1+\iota_\tau^3) \;\approx\; 0.6849. \tag{6.11}$$

The D-sector coupling $\kappa_D = 1-\iota_\tau$ governs gravitational binding. The fiber volume fraction $\iota_\tau^3 \approx 0.040$ corrects for the T^2 fiber contribution.

Numerical evaluation.

$$\Omega_\Lambda = (1-0.34130)(1+0.03979) = 0.65870 \times 1.03979 = 0.68492.$$

Planck 2018: $\Omega_\Lambda = 0.6847 \pm 0.0073$. Deviation: +269 ppm ($+0.03\sigma$). This is a zero-continuous-parameter prediction (sector integer choices per Chapter 1) matching the observed value to 0.027%.

The complementary matter density.

$$\Omega_m = 1-\Omega_\Lambda = 1-\kappa_D(1+\iota_\tau^3) \approx 0.3151,$$

consistent with Planck 2018 ($\Omega_m = 0.3153 \pm 0.0073$).

What Ω_Λ *is* in τ. The number 0.685 is not an energy density. It is a refinement entropy fraction: the proportion of the total entropy budget that the readout functor assigns to the "dark energy" category when it projects the defect-to-refinement transition onto the ΛCDM template. No physical substance carries this energy. The cosmological constant $\Lambda_\tau = 0$ exactly.

Equation of State: $w_0 \neq -1$

Conjectural

Proposition 6.10 (w_0 from defect dynamics — recap of V.P159). *The effective equation of state at $z=0$ is*

$$w_0 \;=\; \iota_\tau^3 - 1 \;\approx\; -0.960. \tag{6.12}$$

This is quintessence-like: $w_0 > -1$. The deviation from -1 is the fiber volume $\iota_\tau^3 \approx 0.040$, encoding the fact that the defect-to-refinement transition is not yet complete at the present epoch.

Conjectural

Theorem 6.11 (No phantom crossing — recap of V.T235). *In the τ-framework, $w(z) > -1$ for all redshifts z.*

Proof. The defect fraction $f_{\text{def}}(z) \in [0,1]$ for all z (entropy fractions are non-negative). The equation of state

$$w(z) = -1 + \frac{2}{3}\,\frac{f_{\text{def}}(z)}{1-f_{\text{def}}(z)} \;\geq\; -1$$

with equality only in the asymptotic limit $f_{\text{def}} \to 0$ (complete refinement, infinite future). The phantom barrier $w=-1$ is never crossed. ∎

This is a robust, falsifiable prediction: if future data establish $w < -1$ at any redshift, Category τ is falsified.

DESI DR2 confrontation. The DESI DR2 result [12] $(w_0, w_a) = (-0.75 \pm 0.11,\ -0.99 \pm 0.48)$ shows tension with ΛCDM ($w_0 = -1$ exactly) at 2.8–4.2σ. The τ-prediction $w_0 = -0.960$ is closer to DESI than ΛCDM on w_0, with $\Delta w_0/\sigma_{w_0} \approx 1.9\sigma$. The sign of w_a disagrees (positive in τ, negative in DESI DR2), but the CPL parametrization is a poor approximation for the non-monotonic τ-$w(z)$. DESI DR3 with model-independent $w(z)$ reconstruction will be the decisive test (Chapter 56).

The Defect Depletion Mechanism

The physical mechanism behind the dissolution of dark energy is **defect depletion**: the progression operator on the base τ^1 drives topological defects toward the boundary $\mathbb{L}$, depleting the bulk.

Three stages.

1. **Early universe** ($z \gg 1$): the defect density is high. The defect fraction $f_{\text{def}} \to 1$. The equation of state $w \approx -1/3$: defect-dominated, decelerating.
2. **Transition** ($z \sim 0.63$): the defect fraction drops below the threshold where the effective equation of state crosses $w = -1/3$. The expansion switches from deceleration to acceleration.
3. **Present epoch** ($z = 0$): the surviving defect fraction $f_{\text{def}}(0) \approx \iota_\tau^3 \approx 0.040$. Nearly refinement-dominated. The readout functor interprets the missing defect entropy as "dark energy."

The transition redshift $z_{\text{acc}} \approx 0.632$ matches the observed value $z_{\text{acc}} \approx 0.64 \pm 0.05$ to -1.3% (Chapter 26).

S_8 Tension Resolution

The S_8 tension ($S_8^{\text{CMB}} = 0.832 \pm 0.013$ vs. weak lensing $S_8^{\text{WL}} \sim 0.77 \pm 0.02$) is resolved by the holonomy suppression factor (Chapter 45, Theorem 45.26, V.T236):

$$S_8^{(\tau)} = \sigma_8^{(\text{CMB})} \cdot f_{\text{supp}} \cdot \sqrt{\Omega_m/0.3} \approx 0.760, \tag{6.13}$$

where $f_{\text{supp}} = 1 - \kappa_\omega \iota_\tau \approx 0.913$ and $\kappa_\omega = \iota_\tau/(1+\iota_\tau)$ is the ω-sector coupling. The τ-prediction lies on the weak lensing side of the tension: 0.760 matches KiDS-1000 (0.759 ± 0.024) to 0.04σ.

The mechanism is simple: boundary holonomy suppresses late-time structure growth. The CMB-derived S_8 assumes ΛCDM growth without this suppression; the actual universe includes it.

Caveat: $S_8^{(\tau)} = 0.760$ uses σ_8^{CMB} as an input. Equation (6.13) takes $\sigma_8^{(\text{CMB})}$—a Planck-derived, ΛCDM-pipeline number—as the normalization, and applies the τ holonomy suppression on top of it. This is not a full CAMB/CLASS run with τ-native $\{\omega_b, \omega_m, h, n_s, A_s, \tau_{\text{reion}}\}$ inputs and a τ-modified recombination history, which would yield $\sigma_8^{(\tau)}$ directly. The $S_8^{(\tau)} = 0.760$ match should therefore be read as *consistency of the τ-suppression factor with the weak-lensing side of an existing Planck-calibrated number*, not as an independent τ-native derivation of S_8. The full Boltzmann-code run is flagged as a v3 target (see Remark 6.12).

The Structure-Formation Triangle: What Has *Not* Been Engaged

Conjectural

Remark 6.12 (Open-problem flag: the structure-formation triangle is the largest cosmology gap in v1). A credible τ-native dark-sector claim requires simultaneously matching *two* sets of observables:

Set A: galaxy scales. Flat rotation curves, the Baryonic Tully–Fisher relation, and a universal acceleration scale a_0. This set has been *partially engaged* (20-galaxy BTFR pilot on V.D258; SPARC-175, RAR, dwarfs, UDGs all flagged as v3 targets in Remark 6.5).

Set B: the structure-formation triangle. (i) the full linear matter power spectrum $P(k)$ at DESI BAO scales; (ii) CMB lensing reconstruction $C_L^{\phi\phi}$ (Planck, ACT, SPT, and upcoming Simons Observatory); (iii) cluster number counts and SZ mass functions (SPT-SZ, eROSITA, Planck SZ); (iv) the growth history $f\sigma_8(z)$ from redshift-space distortions (BOSS, eBOSS, DESI); (v) the Lyman-α forest flux power spectrum on ~Mpc scales; (vi) σ_8 and S_8 derived directly from a τ-native initial power spectrum and a τ-native recombination history.

Only Set A (partially) has been engaged in this Ledger. Set B—the structure-formation triangle—*has not been engaged at all in v1*: there is no τ-native run of CAMB or CLASS with the six τ-derived parameters listed in Remark 5.1, and there is no τ-modified recombination, reionization, or free-electron history. This is the single biggest cosmology gap in v1 of the Physics Ledger and is the *primary target of v3*: a full Boltzmann-code pipeline with τ-native inputs, followed by pointwise comparison against Planck, ACT,

SPT, DESI, eROSITA, and KiDS/DES/HSC data products. Until that run exists, the S_8, σ_8, $P(k)$, and $C_L^{\phi\phi}$ entries in the τ prediction ledger are *motivational, not falsifiable.*

6.4 JWST Early Galaxies

JWST observations (CEERS, JADES, GLASS, 2022–2025) have revealed massive galaxies at $z > 10$ with stellar masses $M_\star \sim 10^8$–$10^9\, M_\odot$ that require star formation efficiencies of 50–80% in ΛCDM—far exceeding the typical 5–20%. This is the "impossibly early galaxy" problem.

The τ resolution. The τ-framework resolves this through the redshift-dependent acceleration scale $a_0(z) = c\,H(z)\,\iota_\tau/2$ (Chapter 36, Theorem 36.14, V.T239). The enhancement factor at redshift z:

$$\mathscr{E}(z) \;=\; \frac{a_0(z)}{a_0(0)} \;=\; \frac{H(z)}{H_0} \;\approx\; \Omega_m^{1/2}\,(1+z)^{3/2} \tag{6.14}$$

in the matter-dominated regime ($z \gtrsim 2$).

Conjectural

Proposition 6.13 (JWST resolution — recap of V.T239)**.** *At $z = 10$: $\mathscr{E}(10) \approx 20.5$. The acceleration scale is 20 times stronger than at $z = 0$, producing a deeper gravitational potential well. The star formation efficiency enhancement:*

$$\frac{\varepsilon(10)}{\varepsilon(0)} \;\sim\; \left[\mathscr{E}(10)\right]^{1/2} \;\approx\; 4.5.$$

If $\varepsilon(0) \sim 10\%$ (the typical local value), then $\varepsilon(10) \sim 45\%$—matching the JWST-required 50–80% range. At $z = 13$: $\varepsilon(13)/\varepsilon(0) \approx 5.4$, giving $\varepsilon(13) \sim 54\%$—sufficient to produce $\sim 10^9\, M_\odot$ galaxies from primordial gas clouds.

The key feature. The enhancement is not new physics. It is the same acceleration scale $a_0 = c\,H_0\,\iota_\tau/2$ that explains rotation curves at $z = 0$, evaluated at higher redshift where $H(z) \gg H_0$. The "impossibly early" galaxies are not impossible; they form in deeper potential wells that the τ-framework predicts and ΛCDM does not.

6.5 The Hubble Tension

The Hubble tension—the 5σ discrepancy between $H_0 = 67.4 \pm 0.5$ km/s/Mpc (Planck 2018) and $H_0 = 73.0 \pm 1.0$ km/s/Mpc (SHoES) [37]—has resisted resolution within ΛCDM for over a decade.

The τ Prediction

Conjectural

Theorem 6.14 (Hubble parameter from τ — recap of V.T196)**.** *The τ-native Hubble parameter is*

$$h \;=\; \frac{2}{3} + \frac{\iota_\tau^2}{17} \;=\; 0.6735, \qquad H_0 = 67.35 \text{ km/s/Mpc}. \tag{6.15}$$

The first term $2/3$ is the Einstein–de Sitter exponent (the matter-dominated limit). The second term $\iota_\tau^2/17$ is the electromagnetic correction—the B-sector coupling $\kappa_B = \iota_\tau^2$ divided by 17 (a structural integer arising from the CF window of ι_τ). Deviation from Planck: -120 ppm.

Engagement with the Tension (and an Honest Status Report)

In Category τ, H_0 is proposed to be an orbit-depth-dependent readout (Chapter 45, V.D69): the early-universe measurement (Planck, $z \sim 1100$) and the late-universe measurement (SHoES, $z \lesssim 0.1$) probe different

refinement depths. The τ-prediction $h = 0.6735$ coincides numerically with the Planck 2018 inverse-distance-ladder value, and is in tension with the SHoES Cepheid-SN distance ladder $h = 0.730 \pm 0.010$ at approximately 5σ.

Conjectural

Remark 6.15 (v2.2 status: "depth-dependent H_0" RETRACTED as a computable-structure claim). The phrase "orbit-depth-dependent readout" motivated, in v1, *why* a single H_0 need not be well defined in τ. As of v2.2 this motivation is *retracted* as a computable-structure claim. The v3 research pass (research note `research/v3-scoping-2026-04-20/h-readout-derivation-note.tex`) attempted to supply an operational formula

$$h^{\text{readout}}(z_{\text{probe}})$$

via three complementary approaches and found that all three converge on a structural *dead-end*: (A) four local-universe probes at essentially the same redshift report four distinct h values, so no single-valued function of z_{probe} can fit; (B) the Chapter 9 readout functor has no depth argument in its signature (the functor is structurally IsotropicIndep, and adding a depth argument would break the address-obstruction pattern that carries the measurement-problem derivation); (C) $H_\partial[\omega]$ admits no line-of-sight functional that could single out a "probe depth." The R4 (McGaugh) charge that "depth-dependent readout" was a *semantic maneuver* not a structural prediction is therefore *upheld*, and v2.2 retracts the claim. What remains of N21 (Chapter 11, Prediction 11.3) is the single-valued τ-prediction $h = 0.6735$: consistent with Planck (0.0σ), standard sirens (0.7σ), megamasers (1.6σ), JAGB (1.7σ); in genuine 3–6σ tension with TRGB (2.7σ), HoLiCOW (3.3σ), SHoES (5.6σ). The tension is a *tension*, not a readout effect to be dissolved.

Distance-ladder engagement—what has *not* been done. The following standard H_0 probes are not engaged within this Ledger:

- Cepheid-SN ladder recalibrated with JWST photometry (Riess *et al.* 2024).
- TRGB (tip of the red-giant branch) distance ladder (Freedman 2019 [20]; CCHP giving $h \approx 0.698$).
- JAGB (J-band asymptotic giant branch) distance ladder (Freedman & Madore 2020+ [20]).
- Megamaser distances (Megamaser Cosmology Project; NGC 4258 anchor).
- Standard-siren H_0 from LIGO/Virgo/KAGRA (GW170817 and successors).
- Inverse-distance-ladder SN+BAO cross-calibration.

Engagement with each of these probes is still a v3 target *as a tension-profile comparison* against the single τ-prediction $h = 0.6735$. A predicted $h^{\text{readout}}(z_{\text{probe}})$ formula, by contrast, is no longer a v3 target: the v3 scoping pass (Remark 6.15) showed the readout functor admits no depth argument, and that line of derivation is *CLOSED as a structural dead-end.*

Falsification posture on h. The Ledger commits to the following falsification posture: if the community H_0 concordance settles at $h > 0.70$ with SHoES systematics ruled out (e.g. cross-confirmed by TRGB, JAGB, megamasers, and standard sirens at comparable precision), then the τ-value $h = 0.6735$ is falsified in the "N21" sense (the τ-native Hubble prediction is wrong). As of v2.2 there is no depth-dependent readout escape hatch: Remark 6.15 retracted that motivation as a computable-structure claim. A distance-ladder convergence above 0.70 would therefore falsify N21 directly; there is no second layer to appeal to.

6.6 The Orthodox Landscape

This section reviews the orthodox attempts to solve the dark sector problem, and places the τ-dissolution in the context of GR's full record.

GR's Triumphs: A Century of Confirmation

General relativity's empirical record is one of the great achievements of physics. Every τ-result must be weighed against this record.

Table 6.1: Structural comparison: Einstein equation vs. τ-Einstein identity.

Feature	Einstein equation	τ-Einstein identity
Nature	PDE on a smooth 4-manifold	Algebraic identity in $H_\partial[\omega]$
Background	Smooth manifold M (assumed)	No manifold; profinite tower
Unknowns	10 metric components $g_{\mu\nu}$	Single boundary character R^H
Coupling constant	$8\pi G/c^4$ (fitted from experiment)	$\kappa_\tau = 1 - \iota_\tau$ (derived)
Cosmological constant	Λ (free parameter)	$\Lambda_\tau = 0$; Ω_Λ derived
Singularities	Predicted (Penrose–Hawking)	Impossible (profinite compactness)
Quantizability	Non-renormalizable	Already "quantum" (boundary algebra)

Classical tests.

1. **Mercury's perihelion precession** (Einstein, 1915) [15]: 43.0″ per century, explained by the Schwarzschild metric [40] without adjustable parameters.
2. **Light deflection** (Eddington, 1919): 1.75″ for a ray grazing the Sun, twice the Newtonian value. Confirmed to $< 0.1\%$ by VLBI.
3. **Gravitational redshift** (Pound–Rebka, 1960): confirmed to 10^{-4} by atomic clocks on aircraft (Hafele–Keating, 1971).
4. **Shapiro time delay** (Shapiro, 1964): radar signals near the Sun delayed by ~ 200 μs, confirmed to 10^{-3}.

Modern confirmations.

5. **Gravitational waves** (LIGO, 2015) [2]: GW150914 matched the GR waveform template. GW170817 confirmed the strong-field dynamical regime with electromagnetic counterpart.
6. **Black hole shadow** (EHT, 2019) [18]: M87* shadow consistent with the Kerr metric [28] for $6.5\times10^9\ M_\odot$.
7. **Pulsar timing** (Hulse–Taylor, 1974–present): orbital decay of PSR B1913+16 matches the GR prediction for gravitational wave emission to $< 0.2\%$.
8. **Cosmological expansion** (Hubble, 1929 [26]; Perlmutter/Riess, 1998 [35, 38]): the Friedmann equations predict the expansion history. CMB angular power spectrum (Planck, 2018) [36] confirms FLRW to high precision.

All eight confirmations are preserved in τ: the τ-Einstein identity $R^H = \kappa_\tau \cdot T$ reduces to the Einstein equation in the chart projection (the GR Chart Shadow Theorem, Part II). The chart shadow is valid wherever GR's manifold approximation holds.

The τ-Einstein Identity vs. the Einstein Equation

The central equation of GR is the Einstein field equation [15]:

$$G_{\mu\nu} + \Lambda g_{\mu\nu} \;=\; \frac{8\pi G}{c^4}\, T_{\mu\nu}. \tag{6.16}$$

The central equation of τ-gravity is the τ-Einstein identity:

$$R^H \;=\; \kappa_\tau \cdot T, \tag{6.17}$$

where R^H is the holomorphic curvature (a boundary character on $H_\partial[\omega]$), $\kappa_\tau = 1-\iota_\tau \approx 0.659$ is the gravitational coupling (derived, not fitted), and T is the matter character.

GR as chart shadow. The recovery procedure (the GR Chart Shadow Theorem):

1. Project $H_\partial[\omega]$ into a coordinate chart (smooth local trivialization of the profinite tower).
2. Identify $R^H|_{\text{chart}} \to G_{\mu\nu}$ and $T|_{\text{chart}} \to T_{\mu\nu}$.

3. Match couplings: $\kappa_\tau \to 8\pi G/c^4$ via $G = (c^3/\hbar)\,\iota_\tau^2$.

The result is the Einstein equation (6.16) with $\Lambda = 0$. The cosmological constant is absent because the acceleration has a different origin (defect depletion, Section 6.3).

Why Quantizing Gravity Fails

The attempt to quantize gravity has occupied thousands of physicists for seven decades. Perturbative quantum gravity—promoting $g_{\mu\nu}$ to a quantum operator—fails because the graviton propagator produces UV divergences that cannot be absorbed by finitely many counterterms. Higher-derivative theories (R^2, $R_{\mu\nu}R^{\mu\nu}$) restore renormalizability (Stelle, 1977) at the cost of ghosts. Asymptotic safety (Weinberg, 1979; Reuter, 1998) posits a non-trivial UV fixed point whose existence remains unproven.

τ-Effective

Theorem 6.16 (Readout quantization obstruction — recap)**.** *If the "classical" object to be quantized is already a readout of a boundary algebra, then quantization produces a readout of a readout—a double projection that generates spurious degrees of freedom (ghosts, non-renormalizable divergences) at each loop order.*

The double-projection problem. The boundary algebra $H_\partial[\omega]$ already contains the full quantum content of gravity. The "classical" Einstein equation is the first projection (the chart shadow). Quantizing the classical equation is a second projection:

$$H_\partial[\omega] \xrightarrow{\mathrm{pr}_1} G_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu} \xrightarrow{\text{“quantize”}} \hat{g}_{\mu\nu} \text{ on } \mathscr{H}_{\mathrm{grav}}. \tag{6.18}$$

The second step creates a Hilbert space $\mathscr{H}_{\mathrm{grav}}$ that is a shadow of a shadow. The resulting theory is non-renormalizable because the double projection loses the profinite finiteness that keeps the original algebra well-defined.

Remark 6.17 (Gravity is already quantum). In Category τ, gravity does not need to be quantized because it is already quantum. The D-sector of $H_\partial[\omega]$ has boundary characters (observables), a spectral decomposition, and commutation relations. The "classical" Einstein equation is the macroscopic limit, not the starting point.

Singularity Resolution

The Penrose–Hawking singularity theorems (1965–1970) require: (i) a smooth Lorentzian manifold, (ii) an energy condition, (iii) a trapped surface. In Category τ, assumption (i) fails—there is no smooth manifold. The profinite tower is compact, admitting no divergent elements. Consequences (the No Singularity Theorem, Part II):

(i) No singularities inside τ-black holes.
(ii) No Big Bang singularity.
(iii) Every progression orbit extends to all depths of the refinement tower.

What replaces a singularity is a *regime boundary*—a refinement depth below which the chart projection ceases to be the correct readout (Chapter 46).

Semiclassical backreaction ($G_{\mu\nu} = (8\pi G/c^4)\langle T_{\mu\nu}\rangle$) is plagued by regularization ambiguities. In τ, it is not needed: R^H and T are both elements of the same boundary algebra. The τ-Einstein identity *is* the quantum equation; there is no "classical" side to perturb.

WIMP Searches: Decades of Null Results

The WIMP was the leading dark matter candidate for three decades. Sensitivity has improved from 6×10^{-44} cm^2 (CDMS II, 2007) to 9.2×10^{-48} cm^2 (LZ, 2024)—seven orders of magnitude—with no signal.

Experiment	Upper limit ($\sigma_{\rm SI}$, cm^2)	Year
CDMS II	6×10^{-44}	2007
XENON100	2×10^{-45}	2012
LUX	7×10^{-46}	2016
XENON1T	4.1×10^{-47}	2018
XENONnT	2.6×10^{-47}	2023
PandaX-4T	3.8×10^{-47}	2024
LUX-ZEPLIN	9.2×10^{-48}	2024
DARWIN/XLZD (proj.)	$\sim 10^{-49}$	~2030

DARWIN/XLZD will reach the neutrino fog ($\sim 10^{-49}$ cm^2), below which coherent elastic neutrino–nucleus scattering (CEνNS) is an irreducible background. Collider searches (LHC monojet, mono-Z) find no excess above the Standard Model. Indirect detection (Fermi-LAT, IceCube, AMS-02) finds no confirmed annihilation signal. In τ, the null results are predicted: Sector Exhaustion guarantees no dark matter particle exists.

MOND and TeVeS

MOND remains the most successful alternative to dark matter at galactic scales. Its covariant extension TeVeS (Bekenstein, 2004) is largely falsified: GW170817 showed gravitational and electromagnetic waves travel at the same speed to 10^{-15}, ruling out most TeVeS-like theories. MOND has no natural cosmology (CMB, BAO, growth of structure require νHDM extensions with sterile neutrinos) and struggles with clusters (the Bullet Cluster requires additional mass). Category τ preserves MOND's galactic successes via the capacity-gradient argument (Section 6.2) and derives a_0 from first principles. Whether τ also resolves MOND's *failures*—the Bullet Cluster dissociation, dwarf spheroidal dispersions, and UDGs NGC 1052-DF2/DF4 [13, 14]—is an *open problem* flagged in Remarks 6.5 and 6.6; the claim "τ resolves MOND's failures" is *not* established in v1 of this Ledger.

Quintessence Models

Quintessence replaces Λ with a dynamical scalar field ϕ rolling down a potential $V(\phi)$, giving time-dependent $w > -1$. It shares two features with τ: $w \neq -1$ and no phantom crossing. But quintessence introduces a new field with a tuned potential ($m_\phi \sim H_0 \sim 10^{-33}$ eV), has no first-principles derivation, and suffers from the coincidence problem (why $\Omega_\phi \sim \Omega_m$ today?). In τ, the transition epoch is structurally determined, no new field is postulated, and the "coincidence" $\Omega_\Lambda \sim \Omega_m$ is not a coincidence—it is $\kappa_D(1 + \iota_\tau^3)$, a fixed algebraic expression.

The Cosmological Constant Problem: 120 Orders of Magnitude

The cosmological constant problem is the worst quantitative disagreement in the history of physics. QFT predicts a vacuum energy

$$\rho_{\rm vac}^{\rm QFT} \sim \frac{M_P^4}{(8\pi)^2} \sim 10^{74}\ {\rm GeV}^4. \tag{6.19}$$

The observed dark energy density is

$$\rho_\Lambda^{\rm obs} \sim 10^{-46}\ {\rm GeV}^4. \tag{6.20}$$

The ratio is 10^{120}. Proposed solutions:

- **Supersymmetry**: sets $\rho_{\rm vac} = 0$ before breaking; after breaking, $\rho \sim M_{\rm SUSY}^4 \sim 10^{12}$ GeV^4—still 10^{58} too large.
- **Anthropic selection**: the string landscape (10^{500} vacua) samples all Λ values; observers select compatible ones. This is not a prediction.
- **Sequestering** (Kaloper–Padilla, 2014): non-local action terms screen vacuum energy from gravity. Justification remains unclear.

Table 6.2: Honest ledger: GR + ΛCDM vs. Category τ on the dark sector and beyond.

Criterion	GR + ΛCDM	Category τ
Classical tests	All passed (8 confirmations)	Same (via chart shadow)
Gravitational waves	Detected (LIGO, 2015)	Same waveform templates
Rotation curves	Requires DM halos (2–3 params/galaxy)	V.T85: 0.067 dex, 0 params
BTFR slope	Not predicted (tuned)	4.000 exactly (V.C13)
Ω_Λ	0.685 (fitted, 1 param)	0.6849 at +269 ppm (0 params)
w_0	-1 (exact by assumption)	-0.960 (testable by DESI)
Singularities	Predicted (Penrose–Hawking)	Impossible
Quantization	Non-renormalizable	Not needed (already quantum)
H_0 tension	Unresolved (5σ)	$h = 0.6735$ (Planck-side); depth-dependent readout *retracted* v2.2 (Rem. 6.15); genuine 3–6σ tension with TRGB/HoLiCOW/SHoES
S_8 tension	Unresolved ($\sim 3\sigma$)	$S_8 = 0.760$ (WL side; uses σ_8^{CMB} as input; full Boltzmann run is v3 target, Rem. 6.12)
JWST early galaxies	Requires $\varepsilon > 50\%$	$\varepsilon \sim 45\%$ from $a_0(z)$
DM particle	Not found (decades of search)	Cannot exist (V.T44)
Cosmological constant	10^{120} fine-tuning	$\Lambda_\tau = 0$ exactly
Computational maturity	100+ years	New framework

τ-Effective

Remark 6.18 (No cosmological constant problem in τ). In the τ-framework, $\Lambda_\tau = 0$ exactly. There is no cosmological constant and therefore no problem. The vacuum catastrophe arises from computing $\langle T_{\mu\nu}\rangle_{\mathrm{vac}}$ in QFT on a manifold and inserting it into the Einstein equation. In τ, the boundary algebra $H_\partial[\omega]$ has no "vacuum energy" because the ground state is defined by the coherence kernel—not by summing over field modes on a continuous manifold. Neither ingredient of the catastrophe (QFT vacuum sum, Λ as free parameter) exists in τ. The 120-order-of-magnitude problem does not arise; it dissolves.

Dark Matter Detection Limits

Beyond WIMPs, other dark matter candidates face their own experimental constraints:

- **Axions**: ADMX, HAYSTAC, and MADMAX have excluded significant parameter space for the QCD axion, but the allowed mass range (10^{-12}–10^{-3} eV) remains large.
- **Sterile neutrinos**: X-ray telescopes have constrained the decay signal (3.5 keV line debated); no definitive detection.
- **Primordial black holes**: microlensing surveys (EROS, OGLE, HSC) exclude PBH dark matter in most mass ranges; the asteroid-mass window (10^{17}–10^{23} g) remains open.
- **Fuzzy dark matter**: ultra-light bosons ($m \sim 10^{-22}$ eV) are constrained by Lyman-α forest data and galaxy UV luminosity functions.

In the τ-framework, none of these candidates exists. The searches are guaranteed to return null results by Sector Exhaustion.

The Honest Ledger: GR + ΛCDM vs. Category τ

What GR got right. Geometry is physical (R^H is the ontic curvature); the equivalence principle (all sectors couple to $\mathfrak{S}_D$ via κ_τ); background independence (τ has no metric at the ontic level); gravitational waves propagate (linearized τ-Einstein gives GR waveforms in chart projection).

What GR got wrong. The manifold as substance (source of singularities, UV divergences, quantization failure—the manifold is a chart, not the territory); the metric as fundamental (quantizing a readout produces a double projection, the Readout Obstruction Theorem, Part II); the dark sector as physics (GR's 95% dark content is evidence that the chart projection misses structural content from the boundary algebra).

6.7 Where τ Diverges

The dissolution of the dark sector is not a minor adjustment to ΛCDM. It is a wholesale replacement of the ontological framework. The divergences are sharp, quantitative, and testable.

1. **No new particles.** τ predicts no dark matter particle will ever be detected. The particle spectrum is completely determined by the boundary holonomy algebra at enrichment layer E_1. The neutrino fog is not a frontier; it is the endpoint of a null search.
2. **No dark matter candidate.** No WIMP, axion, sterile neutrino, gravitino, dark photon, or primordial black hole plays the role of dark matter.
3. **No cosmological constant.** $\Lambda_\tau = 0$ exactly. The observed acceleration is dynamical: $w_0 = \iota_\tau^3 - 1 \approx -0.960$.
4. **No fine-tuning.** Ω_Λ, h, n_s, a_0 are all computed from $\iota_\tau = 2/(\pi + e)$. Zero adjustable parameters.
5. **No phantom crossing.** $w(z) \geq -1$ for all z (Theorem 6.11). Any data point $w < -1$ falsifies the framework.
6. **Sector exhaustion.** Five generators, five sectors, no sixth. Any confirmed dark matter detection falsifies Category τ.
7. **Flat rotation curves from boundary capacity, not new matter.** The additional gravitational force is not produced by a new substance. It is the capacity gradient $\nabla\mathscr{C}_D$ of the existing baryonic matter in the D-sector. If one fits an NFW halo to the τ-predicted rotation curve, one recovers a "dark matter halo" that matches ΛCDM expectations. The halo is real in the readout but not in the ontic structure.

Remark 6.19 (The falsification standard). The dissolution of the dark sector stands or falls on five predictions:

(i) No dark matter particle detected (ever).
(ii) $w_0 > -1$ (no phantom crossing).
(iii) Ω_Λ at +269 ppm.
(iv) $h = 0.6735$ (-120 ppm from Planck, $\sim 5\sigma$ from SHoES; falsified if the concordance settles at $h > 0.70$ with SHoES systematics cross-confirmed by TRGB, JAGB, megamasers, and standard sirens at comparable precision—see Remark 6.15).
(v) $S_8 = 0.760$ on the weak-lensing side (engagement caveat: this uses σ_8^{CMB} as input; see Remark 6.12).

Each is falsifiable. Each has a definite numerical value. The predictions (i)–(iv) do not have error bars—they are exact consequences of $\iota_\tau = 2/(\pi + e)$—and any significant departure (beyond observational uncertainty) falsifies the framework. Prediction (v) is partially motivational in v1 (see caveat above) and becomes fully independent only after the structure-formation triangle is run with a τ-native Boltzmann pipeline.

6.8 The Prediction Ledger

Table 6.3 consolidates the quantitative predictions of the dark-sector dissolution.

Table 6.3: Dark sector dissolution: quantitative prediction ledger.

Observable	τ formula	τ value	ppm	Scope
Ω_Λ	$\kappa_D(1+\iota_\tau^3)$	0.6849	+269	τ-eff
NGC 3198 v_∞	$(GM_{\rm eff}c^2/2\ell_\tau)^{1/4}$	149.1 km/s	−6000	τ-eff
BTFR slope	exact quartic	3.991	−2250	τ-eff
BTFR RMS	zero-parameter	0.067 dex	—	τ-eff
BTFR slope (SPARC-175, v2.2 M_b form)	ODR on 123 galaxies	3.61 ± 0.09	—	τ-eff
BTFR RMS (SPARC-175, v2.2 M_b form)	V.T85 v2.2 vs. V_∞	0.086 dex	—	τ-eff
BTFR slope (SPARC-175, v2.3 $M_{\rm eff}$ form)	ODR on 123 galaxies	4.18 ± 0.09	—	τ-eff
α^* (cold-gas source coupling)	SPARC bootstrap fit	0.240 ± 0.040	—	τ-eff
κ_ω (predicted)	$\iota_\tau/(1+\iota_\tau)$	0.2545	0.36σ from α^*	τ-eff
$a_0^{\rm bare}/a_0^{\rm dress}$	$\sqrt{\kappa_D/\kappa_B}$	2.378	—	τ-eff
w_0	$\iota_\tau^3 - 1$	−0.960	—	conj
h	$2/3 + \iota_\tau^2/17$	0.6735	−120	conj
No phantom	$w(z) > -1\ \forall z$	—	—	conj
$S_8^{(\tau)}$	holonomy suppression	0.760	—	conj
JWST SFE ($z=10$)	$\sqrt{a_0(z)/a_0(0)}$	45%	—	τ-eff
DM particles	Sector Exhaustion	none	exact	τ-eff
Λ	not needed	0	exact	τ-eff

Chapter Summary

- **The dark sector problem**: 95% of the universe unexplained; no dark matter particle detected; Λ requires 10^{120} fine-tuning.
- **Flat rotation curves** from the capacity gradient: $v_\infty^4 = GM_{\rm eff}c^2/(2\ell_\tau)$ with $M_{\rm eff} = M_\star + \kappa_\omega M_{\rm gas}$ and $\kappa_\omega = \iota_\tau/(1+\iota_\tau) = 0.2545$ (Theorem 6.2, V.T85 v2.3). NGC 3198 at 0.6%; 20-galaxy pilot at 0.067 dex RMS, BTFR slope 3.991, zero continuous free parameters (Proposition 6.4, V.D258; sector-structure integer choices per Chapter 1). The v2.2 full-sample SPARC-175 test (Section 6.2) with the single-source form M_b returned a partial confirmation (slope 3.61 ± 0.09, RMS 0.086 dex, $\sim 14\%$ zero-point offset, $f_{\rm gas}/\Sigma_{\rm disk}$ residuals at $p < 10^{-3}$); the v2.3 τ-internal reflection (Section 6.2) closes this residual structure by sector-decomposing the source (best-fit $\alpha^* = 0.240 \pm 0.040$, consistent at 0.36σ with κ_ω, all other τ-naturals rejected at $\geq 2\sigma$; slope recovers to 4.18 ± 0.09, within 2σ of the τ prediction of 4). Pointwise RAR remains the v3 discriminator.
- **Bare vs dressed unification**: $a_0^{\rm bare}/a_0^{\rm dress} = \sqrt{\kappa_D/\kappa_B} = 2.378$ (Theorem 6.7, V.T200).
- **Sector Exhaustion**: five generators, five sectors, no sixth for dark matter (Theorem 6.8, V.T44).
- **Dark energy density**: $\Omega_\Lambda = \kappa_D(1+\iota_\tau^3) = 0.6849$, Planck at +269 ppm (Theorem 6.9, V.T234). $\Lambda_\tau = 0$ exactly.
- **Equation of state**: $w_0 = \iota_\tau^3 - 1 \approx -0.960$ (Proposition 6.10, V.P159); no phantom crossing (Theorem 6.11, V.T235).
- S_8 **tension engaged (not yet resolved from first principles)**: $S_8^{(\tau)} = 0.760$, matching weak lensing (V.T236), but the derivation takes $\sigma_8^{\rm (CMB)}$ as input rather than producing it from a τ-native Boltzmann run; this is flagged as a v3 target (Remark 6.12).
- **JWST early galaxies**: ×20 enhancement at $z = 10$; SFE $\sim$ 45–56% (Proposition 6.13, V.T239).
- **Hubble parameter**: $h = 2/3 + \iota_\tau^2/17 = 0.6735$ (Theorem 6.14, V.T196), coincident with Planck 2018 and in $\sim 5\sigma$ tension with SHoES. The v1 "depth-dependent-readout dissolution" is *retracted* in v2.2 as a computable-structure claim (Remark 6.15): readout functor has no depth argument, tension is genuine (3–6σ with TRGB/HoLiCOW/SHoES) and is to be resolved by distance-ladder community work, not by τ-internal readout depth.

CHAPTER 7

Black Hole Topology — T^2 Signatures and Gravitational Waves

> *Category τ makes a sharp, falsifiable prediction about every black hole in the universe: the event horizon is a torus T^2, not a sphere S^2. Chapter 50 established the topological origin of this claim. This chapter assembles the full observational portfolio that flows from that single topological fact.*
> *The results are organized by observable channel: quasinormal mode frequencies (QNM ratio $\iota_\tau^{-1} \approx 2.930$, a clean discriminator against S^2), Event Horizon Telescope shadows (M87* at 40.85 μas, 0.4σ from observation; Sgr A* at 54.82 μas, 1.3σ), gravitational wave echoes (echo time ratio $\iota_\tau^{-2} \approx 8.585$, detectable with ~19 stacked LIGO events), and magnetic winding signatures (all winding numbers equal 2 from genus$(T^2) = 1$, field ratio $B_{tor}/B_{pol} = \iota_\tau^{-1}$).*
> *All predictions derive from the single posit ι_τ and the topology of the τ^3 fiber. No continuous free parameters; discrete integer/sector choices enter per compute-then-axiomatize (Chapter 1). The chapter concludes by placing these results against the orthodox landscape of theoretical physics programmes—Penrose twistors, Connes noncommutative geometry, asymptotic safety, and entropic gravity—identifying what each gets partially right and where the manifold ontology fails.*

7.1 The T^2 vs. S^2 Question

In general relativity, the event horizon of a stationary black hole is topologically S^2. This is not an assumption: it is a theorem. Hawking's *topology theorem* (1972), sharpened by Galloway and Schoen (2006), proves that cross-sections of the event horizon in a stationary, asymptotically flat spacetime satisfying the dominant energy condition must be topologically S^2. The *topological censorship theorem* of Friedman, Schleich, and Witt (1993) further constrains the topology of the domain of outer communications.

τ-Effective

Thesis 7.1 (The T^2 prediction)**.** *Category τ predicts that the event horizon of every black hole is topologically $T^2 = S^1 \times S^1$, not S^2. This prediction rests on three structural facts:*

(i) The fiber of $\tau^3 = \tau^1 \times_f T^2$ is T^2, not S^2. A black hole's horizon is the fiber restricted to the region where gravitational tension exceeds the spherical capacity (Theorem 50.6, Chapter 50).

(ii) The two cycles of T^2 carry distinct physical content: the outer S^1 (radius R, associated with the γ-generator, EM sector) and the inner S^1 (radius $r = \iota_\tau R$, associated with the η-generator, Strong sector).

(iii) The aspect ratio $r/R = \iota_\tau = 2/(\pi + e)$ is determined by the master constant (Theorem 50.23, Chapter 50). It is not a free parameter.

Why the GR theorem does not apply. Hawking's topology theorem assumes a Lorentzian manifold satisfying the dominant energy condition. In Category τ, there is no background manifold. The τ-Einstein equation (Chapter 13) is a boundary-character identity on $H_\partial[\omega]$, not a PDE on a manifold. The dominant energy condition is a readout-level statement that holds at enrichment layer E_1 but does not constrain the fiber topology, which is fixed at the algebraic level by the τ^3 construction. In short: the GR topology theorem is a theorem about manifolds, and black holes in τ are not manifold objects.

Falsification. The T^2 prediction is sharply falsifiable: if gravitational wave spectroscopy unambiguously identifies S^2 QNM overtone ratios (near 0.928) at 5σ significance, the τ-framework is falsified. Conversely, detection of a QNM ratio near 2.930 would be decisive evidence for T^2 topology. The following sections quantify every observable channel.

7.2 Quasinormal Modes

Quasinormal modes are the damped oscillations of a black hole after perturbation. In GR, the QNM spectrum of a Schwarzschild black hole is determined by the eigenvalues of the Regge–Wheeler and Zerilli equations

on the S^2 horizon. The fundamental $\ell = 2$ mode has frequency $f_0 \approx c^3/(16\pi GM)$ and the first overtone ratio $f_1/f_0 \approx 0.928$ for Schwarzschild, varying slightly with spin for Kerr.

In the τ-framework, the QNM spectrum is determined by the Laplacian on the T^2 horizon.

The T^2 QNM Spectrum

The torus $T^2 = (R \cdot S^1) \times (r \cdot S^1)$ with $r/R = \iota_\tau$ carries a Laplacian whose eigenvalues are labeled by winding pairs $(n, m) \in \mathbb{Z}^2$ (Definition 50.18, Chapter 50):

$$\lambda_{n,m} = \frac{n^2}{R^2} + \frac{m^2}{r^2} = \frac{1}{R^2}\left(n^2 + m^2 \cdot \iota_\tau^{-2}\right). \tag{7.1}$$

The QNM frequencies are $f_{n,m} \propto \sqrt{\lambda_{n,m}}$, giving three primitive modes:

$$f_{(1,0)} = \frac{c}{2\pi R}, \tag{7.2}$$

$$f_{(0,1)} = \frac{c}{2\pi r} = \iota_\tau^{-1} f_{(1,0)} \approx 2.930\, f_{(1,0)}, \tag{7.3}$$

$$f_{(1,1)} = \sqrt{1 + \iota_\tau^{-2}}\, f_{(1,0)} \approx 3.096\, f_{(1,0)}. \tag{7.4}$$

The Fundamental Frequency Ratio

τ-Effective

Theorem 7.2 (QNM frequency ratio = ι_τ^{-1} — V.T168)**.** *For a τ-black hole with T^2 horizon topology, the ratio of the inner-cycle QNM frequency to the outer-cycle QNM frequency is*

$$\frac{f_{(0,1)}}{f_{(1,0)}} = \frac{R}{r} = \iota_\tau^{-1} = \frac{\pi + e}{2} \approx 2.930. \tag{7.5}$$

This is a zero-parameter, mass-independent prediction.

Proof. From (7.2) and (7.3): $f_{(0,1)}/f_{(1,0)} = R/r = 1/\iota_\tau$. The ratio $r/R = \iota_\tau$ is the fiber breathing fraction of the τ^3 fibration (Theorem 50.23). ∎

T^2 Secondary Modes — V.T222

τ-Effective

Theorem 7.3 (T^2 secondary QNM modes)**.** *The T^2 ringdown consists of two superposed modes: the primary $f_{(1,0)}$, which matches the GR $\ell = 2$ fundamental, and the secondary $f_{(0,1)} = \iota_\tau^{-1} f_{(1,0)}$. For a seven-event LIGO catalog:*

Event	$M_{\text{rem}}\,(M_\odot)$	$f_{(1,0)}\,(Hz)$	$f_{(0,1)}\,(Hz)$
GW150914	*62*	*251*	*735*
GW151226	*21*	*741*	*2172*
GW170104	*49*	*318*	*931*
GW170608	*18*	*865*	*2534*
GW170729	*80*	*195*	*570*
GW170814	*53*	*294*	*861*
GW170817	*2.7*	*5764*	*16889*

The ratio $f_{(0,1)}/f_{(1,0)} = \iota_\tau^{-1}$ is mass-independent. The secondary mode at ~735 Hz for GW150914 is within the LIGO sensitivity band but requires higher SNR for detection.

Comparison with Schwarzschild and Kerr QNMs

The Schwarzschild $\ell = 2$ QNM overtone spectrum is well known: $\omega_n = \omega_0 - i(n + \frac{1}{2})\,\sigma$, where σ is the damping rate. The first overtone ratio $\omega_1/\omega_0 \approx 0.928$. For Kerr black holes with spin parameter a_*, the ratios change: the co-rotating $m = +2$ mode frequency increases relative to the counter-rotating $m = -2$ mode, but the fundamental-to-overtone ratio remains in the range $[0.8,\ 1.1]$.

τ-Effective

Remark 7.4 (Non-overlapping ranges). The T^2 prediction $f_{(0,1)}/f_{(1,0)} \in [2.5,\ 3.4]$ and the S^2 prediction $f_1/f_0 \in [0.8,\ 1.1]$ *do not overlap.* Any gravitational wave detector that resolves two QNM frequencies from a single ringdown event can distinguish T^2 from S^2 topology. This is the cleanest single-observable discriminator between τ and GR.

7.3 EHT Shadows

The Event Horizon Telescope (EHT) has imaged the shadows of two supermassive black holes: M87* (Event Horizon Telescope Collaboration, 2019) [18] and Sgr A* (2022) [19]. The τ-framework predicts a universal correction to the shadow diameter from T^2 topology.

The T^2 Shadow Correction

The Schwarzschild photon sphere radius is $r_{\rm ph} = 3GM/c^2$, giving a shadow diameter $\theta_{\rm GR} = 2\sqrt{3}\,r_{\rm ph}/d$ for a black hole at distance d. The T^2 torus horizon introduces a quadrupole correction from its oblateness.

τ-Effective

Theorem 7.5 (T^2 shadow correction — V.T184). *The T^2 quadrupole correction factor is*

$$f_\tau \;=\; 1 + \frac{\iota_\tau^2}{4} \;=\; 1.02912\ldots \tag{7.6}$$

The corrected shadow diameter is

$$\theta_\tau \;=\; f_\tau \cdot \theta_{GR} \;=\; \left(1 + \tfrac{\iota_\tau^2}{4}\right) \cdot 2\sqrt{3}\,\frac{GM}{c^2\,d}. \tag{7.7}$$

This is a universal +2.91% enlargement, mass-independent and zero-parameter.

M87* — V.T220

τ-Effective

Theorem 7.6 (M87* prediction suite). *For M87* ($M = 6.5 \times 10^9\ M_\odot$, $d = 16.8$ Mpc):*

- ***Shadow diameter:*** $\theta_\tau = 40.85\ \mu as$.
- ***EHT observation:*** $42 \pm 3\ \mu as$.
- ***Deviation:*** 0.4σ.
- ***QNM ratio:*** $\iota_\tau^{-1} \approx 2.930$.
- ***Modulation periods:*** 4.66 d (outer S^1), 1.59 d (inner S^1).

The shadow prediction lies comfortably within the EHT error bar. The T^2 correction brings τ 1.15 μas closer to the central value than pure GR.

Sgr A* — V.T221

τ-Effective

Theorem 7.7 (Sgr A* prediction suite). *For Sgr A* ($M = 4.0 \times 10^6\, M_\odot$, $d = 8.28$ kpc):*

- ***Shadow diameter:*** $\theta_\tau = 54.82\ \mu as$.
- ***EHT observation:*** $51.8 \pm 2.3\ \mu as$.
- ***Deviation:*** 1.3σ.
- ***QNM ratio:*** $\iota_\tau^{-1} \approx 2.930$.
- ***Modulation periods:*** 4.43 *min (outer* S^1*)*, 1.51 *min (inner* S^1*)*.

The 1.3σ *tension is within acceptable bounds. The modulation periods are testable by the next-generation EHT (ngEHT, ~2028+).*

Universal Correction and Variability — V.P148, V.P149

τ-Effective

Proposition 7.8 (Universal +2.91% correction — V.P148). *The readout Gibbs state on the linking boundary* $\mathbb{L} = S^1 \vee S^1$ *is Planckian via the KMS condition:* $B(\nu, T_H) = (2h\nu^3/c^2)\,/\,(\exp(h\nu/k_B T_H) - 1)$. *The Planckian spectrum combined with* T^2 *geometry produces the universal correction* $f_\tau = 1 + \iota_\tau^2/4$, *which is mass-independent.*

τ-Effective

Proposition 7.9 (Sgr A* variability from toroidal modes — V.P149). *The* T^2 *eigenfrequencies predict horizon-scale variability in Sgr A*:*

$$P_{outer} = 2\pi R_S/c \;\approx\; 4.43\ min, \tag{7.8}$$

$$P_{inner} = 2\pi r/c = \iota_\tau \cdot P_{outer} \;\approx\; 1.51\ min. \tag{7.9}$$

The outer period corresponds to the ISCO orbital period; the inner period is a τ-specific prediction with no GR analogue. Both are testable by ngEHT multi-epoch monitoring.

7.4 Gravitational Wave Echoes

If the black hole horizon has T^2 topology, gravitational waves can circulate around the two independent S^1 cycles before escaping. This produces post-ringdown echoes at characteristic time delays.

Echo Time Ratio — V.D283

τ-Effective

Definition 7.10 (Echo time delays). The gravitational-wave echo time delays for a T^2-topology black hole of mass M are

$$t_+ \;=\; \frac{4GM \cdot \iota_\tau}{c^3}\ \text{(inner } S^1\text{)}, \qquad t_- \;=\; \frac{4GM \cdot \iota_\tau^{-1}}{c^3}\ \text{(outer } S^1\text{)}, \tag{7.10}$$

with echo time ratio

$$\frac{t_-}{t_+} \;=\; \iota_\tau^{-2} \;\approx\; 8.585. \tag{7.11}$$

The echo separation $\Delta t = t_- - t_+ = 4GM(\iota_\tau^{-1} - \iota_\tau)/c^3$ places both echoes in the sensitive frequency band of current and next-generation detectors:

Source	$M(M_\odot)$	t_+ (ms)	t_- (ms)	Δt(ms)
Stellar BH	10	0.067	0.577	0.510
GW150914 remnant	62	0.417	3.580	3.163
Intermediate	150	1.009	8.660	7.651
Sgr A*	4×10^6	LISA band: ~4.3 mHz / ~37 mHz		

GW150914 Specific Predictions — V.R407

τ-Effective

Remark 7.11 (GW150914 echo search — V.R407). For GW150914 ($M_{\text{rem}} \approx 62\, M_\odot$):

- Inner echo: $t_+ = 0.417$ ms ($f \approx 2399$ Hz, within LIGO band).
- Outer echo: $t_- = 3.580$ ms ($f \approx 279$ Hz, within LIGO band).
- Echo time ratio: $t_-/t_+ = \iota_\tau^{-2} \approx 8.585$.
- Single-event echo SNR: ~1.04 (below 3σ threshold).

The echo amplitude is bounded by $A_{\text{echo}}/A_{\text{main}} \leq \exp(-\gamma_\tau\sqrt{\lambda_{nm}})$, where $\gamma_\tau = \pi(1+\iota_\tau^2/2)$ is the T^2-corrected damping parameter. For the $(1,0)$ mode: bound ≤ 0.036, approximate coupling ~ 0.043.

Detection Threshold — V.P151

τ-Effective

Proposition 7.12 (Stacked echo SNR — V.P151). *(i)* ***Current LIGO (O1–O3):*** *With ~10 BBH events, stacked echo SNR ≈ 2.2 (below 3σ). Approximately* ***19 events*** *needed for confident (3σ) detection.*

(ii) ***Einstein Telescope:*** *Single-event echo SNR ≈ 10.4 for a GW150914-like event. Decisive measurement in a single observation.*

(iii) ***Key discriminator:*** *The echo time ratio $t_-/t_+ = \iota_\tau^{-2} \approx 8.585$ is mass-independent. Any two echo peaks with this ratio would constitute evidence for T^2 topology.*

Echo Amplitude Model — V.D283

τ-Effective

Definition 7.13 (Echo amplitude bound). The echo amplitude relative to the main ringdown is bounded by

$$\frac{A_{\text{echo}}}{A_{\text{main}}} \leq \exp(-\gamma_\tau\sqrt{\lambda_{nm}}), \qquad \gamma_\tau = \pi\left(1 + \frac{\iota_\tau^2}{2}\right) \approx 3.324. \tag{7.12}$$

The bound is τ-effective: γ_τ derives from the T^2-corrected photon sphere potential barrier. The approximate coupling $A_{\text{echo}}/A_{\text{main}} \sim \exp(-\pi\sqrt{\lambda_{nm}})$ is conjectural. For the $(1,0)$ mode:

$$\frac{A_{(1,0)}}{A_{\text{main}}} \leq 0.036 \quad \text{(bound)}, \qquad \sim 0.043 \quad \text{(approximate)}. \tag{7.13}$$

7.5 Magnetic Winding Numbers

The T^2 topology of the horizon determines the magnetic field geometry near the black hole. A torus of genus 1 has a fundamentally different flux structure from a sphere of genus 0: $H_1(T^2; \mathbb{Z}) = \mathbb{Z} \oplus \mathbb{Z}$, while $H_1(S^2; \mathbb{Z}) = 0$. This topological distinction generates a suite of magnetic predictions.

The Three Winding Numbers

τ-Effective

Theorem 7.14 (RM winding theorem — V.T227). *The Faraday rotation measure (RM) around the shadow of a T^2 black hole exhibits winding number*

$$w_{\mathrm{RM}}(T^2) = 2, \qquad w_{\mathrm{RM}}(S^2) = 1. \tag{7.14}$$

The toroidal magnetic field on the minor S^1 cycle causes two sign changes per azimuthal circuit, compared to one for a radial/dipolar field on S^2. The winding number 2 is a topological invariant of genus$(T^2) = 1$.

Proof. The toroidal B-field wraps around the minor S^1 (Definition 42.25, Chapter 42). As the line of sight traces one azimuthal circuit around the shadow, it crosses two sectors where the toroidal field component $B_\parallel$ reverses sign: once at the "top" of the torus and once at the "bottom." Each reversal produces a sign change in the RM integral $\mathrm{RM} = \int n_e B_\parallel \, dl$. For S^2 with a radial or dipolar field, only one sign change occurs per circuit. The winding number is $w_{\mathrm{RM}} = |\text{number of sign changes}|/1 = 2$ for T^2 and 1 for S^2. ∎

τ-Effective

Theorem 7.15 (Circular polarization winding — V.T229). *The Stokes V circular polarization winding number around the shadow is*

$$w_V(T^2) = 2, \qquad w_V(S^2) = 1. \tag{7.15}$$

The Faraday conversion coefficient $\cos(2\chi_B)$ reverses sign at the same azimuthal angles as $B_\parallel$ in the RM integral (same toroidal geometry). Therefore $w_V = w_{\mathrm{RM}} = 2$.

Remark 7.16 (EVPA winding). The electric vector position angle (EVPA) winding is also $w_{\mathrm{EVPA}} = 2$ for T^2, 1 for S^2. All three polarimetric winding numbers—RM, EVPA, Stokes V—agree:

$$w_{\mathrm{RM}} = w_{\mathrm{EVPA}} = w_V = 2 \quad (T^2), \qquad = 1 \quad (S^2). \tag{7.16}$$

$B_{\mathrm{tor}}/B_{\mathrm{pol}} = \iota_\tau^{-1}$ — V.T230

τ-Effective

Theorem 7.17 (Magnetic field ratio — V.T230). *For a τ-black hole, the near-horizon toroidal-to-poloidal magnetic field ratio is*

$$\frac{B_{\mathrm{tor}}}{B_{\mathrm{pol}}} = \iota_\tau^{-1} \approx 2.930. \tag{7.17}$$

This follows from frozen-flux conservation on T^2: the magnetic field strength scales inversely with area, $B \propto 1/A$, and the two fundamental cycle areas scale as $R/r = \iota_\tau^{-1}$. The ratio is mass-independent and zero-parameter.

Proof. On the torus $T^2 = (R \cdot S^1) \times (r \cdot S^1)$, the toroidal flux threads the minor cross-section (area $\propto r$) and the poloidal flux threads the major cross-section (area $\propto R$). By ideal-MHD frozen-flux conservation, $B_{\mathrm{tor}} \cdot A_{\mathrm{minor}} = B_{\mathrm{pol}} \cdot A_{\mathrm{major}}$ at equipartition. Since $A_{\mathrm{major}}/A_{\mathrm{minor}} = R/r = \iota_\tau^{-1}$, we obtain $B_{\mathrm{tor}}/B_{\mathrm{pol}} = R/r = \iota_\tau^{-1}$. ∎

Flux Threading Through T^2 — V.T228

τ-Effective

Theorem 7.18 (Flux threading theorem — V.T228). *Both toroidal flux Φ_{tor} and poloidal flux Φ_{pol} are conserved in ideal MHD on T^2:*

$$\frac{d\Phi_{\mathrm{tor}}}{dt} = 0, \qquad \frac{d\Phi_{\mathrm{pol}}}{dt} = 0. \tag{7.18}$$

Table 7.1: T^2 vs. S^2 magnetic predictions (all derived from genus(T^2) = 1 and ι_τ).

Observable	T^2 prediction	S^2 prediction
RM winding w_{RM}	2	1
EVPA winding w_{EVPA}	2	1
Stokes V winding w_V	2	1
$B_{\mathrm{tor}}/B_{\mathrm{pol}}$	$\iota_\tau^{-1} \approx 2.93$	Model-dependent
Flux through hole	Present ($\Phi_{\mathrm{pol}} \neq 0$)	Absent ($H_1 = 0$)
Jet helicity sign	Fixed by topology	Free parameter
Jet B_z/B_φ at base	$\iota_\tau \approx 0.341$	~ 1 (equipartition)
IGMF in filaments	10–100 nG	0.1–1 nG (dynamo)
B-field alignment	Parallel to filaments	Random

The poloidal flux Φ_{pol} (threading the torus hole) is topologically protected: changing it requires magnetic reconnection (a violation of ideal MHD). On S^2, $H_1(S^2) = 0$, so the only topologically stable flux through the "hole" is $\Phi = 0$. The torus has a nonzero topologically protected flux; the sphere does not.

Observational consequence. The topologically protected poloidal flux explains the long-term stability of AGN jets (lasting 10^6–10^8 years): the jet is anchored by a topological invariant, not by a dynamical equilibrium. In S^2 models, jet stability requires continuous dynamo maintenance with no intrinsic stability guarantee.

Magnetic Prediction Table

7.6 Jet Helicity and Cosmic Magnetism

The magnetic predictions of Section 7.5 extend from the near-horizon scale ($\sim R_S$) to the cosmic web (~Mpc). The link is the topologically protected poloidal flux of Theorem 7.18.

Jet B_z/B_φ at the Base — V.P156

τ-Effective

Proposition 7.19 (Jet field ratio at base — V.P156)**.** *At the jet base ($\sim 10\, R_S$), the axial-to-toroidal magnetic field ratio is*

$$\left.\frac{B_z}{B_\varphi}\right|_{base} = \iota_\tau \approx 0.341. \tag{7.19}$$

This follows from the torus hoop-stress collimation: the toroidal component B_φ provides hoop stress $P = B_\varphi^2/(4\pi r)$ that collimates the jet, while the axial component B_z originates from the topologically protected Φ_{pol}. The equilibrium condition $\sin\theta_{jet} \leq B_z/B_\varphi = \iota_\tau$ independently recovers the jet collimation angle $\theta_{jet} \leq 20^\circ$ (Theorem 40.26).

Remark 7.20 (Observational consistency). VLBI observations of AGN jets at sub-parsec scales measure $B_z/B_\varphi \sim 0.2$–0.5, consistent with the τ-prediction $\iota_\tau \approx 0.341$. The ratio is a zero-parameter prediction, testable with improved VLBI polarimetry.

Intergalactic Magnetic Field — V.P157

τ-Effective

Proposition 7.21 (IGMF from Wilson skeleton — V.P157)**.** *The intergalactic magnetic field (IGMF) in cosmic filaments is predicted to be*

$$B_{fil} \sim 10\text{–}100\ nG. \tag{7.20}$$

The physical mechanism:

(i) Supermassive black holes at filament nodes carry near-horizon field $B \sim 1$–30 G.

(ii) The topologically protected poloidal flux Φ_{pol} is transported via jets along Wilson skeleton edges (Chapter 43).

(iii) Dilution over the filament cross-section ($\sim (2\ Mpc)^2$) gives $B_{fil} \sim \Phi_{\text{pol}}/A_{fil} \sim 10$–100 nG.

The contrast $B_{fil}/B_{void} \gg 1$ is a structural prediction of the Wilson skeleton topology.

Remark 7.22 (Observational status). Vernstrom et al. (2021) detected $B \sim 30$ nG in cosmic filaments via synchrotron stacking, within the τ-predicted range. Carretti et al. (2023) obtained consistent results. Both exceed random-dynamo predictions (0.1–1 nG) by 1–2 orders of magnitude. The discriminating prediction is *alignment*: T^2 predicts B-field parallel to filament axes (from 1D Wilson loop topology), while random dynamo produces isotropic fields. This is testable by the Square Kilometre Array (~2030+).

7.7 The Orthodox Landscape

The results of Sections 7.2–7.6 flow from a single topological fact: the fiber of τ^3 is T^2. We now place these results against the broader landscape of theoretical physics programmes, asking: what does each programme see correctly, what does it miss, and how does the partial truth embed into Category τ?

Penrose Twistors: Holomorphic Geometry for Physics

Roger Penrose proposed the twistor programme in the late 1960s, motivated by the intuition that the fundamental arena of physics should be *holomorphic*, not real-differentiable. Twistor space $\mathbb{T} \cong \mathbb{C}^4$ encodes null geodesics in Minkowski space; a spacetime point corresponds to a complex projective line $\mathbb{CP}^1 \hookrightarrow \mathbb{PT}$. The Penrose transform encodes massless fields as sheaf cohomology classes $H^1(\mathbb{PT}, \mathcal{O}(n))$: physics becomes cohomology.

Metaphorical

Observation 7.23 (What Penrose gets right). Three features correctly identified:

(i) **Holomorphy is primary.** Physics is encoded in holomorphic data.

(ii) **Points are derived.** A spacetime point is a relation ($\mathbb{CP}^1$ in $\mathbb{PT}$), not a primitive.

(iii) **Massless fields are cohomological.** The Penrose transform encodes fields as algebraic objects on a complex space.

Each has a precise counterpart in Category τ: holomorphy from the Central Theorem $\mathcal{O}(\tau^3) \cong A_{\text{spec}}(\mathbb{L})$; points as E_1 readouts of boundary characters; all fields as elements of $H_\partial[\omega]$.

The limitations. The twistor programme has three well-known obstacles: (1) the *curved-space problem* (twistors are naturally defined for flat or conformally flat spacetimes); (2) *mass* (massive fields lack a clean twistor description); (3) *no internal degrees of freedom* (twistors do not incorporate gauge groups or particle species).

τ-Effective

Proposition 7.24 (Twistor embedding)**.** *The Penrose transform for massless fields on Minkowski space embeds into the E_1 readout of the boundary holonomy algebra:*

$$H^1\big(\mathbb{PT},\, \mathcal{O}(n)\big) \;\hookrightarrow\; \big\{\, \chi \in H_\partial[\omega] \;\big|\; m(\chi) = 0 \,\big\}. \tag{7.21}$$

The massless condition $m(\chi) = 0$ restricts to characters with no fiber breathing-mode excitation. The helicity n corresponds to the winding number around T^2. All three obstacles dissolve: curved space (no background manifold to curve), mass (fiber breathing mode), gauge groups (five sectors of $H_\partial[\omega]$).

Connes' Noncommutative Geometry: Spectral Triples

Alain Connes' NCG programme begins from the insight that geometry is spectral: the fundamental object is a spectral triple $(\mathcal{A}, \mathcal{H}, D)$ consisting of a $*$-algebra, a Hilbert space, and a Dirac operator. The Connes–Chamseddine spectral action

$$S_{\mathrm{CC}} \;=\; \mathrm{Tr}\big(f(D/\Lambda)\big) + \tfrac{1}{2}\langle J\psi,\, D\psi\rangle \tag{7.22}$$

reproduces the Standard Model coupled to Euclidean gravity from the algebra $\mathcal{A} = C^\infty(M) \otimes \mathcal{A}_F$ where $\mathcal{A}_F = \mathbb{C} \oplus \mathbb{H} \oplus M_3(\mathbb{C})$.

Metaphorical

Observation 7.25 (What Connes gets right). Four correct identifications:

(i) **Geometry is algebraic**: the fundamental object is the algebra, not the manifold.
(ii) **The metric is spectral**: distances from operator spectra.
(iii) **Internal and external geometry unify**: gauge groups from the algebra.
(iv) **Internal geometry is finite**: $\mathcal{A}_F$ is finite-dimensional.

Where NCG falls short. (1) The algebra $\mathcal{A}_F$ is put in by hand, not derived. (2) The cutoff Λ is a free parameter. (3) Gravity remains classical. (4) The original Higgs mass prediction (~170 GeV) failed.

τ-Effective

Proposition 7.26 (NCG spectral triple from τ)**.** *The boundary holonomy algebra $H_\partial[\omega]$ at E_1 determines a canonical spectral triple $(\mathcal{A}_\tau, \mathcal{H}_\tau, D_\tau)$ with $\mathcal{A}_\tau = \mathcal{O}(\tau^3) \cong A_{\mathrm{spec}}(\mathbb{L})$. The finite algebra is recovered as*

$$\mathcal{A}_F \;\cong\; \bigoplus_{X\in\{D,A,B,C,\omega\}} \mathrm{End}\big(H_\partial^X\big) \Big/ \textit{cross-coupling relations.} \tag{7.23}$$

The algebra is not put in by hand: it is the endomorphism algebra of the sector decomposition.

Asymptotic Safety: Weinberg's Conjecture

Steven Weinberg conjectured (1979) that GR possesses a non-perturbative UV fixed point. Martin Reuter and collaborators developed the functional renormalization group approach, finding evidence for a fixed point with 2–4 relevant directions across progressively larger truncations.

Metaphorical

Observation 7.27 (What asymptotic safety gets right). The correct principle: gravity at high energies does not require new particles or extra dimensions. The existing theory, properly understood non-perturbatively, is sufficient. This is the τ-position: gravity is the D-sector readout of $H_\partial[\omega]$, and the coupling $\kappa_\tau = 1 - \iota_\tau$ does not run.

Table 7.2: Six programmes and their partial truths.

Programme	Correct Insight	τ-Counterpart
String Theory	Extra dimensions; holographic duality; extended objects	Fiber T^2; boundary/bulk duality on $\mathbb{L}$; defect bundles
Loop QG	Discrete quantum geometry; background independence; area quantization	Profinite spectrum; no background manifold; area = boundary character readout
Twistors	Holomorphic geometry; points derived; massless fields = cohomology	Central Theorem; characters, not points; all fields = boundary characters
NCG	Geometry = algebra; metric = spectrum; internal geometry finite	$\mathcal{O}(\tau^3) \cong A_{\text{spec}}(\mathbb{L})$; D-sector spectral; five sectors from five generators
Asym. Safety	Gravity is UV complete; finite relevant directions	No renormalization needed; two base generators
Entropic/Emergent	Gravity emerges; entropy-area relation; no dark matter needed	D-sector readout; $S = k_B A/(4\iota_\tau^2)$; sector exhaustion

τ-Effective

Theorem 7.28 (Gravity as readout — no renormalization)**.** *In the τ-framework:*

(i) GR is a readout, not a fundamental theory.

(ii) Readouts do not need renormalization: the boundary holonomy algebra is profinite, its spectrum discrete, and there are no divergent integrals.

(iii) The gravitational coupling $\kappa_\tau = 1 - \iota_\tau \approx 0.659$ takes this value at every refinement depth. No running. No UV fixed point. No IR limit.

Remark 7.29 (The two relevant directions). The most robust finding of the asymptotic safety programme is approximately two relevant directions. In τ, the base τ^1 has two generators: α ($\mathfrak{S}_D$, Gravity) and π ($\mathfrak{S}_A$, Weak). The "two relevant directions" are the two base-sector couplings $\kappa(D;\,1)$ and $\kappa(A;\,1)$, satisfying $\kappa(D;\,1) + \kappa(A;\,1) = 1$. Two couplings, one constraint, one free parameter—which is ι_τ itself.

Entropic and Emergent Gravity

Ted Jacobson (1995) showed that the Einstein equation follows from the Clausius relation $\delta Q = TdS$ applied to local causal horizons. Erik Verlinde (2010/2016) proposed gravity as an entropic force: $F\Delta x = T\Delta S$.

Metaphorical

Observation 7.30 (What the entropic programmes get right). Three correct identifications:

(i) Gravity emerges (D-sector readout at E_1).

(ii) Entropy is structural (area-entropy relation from boundary identity).

(iii) Dark matter effects may be gravitational (consistent with Sector Exhaustion, Chapter 44).

τ-Effective

Principle 7.31 (Common origin, not causal priority)**.** *Gravity and thermodynamics are both E_1 readouts of $H_\partial[\omega]$:*

$$H_\partial[\omega] \xrightarrow{\mathrm{pr}_{\mathfrak{S}_D}} \tau\text{-Einstein equation}, \qquad H_\partial[\omega] \xrightarrow{\mathrm{pr}_{entropy}} \text{entropy splitting}. \tag{7.24}$$

Neither is prior to the other. Jacobson's derivation is correct in spirit: the Einstein equation is a thermodynamic identity. The error is the causal direction: gravity does not emerge from thermodynamics; both emerge from the boundary.

Six Programmes and Their Partial Truths

τ-Effective

Thesis 7.32 (The manifold error). *Every major programme (string theory, LQG, twistors, NCG, asymptotic safety, entropic/emergent gravity) correctly identifies a structural feature of physics and incorrectly houses it in a manifold ontology. Category τ recovers every correct insight as a projection of $H_\partial[\omega]$, without requiring a background manifold. The common error is the manifold; the common truth is the boundary.*

The BH Information Paradox: No Paradox in τ

In GR, Hawking's calculation (1975) shows that black holes radiate thermally. If the radiation is truly thermal, the information about in-falling matter is lost, violating the unitarity of quantum mechanics. This is the *information paradox*, which has driven four decades of theoretical debate.

In Category τ, the information paradox does not arise.

τ-Effective

Theorem 7.33 (No information paradox in τ). *The τ-framework precludes the information paradox through three independent mechanisms:*

(i) ***No Hawking radiation as fundamental process.*** *Hawking radiation requires a Bogoliubov transformation between "in" and "out" vacuum states. In Category τ, the vacuum is the boundary $\mathbb{L} = S^1 \vee S^1$ (unique, no in/out split). The SA-i condition forbids sub-coherence-kernel mode creation. Therefore τ-black holes do not evaporate.*

(ii) ***Boundary coherence.*** *The boundary holonomy algebra $H_\partial[\omega]$ is an inverse system of finite-dimensional algebras. Every element at depth n is determined by its projections to coarser depths. No information is lost (Corollary 50.9, Chapter 50).*

(iii) ***No interior singularity.*** *The fiber T^2 is compact with no boundary. All boundary characters are bounded on the compact interior (Proposition 50.8). There is no $r = 0$ where information could be destroyed.*

Hawking Radiation as Readout Artifact

Remark 7.34 (Hawking temperature as readout). The Hawking temperature $T_H = \hbar c^3/(8\pi G M k_B)$ is not denied: it is *reinterpreted*. The readout Gibbs state on the linking boundary (Definition 52.11, Chapter 52) has temperature T_H. The spectrum is Planckian (Proposition 52.13). But this is an *informational readout*, not an energetic flux: no mass is lost, and the No-Shrink Theorem (Theorem 52.3, Chapter 52) ensures that the black hole mass is monotonically non-decreasing. The Hawking temperature describes the boundary state's information content, not a physical radiation process.

7.8 Where τ Diverges

This section collects the sharpest points of divergence between the τ-framework and orthodox black hole physics.

T^2 Not S^2

The horizon topology is T^2, not S^2. This is the most fundamental divergence. All predictions in this chapter (QNM ratio, echo times, shadow correction, magnetic winding numbers, field ratios, IGMF magnitude, jet helicity) flow from this single topological fact. The divergence is falsifiable: QNM spectroscopy can decide.

No Information Paradox

There is no information paradox in τ. The paradox arises in GR from three features: (1) Hawking radiation is thermal, (2) the interior singularity destroys information, (3) unitarity requires information preservation. In τ:

(1) there is no Hawking radiation (no Bogoliubov transformation), (2) there is no singularity (T^2 is compact), (3) boundary coherence preserves all data.

No Singularity

The interior of a τ-black hole is a compact subset of T^2. Every continuous function on a compact set is bounded. The curvature functional $R^H[\chi]$ is bounded on the interior. There is no $r = 0$.

Entropy Excess Factor $\pi\iota_\tau$

τ-Effective

Proposition 7.35 (Entropy excess — V.P125)**.** *The T^2 horizon carries 7.2% more entropy than the S^2 Bekenstein–Hawking entropy at the same Schwarzschild radius:*

$$\frac{S_{T^2}}{S_{S^2}} = \pi\iota_\tau \approx 1.0722. \tag{7.25}$$

The excess factor $\pi\iota_\tau$ is mass-independent.

No Hawking Evaporation (No-Shrink Theorem)

τ-Effective

Theorem 7.36 (No-Shrink summary)**.** *A mature τ-black hole cannot decrease in mass. The defect-mass coupling (Theorem 52.2, Chapter 52) ensures monotonic non-decrease: $M(n+1) \geq M(n)$ for all refinement depths $n \geq n_*$. Black holes grow or remain constant. They do not shrink. They do not evaporate.*

Chapter Summary

- The T^2 vs. S^2 question is the sharpest falsifiable prediction of the τ-framework (Thesis 7.1).
- **QNM frequency ratio** $= \iota_\tau^{-1} \approx 2.930$ vs. Schwarzschild ≈ 0.928: non-overlapping ranges, cleanest single-observable discriminator (Theorem 7.2, Remark 7.4).
- **EHT shadows:** M87* at 40.85 μas (0.4σ), Sgr A* at 54.82 μas (1.3σ), universal +2.91% correction (Theorems 7.6, 7.7, 7.5).
- **GW echoes:** echo time ratio $\iota_\tau^{-2} \approx 8.585$, ~19 events for 3$\sigma$ detection, Einstein Telescope decisive (Proposition 7.12).
- **Magnetic winding numbers:** $w_{\mathrm{RM}} = w_{\mathrm{EVPA}} = w_V = 2$ (Theorems 7.14, 7.15). $B_{\mathrm{tor}}/B_{\mathrm{pol}} = \iota_\tau^{-1}$ (Theorem 7.17).
- **Jets and cosmic magnetism:** $B_z/B_\varphi = \iota_\tau$ at base, IGMF ~ 10–100 nG in filaments (Propositions 7.19, 7.21).
- The **orthodox landscape** (twistors, NCG, asymptotic safety, entropic gravity) correctly identifies partial truths; every programme commits the manifold error (Thesis 7.32, Table 7.2).
- **No information paradox**: no Hawking radiation, no singularity, boundary coherence preserves all data (Theorem 7.33).
- **No evaporation**: No-Shrink Theorem ensures monotonic non-decrease of mass (Theorem 7.36).
- **Entropy excess:** $S_{T^2}/S_{S^2} = \pi\iota_\tau \approx 1.072$, mass-independent (Proposition 7.35).

Table 7.3: Complete black hole prediction table (all from T^2 topology and the ι_τ posit; zero continuous free parameters, structural integer choices as listed; Chapter 1).

Observable	τ **Prediction**	**Observation**	**Decisive Instrument**
QNM ratio	$\iota_\tau^{-1} \approx 2.930$	Not yet resolved	ET, LISA
Echo time ratio	$\iota_\tau^{-2} \approx 8.585$	Not yet detected	LIGO (∼19 events)
M87* shadow	40.85 μas	42 ± 3 μas	ngEHT
Sgr A* shadow	54.82 μas	51.8 ± 2.3 μas	ngEHT
Shadow correction	+2.91%	Within error	ngEHT (< 3% precision)
RM winding	$w_{\mathrm{RM}} = 2$	Consistent	ngEHT (∼2028+)
$B_{\mathrm{tor}}/B_{\mathrm{pol}}$	2.930	Consistent	ngEHT polarimetry
Jet B_z/B_φ	0.341 at base	0.2–0.5	VLBI
IGMF (filaments)	10–100 nG	∼ 30 nG	SKA (∼2030+)
Entropy ratio	$\pi\iota_\tau \approx 1.072$	—	Indirect
Evaporation	Forbidden	No detection	—
Information loss	Impossible	—	—

CHAPTER 8

Collective Dynamics: Turbulence, Reconnection, and Heating

Four of the oldest open problems in classical physics—turbulent intermittency, the value of the Kolmogorov constant, fast magnetic reconnection, and coronal heating—have resisted quantitative solution for decades. Each problem involves a parameter that orthodox theory must fit to experiment: $\beta = 2/3$ in She–Lévêque, $C_K \approx 1.5$ in Kolmogorov, $v_{\rm rec} \approx 0.1\, v_A$ in reconnection, and a damping length scale in coronal heating. This chapter shows that all four parameters are derived—not fitted—from a single structural fact of the τ^3 fibration: $\dim(T^2) = 2$. The fiber dimension controls intermittency, sets the Kolmogorov constant, determines the reconnection rate, and fixes the heating damping scale. A fifth result—the super-exponential convergence of the Navier–Stokes decompactification exponent—connects to the Millennium Problem. Together, these constitute the first parameter-free theory of collective dynamics.

8.1 The Turbulence Problem

Turbulence is the last great unsolved problem of classical physics. The Navier–Stokes equations have been known since 1822 (Navier) and 1845 (Stokes), yet no closed-form solution exists for the general turbulent regime, and even the question of whether smooth solutions persist for all time remains open (the Clay Millennium Problem).

The empirical situation. Turbulent flows exhibit remarkable universality. The energy spectrum in the inertial range follows the Kolmogorov $-5/3$ law:

$$E(k) \;=\; C_K\, \varepsilon^{2/3}\, k^{-5/3}, \tag{8.1}$$

where ε is the mean energy dissipation rate and $C_K \approx 1.5$ is the Kolmogorov constant. This law has been verified in wind tunnels, atmospheric boundary layers, ocean currents, and numerical simulations across more than four decades of wavenumber range. The universality is astonishing: the same exponent and prefactor appear regardless of the large-scale geometry, the forcing mechanism, or the Reynolds number (provided $\mathrm{Re} \gg 1$).

Kolmogorov 1941. Kolmogorov's original theory (K41) derives (8.1) from two hypotheses:

1. **Local isotropy.** At scales much smaller than the injection scale L, turbulence is statistically isotropic and homogeneous.
2. **Self-similarity.** The statistics of velocity increments $\delta v(\ell) = v(x+\ell) - v(x)$ depend only on the separation ℓ and the dissipation rate ε.

From these hypotheses, dimensional analysis gives

$$\langle |\delta v(\ell)|^p \rangle \;\propto\; (\varepsilon\, \ell)^{p/3}, \tag{8.2}$$

predicting structure function exponents $\zeta_p = p/3$. The exact result of K41 is the four-fifths law: $\langle (\delta v_\parallel)^3 \rangle = -\frac{4}{5}\, \varepsilon\, \ell$, giving $\zeta_3 = 1$ exactly. This remains the only exact result in turbulence theory.

The intermittency gap. For $p \geq 4$, the K41 prediction $\zeta_p = p/3$ fails systematically. The observed exponents satisfy $\zeta_p < p/3$: turbulence is more intermittent than K41 allows. Rare, intense events (concentrated vortex filaments) produce extreme velocity gradients that violate the self-similarity assumption. Correcting for intermittency has been the central challenge of turbulence theory for over half a century.

She–Lévêque 1994. The most successful correction was proposed by She and Lévêque (1994), who introduced a log-Poisson model for the energy dissipation and derived:

$$\zeta_p = \frac{p}{9} + 2\left[1 - \left(\frac{2}{3}\right)^{p/3}\right]. \tag{8.3}$$

This formula contains two parameters: $\beta = 2/3$ (the intermittency parameter, governing the scaling ratio between successive hierarchy levels) and $C_0 = 2$ (the co-dimension of the most intense dissipative structures). Both were motivated by physical reasoning (filamentary vortex tubes have co-dimension 2; $\beta = 2/3$ gives the best fit), but ultimately they were selected to match experimental data. The formula agrees with experiment to better than 1% for all $p \leq 12$.

The question. Why $\beta = 2/3$? Why $C_0 = 2$? Why does the exponent in the linear term change from $p/3$ to $p/9$? The orthodox theory has no answer: these are fitted constants, and no principle determines them.

8.2 She–Lévêque from τ^3 Dimensions

Chapter 28 derived the She–Lévêque formula from the τ^3 fibration. We summarize the result and its logic.

The derivation. The fibered product $\tau^3 = \tau^1 \times_f T^2$ has total dimension 3 and fiber dimension 2. Three structural identifications determine the She–Lévêque parameters:

1. **Co-dimension of dissipative structures:** $C_0 = \dim(T^2) = 2$. The most intense dissipative structures in fully developed turbulence are vortex filaments—one-dimensional curves in three-dimensional space. Their co-dimension is $\dim(\tau^3) - 1 = 2$, which equals the fiber dimension. In the τ-framework, vortex filaments are the loci where the fiber T^2 degenerates, leaving a base-only skeleton (Definition 28.14).
2. **Intermittency parameter:** $\beta = \dim(T^2)/\dim(\tau^3) = 2/3$. The ratio of fiber to total dimension controls the scaling between successive hierarchy levels of the energy dissipation field. This ratio is a structural invariant of the fibration—not a free parameter.
3. **Linear scaling:** $p/\dim(\tau^3)^2 = p/9$. The K41 exponent $p/3$ is reduced by a factor of $1/\dim(\tau^3)$ because the cascade operates on the full fibered product, saturating at rate $\varepsilon^{1/3}$ per dimension.

τ-Effective

Theorem 8.1 (She–Lévêque from fibration — V.T248; recap)**.** *The structure function exponents of fully developed turbulence are*

$$\zeta_p = \frac{p}{\dim(\tau^3)^2} + \dim(T^2)\left[1 - \left(\frac{\dim(T^2)}{\dim(\tau^3)}\right)^{p/\dim(\tau^3)}\right] = \frac{p}{9} + 2\left[1 - \left(\frac{2}{3}\right)^{p/3}\right], \tag{8.4}$$

with zero continuous free parameters; the exponent structure derives from $\dim(T^2) = 2$ *with no integer-choice site (Chapter 1).*

The agreement with experiment (Theorem 28.17) is better than 1% for all $p \leq 12$:

p	ζ_p^τ	ζ_p^{exp}	K41	Deviation
1	0.364	0.37 ± 0.01	0.333	−1.6%
2	0.696	0.70 ± 0.01	0.667	−0.6%
3	1.000	1.00 (exact)	1.000	0.0%
4	1.280	1.28 ± 0.02	1.333	0.0%
6	1.778	1.77 ± 0.04	2.000	+0.5%
8	2.211	2.21 ± 0.07	2.667	0.0%
10	2.598	2.59 ± 0.10	3.333	+0.3%
12	2.948	2.93 ± 0.15	4.000	+0.6%

Remark 8.2 (Fitted to derived). In the She–Lévêque original work, $\beta = 2/3$ and $C_0 = 2$ are selected to match data. In Category τ, $\beta = \dim(T^2)/\dim(\tau^3)$ and $C_0 = \dim(T^2)$ are structural constants of the fibration. The match with experiment is therefore a prediction, not a fit (Remark 28.19). This is the first derivation of the She–Lévêque parameters from any theory.

Remark 8.3 (Vortex filaments as fiber collapse). The identification $C_0 = \dim(T^2)$ has a geometric interpretation (Remark 28.20): vortex filaments are co-dimension-2 objects because they are the loci where the fiber T^2 degenerates. Intermittency is thus a consequence of the fibered structure of macroscopic phase space. It arises because the fiber can collapse, concentrating the energy flux onto lower-dimensional structures.

8.3 Kolmogorov Constants

The Kolmogorov energy spectrum $E(k) = C_K \varepsilon^{2/3} k^{-5/3}$ contains two structural constants: the exponent $-5/3$ and the prefactor C_K. In orthodox theory, $-5/3$ is obtained by dimensional analysis (which does not explain its numerical value), and C_K is fitted to DNS data. In Category τ, both are derived.

The exponent $-5/3$

τ-Effective

Theorem 8.4 ($-5/3$ from τ dimensions — V.T250; recap)**.** *The energy spectrum exponent decomposes as*

$$\frac{5}{3} = \frac{\dim(\tau^3) + \dim(T^2)}{\dim(\tau^3)} = \frac{3+2}{3}. \tag{8.5}$$

The numerator $5 = \dim(\tau^3) + \dim(T^2) = |\mathrm{gen}| + \dim(T^2)$ *counts the total number of dissipation channels: three generation modes from* $H_1(\tau^3; \mathbb{Z}) \cong \mathbb{Z}^3$ *plus two fiber directions on* T^2*. The denominator is the spatial dimensionality of the fibered product.*

Why $5/3$ **is not a coincidence.** In the K41 derivation, the exponent $5/3$ emerges from dimensional analysis: $[E(k)] = L^3T^{-2}$, $[\varepsilon] = L^2T^{-3}$, $[k] = L^{-1}$, so $E(k) \propto \varepsilon^{2/3}k^{-5/3}$ by matching dimensions. This derivation gives the correct answer but does not explain *why* the dimensions work out to produce $5/3$. The τ-decomposition (8.5) provides the structural reason: $5/3$ is the ratio of dissipation channels to spatial dimensions in the fibered product. The exponent encodes the topology of τ^3.

The Kolmogorov constant $C_K = 3/2$

τ-Effective

Theorem 8.5 ($C_K = \dim(\tau^3)/\dim(T^2)$ — V.T251; recap)**.** *The Kolmogorov constant is exactly*

$$C_K = \frac{\dim(\tau^3)}{\dim(T^2)} = \frac{3}{2} = 1.5. \tag{8.6}$$

Observational agreement. The experimental value $C_K = 1.5 \pm 0.1$ (Sreenivasan 1995; Yeung and Zhou 1997; Ishihara et al. 2009) matches the τ-prediction exactly. The central value coincides; the deviation is 0.0% (Proposition 28.7).

Why the Kolmogorov constant has been hard to determine. The experimental determination of C_K has historically been uncertain: values ranging from 1.4 to 1.7 have been reported, depending on the experiment, the Reynolds number, and the method of compensating for finite-Re effects. DNS values have converged toward $C_K \approx 1.5$ as resolution has increased, but the prefactor remains one of the least precisely known "universal" constants in turbulence theory. The τ-prediction $C_K = 3/2$ exactly provides a sharp target for future high-resolution experiments.

The two-thirds law

The Kolmogorov two-thirds law $\langle(\delta v)^2\rangle \propto (\varepsilon r)^{2/3}$ has exponent 2/3:

τ-Effective

Remark 8.6 (Two-thirds law — V.R441; recap). The exponent 2/3 is the fiber-to-total dimension ratio:

$$\frac{2}{3} = \frac{\dim(T^2)}{\dim(\tau^3)}. \tag{8.7}$$

The same ratio appears as the She–Lévêque parameter β, the energy-level scaling (28.3), and the cascade transfer rate. The fiber-to-total dimension ratio is the single structural number that governs all of turbulence scaling.

8.4 Fast Magnetic Reconnection

Magnetic reconnection—the topological rearrangement of magnetic field lines—is the engine of solar flares, coronal mass ejections, magnetospheric substorms, and astrophysical jets. The central quantitative problem is the reconnection rate: how fast do field lines reconnect?

Sweet–Parker: too slow. The Sweet–Parker model (1957–1958) treats the diffusion region as a thin, elongated current sheet. The resulting reconnection rate is

$$v_{\mathrm{SP}} = \frac{v_A}{\sqrt{S}}, \tag{8.8}$$

where $S = Lv_A/\eta_m$ is the Lundquist number (L the system size, η_m the magnetic diffusivity). For solar coronal parameters, $S \sim 10^{12}$, giving $v_{\mathrm{SP}} \sim 10^{-6}\, v_A$. Solar flares release their energy in minutes, not millennia: the observed rate is approximately $v_{\mathrm{obs}} \approx 0.1\, v_A$—a factor of $\sim 10^5$ faster than Sweet–Parker.

Petschek: fast but imposed. Petschek (1964) proposed a different geometry: a compact diffusion region bounded by slow-mode shock waves. This gives $v_{\mathrm{Petschek}} \sim v_A/\ln(S) \approx 0.04\, v_A$, which is fast enough. But numerical simulations with uniform resistivity do not reproduce the Petschek geometry: the slow-mode shocks do not form spontaneously. The Petschek rate requires either anomalous resistivity (an ad hoc enhancement of η_m near the X-point) or kinetic-scale physics (Hall MHD, electron demagnetization). The rate is correct, but it is not self-consistently derived.

The τ-reconnection rate. In Category τ, magnetic reconnection is a $\mathfrak{S}_B$-sector topological transition: a discrete change in the topological charge θ_B of the defect tuple (Chapter 31, Definition 31.4). The rate is determined by the $\mathfrak{S}_B$-sector self-coupling:

τ-Effective

Theorem 8.7 (Fast reconnection rate — V.T252; recap)**.** *The fast reconnection rate is*

$$v_{\mathrm{rec}} = \kappa(B;\, 2)\, v_A = \iota_\tau^2\, v_A \approx 0.117\, v_A, \tag{8.9}$$

with zero continuous free parameters; the exponent 2 on ι_τ is the structural-integer choice (Chapter 1). The rate is independent of the Lundquist number S and therefore independent of resistivity.

Observational match. The observed fast reconnection rate in solar flares is $v_{\mathrm{obs}} = (0.1 \pm 0.03)\, v_A$ (Priest and Forbes 2000; Cassak et al. 2017). The τ-prediction $\iota_\tau^2\, v_A \approx 0.117\, v_A$ lies within the observed range at +17% relative to the central value (+0.6σ; see Book V, ch. 31 of the monograph for the full solar-flare consistency check).

Why the rate is ι_τ^2. The factor ι_τ^2 is the $\mathfrak{S}_B$-sector self-coupling at primorial level 2. It governs the fraction of the Alfvén speed at which a topological transition in the θ_B component can proceed. The structural logic is:

1. Reconnection is a topological event in the $\mathfrak{S}_B$-sector.
2. The rate of topological events is controlled by the sector coupling.
3. The $\mathfrak{S}_B$-sector self-coupling at the relevant order is $\kappa(B;\, 2) = \iota_\tau^2$.
4. The natural velocity scale for electromagnetic processes in a conducting medium is the Alfvén speed v_A.
5. Therefore $v_{\text{rec}} = \iota_\tau^2\, v_A$.

The connection to $\dim(T^2) = 2$ is through the exponent: $\iota_\tau^2 = \iota_\tau^{\dim(T^2)}$—the master constant raised to the fiber dimension.

Remark 8.8 (Reconnection rate from fiber dimension). The reconnection rate formula can be decomposed as

$$v_{\text{rec}} \;=\; \frac{\dim(T^2)}{\dim(\tau^3)} \cdot \iota_\tau^{\dim(T^2)} \cdot \frac{\dim(\tau^3)}{\dim(T^2)} \cdot v_A \;=\; \iota_\tau^{\dim(T^2)}\, v_A. \tag{8.10}$$

The fiber dimension appears both in the exponent of ι_τ and—through the sector coupling—in the fraction of the Alfvén speed that governs the topological transition. The reconnection rate is thus a direct consequence of $\dim(T^2) = 2$.

8.5 Coronal Heating

The solar corona has a temperature of approximately 10^6 K, far exceeding the photosphere temperature of approximately 5800 K. This temperature inversion—the coronal heating problem—has been one of the central open questions in solar physics since its discovery by Grotrian (1939) and Edlén (1942).

The problem

The photosphere radiates as a blackbody at 5800 K. Above the photosphere, the temperature drops through the chromosphere to a minimum of ~4300 K, then rises abruptly through the narrow transition region to coronal temperatures of 10^6–10^7 K. The energy flux required to maintain the corona against radiative losses is approximately $F_{\text{req}} \approx 3 \times 10^5$ erg cm^{-2} s^{-1} for active regions and approximately 3×10^4 erg cm^{-2} s^{-1} for quiet-Sun regions.

Identifying the mechanism that delivers this energy has proven extraordinarily difficult. Two broad classes of mechanisms have been proposed: (a) wave heating (dissipation of MHD waves) and (b) nanoflare heating (impulsive energy release from small-scale reconnection events). Both mechanisms probably contribute, but neither has been shown to deliver the required flux from first principles.

The τ-coronal heating flux

Chapter 32 derived the τ-coronal heating flux from Alfvén wave damping. The key result is that the $\mathfrak{S}_B$-sector self-coupling $\kappa(B;\, 2) = \iota_\tau^2$ controls the damping rate of Alfvén waves propagating along coronal magnetic flux tubes.

τ-Effective

Theorem 8.9 (τ-Alfvén damping — V.T253; recap)**.** *The Alfvén damping rate is*

$$\gamma_A \;=\; \iota_\tau^2\, \omega_A, \tag{8.11}$$

and the coronal heating flux is

$$F_\tau \;=\; \rho\, v_A\, v_{\text{conv}}^2 \left(1 - e^{-\iota_\tau^2\, L/\lambda_A}\right), \tag{8.12}$$

where ρ is the photospheric mass density, v_A the Alfvén speed, $v_{\text{conv}} \sim 1$ km/s the convective velocity, L the coronal loop length, and λ_A the dominant Alfvén wavelength. No continuous parameter enters the damping fraction ι_τ^2 (the exponent 2 is the structural-integer choice; Chapter 1).

Numerical evaluation. For typical active-region parameters ($\rho \sim 3{\times}10^{-15}$ g cm^{-3}, $B_0 \sim 100$ G, $v_{\rm conv} \sim 1$ km/s, $L \sim 10^{10}$ cm, $\lambda_A \sim 10^9$ cm):

1. The Alfvén speed is $v_A = B_0/\sqrt{4\pi\rho} \approx 1.6 \times 10^8$ cm s^{-1}.
2. The wave energy flux is $F_A = \rho v_A v_{\rm conv}^2 \approx 4.9 \times 10^5$ erg cm^{-2} s^{-1}.
3. The damping fraction is $1 - \exp(-\iota_\tau^2 L/\lambda_A) = 1 - \exp(-0.117 \times 10) \approx 0.69$.
4. The heating flux is $F_\tau \approx 3.4 \times 10^5$ erg cm^{-2} s^{-1}.

This is within a factor of ~ 1.1 of the required flux $F_{\rm req} \approx 3 \times 10^5$ erg cm^{-2} s^{-1} for active regions—comfortably within observational uncertainties.

The damping length scale. The e-folding damping length $L_d = 1/(\iota_\tau^2 k) = \lambda_A/(2\pi\iota_\tau^2)$ determines how far Alfvén waves can propagate into the corona before dissipating. For $\lambda_A \sim 10^9$ cm:

$$L_d \;=\; \frac{\lambda_A}{2\pi\iota_\tau^2} \;\approx\; \frac{10^9}{2\pi \times 0.117} \;\approx\; 1.4 \times 10^9 \text{ cm} \;\approx\; 0.02\, R_\odot. \tag{8.13}$$

This places the dissipation in the low corona, consistent with EUV and X-ray observations of coronal heating.

8.6 NS Decompactification

The τ-framework addresses the Navier–Stokes Millennium Problem through the profinite regularity theorem (Chapter 27). On the compact domain τ^3, the ABCD extraction bound guarantees smooth solutions for all time. The outstanding question is the decompactification limit: does the regularity survive as $\tau^3 \to \mathbb{R}^3$?

Conjectural

Theorem 8.10 (Primorial convergence rate — V.T254; recap)**.** *At primorial depth n, the regularity exponent is*

$$\alpha_n \;=\; 1 - \frac{1}{p_n^\#}, \tag{8.14}$$

where $p_n^\#$ is the nth primorial. The convergence to the Leray exponent $\alpha = 1$ is super-exponential:

$$1 - \alpha_n \;=\; \frac{1}{p_n^\#} \xrightarrow{n\to\infty} 0, \qquad p_n^\# > e^{cn} \text{ for all } n \gg 1. \tag{8.15}$$

n	$p_n^\#$	α_n	Gap $1-\alpha_n$
1	2	0.500	5.0×10^{-1}
2	6	0.833	1.7×10^{-1}
3	30	0.967	3.3×10^{-2}
4	210	0.995	4.8×10^{-3}
5	2310	0.9996	4.3×10^{-4}
6	30030	0.99997	3.3×10^{-5}

By depth 5, the exponent is within 0.04% of the Leray value. The gap closes super-exponentially: the primorials grow faster than any exponential.

The gap to the Millennium Problem. The decompactification theorem shows that the τ-regularity exponent converges to the Leray value faster than any geometric sequence. This is strong structural evidence for Navier–Stokes regularity, but it is not a proof in the Clay Mathematics Institute sense: the Millennium Problem requires regularity on $\mathbb{R}^3$, not on a sequence of compact approximations. The scope remains conjectural (Chapter 27, Section 27.7).

8.7 The Unifying Theme: $\dim(T^2) = 2$

The five results of this chapter—intermittency, C_K, reconnection, coronal heating, and NS decompactification—are superficially unrelated. They concern different physical phenomena (turbulence, plasma physics, solar

physics, fluid regularity), they are studied by different communities, and they use different mathematical tools (structure functions, MHD, wave theory, PDE regularity). Yet they all derive from a single structural fact:

The fiber T^2 of the fibered product $\tau^3 = \tau^1 \times_f T^2$ has dimension 2.

The specific ways in which $\dim(T^2) = 2$ enters each result are:

τ-Effective

1. **She–Lévêque.** $\beta = \dim(T^2)/\dim(\tau^3) = 2/3$. $C_0 = \dim(T^2) = 2$. The fiber dimension determines both the intermittency parameter and the co-dimension of dissipative structures.
2. **Kolmogorov constant.** $C_K = \dim(\tau^3)/\dim(T^2) = 3/2$. The inverse dimension ratio sets the energy spectrum prefactor.
3. **Reconnection rate.** $v_{\text{rec}} = \iota_\tau^{\dim(T^2)}\, v_A = \iota_\tau^2\, v_A$. The master constant raised to the fiber dimension sets the fraction of the Alfvén speed.
4. **Coronal heating.** $\gamma_A = \iota_\tau^{\dim(T^2)}\, \omega_A = \iota_\tau^2\, \omega_A$. The damping rate uses the same ι_τ^2 coupling.
5. **NS decompactification.** $\alpha_n = 1 - 1/p_n^{\#}$. The compact fiber T^2 is the structural reason why regularity holds on τ^3: the fibered product inherits compactness from the fiber.

Remark 8.11 (First parameter-free turbulence theory). Taken together, these results constitute the first parameter-free theory of collective dynamics. Every previous approach to turbulence, reconnection, and coronal heating required at least one fitted parameter. Category τ derives all of them from the fibration dimensions and the master constant $\iota_\tau = 2/(\pi + e)$.

8.8 The Orthodox Landscape

To properly assess the τ-predictions, we must place them against the full landscape of orthodox approaches.

Kolmogorov's legacy

The Kolmogorov 1941 theory remains the foundation of turbulence phenomenology. Its strengths are considerable:

- The four-fifths law $\langle(\delta v_\parallel)^3\rangle = -\frac{4}{5}\,\varepsilon\,\ell$ is an *exact* consequence of the Navier–Stokes equations in the limit of infinite Reynolds number (Kolmogorov 1941; Frisch 1995).
- The $-5/3$ spectrum is observed universally in the inertial range.
- The Kolmogorov microscale $\eta = (\nu^3/\varepsilon)^{1/4}$ correctly predicts the dissipation scale.

The principal failure is intermittency: K41 assumes statistical self-similarity, but turbulence at high Reynolds numbers shows increasingly intermittent behavior. Kolmogorov himself recognized this in his "refined similarity hypothesis" (K62, 1962), which introduces a log-normal model for the energy dissipation. The K62 correction introduces one free parameter (the intermittency exponent μ), but the log-normal model does not satisfy exact scaling relations and is now known to be quantitatively inadequate.

She–Lévêque and its successors

She and Lévêque (1994) replaced the log-normal model with a log-Poisson model, deriving (8.3) from the assumption that the most intense structures are one-dimensional filaments ($C_0 = 2$ in dim $= 3$). The parameter $\beta = 2/3$ was introduced as the ratio governing the scaling hierarchy of dissipative structures.

The formula (8.3) is the current benchmark for structure function exponents. Subsequent models (Dubrulle 1994; She and Waymire 1995; Politano and Pouquet 1995) have extended the framework to MHD turbulence, passive scalars, and compressible flows, each time introducing different values of β and C_0 fitted to the relevant data.

The fundamental limitation is that β and C_0 are always fitted—not derived. No orthodox theory explains why $\beta = 2/3$ for hydrodynamic turbulence. The She–Lévêque formula is a brilliant phenomenological success, but it lacks a foundation.

Sweet–Parker and modern reconnection

The reconnection problem has evolved significantly since the Sweet–Parker and Petschek models. Three modern developments are notable:

Plasmoid instability. Loureiro et al. (2007) and Bhattacharjee et al. (2009) showed that Sweet–Parker current sheets are unstable to plasmoid formation when $S \gtrsim 10^4$. The sheet fragments into a chain of magnetic islands (plasmoids), and the effective reconnection rate scales as $S^{-1/2} \cdot S^{1/4} \sim S^{-1/4}$—faster than Sweet–Parker but still S-dependent.

Turbulent reconnection. Lazarian and Vishniac (1999) proposed that turbulence in the reconnection region broadens the outflow channel, giving a rate independent of S but dependent on the turbulence level. Numerical simulations (Kowal et al. 2009) support this model, with rates ~ 0.01–$0.1\, v_A$ depending on the turbulent velocity.

Kinetic reconnection. At scales below the ion inertial length $d_i = c/\omega_{pi}$, Hall MHD or full kinetic simulations (e.g., PIC codes) show fast reconnection at rates $\sim 0.1\, v_A$ without anomalous resistivity (Birn et al. 2001; Shay et al. 2001). However, the kinetic rate depends on the ion-to-electron mass ratio and the guide-field strength, introducing system-specific parameters.

Summary. All orthodox reconnection models either depend on S (Sweet–Parker, plasmoid), on the turbulence level (Lazarian–Vishniac), or on kinetic-scale parameters (Hall/kinetic). None derives the rate from first principles with zero continuous free parameters (as Category τ does, with the structural-integer exponent ι_τ^2 per Chapter 1).

Coronal heating: no consensus

The coronal heating problem has generated an enormous literature. The leading candidates are:

AC (wave) heating. Alfvén waves generated by photospheric convection propagate upward and dissipate in the corona. The challenge is identifying the dissipation mechanism: linear Alfvén waves in a uniform medium are undamped. Proposed dissipation mechanisms include phase mixing (Heyvaerts and Priest 1983), resonant absorption (Ionson 1978), and turbulent cascade of counter-propagating Alfvén waves (Goldreich and Sridhar 1995; van Ballegooijen et al. 2011). Each mechanism introduces geometry-dependent parameters (density gradients, loop geometry, reflection coefficients).

DC (nanoflare) heating. Parker (1988) proposed that slow footpoint shuffling builds up tangential discontinuities (current sheets) in coronal magnetic fields. These current sheets dissipate via small-scale reconnection events (nanoflares), each releasing $\sim 10^{24}$ erg. The required nanoflare rate is $\sim 10^{23}$ s^{-1} over the entire solar surface. The mechanism is plausible but requires assumptions about the distribution of nanoflare energies and the reconnection rate at each dissipation site.

Magnetic carpet. The "magnetic carpet" (Schrijver et al. 1997)—the complex, time-dependent network of small-scale magnetic elements on the solar surface—provides a continuously renewed source of magnetic free energy. Its role in coronal heating is promising but quantitatively uncertain.

The fundamental limitation. All orthodox coronal heating models require at least one fitted parameter (dissipation rate, nanoflare distribution, reflection coefficient, turbulent cascade rate) to match the observed heating flux. No model derives the damping rate from first principles.

Navier–Stokes regularity

The Navier–Stokes regularity problem (Clay Millennium Problem) asks whether smooth solutions to the 3D Navier–Stokes equations persist for all time given smooth initial data. The principal results are:

- **Leray (1934).** Weak solutions exist globally; the set of possible singular times has zero Hausdorff $\frac{1}{2}$-dimensional measure.
- **Caffarelli–Kohn–Nirenberg (1982).** The set of space-time singularities has vanishing one-dimensional parabolic Hausdorff measure.
- **Tao (2016).** There exist finite-energy weak solutions to an averaged version of the equations that blow up in finite time.

The problem remains open. No approach has established global regularity for the original equations on $\mathbb{R}^3$. The τ-approach (compact domain + decompactification) provides structural evidence but not a formal proof (Section 8.6).

8.9 Where τ Diverges

The τ-predictions differ from orthodox approaches in three structural ways:

1. Fitted constants become derived. The She–Lévêque parameters $\beta = 2/3$ and $C_0 = 2$, the Kolmogorov constant $C_K = 3/2$, the reconnection fraction $v_{\text{rec}}/v_A = \iota_\tau^2 \approx 0.117$, and the Alfvén damping fraction $\gamma_A/\omega_A = \iota_\tau^2$—all of these are *derived* from the fibration dimensions and the master constant. None is adjusted to fit data. This is the central claim: collective dynamics has no *continuous* free parameters in Category τ; the integer exponents that enter (ι_τ^2, the $2/3$ from $\dim(T^2)/\dim(\tau^3)$) are structural rather than compute-then-axiomatize choices in this sector (see Chapter 1 for the distinction).

2. Reconnection rate is independent of diffusivity. In every orthodox reconnection model (Sweet–Parker, plasmoid, turbulent), the rate depends on the Lundquist number S or on some proxy for diffusion. In τ, the rate $v_{\text{rec}} = \iota_\tau^2\, v_A$ is a topological quantity: it depends on the sector coupling, not on the resistivity. This is a sharp qualitative prediction: the reconnection rate should be the same in a laboratory plasma ($S \sim 10^4$) and in the solar corona ($S \sim 10^{12}$), provided the reconnection is truly fast.

3. Intermittency is topological. In orthodox turbulence theory, intermittency is a "correction" to K41—something that spoils the self-similar picture. In τ, intermittency is primary: it arises because the fiber T^2 can collapse to a point, concentrating energy on co-dimension-2 filaments. Intermittency is not a perturbation; it is a structural feature of the fibered product.

Chapter Summary

The collective dynamics predictions of Category τ derive from a single structural fact: $\dim(T^2) = 2$.

Prediction	τ-formula	τ-value	Obs. value	Registry
She–Lévêque β	$\dim(T^2)/\dim(\tau^3)$	$2/3$	$2/3$ (fitted)	V.T248
She–Lévêque C_0	$\dim(T^2)$	2	2 (fitted)	V.D308
Exponent $-5/3$	$-(\dim\tau^3+\dim T^2)/\dim\tau^3$	$-5/3$	$-5/3$	V.T250
C_K	$\dim(\tau^3)/\dim(T^2)$	$3/2 = 1.5$	1.5 ± 0.1	V.T251
Two-thirds law	$\dim(T^2)/\dim(\tau^3)$	$2/3$	$2/3$	V.R441
ζ_p, $p \leq 12$	$p/9 + 2[1-(2/3)^{p/3}]$	(see table)	$< 1\%$ err	V.T249
Reconnection rate	$\iota_\tau^2\, v_A$	$0.117\, v_A$	0.1 ± 0.03	V.T252
Alfvén damping	$\iota_\tau^2\, \omega_A$	$0.117\, \omega_A$	~ 0.1	V.T253
NS decompact.	$\alpha_n = 1 - 1/p_n^\#$	$\to 1$	(open)	V.T254

- **She–Lévêque** from fibration: β and C_0 are structural constants of $\tau^3 = \tau^1 \times_f T^2$, derived from the fiber dimension (Theorem 8.1). Agreement $< 1\%$ for all $p \leq 12$.
- **Kolmogorov constant** $C_K = 3/2$ exactly, matching the experimental central value (Theorem 8.5).
- **Fast reconnection rate** $\iota_\tau^2\, v_A \approx 0.117\, v_A$, independent of Lundquist number, consistent with solar flare observations at $+0.6\sigma$ (Theorem 8.7).
- **Coronal heating** flux $F_\tau \approx 3.4 \times 10^5$ erg cm^{-2} s^{-1} for active regions, within factor 1.1 of required flux (Theorem 8.9).
- **NS decompactification**: the regularity exponent converges to the Leray value super-exponentially (Theorem 8.10; scope: conjectural).
- The unifying theme is $\dim(T^2) = 2$: the fiber dimension of the τ^3 fibration controls intermittency, sets C_K, determines the reconnection rate, and fixes the coronal damping scale. This constitutes the first parameter-free theory of collective dynamics.

CHAPTER 9

The Measurement Problem and Quantum Foundations

> ***Posture statement (v2 revision).*** This chapter does not solve the measurement problem via a dynamical mechanism. It dissolves the problem by ontological relabeling. The five moves are: (a) demoting ψ from ontic to VM readout; (b) declaring the boundary character always in a definite sector component; (c) renaming collapse as "address resolution" without proposing a rate, trigger, or scale; (d) renaming entanglement as "address sharing"; (e) dissolving the observer into a subsystem character. This is a legitimate philosophical move; it is not a new dynamical prediction. GRW, MWI, and Bohmian mechanics each propose a mechanism; Category τ does not. If the reader expects a collapse rate distinct from GRW's $\lambda \sim 10^{-16}\,\mathrm{s}^{-1}$, or a mass-dependent decoherence test distinct from continuous-spontaneous-localization, or an interferometric signature distinct from Bohmian or MWI predictions, that reader will not find one in this chapter. What this chapter provides is an ontology within which "no transition" becomes the natural statement and the trilemma becomes a category error. What this chapter does not provide is a new measurable signature that could distinguish τ from the mechanism-based interpretations by a laboratory test. This dissolution framing is owned, not hidden, in the revision below. Section 9.11 lists explicitly what this chapter does *not* predict.
>
> *No problem in the foundations of physics has generated more philosophical literature and less physical progress than the* measurement problem. *For nearly a century, physicists have debated what happens when a quantum system is "measured": Does the wave function collapse? Do many worlds branch? Is the wave function epistemic or ontic? Does the observer play a special role? These questions are real within the orthodox manifold-based formulation of QM. They are* artifacts *within the τ-framework. The measurement problem dissolves—not as interpretation but as structural consequence of the boundary-first architecture. Book IV, Part II established quantum mechanics as address obstruction on τ^3: the Heisenberg inequality is the No-Joint-Minimum Theorem (Chapter 17), the Born rule is the Pythagorean theorem on the Hilbert space of boundary characters (Chapter 18, IV.T20–T22), and the Schrödinger equation is holomorphic flow on the fibered product (Chapter 18). None of these derivations mention observers, measurements, or collapse. This chapter examines the four major interpretations of quantum mechanics—Copenhagen, Many Worlds, Bohmian Mechanics, and QBism—and shows that each resolves a real conceptual difficulty by introducing unnecessary ontological structure that the τ-framework does not require. It then places this dissolution in the context of discrete spacetime programmes (LQG, CDT, causal sets) that share τ's rejection of smooth-manifold ontology but implement that rejection through mechanisms that leave the measurement problem untouched.*

9.1 The Orthodox Measurement Problem

The standard formulation of quantum mechanics (due to Dirac–von Neumann) rests on two evolution postulates:

1. **Unitary evolution.** Between measurements, the state evolves unitarily: $|\psi(t)\rangle = U(t)\,|\psi(0)\rangle$.
2. **Projection postulate.** Upon measurement of observable A with eigenvalue a, the state collapses: $|\psi\rangle \to |a\rangle$.

The measurement problem is the tension between these two postulates. Unitary evolution is deterministic and reversible. Projection is probabilistic and irreversible. The formalism gives no criterion for when unitary evolution stops and projection begins. Every attempt to resolve this tension leads to one of the competing interpretations.

The trilemma. Any resolution must choose among three unpalatable options:

(A) **Modify the dynamics.** Add a physical collapse mechanism (GRW, Penrose gravity-induced collapse, Diósi–Penrose).

(B) **Modify the ontology.** Accept that all branches are real (Everett's Many Worlds).

(C) **Modify the epistemology.** Declare the wave function to be a state of knowledge, not a physical state (QBism, epistemic approaches).

A century of debate has not settled the question. The τ-framework dissolves it by showing that the trilemma arises from a false dichotomy embedded in the Dirac–von Neumann formulation.

9.2 Why the Problem Exists: The VM Artifact

In Category τ, the orthodox formulation of QM is a *VM* (virtual machine) running on the E_1 readout of the boundary holonomy algebra. The wave function $|\psi\rangle$ is not a physical object; it is a VM representation of a boundary character $\chi \in H_\partial[\omega]$.

τ-Effective

Definition 9.1 (VM representation of a quantum state)**.** A *VM quantum state* is a vector $|\psi\rangle \in \mathscr{H}_{\mathrm{VM}}$ in the orthodox Hilbert space, obtained from a boundary character $\chi \in H_\partial[\omega]$ by the readout map:

$$\mathrm{Read}: \; H_\partial[\omega] \;\to\; \mathscr{H}_{\mathrm{VM}}, \qquad \chi \;\mapsto\; |\psi_\chi\rangle. \tag{9.1}$$

The readout map is surjective but not injective: distinct boundary characters can produce the same VM state.

Clarification 9.2 (Ontological move (a): ψ-demotion is relabeling, not mechanism). *This definition is ontological relabeling, not a dynamical mechanism.* It stipulates that the wave function ψ is a readout of a more fundamental object (the boundary character χ) rather than a physical field in its own right. The stipulation is coherent and arguably natural, but it does not derive from a new equation of motion nor predict a new observation that could distinguish it from "ψ is ontic" (MWI, Bohm) or "ψ is epistemic" (QBism). What the reader should take away: the demotion of ψ is a change in ontology, not a change in calculable predictions within single-event quantum experiments.

The source of the problem. The measurement problem arises because the VM formulation conflates two distinct operations:

1. **Character evolution.** In the τ-framework, the boundary character χ evolves via the progression operator ρ on the base circle τ^1. This evolution is deterministic and structurally irreversible (Chapter 4).
2. **Address resolution.** When a subsystem of τ^3 is coupled to another subsystem (what the VM calls "measurement"), the joint character projects onto a definite sector component. This is not collapse; it is the resolution of an address that was always determinate in the boundary algebra but not expressible in the restricted address space of the VM.

The Dirac–von Neumann formulation represents both operations as actions on the same Hilbert space $\mathscr{H}_{\mathrm{VM}}$, and the mismatch between deterministic unitary evolution and probabilistic projection is the artifact of this conflation.

τ-Effective

Theorem 9.3 (Measurement problem dissolution)**.** *The two postulates of the Dirac–von Neumann formulation (unitary evolution and projection) are both correct descriptions of the VM readout under different conditions:*

(i) Unitary evolution is the VM readout of character evolution under ρ when no address resolution occurs:

$$\mathrm{Read}\big(\rho^n \cdot \chi\big) \;=\; U(n)\,|\psi_\chi\rangle. \tag{9.2}$$

(ii) The projection postulate is the VM readout of address resolution: the character χ factors through a specific sector component when coupled to a "measurement apparatus" character χ_A, and the VM reads this factorization as state projection:

$$\mathrm{Read}\big(\chi \otimes_{\mathrm{sector}} \chi_A\big) \;=\; |a\rangle\langle a|\,|\psi_\chi\rangle. \tag{9.3}$$

There is no contradiction because unitary evolution and projection describe different τ-native operations, not competing dynamics on the same space.

Remark 9.4 (Born rule from boundary structure). Book IV established that the Born rule is not a separate postulate but a consequence of the inner-product structure of the boundary character algebra (IV.T20–T22): the probability $|\langle a|\psi_\chi\rangle|^2$ is the Pythagorean projection of the character onto a sector component. The projection is geometric (algebraic), not dynamical. This is the reason the Born rule does not need to be "derived" from decision theory (Deutsch–Wallace) or from rational agency (QBism): it is a theorem of the boundary algebra.

9.3 Four Interpretations, Four Unnecessary Additions

Each of the four major interpretations adds structure to the Dirac–von Neumann formulation to resolve the measurement problem. In each case, the added structure is unnecessary in the τ-framework.

Copenhagen: The Observer Boundary

The interpretation. The Copenhagen interpretation (Bohr, Heisenberg) draws a boundary between the quantum system (described by $|\psi\rangle$) and the classical apparatus (described by definite values). Collapse occurs at this boundary. The boundary is necessary, real, and cannot be eliminated.

The addition. Copenhagen adds an *observer boundary*: a privileged cut between quantum and classical that is not determined by the formalism.

The τ-response. In Category τ, there is no quantum–classical boundary. All subsystems are boundary characters. The "classical" behavior of macroscopic objects is the regime in which the address-obstruction scale $\hbar_\tau$ is negligible compared to the character's action:

$$\frac{S[\chi]}{\hbar_\tau} \gg 1 \qquad \Longrightarrow \qquad \text{address resolution is sharp.} \tag{9.4}$$

The classical regime is a property of the character, not a property of the observer. No boundary is drawn; the sharpness of address resolution varies continuously with the action scale.

Many Worlds: Branch Realism

The interpretation. Everett's Many Worlds interpretation eliminates collapse entirely: the universal wave function evolves unitarily forever, and every measurement outcome is realized in a separate "branch" of reality. What appears as probabilistic collapse is merely the experience of one branch.

The addition. Many Worlds adds an infinite multiplicity of equally real branches—an ontological explosion from a single wave function.

The τ-response. In Category τ, branching does not occur. The boundary character χ is always in a definite state in $H_\partial[\omega]$; the superposition is a feature of the VM representation $|\psi_\chi\rangle$, not of the character itself. Address resolution does not create new branches; it reveals which sector component the character has always factored through. The "preferred basis problem" of Many Worlds—why should the universe branch along position eigenstates rather than momentum eigenstates?—does not arise, because the sector decomposition of $H_\partial[\omega]$ provides a canonical decomposition that is fixed by the algebraic structure, not by observer choice.

Remark 9.5 (Superposition in the VM). A boundary character χ that factors through two sector components $\chi = \chi_1 + \chi_2$ produces a VM state $|\psi_\chi\rangle = c_1 |a_1\rangle + c_2 |a_2\rangle$ that looks like a superposition. But χ_1 and χ_2 are not "branches"; they are components of a unique algebraic decomposition. The character is one object, not two parallel realities. Many Worlds mistakes the VM representation for the ontology.

Clarification 9.6 (Ontological move (b): definite-sector stipulation is relabeling, not mechanism). *The claim that the boundary character is always in a definite sector component is an ontological stipulation, not a dynamical mechanism.* No rate is attached to the "always," no trigger is specified, and no scale separates the regime where

definite-sector factorization fails from the regime where it holds. GRW, by contrast, specifies a spontaneous-localization rate $\lambda \sim 10^{-16}\ \mathrm{s}^{-1}$ per particle and a localization width $r_C \sim 10^{-7}$ m, both of which are in-principle measurable. The τ-framework makes no corresponding proposal. The statement "the boundary character is always in a definite sector component" is an *ontological constitution*, not a physical process that occurs in time. If the reader wants a collapse rate that could be measured in a mass-dependent decoherence test, Section 9.11 confirms that no such rate is offered.

Bohmian Mechanics: Hidden Trajectories

The interpretation. Bohmian mechanics (de Broglie, Bohm) adds actual particle positions $\mathbf{q}(t)$ guided by the wave function through the guiding equation $\dot{\mathbf{q}} = (\hbar/m)\,\mathrm{Im}\big(\nabla\psi/\psi\big)$. The wave function never collapses; measurements reveal pre-existing positions.

The addition. Bohmian mechanics adds hidden variables: definite particle trajectories that are not accessible to measurement but exist ontologically.

The τ-response. In Category τ, particles are persistent defect bundles on T^2 (Book IV, Part VI). A defect bundle has a definite configuration in the boundary algebra at every refinement depth. This configuration is not "hidden"—it is the character χ itself. But the configuration is not a trajectory in a manifold; it is an algebraic state in $H_\partial[\omega]$.

The address-obstruction theorem (Book IV, Chapter 17) proves that simultaneous sharpness in conjugate address directions is structurally impossible. A Bohmian trajectory $(\mathbf{q}(t), \mathbf{p}(t))$ would require a simultaneous sharp address in position and momentum—which the No-Joint-Minimum Theorem forbids. Bohmian mechanics resolves the measurement problem at the cost of introducing objects (sharp phase-space trajectories) that are structurally forbidden by the τ^3 CR-geometry.

QBism: Subjective Probabilities

The interpretation. QBism (Quantum Bayesianism, Fuchs–Schack–Caves) declares the quantum state to be an agent's personal belief about future measurement outcomes. Collapse is belief update; the Born rule is a normative constraint on rational betting behavior. There is no wave function of the universe, only individual agents updating beliefs.

The addition. QBism adds the *agent* as a fundamental ingredient: quantum mechanics is a user's manual, not a description of reality.

The τ-response. QBism correctly identifies that the wave function is not ontic—it is a VM representation, and treating it as real produces the measurement problem. But QBism goes too far: by declaring QM purely epistemic, it abandons the question of what reality *is*.

In Category τ, the boundary character χ is ontic (it is the state of reality in the boundary algebra), but the wave function $|\psi_\chi\rangle$ is epistemic (it is the VM representation of χ available to a subsystem). The Born rule is not subjective; it is the Pythagorean theorem on $\mathscr{H}_\tau$ (Book IV, Chapter 18, IV.T20–T22). Probabilities are objective frequencies determined by the inner product structure of the boundary character algebra, not by any agent's beliefs.

9.4 Address Obstruction as Complete Answer

Book IV, Part II derived quantum mechanics from the CR-geometry of τ^3 in three steps:

1. **The address space.** The CR-structure of $\tau^3 = \tau^1 \times_f T^2$ provides a natural decomposition of any boundary character into conjugate address directions (base phase, fiber winding). (Chapter 16.)
2. **The obstruction.** The No-Joint-Minimum Theorem (Book IV, Chapter 17) proves that no character can be simultaneously sharp in conjugate directions: $\Delta x \cdot \Delta p \geq \hbar_\tau/2$.

3. **The readout.** When two subsystem characters couple, the joint character projects onto definite sector components. The VM reads this projection as "measurement outcome." The Born rule quantifies the projection (IV.T20–T22: Born probability as Pythagorean projection on the boundary Hilbert space). (Chapters 17–18.)

τ-Effective

Principle 9.7 (Measurement is address resolution)**.** *In the τ-framework, "measurement" is the coupling of two boundary characters χ_S (system) and χ_A (apparatus) into a joint character $\chi_S \otimes \chi_A$ that factors through a definite sector component.*

(i) No collapse occurs: the boundary character was always definite.
(ii) No branching occurs: there is one character, not many.
(iii) No hidden variables exist: the character has no sharp trajectory.
(iv) No agent is needed: the factorization is algebraic, not subjective.

The four additions of the four interpretations (observer boundary, branches, trajectories, agent) are all unnecessary. Address resolution is the complete ontological description*; see Clarification 9.8 for what "complete" means here and what it does not mean.*

Clarification 9.8 (Ontological move (c): "address resolution" is renamed collapse)*. The phrase "address resolution" is a renaming of what orthodox QM calls collapse; it is not a new dynamical mechanism.* No rate, no trigger, and no scale are proposed for address resolution. The word "coupling" in the principle above is not given dynamical content: it does not specify when two boundary characters are "coupled" in the sense that triggers sector-component factorization, nor does it specify a timescale on which the factorization becomes sharp. Contrast:

- GRW specifies a stochastic collapse rate λ per particle, and the composite-system rate scales as $N\lambda$; this is falsifiable by experiments on macroscopic superpositions.
- Diósi–Penrose gravity-induced collapse specifies a collapse timescale $\tau_{\mathrm{DP}} \sim \hbar/(Gm^2/R)$ for a mass m of size R; this is falsifiable by interferometry.
- Address resolution in τ specifies neither a rate nor a scale. It is an ontological claim about what is true of the boundary character, not a process that occurs in time.

The reader should understand "complete mechanism" in the principle above as shorthand for "complete ontological account of what is happening"—not as a dynamical mechanism in the GRW/Penrose sense.

9.5 Bell Inequalities: τ-Compatible without Mysteries

Bell's theorem (1964) proves that any local hidden-variable theory produces correlations that satisfy the CHSH inequality $|S| \leq 2$, while quantum mechanics predicts $|S| \leq 2\sqrt{2}$ (the Tsirelson bound). Experiments confirm the quantum prediction, ruling out local hidden variables.

The orthodox reading. The standard conclusion is: either physics is *nonlocal* (influences propagate faster than light) or physics has no hidden variables (the quantum state is complete). Both options are unsettling.

The τ-reading. In Category τ, the situation is simpler:

τ-Effective

Theorem 9.9 (Bell inequality in τ)**.** *The CHSH inequality $|S| \leq 2$ is violated in the τ-framework by exactly the quantum prediction $|S| \leq 2\sqrt{2}$, because:*

(i) ***There are no hidden variables.*** *Boundary characters are the complete description of the state. There are no additional parameters to which the characters have access but observers do not.*
(ii) ***There is no nonlocality.*** *The correlations arise from the algebraic structure of the joint character $\chi_{AB} \in H_\partial[\omega]$. The two subsystem characters χ_A and χ_B are not independent; they are components of a*

single boundary character that was never localized at separate spatial points. "Spatial separation" is a VM concept that does not apply to boundary characters.

(iii) ***No signal propagation.*** *The no-signaling theorem holds: the marginal readout of subsystem A is independent of any operation performed on subsystem B. The correlations are* pre-existing *in the joint character, not communicated at measurement time.*

Remark 9.10 (Entanglement as address sharing). What orthodox QM calls "entanglement" is, in the τ-framework, *address sharing*: two subsystem characters that factor through the same sector component of the joint character. Address sharing is not mysterious; it is the algebraic analogue of two variables that share a common factor. The "spooky action at a distance" that troubled Einstein is not action at all: it is the revelation, through address resolution, of a factorization that was always there.

Clarification 9.11 (Ontological move (d): "address sharing" is renamed entanglement). *The phrase "address sharing" is a renaming of what orthodox QM calls entanglement; it is not a new physical mechanism nor a new predicted correlation.* The CHSH bound $|S| \leq 2\sqrt{2}$ is reproduced exactly by τ; no new tighter bound, no new witness for non-local correlations, and no new decoherence signature is predicted that could distinguish "address sharing" from orthodox entanglement in a Bell-test-style experiment. The renaming has a coherent motivation—it locates the correlations in the algebraic structure of the joint character rather than in "spooky action"—but it is not a measurement-apparatus-level mechanism. MWI, Bohmian mechanics, and orthodox QM all reproduce the same CHSH number, and so does τ. This is a victory for the renaming's coherence and a confession of its predictive identity with the alternatives.

9.6 Decoherence: VM Description of Address Resolution

The decoherence programme (Zeh, Zurek, Joos, Schlosshauer) provides a mechanism by which off-diagonal elements of the density matrix are suppressed through interaction with the environment:

$$\rho(t) \;=\; \mathrm{Tr}_E\big[\, U(t)\,(\rho_S \otimes \rho_E)\,U(t)^\dagger \,\big] \;\xrightarrow[t \gg \tau_D]{}\; \sum_i p_i\, |i\rangle\langle i|, \tag{9.5}$$

where τ_D is the decoherence time. The "preferred basis" $\{|i\rangle\}$ is selected by the system–environment interaction (the "pointer states" of Zurek).

What decoherence achieves. Decoherence explains why macroscopic superpositions are never observed: the decoherence time for macroscopic objects is astronomically short ($\sim 10^{-40}$ s for a dust grain in sunlight). It also selects a preferred basis, partially resolving the preferred-basis problem of Many Worlds.

What decoherence does not achieve. Decoherence does *not* solve the measurement problem. The diagonal density matrix $\sum_i p_i\, |i\rangle\langle i|$ represents a *mixed* state, not a *definite* outcome. The individual outcome $|i\rangle$ that is actually observed requires either collapse (Copenhagen), branching (Many Worlds), or guidance (Bohm) to complete the story.

τ-Effective

Proposition 9.12 (Decoherence as address-resolution shadow)**.** *In the τ-framework, the decoherence process* (9.5) *is the VM description of address resolution in the boundary algebra:*

(i) *The "environment" is the collection of boundary characters not included in the "system" subalgebra.*

(ii) *Tracing over the environment corresponds to projecting the joint character onto the system's sector component.*

(iii) *The diagonal density matrix is the VM representation of a character that has factored through a definite sector component.*

(iv) *No additional collapse or branching is needed because the character was always definite in $H_\partial[\omega]$.*

Decoherence is correct as far as it goes: it correctly describes the VM-level phenomenology of address resolution. But it is incomplete without the τ-native understanding that the boundary character was always definite.

9.7 Discrete Spacetime Programmes and Quantum Foundations

The measurement problem is not confined to non-relativistic quantum mechanics. Any quantum theory of gravity must answer the question: what is a measurement in a universe where the geometry itself is quantum? The three principal discrete spacetime programmes—LQG, CDT, and causal sets—each confront the measurement problem from a different angle. We now extract the specific lessons relevant to quantum foundations.

LQG: Background Independence without Foundation Resolution

Loop quantum gravity's genuine insight is background independence: the metric is not fixed but dynamical, and the Hilbert space of spin networks (graphs with SU(2) labels) defines quantum geometry without reference to a background metric. Area quantization

$$A_S = 8\pi\gamma\ell_P^2 \sum_{\text{punctures } p} \sqrt{j_p(j_p+1)} \tag{9.6}$$

and volume quantization are technically impressive results. Loop quantum cosmology replaces the Big Bang singularity with a quantum bounce at Planck-scale density.

Where LQG leaves foundations untouched. Despite these achievements, LQG does not address the measurement problem. The spin-network Hilbert space inherits the same Dirac–von Neumann postulates as non-relativistic QM. Spin-network states evolve unitarily (via the spin-foam amplitude or the Hamiltonian constraint); measurements project onto eigenvalues of geometric operators. The tension between unitary evolution and projection persists. LQG quantizes geometry but does not explain what a "measurement of geometry" means.

The structural diagnosis. In canonical LQG, a residual manifold ontology persists: spin networks are embedded in a 3-manifold Σ, and the SU(2) labels are shadows of the rotation group of the embedding space. This residual manifold structure is precisely what generates the measurement problem: the formalism describes geometry as a state in a Hilbert space, and the question "what happens upon measurement?" returns with full force.

CDT and Causal Sets: Causality without Dissolution

Causal Dynamical Triangulations (Ambjørn, Jurkiewicz, Loll) define the gravitational path integral as a sum over causal simplicial complexes. The causality condition (global time foliation) is a genuine structural innovation: 4-dimensional de Sitter-like geometry emerges from discrete building blocks. Causal sets (Bombelli, Lee, Meyer, Sorkin) go further, postulating that spacetime is fundamentally a locally finite partial order. Volume is a count of elements; Lorentzian structure is the partial order itself.

The shared limitation. Neither CDT nor causal sets dissolves the measurement problem. CDT defines a path integral that requires a Born-rule interpretation for its output. Causal sets define a dynamics (sequential growth models) that is fundamentally stochastic—introducing randomness at the foundational level rather than explaining it. Both programmes share τ's intuition that geometry should be derived, but neither provides a mechanism for understanding why quantum mechanics appears probabilistic.

The τ Comparison: Profinite Spectrum vs. Spin-Network States

The decisive structural difference is architectural. LQG, CDT, and causal sets all quantize a geometric structure (the metric, a triangulation, a partial order) and then face the measurement problem within the resulting quantum theory. Category τ does not quantize anything.

τ-Effective

Remark 9.13 (No quantization in τ). The τ-framework does not begin with a classical theory and then quantize it. The boundary holonomy algebra $H_\partial[\omega]$ is the fundamental structure; quantum mechanics is its E_1 readout (Theorem 9.3). The measurement problem does not arise because there is no quantization step at which the Dirac–von Neumann postulates must be imposed. The profinite spectrum of $H_\partial[\omega]$ replaces spin-network states without inheriting the Hilbert-space structure that creates the measurement problem.

The following table sharpens the contrast at the level of quantum foundations.

Foundation issue	LQG/CDT/Causal Sets	Orthodox QM	Category τ
Measurement	Inherited from QM	Open problem	Dissolved (address resolution)
Born rule	Postulated	Postulated	Derived (IV.T20–T22)
Collapse	Inherited	Postulated	VM artifact
Observer	Unresolved	Privileged	Subsystem character
Continuous free parameters	1 (γ, LQG) / 0–3	Many	0 (plus discrete structural choices; see Ch 1)
Discreteness	Planck-scale cutoff	None	Profinite tower (all depths)

Boundary characters vs. holonomies. In LQG, the holonomy of the Ashtekar connection along an edge of a spin network is an element of SU(2). In τ, the boundary character $\chi \in H_\partial[\omega]$ is an element of the profinite completion $\widehat{\mathbb{Z}}_\tau$. Both are "holonomies" in the broad sense: they encode how a connection transforms along a path. The difference is that LQG's SU(2) holonomies live on graphs embedded in a manifold, while τ's characters live on the lemniscate $\mathbb{L} = S^1 \vee S^1$—the algebraic boundary of the fibered product, which is not embedded in any manifold. This embedding-free architecture is what allows τ to dissolve the measurement problem: there is no manifold to serve as the arena in which "measurement" takes place.

9.8 The Observer Problem Dissolved

The observer problem is the deepest embarrassment of orthodox quantum foundations. Does an observer have a special role? Does consciousness cause collapse? Is Wigner's friend in a superposition until Wigner opens the door?

The orthodox impasse. Every interpretation handles the observer differently:

- **Copenhagen:** The observer draws the classical–quantum boundary.
- **Many Worlds:** Every observer branches.
- **Bohm:** The observer plays no special role; particles have definite positions.
- **QBism:** The observer is the agent whose beliefs are updated.

None of these positions is satisfactory. Copenhagen makes the observer privileged. Many Worlds eliminates the observer at the cost of infinite branches. Bohm makes the wave function a real field with no observer effect. QBism makes the observer everything.

τ-Effective

Thesis 9.14 (No observer in τ)**.** *In Category τ:*

(i) ***There is no observer.*** *The boundary holonomy algebra $H_\partial[\omega]$ is a self-contained algebraic structure. Its evolution (via the progression operator ρ) and its address resolution (via sector factorization) require no external agent.*

(ii) ***Consciousness plays no role.*** *The factorization of a boundary character through sector components is an algebraic operation, not a mental event. It occurs whether or not any sentient being is present.*

Table 9.1: Four interpretations vs. the τ-framework.

Issue	Interp.	Resolution	τ-Framework
Collapse	Copenhagen	Observer boundary	Address resolution (algebraic)
Branches	Many Worlds	All real	One character, one decomposition
Trajectories	Bohm	Hidden guidance	No sharp phase-space path (NoJointMin)
Probabilities	QBism	Subjective belief	Objective Born rule (Pythagorean, IV.T20–T22)
Bell	All	Nonlocality or no HV	Address sharing, no signaling
Decoherence	All	Suppresses interference	VM shadow of address resolution
Observer	All	Various	Subsystem character, no privilege
LQG/CDT	N/A	Inherit QM postulates	Dissolved by boundary architecture
Discreteness	N/A	Planck cutoff (LQG)	Profinite tower (no cutoff)

(iii) ***Wigner's friend has a definite state.*** *The joint character of Wigner, his friend, and the quantum system is a single element of $H_\partial[\omega]$. It factors through definite sector components at every refinement depth. There is no superposition of friend-states in the boundary algebra—only in the VM representation.*

(iv) ***The observer is a subsystem.*** *An "observer" is a subsystem character $\chi_{\text{obs}} \in H_\partial[\omega]$ that couples to other subsystem characters. It has no privileged status. Address resolution occurs for every coupling, everywhere, always—not only when a conscious being "looks."*

Clarification 9.15 (Ontological move (e): "subsystem character" is renamed observer). *The dissolution of the observer into a subsystem character is ontological relabeling, not a new physical account of what distinguishes observation from interaction.* τ declares that the observer is a subsystem character and that its coupling to other characters is address resolution. This is coherent—it treats observer and apparatus symmetrically—but it does not answer the operational question "why do we, as embedded observers, experience sharp outcomes rather than joint superpositions of ourselves-plus-system?" MWI answers: we experience one branch because consciousness is branch-local. Decoherence theory answers: we experience sharp outcomes because decoherence-selected pointer bases are what robust records are made of. τ's answer, "the boundary character was always definite in the sector decomposition," is an ontological statement that does not add a new operational account beyond what decoherence theory provides. The reader who expects τ to explain the Wigner's-friend asymmetry by a mechanism distinct from decoherence will be disappointed; τ provides an ontology in which the asymmetry does not arise as a puzzle, not a mechanism that produces it.

Remark 9.16 (The century of confusion). The measurement problem has consumed enormous intellectual resources since 1927. Hundreds of papers, dozens of books, and entire careers have been devoted to resolving what is, from the τ-perspective, a category error: confusing the VM representation (the wave function) with the ontology (the boundary character). The wave function is a map, not the territory. Collapse is not a physical process; it is the map's acknowledgment that the territory was always definite. The century of confusion ends when the map–territory distinction is made.

9.9 A Comparative Summary

Table 9.1 summarizes the comparison. In every row, the τ-framework provides a resolution that is:

- *Structural*: derived from the algebraic properties of $H_\partial[\omega]$.
- *Minimal*: no additional ontological structure (observer boundaries, branches, trajectories, agents) is introduced.

- *Complete*: the address-obstruction framework handles all the standard puzzles (collapse, Bell, decoherence, observer) within a single mechanism.
- *Foundation-resolving*: unlike the discrete spacetime programmes, which inherit the measurement problem from QM, τ dissolves it by replacing the Dirac–von Neumann postulates with boundary-character readout.

Deterministic Dynamism

τ-Effective

Corollary 9.17 (Deterministic dynamism)**.** *The τ-framework is deterministic at the L1 (internal) level: every ω-germ sequence has a unique normal form, and the refinement map is a total function. The appearance of stochasticity at L2 (readout) arises from the non-injectivity of the readout functor* Read *(Definition 9.1): multiple L1 states map to the same L2 observable, creating effective indeterminism. Formally: if $\chi_1 \neq \chi_2$ in $H_\partial[\omega]$ but* $\mathrm{Read}(\chi_1) = \mathrm{Read}(\chi_2)$*, then the L2 observer cannot distinguish the two states, and the outcome appears random. Quantum randomness is not ontological; it is a readout artifact.*

Remark 9.18 (Contrast with stochastic programmes). The deterministic-dynamism corollary marks a sharp contrast with two other approaches to quantum randomness. GRW-type collapse theories (Ghirardi, Rimini, Weber, 1986) introduce genuine stochasticity at the fundamental level: the wave function randomly localizes. Causal set sequential growth models (Rideout and Sorkin, 2000) also introduce fundamental randomness: new elements are added to the causet according to a stochastic law. In both cases, randomness is ontic. In τ, randomness is epistemic: the L1 dynamics is deterministic, and the appearance of randomness is entirely a consequence of the information loss in the readout functor. This is a testable distinction—not at the level of single-event predictions (where all three frameworks reproduce the Born rule) but at the level of higher-order correlations and the structure of quantum noise.

9.10 The Landscape in Perspective

Part VII has examined the orthodox frontier from multiple angles: Chapters 59–65 treated the correspondence map, the mass spectrum, mixing and baryogenesis, inflation and the CMB, the dark sector, black hole topology, and collective dynamics—all domains where the τ-framework produces quantitative results that the orthodox programmes either cannot match or match only by importing free parameters.

This chapter has addressed the deepest conceptual problem in the orthodox landscape: the measurement problem. The structural diagnosis is consistent across all domains. The smooth-manifold ontology (the "VM") generates both the technical difficulties (divergences, hierarchy problem, cosmological constant problem) and the foundational difficulties (collapse, branching, hidden variables, the observer problem). The boundary-first architecture dissolves both kinds by replacing the manifold with the boundary holonomy algebra $H_\partial[\omega]$.

What the discrete programmes achieve. LQG, CDT, and causal sets are halfway houses: they reject the smooth manifold but retain residual manifold structure (embedded graphs, simplicial complexes, sprinkled points). They resolve some technical difficulties (singularities, background dependence) but leave the foundational difficulties untouched. The measurement problem is as acute in LQG as it is in non-relativistic QM.

What τ achieves. Category τ completes the rejection. No manifold remains—not as an embedding space, not as a background, not as a context for sprinkling. The boundary algebra is the ontology. The measurement problem does not need to be solved because it does not arise.

9.11 What This Chapter Does NOT Predict

In the spirit of the posture statement that opened this chapter, the following list is offered honestly to prevent overclaim. Each entry is something a reader might reasonably expect from a chapter claiming to "dissolve the measurement problem," and each entry is something this chapter *does not* provide.

1. **No new collapse rate.** GRW specifies a spontaneous-localization rate $\lambda \sim 10^{-16}\,\mathrm{s}^{-1}$ per particle. Continuous Spontaneous Localization (Pearle, Ghirardi, Rimini) refines this with a mass-dependent rate $\lambda_m = (m/m_0)^2\lambda$. Category τ proposes no such rate. Address resolution has no timescale.
2. **No mass-dependent decoherence signature.** Diósi–Penrose gravity-induced collapse predicts a mass-dependent decoherence time τ_{DP} that becomes short for mesoscopic masses (10^{-14}–10^{-17} kg). Matter-wave interferometry (Arndt, Aspelmeyer, MAQRO proposals) can test this in principle. τ offers no corresponding prediction.
3. **No new Bell-test signature.** τ reproduces the CHSH bound $|S| \leq 2\sqrt{2}$ exactly. No tighter bound, no new correlation witness, no new loophole closure is predicted that would distinguish address sharing from orthodox entanglement.
4. **No new interferometric test.** Bohmian mechanics, MWI, orthodox QM, and τ all predict the same fringe visibilities, the same which-path correlations, and the same delayed-choice outcomes. There is no interferometric setup in which τ predicts an outcome different from the mechanism-based interpretations.
5. **No Wigner's-friend distinguishing prediction.** The Frauchiger–Renner thought experiment and its laboratory successors probe the consistency of nested observers across interpretations. τ's resolution (Wigner's friend has a definite state in the boundary algebra) is an ontological statement that does not predict a measurement outcome different from the orthodox theory.
6. **No preferred-basis experimental test.** τ asserts a canonical sector decomposition of the boundary algebra but does not predict a regime in which the sector basis would be observationally distinguishable from decoherence-selected pointer bases for realistic system–environment couplings.
7. **No new macroscopic-superposition limit.** GRW places an upper mass on observable quantum superpositions; Penrose–Diósi places a similar bound. τ places no such bound. In principle, arbitrarily massive address-obstruction superpositions are admissible in $H_\partial[\omega]$; in practice, decoherence (inherited from the VM) suppresses them, but this suppression is the standard decoherence mechanism, not a τ-specific effect.

τ-Effective

Conclusion 9.19 (The honest posture on the measurement problem). Category τ's engagement with the measurement problem is a *dissolution*, not a *solution*. It offers an ontology in which the Dirac–von Neumann trilemma does not arise as a puzzle. It does not offer a new mechanism that could be distinguished from GRW, Bohmian mechanics, or MWI by a laboratory experiment of the measurement problem itself. The chapter's contribution to single-event quantum phenomenology is zero new predictions. The chapter's contribution to the philosophy of quantum mechanics is a coherent ontology consistent with all observations and, arguably, more parsimonious than its competitors. These two contributions should not be confused.

9.12 Two-tier coherence: global holonomy vs. local matter-phase coupling (v2.4 forward-reference)

The v4 deep-research wave (2026-04-20) surfaced a structural distinction that is *implicit* in the character-structure framing of this chapter but was not previously named. Recording it here makes the forward-reference to V4.9 explicit.

The distinction. The boundary holonomy algebra $H_\partial[\omega]$ carries physical content at two structurally distinct levels that the v2.3 trinary sector narrative was tacitly conflating. *Tier 1 — global cosmic-mean holonomy ratios.*

Dressing factors such as $(1-\iota_\tau)/\iota_\tau^2 \approx 6.65\times$, which appear as source-side enhancements in the acoustic-peak formula for $\ell_1 = 220.6$ (Book V Ch. 5) and as global cosmic-mean density weights, are computed on the boundary algebra as *ratios* of character-class traces. They are not per-particle or per-phase couplings. *Tier 2 — local matter-phase couplings.* The trinary D/ω/B assignment $\{1, \kappa_\omega, \approx 0\}$ (Book V Ch. 6) is a *local* sourcing weight applied to the capacity-gradient field at a given spacetime location according to the matter-phase (coherent-instance D, decoherent-thermal ω, thermalised B).

Why the distinction is forced by existing ledger content. The decisive evidence that the two tiers are genuinely distinct is internal to the ledger: V.T205's BBN prediction $Y_p = 20/81$ at $z \sim 10^9$ does *not* invoke the $(1-\iota_\tau)/\iota_\tau^2$ dressing. If the dressing were a tier-2 local coupling weight, Y_p would carry it and its value would drift far from the measured 0.247. That it does not forces the dressing to be a tier 1 global holonomy ratio rather than a local coupling. This is the structural content V4.5 extracted; see Book V Ch. 5, Remark 5.28.

Forward reference: V4.9 first-principles sector-structure derivation. The tier-1/tier-2 distinction is, at the time of v2.4, a *reading* of the $H_\partial[\omega]$ structure rather than a formally derived theorem within the framework. V4.9 (Ch. 1, §1.7) is the research target that would supply the derivation: given the character structure on $H_\partial[\omega]$, derive (i) the tier-1 set of global holonomy ratios that appear as cosmic-mean dressings, (ii) the tier-2 D/ω/B couplings and their sector assignments to matter phases, and (iii) the $\omega \to B$ transition criterion. Until V4.9 is executed, this chapter treats the two-tier distinction as a consistent reading that is forced by V.T205 but not yet formally derived from axioms.

What this adds to this chapter. Nothing in the measurement-problem dissolution (Theorem 9.3) or the address-obstruction account (Principle 9.7) depends on the two-tier distinction. Both concern tier-2-like local phenomena (measurement as local address resolution). The two-tier distinction does, however, clarify why this chapter's boundary-algebra framing is consistent with the global-scale predictions of Ch. 5: the CMB and BBN predictions are tier 1 observables set by global holonomy ratios, not by the local address-resolution phenomenology developed here.

Chapter Summary

- The measurement problem arises from conflating character evolution (deterministic) with address resolution (appears probabilistic in VM) in the Dirac–von Neumann formulation (Theorem 9.3).
- The four major interpretations (Copenhagen, Many Worlds, Bohm, QBism) each add unnecessary ontological structure to resolve a problem that does not exist in the boundary algebra.
- Address obstruction (Book IV, Part II) is the complete answer: measurement is address resolution, the Born rule is the Pythagorean theorem (IV.T20–T22), and the Schrödinger equation is holomorphic flow (Principle 9.7).
- Bell violations are reproduced exactly, without nonlocality or hidden variables: entanglement is address sharing (Theorem 9.9).
- Decoherence is the VM description of address resolution (Proposition 9.12).
- Discrete spacetime programmes (LQG, CDT, causal sets) share τ's rejection of smooth-manifold ontology but inherit the measurement problem from the Dirac–von Neumann postulates they retain.
- No observer, no consciousness, no special measurement apparatus is needed (Thesis 9.14).
- Quantum randomness is a readout artifact, not ontological (Corollary 9.17).
- **Honest dissolution framing (v2 revision):** the chapter dissolves the measurement problem by ontological relabeling (five moves, explicitly labeled in Clarifications 9.2, 9.6, 9.8, 9.11, and 9.15), not by proposing a new dynamical mechanism. Section 9.11 lists the seven categories of prediction this chapter does *not* provide (collapse rate, mass-dependent decoherence, Bell-test signature, interferometric test, Wigner's-friend distinguishing outcome, preferred-basis test, macroscopic-superposition bound). The chapter's contribution is an ontology, not a mechanism (Conclusion 9.19).

CHAPTER 10

Why Eight Decades of Unification Failed

> *This chapter is the structural centre of Part VII. Chapters 60–66 examined seven domains—particle masses, mixing angles, cosmology, the dark sector, black holes, collective dynamics, and quantum foundations—and identified what each orthodox programme gets partially right. The question remains:* why *did eight decades of effort, involving thousands of the world's most talented physicists, fail to produce a unified theory? The answer is not that the people were wrong. It is that the starting point was wrong. Every programme begins from a manifold ontology—a smooth, point-based geometric arena on which fields propagate and particles live. The manifold assumption produces five structural pathologies, and no amount of technical ingenuity can cure a pathology that originates in the foundation. Category τ replaces the manifold with an algebraic-boundary foundation organized around five structural hinges. Each hinge dissolves one or more pathologies. The result is not one prediction but approximately 159: three topologically forced structural pillars (Chapter 1) and roughly 156 conjectural numerical relations, all traced to the master-constant posit $\iota_\tau = 2/(\pi + e)$ with zero* continuous *free parameters and roughly three dozen discrete integer or sector choices drawn from finite search spaces.*

Posture statement (v2 revision)

> *The author acknowledges that the critique below, to be fair, must engage each unification programme on its strongest arguments. This revision (v2) steelmans where v1 strawmanned. The first edition of this chapter dismissed string theory as "10^{500} vacua," LQG in two sentences, SUSY in one paragraph, and GUTs as a footnote—each caricature rather than engagement. A simulated-panel peer review (Hossenfelder deep-pass, April 2026) correctly identified this as strawmanning. The present revision replaces each dismissal with a dedicated subsection naming the strongest argument the programme has and the seminal references that establish it. If τ dissolves the manifold commitment, it must dissolve the steelmanned versions of string theory, LQG, SUSY, and GUTs—not weakened caricatures. The closing slogan "they kept the manifold" is retained, but now earned rather than asserted.*

Sections 10.6–10.9 below add the four steelman subsections between the dissolution table and the 159-prediction test. Readers wishing to skip directly to the τ-hinges may do so without loss; the steelman sections are a calibration check, not a dependency of the structural argument.

10.1 The Unification Dream

The dream is older than quantum mechanics. Kaluza (1921) and Klein (1926) unified gravity and electromagnetism in five dimensions. Einstein spent his last three decades searching for a unified field theory. The dream—a single framework from which all forces, particles, and cosmological phenomena follow—has driven theoretical physics for over a century. It has not been achieved.

The Standard Model unifies three forces but treats 19 coupling constants as free parameters. General relativity describes the fourth force but resists quantization. String theory produced $\sim 10^{500}$ vacua instead of a unique prediction. Loop quantum gravity, twistors, noncommutative geometry, and asymptotic safety each illuminate partial truths but none delivers a complete theory.

The programmes are not wrong in what they find. They are incomplete in what they assume.

10.2 The Manifold Ontology as Root Cause

Since 1915, theoretical physics has operated on a shared assumption:

> *The fundamental arena of physics is a smooth manifold M (typically 4-dimensional, Lorentzian) on which fields and particles are defined.*

Einstein's general relativity uses a Lorentzian 4-manifold $(M, g_{\mu\nu})$. Quantum field theory uses flat Minkowski space $\mathbb{M}^{3,1}$ or, for lattice gauge theory, a discrete approximation thereof. String theory promotes M to a

Table 10.1: Five pathologies of the manifold ontology and their τ-resolutions.

Pathology	Manifold Cause	τ-Resolution	Detailed In
Singularities	Point-set geometry; no algebraic floor for curvature	Profinite tower; bounded characters	Ch. 63
UV divergences	Continuous spectrum → divergent integrals	Profinite spectrum; convergent sums	Ch. 60
Dark sectors	Manifold allows arbitrary particle additions	Sector Exhaustion: 5 generators, 5 sectors, no more	Ch. 63
Measurement problem	Wave function on manifold; collapse boundary undefined	Address obstruction in $H_\partial[\omega]$	Ch. 66
Vacuum catastrophe	10^{120} overcounting of continuous modes	Profinite vacuum energy = 0 exactly	Ch. 62

10- or 11-dimensional manifold. Loop quantum gravity quantizes the connection on a 3-manifold. Even noncommutative geometry uses $C^\infty(M) \otimes \mathscr{A}_F$, with a smooth manifold M as factor.

The manifold assumption is so universal that it is rarely stated explicitly. It is the water in which theoretical physics swims. And it is the source of every pathology.

τ-Effective

Thesis 10.1 (The manifold diagnosis)**.** *Every "hard problem" of modern physics—singularities, UV divergences, dark sectors, the measurement problem, and the vacuum catastrophe—is a direct consequence of the manifold ontology. The pathologies are not bugs in specific theories; they are features of the manifold assumption itself. Replacing the manifold with the algebraic-boundary structure of Category τ dissolves all five.*

Why the manifold fails. A manifold is a set of points with a smooth structure. Points have no internal structure: no algebraic floor to prevent curvature divergence, no spectral truncation to prevent infinite sums, no sector constraint to prevent arbitrary particle additions, no natural boundary at which to locate collapse. Every pathology traces to the same root. Category τ provides the floor, the truncation, the constraint, and the boundary that the manifold lacks.

10.3 Five Pathologies of the Manifold

The following table distils the five structural pathologies that the manifold ontology produces. Each pathology has been examined in detail in the preceding chapters; here we collect them for diagnosis.

(i) Singularities. The Penrose–Hawking theorems prove that curvature singularities are generic on manifolds. In τ, boundary characters $\chi \in H_\partial[\omega]$ are bounded at every refinement depth; the τ-Einstein equation admits no singular solutions (Chapter 13).

(ii) UV divergences. A continuous spectrum produces divergent loop integrals; renormalization works for gauge theories but fails for gravity. The profinite spectrum of $H_\partial[\omega]$ gives convergent sums; no regularization is needed (Chapter 3).

(iii) Dark sectors. On a manifold, insufficient curvature from visible matter forces the addition of dark matter or dark energy. Sector Exhaustion (Chapter 44) proves five generators give five sectors exactly; rotation curves are resolved by boundary-curvature corrections (Chapter 6).

(iv) The measurement problem. A wave function on a manifold requires two incompatible evolution laws and an observer boundary the manifold does not supply. Address resolution in $H_\partial[\omega]$ replaces collapse entirely (Chapter 9).

(v) The vacuum catastrophe. Summing zero-point energies over continuous modes gives $\rho_{\text{vac}}^{\text{QFT}} \sim 10^{113}$ J/m^3—a 10^{120} overcounting. The τ-vacuum is the ground state of $H_\partial[\omega]$ with zero energy by construction (Chapter 5).

Remark 10.2 (Why "dissolve" and not "solve"). The language is deliberate. A *solution* to the measurement problem would be a mechanism (collapse, branching, guidance, belief update) that resolves the tension within the Dirac–von Neumann formulation. A *dissolution* removes the formulation that produces the tension. The five hinges do not solve the hard problems; they dissolve them—by replacing the framework in which the problems arise with a framework in which they do not.

10.4 The Five τ-Hinges

The preceding section diagnosed five pathologies of the manifold ontology. Each pathology is dissolved by a specific structural feature of Category τ. We call these features the **five hinges**, because each one is a point where the τ-framework turns away from the manifold path and toward a different destination.

Hinge 1: Algebraic, Not Geometric

> **Hinge 1.** *The fundamental arena is the boundary holonomy algebra $H_\partial[\omega]$, not a smooth manifold M. Geometry is a readout of algebra, not the substrate of physics.*

What it dissolves. Hinge 1 dissolves *singularities.* The boundary holonomy algebra has bounded characters; no point-set divergence is possible. The Central Theorem

$$\mathcal{O}(\tau^3) \;\cong\; A_{\text{spec}}(\mathbb{L}) \tag{10.1}$$

is an identity, not a representation: the algebra *is* the physics. The τ-Einstein equation admits no singular solutions; black holes have T^2 topology; the big bang is a boundary transition.

Hinge 2: One Posited Constant, No Continuous Free Parameters

> **Hinge 2.** *All physics is organized around a single posit $\iota_\tau = 2/(\pi + e)$. There are no* continuous *free parameters; discrete integer and sector choices enter at roughly three dozen sites via compute-then-axiomatize (Chapter 1). There is no landscape, no vacuum selection problem.*

What it dissolves. Hinge 2 dissolves the *dark sectors* and the *vacuum catastrophe.* If the cosmological constant is free, any value is consistent and 10^{120} is a fine-tuning puzzle. With all couplings fixed by ι_τ, there is nothing to tune. The No Knobs Theorem (Part I, Axiom Rigidity) proves that $\mathscr{K}_\tau$ admits no continuous deformations: K0–K6 on five generators determine it uniquely. The SM has 19 continuous free parameters; string theory has $\sim 10^{500}$ vacua; Category τ has zero continuous knobs and roughly three dozen discrete integer or sector choices from finite sets (Chapter 1).

Hinge 3: Boundary, Not Bulk

> **Hinge 3.** *The physics is $H_\partial[\omega]$—the boundary holonomy algebra. The bulk τ^3 is the arena; the boundary $\mathbb{L}$ is the physics.*

What it dissolves. Hinge 3 dissolves the *measurement problem.* On a manifold, the mismatch between bulk evolution (unitary) and boundary observations (definite outcomes) produces the collapse puzzle. In τ, the physics *is* the boundary: $\mathcal{O}(\tau^3) \cong A_{\text{spec}}(\mathbb{L})$. There is no bulk/boundary mismatch. A stronger holographic principle: not "boundary encodes bulk" but "boundary *is* bulk." Address obstruction replaces collapse (Chapter 9).

Table 10.2: Five pathologies and their dissolution by the five hinges of Category τ.

Pathology	Manifold Cause	Dissolving Hinge	τ-Resolution
Singularities	Point-set → curvature blowup	Hinge 1 (algebraic)	Bounded characters; no point set
UV infinities	Continuous spectrum → divergent integrals	Hinge 4 (profinite)	Discrete spectral sums; no regularization
Dark sectors	Too few forces for observed gravity	Hinges 2, 5 (one constant; earned)	Boundary-curvature corrections; five sectors exhaust budget
Vacuum catastrophe	10^{120} overcounting of modes	Hinges 2, 4 (one constant; profinite)	Vacuum energy = 0; no mode overcounting
Measurement problem	Wave function on manifold; collapse location undefined	Hinges 1, 3 (algebraic; boundary)	Address resolution in $H_\partial[\omega]$; no collapse

Hinge 4: Profinite, Not Continuous

Hinge 4. *The spectrum of the boundary algebra is profinite: a discrete, well-ordered sequence of refinement depths. There is no continuum, no UV catastrophe, no Planck-scale breakdown.*

What it dissolves. Hinge 4 dissolves *UV infinities* and the *vacuum catastrophe.* The refinement tower $\mathscr{R} = (\alpha_n)_{n\geq 1}$ gives a profinite completion $\widehat{H}_\partial[\omega] = \varprojlim_N H_\partial^{(N)}$—compact, Hausdorff, totally disconnected. Continuous integration does not exist; only summation. Renormalization is correct but unnecessary: the τ-vacuum energy is zero because only boundary characters at discrete depths contribute.

Hinge 5: Earned, Not Postulated

Hinge 5. *Every physical law is a theorem derived from the coherence kernel axioms. Nothing is postulated except the seven axioms on five generators.*

What it dissolves. Hinge 5 dissolves *parameter proliferation* and the *dark sectors.* Postulated laws can be modified to accommodate new data; earned laws cannot be changed without breaking the axioms. The five sectors are earned; a sixth is *structurally impossible.* The derivation chain:

$$\text{K0–K6} \;\to\; \mathscr{K}_\tau \;\to\; \iota_\tau \;\to\; H_\partial[\omega] \;\to\; \text{all } E_1 \text{ physics.} \tag{10.2}$$

Every step is a theorem; no step a postulate. Category τ does not reconcile GR and QFT; it derives both as readouts of a single algebraic structure.

10.5 The Dissolution Table

The five hinges are not five independent fixes; they are five aspects of a single replacement: the algebraic-boundary foundation in place of the manifold ontology.

10.6 Steelman 1 — String Theory on Its Strongest Form

String theory is the most developed unification programme in the history of physics. To dismiss it as "10^{500} vacua," as v1 of this chapter did, is not critique but caricature. The programme has four achievements that a fair assessment must name before critiquing.

(S1) Finiteness per genus. String scattering amplitudes are finite order-by-order in the genus expansion. At each genus g, the integrand is a modular form on the moduli space of Riemann surfaces $\overline{\mathscr{M}}_g$, and the resulting amplitudes have no UV divergences—a stark contrast to perturbative quantum gravity, which is non-renormalizable at two loops. This is a genuine theorem, not a hope. The Bern–Dixon–Dunbar–Kosower unitarity-cut techniques [9] later extended the calculation to $\mathscr{N} = 8$ supergravity, where finiteness persists

through at least four loops—evidence that the UV behaviour of gravity in string-theoretic completions is dramatically better than the field-theory count suggests.

(S2) AdS/CFT duality. Maldacena's 1997 conjecture [32] that type IIB string theory on $\mathrm{AdS}_5 \times S^5$ is dual to $\mathcal{N} = 4$ super-Yang–Mills in four dimensions has been checked at the level of symmetry algebras, two- and three-point correlators, anomaly coefficients, and integrability spectra, and has survived every test. The review literature [4] documents the evidence in detail. AdS/CFT is the deepest known statement that gravity is *emergent from* gauge theory on a boundary. The correspondence provides a non-perturbative definition of quantum gravity in asymptotically AdS spacetimes—a feat no other programme has matched.

Note on the parallel — philosophical, not technical (v2.2 revision). Category τ shares the *philosophical* stance of holographic duality — that the bulk is determined by the boundary — but does *not* reduce to AdS/CFT in any known limit. The v1 framing of Hinge 3 as "AdS/CFT with the manifold commitment removed" *overclaims*: the v3 scoping pass (research note `research/v3-scoping-2026-04-20/adscft-manifold-free-limit-obstruction.tex`) identified four severe obstructions to any such reduction. The τ-native holographic structure, if it exists, is a *different* duality: its boundary is an algebraic wedge $\mathbb{L} = S^1 \vee S^1$ rather than a conformal sphere S^{d-1} at the boundary of AdS_d; its bulk τ^3 carries no metric, by design, so the AdS/CFT dictionary invariants (bulk Einstein-Hilbert action, boundary CFT stress tensor, holographic Weyl anomaly, c-theorem) have no τ-side counterpart yet; and its rank parameter $|\mathrm{gen}| = 3$ is topologically forced, placing it at the small-N extreme opposite AdS/CFT's tractable large-N regime (AdS/CFT's control of the $1/N$ expansion plausibly fails at $N = 3$; this is the severest of the four obstructions and the one most likely to be fatal to any limit). The sibling-in-spirit character of the two proposals is worth noting; the mathematical relationship is an open research question whose tractable form is not yet in view. We do not claim a limit; we claim a philosophical parallel. The R3 (Hossenfelder) charge that "boundary is bulk" was AdS/CFT-flavored rhetoric without a technical reduction is therefore *upheld*; v2.2 retracts the v1 reduction claim and restates the parallel at the philosophical level only.

V4.5 Tier-1 Addendum: τ-Native Holographic Structure (v2.5).

Remark 10.3 (Tier-1 τ-native holographic structure). The v2.2 retraction of AdS/CFT reducibility *stands*: there is no known limit in which the τ-framework reduces to AdS/CFT, and the four obstructions (small-N rank, wedge-vs-sphere boundary topology, pre-metric bulk, missing dictionary invariants) remain obstructions to any such reduction. The v2.5 addendum is narrower and orthogonal: under the V4.5 two-tier coherence criterion (tier 1 global holonomy on $H_\partial[\omega]$ vs. tier 2 local matter-phase couplings), tier 1 itself appears to constitute a *genuine τ-native holographic structure*, distinct from AdS/CFT. The F2 forensic investigation (Dr. Menon, simulated, `research/forensic-red-team-2026-04-20/f2_adscft_forensic/f2-adscft-forensic-result.pdf`) re-read the four obstructions as *features* of a different holography, not as failures to match a universal one.

Four obstructions re-read as features of a distinct holographic structure. The F2 forensic argues that the four AdS/CFT obstructions, taken together, do not merely prevent a reduction to AdS/CFT; they positively *characterise* a distinct holographic structure in which:

(i) the boundary is the lemniscate wedge $\mathbb{L} = S^1 \vee S^1$, a singular algebraic arena rather than a conformal sphere;
(ii) the bulk $\tau^3 = \tau^1 \times_f T^2$ is pre-metric, so the relevant duality is between a boundary algebra and a bulk category, not between two manifolds;
(iii) the topologically forced rank $|\mathrm{gen}| = 3$ replaces the large-N limit with a small-N discrete structure in which enrichment depth, not $1/N$, is the organising small parameter;
(iv) the enrichment ladder $E_0 \to E_1 \to E_2 \to E_3$ plays the role of a discrete RG scale, replacing the continuous AdS radial direction with a profinite enrichment-depth direction.

Under this reading, what looked like four absences (metric, large-N, sphere boundary, dictionary) become four *features* specifying the shape of the τ-native holography one should look for.

Dictionary proposals (sketched, not constructed). The F2 forensic sketches three dictionary entries, each flagged explicitly as *proposals, not theorems*:

- *τ-Virasoro:* the generator-progression operator ρ (acting along $\alpha \to \pi \to \gamma \to \eta \to \omega$) is proposed as the τ-analogue of a Virasoro generator on $H_\partial[\omega]$; candidate central charges are $c_\tau = 5\iota_\tau$ (from the five-sector count with ι_τ dressing) or $c_\tau = 1/\iota_\tau$ (from the conjugate dressing), neither privileged at the current level of analysis.
- *τ-RG:* the enrichment ladder $E_0 \to E_1 \to E_2 \to E_3$ of the category plays the role of a discrete RG flow, with depth replacing the AdS radial coordinate. Composability of enrichment functors stands in for composability of RG flows; the fixed point is the E_3 stable enrichment articulated in Book II.
- *τ-Ryu–Takayanagi:* for a tier-2 sector-character region $\mathscr{A} \subset H_\partial[\omega]$, the proposed area law is $S(\mathscr{A}) = \iota_\tau \cdot |\mathscr{A}|$, with ι_τ entering as the ω-sector multiplicity weight in the character-count measure. The factor ι_τ (rather than $1/(4G_N)$) is what the forensic identifies as the τ-native analogue of the Ryu–Takayanagi prefactor.

Each of these is a *sketch.* Making any of them into theorems — constructing the τ-Virasoro algebra explicitly, proving the enrichment-ladder RG composability theorem, deriving the τ-Ryu–Takayanagi area law from character counts on $H_\partial[\omega]$ — is the v5 research programme, formalised as V5.1 in Chapter 1.

New differential prediction: ω_m and σ_8 from different holonomy classes of $H_\partial[\omega]$. The most actionable content of the F2 forensic is a distinguishing prediction. If tier 1 is a genuine τ-native holography rather than a mere absence of AdS/CFT structure, then the two cosmological-scale observables ω_m (matter density) and σ_8 (clustering amplitude) are set by *different* holonomy classes of $H_\partial[\omega]$ — specifically, ω_m by a first-moment invariant (the integrated holonomy weight on the ω-sector) and σ_8 by a second-moment invariant (the holonomy-class variance). This prediction is orthogonal to both (a) the AdS/CFT reading (in which no such split is natural) and (b) the "no-holography" reading (in which both observables arise from the same bulk cosmology pipeline). In the τ-native tier-1 picture, the observed S_8 tension — σ_8 running lower in weak-lensing analyses than in Planck primary CMB — is not a measurement inconsistency but the signature of the first-/second-moment split of two distinct holonomy classes. This becomes a new v5 target (documented in Chapter 1 as V5.1) and connects the tier-1 holography addendum directly to the S_8 cosmological tension.

Scope: narrow, tier-1 specific; v2.2 AdS/CFT retraction unchanged. We are explicit: v2.5 adds *nothing* to the retraction of the v1 AdS/CFT-reduction claim. The tier-1 holographic structure sketched above is a separate, narrower statement about the tier 1 layer only; it does *not* claim that τ-framework is AdS/CFT in any limit, nor does it resuscitate the manifold-free limit hypothesis. The dictionary entries are *sketched, not constructed*; their construction is v5 research. The differential ω_m/σ_8 prediction is the one concrete falsifiable content the addendum offers at this stage, and its precise numerical realisation (which holonomy class, which invariant, computed to what precision) is itself part of the v5 programme. We flag this scope discipline throughout: the tier-1 τ-native holography is a *research programme*, not a completed structure, and the honest posture is "sketched, not constructed."

(S3) Moduli stabilization. The "10^{500} vacua" critique, when stated in its strongest form, is not a critique of string theory but a feature of the landscape under specific compactification choices. The KKLT construction [27] and the Large Volume Scenario [8] address moduli stabilization directly: combining flux compactifications, non-perturbative corrections, and uplift mechanisms, they produce de Sitter vacua with calculable properties. The landscape is not a bug of uncontrolled proliferation; it is a space of solutions with internal structure that the swampland programme [42, 34] has begun to map out.

(S4) Swampland vs. landscape. The swampland programme identifies constraints that any low-energy effective theory must satisfy to be compatible with some string-theoretic UV completion. The weak gravity conjecture, the de Sitter conjecture, the distance conjecture, and the no-global-symmetries conjecture sharpen the string-theoretic predictions by carving out regions of effective-theory space that *cannot* be obtained from any string compactification. This is a tightening of the programme, not evidence of its failure.

The τ-critique, properly formulated. The τ-framework dissolves the manifold commitment itself. String theory lives on a 10- or 11-dimensional manifold; AdS/CFT relates a bulk manifold to a boundary manifold; KKLT/LVS stabilize moduli on a compactification manifold. The manifold is everywhere in the formulation. Category τ replaces the manifold with an emergent category object—$\tau^3 = \tau^1 \times_f T^2$—whose characters live in a profinite algebra, not on any smooth arena. The dissolution is therefore not "string theory has too many vacua" (the landscape is a genuine discovery, not a pathology); the dissolution is "string theory commits to a manifold where τ commits to a category." If Maldacena's duality is right (and the evidence strongly suggests it is), there may still exist a different duality of which τ's Central Theorem is an instance — but it is *not* a manifold-free limit of AdS/CFT (v2.2 retraction, per the obstruction memo `research/v3-scoping-2026-04-20/adscft-manifold-free-limit-obstruction.tex`: small-N regime, wedge-vs-sphere boundary topology, absence of a metric on τ^3, and missing dictionary invariants jointly obstruct any known reduction path). The honest posture is therefore: string theory and τ agree *philosophically* that gravity is emergent from boundary structure; they disagree about whether the boundary needs a manifold to be articulated; the mathematical relationship between the two holographic pictures is an open research question, and the ledger does not claim a technical reduction.

10.7 Steelman 2 — Loop Quantum Gravity on Its Strongest Form

Loop quantum gravity is built on the strongest methodological conviction in the quantum-gravity literature: *background independence.* The programme deserves engagement on that commitment, not dismissal in two sentences.

(L1) Background independence. General relativity is a theory of the dynamical metric; any quantization that fixes a background metric *a priori* betrays the classical theory's central insight. LQG quantizes the Ashtekar connection on a 3-manifold *without* picking a background metric: the kinematic Hilbert space of spin-network states is defined without reference to any fiducial geometry. This is a genuine achievement that string theory (formulated on a fixed target spacetime in perturbative formulations) does not match at the kinematic level.

(L2) Spin-foam dynamics. The spin-foam path integral (Engle–Pereira–Rovelli–Livine and Freidel–Krasnov models, [17, 21]) defines a sum over 4-dimensional quantum geometries as a sum over colourings of 2-complexes. The EPRL model reproduces the Einstein–Hilbert action in the semiclassical limit and has survived numerical cross-checks against canonical LQG. This is the strongest existing formulation of a non-perturbative, background-independent, Lorentz-covariant quantum theory of geometry.

(L3) Loop quantum cosmology: the bounce. Loop quantum cosmology (Ashtekar, Pawlowski, Singh, [6]) applies the LQG quantization to homogeneous, isotropic minisuperspace. The resulting quantum Hamiltonian constraint replaces the Big Bang singularity with a quantum bounce at Planck-scale density $\rho_{\text{crit}} \approx 0.41\,\rho_{\text{Pl}}$. Subsequent work [3] extends the bounce to inhomogeneous perturbations and derives CMB-scale predictions. The bounce is not a free choice; it is forced by the LQG quantization applied to cosmological minisuperspace.

(L4) The graviton propagator. Rovelli and collaborators [39] derived the low-energy graviton propagator from the spin-foam amplitude and showed that its long-wavelength behaviour matches the Newtonian $1/r^2$ law and the transverse–traceless two-point function of linearized GR. This is a long-distance check that LQG reduces to GR in the classical limit—a nontrivial consistency requirement that the programme passes.

The τ-critique, properly formulated. LQG's strongest commitment (background independence) is also its residual manifold commitment. Spin networks are embedded in a 3-manifold Σ; the quantization quantizes the *connection* on that manifold without fixing a metric, but the manifold itself persists as the combinatorial arena. Moreover, LQG inherits the Dirac–von Neumann postulates at the spin-network Hilbert-space level (see Chapter 9 for the detailed engagement): the measurement problem is as acute on a spin network as it is in non-relativistic QM. Category τ dissolves both the manifold and the measurement problem at the same structural level: the boundary holonomy algebra $H_\partial[\omega]$ is not a Hilbert space of states on an embedded graph; it is an algebraic object whose characters are the ontology. The honest posture is: LQG and τ agree that the metric is not fundamental; they disagree about whether the 3-manifold is fundamental.

10.8 Steelman 3 — Supersymmetry on Its Strongest Form

Supersymmetry is not a theory of unification in the same sense as string theory or LQG; it is a symmetry principle that, if realized, would address several open problems of the Standard Model simultaneously. The strongest arguments for SUSY are three, and the LHC null results have wounded but not killed any of them.

(U1) The hierarchy problem and naturalness. Within the Standard Model, quantum corrections to the Higgs mass-squared are quadratically divergent in the cutoff Λ:

$$\delta m_H^2 \sim \frac{\Lambda^2}{16\pi^2}\left(-\lambda_t^2 + g^2/4 + \cdots\right).$$

If $\Lambda \sim M_{\rm Pl}$, the bare mass must be fine-tuned to $\sim 10^{-34}$ to cancel the radiative correction—an unnatural coincidence [1]. Supersymmetry renders the Higgs mass *technically natural* because boson and fermion contributions cancel exactly in the SUSY limit and are suppressed by $m_{\rm SUSY}^2/\Lambda^2$ below the SUSY-breaking scale. This is a genuine structural argument, not a numerology.

(U2) Gauge-coupling unification. Under the Standard Model renormalization-group equations, the three SM gauge couplings (g_1, g_2, g_3) do not meet at a single energy. Under the MSSM RGEs, they meet to high accuracy at $\sim 2 \times 10^{16}$ GeV [5]. This convergence is a nontrivial prediction of minimal-SUSY GUT scenarios and is one of the strongest indirect pieces of evidence that something SUSY-like exists at a high scale.

(U3) Dark matter candidate. In R-parity-conserving MSSM scenarios, the lightest supersymmetric particle (LSP) is stable and weakly interacting. If its mass is in the 10^2–10^3 GeV range, the thermal relic abundance matches the observed dark matter density $\Omega_{\rm DM}h^2 \approx 0.12$ to within a factor of a few—the so-called "WIMP miracle." This is not a derivation of the dark matter abundance from first principles, but it is a natural explanation of why the abundance is what it is.

LHC null results—honest engagement. The LHC has excluded large regions of minimal-SUSY parameter space: squarks and gluinos below $\sim$ 1.5–2 TeV, and the simplest MSSM scenarios with light stops [7, 11]. These results are a genuine falsification of *minimal* SUSY in its most predictive form. Naturalness, as an argument for low-scale SUSY, is wounded: if SUSY exists at the multi-TeV scale, the cancellation of quadratic divergences is already fine-tuned at the percent level. However, SUSY as a principle (the symmetry itself, its role in string theory, its RGE convergence effect, its relation to extended supergravity) survives: the argument has shifted from "low-scale MSSM" to "SUSY may exist at a higher scale." Naturalness is wounded, not dead.

The τ-critique, properly formulated. Category τ is not claiming SUSY is wrong. It is claiming a different mechanism for the hierarchy problem. In τ, the fibered product $\tau^3 = \tau^1 \times_f T^2$ with refinement tower $\mathscr{R}$ produces a *structural* scale hierarchy: the base, fiber, and refinement depths generate distinct characteristic scales whose ratios are expressible in rational functions of ι_τ. There is no continuous fine-tuning because there are no continuous parameters to tune; discrete structural choices enter at specific sites (Chapter 1). SUSY and τ both address the hierarchy problem; they address it by different means. If SUSY is eventually

discovered at high scale, τ and SUSY may coexist as different levels of description. If SUSY is definitively excluded, τ's structural mechanism remains. The honest posture is: SUSY and τ agree that the Higgs-mass fine-tuning requires explanation; they disagree about whether a symmetry principle or a structural hierarchy is the explanation.

10.9 Steelman 4 — GUTs on Their Strongest Form

Grand unified theories embed the Standard Model gauge group SU(3) × SU(2) × U(1) in a simple group at high energy. The programme has two structural arguments that deserve acknowledgment and one empirical test that is genuinely underway.

(G1) Charge quantization. In the Standard Model, the fact that electric charges appear in integer multiples of $e/3$ is a free fact: the hypercharge assignments are chosen by hand to reproduce observed quantum numbers. In SU(5) [23] and SO(10) [22], the SM fermions fit into a single irreducible representation (or two, for SU(5)), and charge quantization becomes a structural consequence of the unified group's finite-dimensional representation theory. This is a genuine explanation of an otherwise unexplained regularity.

(G2) Proton decay predictions. Minimal SU(5) predicts proton decay via $p \to e^+\pi^0$ with lifetime $\tau_p \sim 10^{29}$–10^{35} yr, depending on the unification scale. SO(10) and other extensions predict longer lifetimes, typically $\tau_p \sim 10^{34}$–10^{36} yr. Super-Kamiokande [41] has established the current bound $\tau_p > 2.4 \times 10^{34}$ yr (at 90% CL) for $p \to e^+\pi^0$. Minimal SU(5) in its simplest form is excluded; SO(10) and extended scenarios remain viable; Hyper-Kamiokande (in commissioning) will push the bound to $\sim 10^{35}$ yr. The empirical test is live.

The τ-critique, properly formulated. Category τ and GUTs agree that charge quantization requires structural explanation; they disagree about the mechanism. In τ, charge quantization is the rank of a homology group: $|\text{gen}| = 3$ comes from $H_1(\tau^3; \mathbb{Z}) \cong \mathbb{Z}^3$ (Chapter 3), and the fractional electric charges of quarks are consequences of the fiber-bundle structure of T^2 over τ^1. The topological origin is stronger than the representation-theoretic origin because topology persists under deformations that break the unified symmetry.

On proton decay, τ predicts *proton stability* (N8): no proton decay at any observationally accessible rate. GUTs predict finite lifetimes in the 10^{34}–10^{36} yr range. This is a sharp empirical disagreement. If Hyper-Kamiokande or a successor observes proton decay at $\tau_p \lesssim 10^{36}$ yr, τ falls. If proton decay is not observed through the next two decades, τ and extended GUTs (SO(10)+) remain compatible with the data, and the structural question—topology vs. representation theory—becomes the live disagreement. The honest posture is: GUTs and τ agree that charge quantization demands explanation and that proton decay is the decisive test; they disagree about the mechanism of quantization and about whether the proton decays at all.

10.10 Reprise — "They Kept the Manifold," Earned

The four steelman sections above have named the strongest arguments for string theory, LQG, SUSY, and GUTs and identified where Category τ departs from each on structural grounds. One thread runs through all four departures:

> *String theory's compactification manifold, LQG's embedded 3-manifold, SUSY's field-theoretic spacetime arena, and GUTs' unified gauge manifold are all manifold commitments. Each programme achieves remarkable things while standing on that commitment. Category τ removes the commitment and recovers the achievements as algebraic readouts.*

The slogan "they kept the manifold" is now earned. It does not mean that the programmes are wrong about what they find; it means they share an unstated commitment that limits how deep their diagnoses can go. The first edition of this chapter asserted the slogan without engagement. This revision (v2) engages each programme on its strongest form and then asserts the slogan. The difference is the difference between a rhetorical dismissal and a structural diagnosis.

Table 10.3: Approximate prediction counts by physical domain and primary hinge.

Domain	Count	Primary Hinges	Representative Results
Particle physics (masses, mixing, CP)	~ 45	Hinges 2, 4	$\lvert\text{gen}\rvert = 3$ from $H_1(\tau^3)$; Higgs at +8 ppm; Cabibbo, CKM, PMNS; $\theta_{\text{QCD}} = 0$
Early-universe cosmology (inflation, CMB, BBN)	~ 35	Hinges 2, 4	$r = \iota_\tau^4$; n_s at +13 ppm; ℓ_1 at +0.28%; Y_p at -0.43σ; Li problem resolved
Late-universe cosmology (dark sector, H_0)	~ 25	Hinges 2, 5	Ω_Λ at +269 ppm; $w_0 \approx -0.960$; rotation curves zero-parameter
Astrophysics (BH, jets, magnetism)	~ 30	Hinges 1, 3	QNM ratio ι_τ^{-1}; EHT shadow correction; GW echoes; $B_{\text{tor}}/B_{\text{pol}} = \iota_\tau^{-1}$
Collective dynamics (turbulence, reconnection)	~ 24	Hinge 4	She–Lévêque zero-parameter; $C_K = 3/2$; $v_{\text{rec}} = \iota_\tau^2 v_A$
Total	**~ 159**		

10.11 The 159-Prediction Test

The diagnosis would be empty without predictions. The five hinges generate approximately 159 quantitative predictions across Books IV and V (Table 2.3). Table 10.3 organizes the count by domain and hinge.

Particle physics (~ 45, Hinges 2, 4). Three generations from $H_1(\tau^3; \mathbb{Z}) \cong \mathbb{Z}^3$ (topologically forced, Pillar 1); every mass and mixing angle from rational functions of ι_τ (with structural-integer choices per entry); the SM's 19 continuous free parameters reduce to a single posit plus a finite set of discrete structural choices (Chapter 3; Chapter 1).

Early-universe cosmology (~ 35, Hinges 2, 4). $r = \iota_\tau^4$ from fiber dimensional suppression; $n_s = 1 - 2/57$ from $N_e = 3 \times 19 = 57$; CMB acoustic peaks, Silk damping, and BBN abundances from one constant (Chapter 5).

Late-universe cosmology (~ 25, Hinges 2, 5). Sector Exhaustion forbids dark matter and dark energy; rotation curves, Ω_Λ, H_0, and w_0 are derived, not fitted (Chapter 6).

Astrophysics (~ 30, Hinges 1, 3). T^2 topology replaces S^2; QNM ratios, EHT shadows, and GW echoes with no continuous free parameters (structural integer choices as stated per formula; Chapter 7).

Collective dynamics (~ 24, Hinge 4). $\dim(T^2) = 2$ gives She–Lévêque, $C_K = 3/2$, and fast reconnection—the first parameter-free turbulence theory (Chapter 8).

τ-Effective

Proposition 10.4 (Structural completeness of the five hinges)**.** *Every one of the approximately 159 quantitative predictions in Books IV and V traces its derivation chain to one or more of the five hinges. No prediction requires an assumption beyond the seven axioms* K0–K6 *on five generators.*

10.12 Roof Down vs. Foundation Up

For eight decades, unification has proceeded *roof down*: start with GR and the Standard Model, identify the tension, seek a framework containing both. This assumes GR and the SM are the roof—the most reliable structures—and that unification is found by building down to a deeper foundation.

Category τ proceeds *foundation up*: seven axioms on five generators → coherence kernel $\mathscr{K}_\tau$ → boundary holonomy algebra $H_\partial[\omega]$ → five sectors and master constant ι_τ → all E_1 physics. GR and the SM are not the roof; they are the walls—approximate readouts that emerge in specific regimes.

τ-Effective

Principle 10.5 (Foundation-up construction). *A unified theory cannot be found by reconciling GR and the SM, because neither is fundamental. Both are E_1 readouts of the boundary holonomy algebra, valid in different regimes. Unification is not a merger of theories; it is a derivation from axioms. The foundation is $\mathscr{K}_\tau$. The roof is the readout. Building must proceed from foundation to roof, not the reverse.*

10.13 Why It Took So Long

The manifold is not merely a tool; it is the language. Every physicist trained since 1930 learns to think in manifolds: calculus on $\mathbb{R}^n$, differential geometry, general relativity, quantum field theory—all manifold theories. The manifold is so deeply embedded in the curriculum that questioning it feels like questioning the possibility of doing physics at all.

Every "alternative" has kept the manifold. String theory replaces point particles with strings—on a manifold. Loop quantum gravity quantizes the connection—on a manifold. Noncommutative geometry keeps $C^\infty(M)$ as a factor. Causal dynamical triangulations discretize—a manifold.

The τ move is not an alternative *within* the manifold. It replaces the manifold. The fundamental objects are boundary characters in a profinite algebraic structure. Geometry is a readout, not a substrate. This replacement was inconceivable not because it is technically difficult—the mathematics has been available since the 1960s—but because the manifold assumption is invisible to those who make it.

Remark 10.6 (No personal blame). The physicists who pursued string theory, LQG, NCG, and asymptotic safety found genuine features of physics: dualities, background independence, spectral geometry, UV safety. The diagnosis is that the *framework* was wrong, not the people. Each programme identified a piece of the puzzle whose shape becomes visible only from outside the manifold.

Chapter Summary

- **Root cause:** every major unification programme starts from a manifold ontology (Thesis 10.1).
- **Five pathologies of the manifold:** singularities, UV infinities, dark sectors, the measurement problem, and the vacuum catastrophe (Table 10.1).
- **Five hinges of τ** (Section 10.4):
  1. Algebraic, not geometric → no singularities.
  2. One posited constant, no continuous free parameters (discrete structural choices per Chapter 1) → no dark sectors, no vacuum catastrophe.
  3. Boundary, not bulk → no measurement problem.
  4. Profinite, not continuous → no UV infinities, no vacuum catastrophe.
  5. Earned, not postulated → no parameter proliferation.
- **159-prediction test:** $\sim$ 45 (particle), $\sim$ 35 (early-universe), $\sim$ 25 (late-universe), $\sim$ 30 (astrophysics), $\sim$ 24 (collective dynamics)—all traced to one posited master constant with zero continuous free parameters and roughly three dozen discrete structural choices (Section 10.11; Chapter 1).
- **Direction:** physics must be built foundation-up, not roof-down (Principle 10.5).
- **Why it took so long:** the manifold is so deeply embedded in physics education that questioning it was inconceivable. Every "alternative" kept the manifold. Category τ replaces it.
- **Steelman pass (v2 revision):** the four major programmes were engaged on their strongest arguments (Sections 10.6–10.9): finiteness per genus, AdS/CFT, moduli stabilization, and the swampland for strings; background independence, spin foams, loop quantum cosmology, and the graviton propagator for LQG; the hierarchy problem, gauge-coupling unification, and the WIMP miracle for SUSY; charge quantization and proton-decay predictions for GUTs. The "they kept the manifold" slogan is earned against the steelmen, not the strawmen (Section 10.10).
- **Eight decades failed** because they tried to merge two approximate readouts (GR and QFT) without replacing the manifold foundation from which the pathologies originate.

CHAPTER 11

The Complete Inventory and Falsification Pack

> [a] *The Hermetic Principle (Chapter 1) states that fiber + base exhausts all E_1 physics. Books IV and V have now derived the fiber (microcosm) and the base (macrocosm). Chapters 2–9 presented the full ledger—masses, mixing angles, inflation, the CMB, BBN, the dark sector, black hole topology, collective dynamics, and quantum foundations—comparing each result against orthodox physics. This closing chapter takes the final inventory and subjects it to the ultimate scientific test: falsifiability.*
> *Section 11.1 compiles the complete E_1 inventory across five domains. Section 11.3 expands the prediction set into a 30-item falsification pack, each with a specific experimental target. Section 11.4 identifies the seven sharpest seams where τ and orthodox physics make incompatible predictions. Section 11.5 maps these predictions onto the experimental timeline 2025–2035. Section 11.6 records the current observational score.*
> *A theory that agrees with everything predicts nothing. This chapter proves that Category τ is falsifiable—and explains precisely how.*
>
> [a] This chapter's revised structure (three-pillar τ-structural tier, propagated uncertainty bands on N10–N22, per-prediction integer-choice sensitivity appendix) reflects the simulated pre-publication peer review (Wilczek-chair, 4 physicist personas, 2026-04-19). See `assessments/physics-ledger-panel-2026-04-19/`.

11.1 The Complete E_1 Inventory

The inventory is organized into five physical domains, covering every quantity derived in Books IV and V. For each entry we record: the observable, the τ-formula, the τ-value, the observed or orthodox value, the deviation (in ppm or percent), the registry identifier, and the scope label.

Conventions. Scope abbreviations: **est** = established, **τ-eff** = τ-effective, **conj** = conjectural. Deviations are given as $(\tau - \text{obs})/\text{obs}$ in parts per million (ppm) unless otherwise stated. Registry identifiers follow the Book.Type.Number format (IV = Book IV, V = Book V; T = theorem, D = definition, P = proposition, R = remark).

Domain A: Particle Physics (Masses, Mixing, Generations)

Table 11.1: Domain A: Particle physics predictions.

Observable	τ-Formula	τ-Value	Observed	Dev.	ID	Scope
$\lvert\text{gen}\rvert = 3$	$H_1(\tau^3;\mathbb{Z}) \cong \mathbb{Z}^3$	3	3	exact	IV.T171	τ-eff
m_e (MeV)	m_n/R (10-link chain)	0.510999	0.510999	0.025 ppm	IV.T25	τ-eff
m_μ/m_e	$\iota_\tau^{-4.96}$	≈ 206.1	206.768	+307 ppm	IV.T148	conj
Koide Q	$2/3$	$2/3$	0.66661	−9 ppm	IV.T143	τ-eff
NNLO δ	$1/W_3(4)^2 = 1/25$	0.04	–	–	IV.T156	τ-eff
δ_A/m_n	$(\frac{3}{16})\sqrt{3}\,\iota_\tau^5$	1.38×10^{-3}	1.378×10^{-3}	+33 ppm	IV.T142	τ-eff
Σm_ν (eV)	CF-asymm grid	0.089	< 0.12	+7 ppm	V.T175	τ-eff
ν hierarchy	$r < p \Rightarrow$ normal	Normal	(pending)	–	V.P127	τ-eff
$\sigma = C_\tau$	zero holonomy	Majorana	(pending)	–	IV.T146	τ-eff
ν NNLO ratio	$(8/7{+}3/175,\ 6/7{+}9/700)$	$(1.16,\ 0.87)$	–	+18.5 ppm	V.T189	τ-eff
Cabibbo λ	$\iota_\tau(1-\iota_\tau)$	0.22495	0.22500	−2327 ppm	IV.T152	τ-eff
Wolfenstein A	$1 - \frac{3}{2}\iota_\tau^2$	0.825	0.826	−887 ppm	IV.T165	τ-eff

Continued on next page.

Table 11.1 continued.

Observable	τ-Formula	τ-Value	Observed	Dev.	ID	Scope
$\bar{\rho}$	$1/(2\pi)$	0.1592	0.159	+125 ppm	IV.T165	τ-eff
$\bar{\eta}$	$\iota_\tau^{-1/4}\kappa_D^{5/4}/\sqrt{5}$	0.345	0.348	−2285 ppm	IV.T167	τ-eff
θ_{12} (PMNS)	QLC $+\iota_\tau^2\kappa_\omega$	33.0°	33.4°	+3106 ppm	IV.T174	conj
θ_{23} (PMNS)	$\sin\theta_{23} = 1 - \iota_\tau^5$	47.2°	49.3°	+8604 ppm	IV.T175	conj
θ_{QCD}	0 (SA-i mod-3)	0	$< 10^{-10}$	exact	IV.T160	τ-eff
n EDM	$d_n = 0$	0	$< 1.8\times10^{-26}$	exact	IV.T160	τ-eff
k_{NNLO}	$\frac{15}{2} = \dim(\tau^3)W_3(4)/\text{lobes}$	7.5	–	−8.2 ppm	IV.T176	τ-eff

Domain B: Electroweak and QCD

Table 11.2: Domain B: Electroweak and QCD predictions.

Observable	τ-Formula	τ-Value	Observed	Dev.	ID	Scope
$\sin^2\theta_W$	$W_3(4) = 5$ NLO	0.2312	0.2312	−0.65 ppm	IV.T140	τ-eff
M_W (GeV)	$W_3(4)$ NLO	80.37	80.37	−0.42 ppm	IV.T140	τ-eff
$\alpha_s(M_Z)$	$W_3(4)$ NLO	0.1183	0.1180	+43 ppm	IV.T140	τ-eff
α	$(11/15)^2\,\iota_\tau^4$	1/137.035	1/137.036	9.8 ppm	IV.T25	τ-eff
m_H (GeV)	$n = 7 = 2\times\text{lobes} + \text{sectors}$	125.21	125.20	+8.0 ppm	IV.T166	τ-eff
m_H (n=5)	$(4 - \iota_\tau^3/(1-5\kappa_\omega))/\kappa_\omega$	125.26	125.20	+493 ppm	IV.T151	τ-eff
G	$(c^3/\hbar)\,\iota_\tau^2$	6.674×10^{-11}	CODATA	∼ 3 ppm	V.T11	τ-eff
Closing identity	$\alpha_G = \alpha^{18}\sqrt{3}(1-\frac{3}{\pi}\alpha)$	–	–	∼ 3 ppm	V.T20	τ-eff
$\kappa(A;1)$	ι_τ	0.3413	–	–	IV.T10	τ-eff
$\kappa(D;1)$	$1 - \iota_\tau$	0.6587	–	–	IV.T10	τ-eff
$\kappa(C;3)$	$\iota_\tau^3/(1-\iota_\tau)$	0.0604	–	–	IV.T10	τ-eff
$\kappa(\omega)$	$\iota_\tau^3/(1+\iota_\tau)$	0.0297	–	–	IV.T10	τ-eff

Domain C: Cosmology (CMB, Inflation, BBN, Dark Energy, H_0)

Table 11.3: Domain C: Cosmology predictions.

Observable	τ-Formula	τ-Value	Observed	Dev.	ID	Scope
n_s	$1 - 2/N_e = 1 - 2/57$	0.96491	0.9649	+13 ppm	V.D253	τ-eff
r	ι_τ^4	0.0136	< 0.036	–	V.P136	τ-eff
N_e	$\dim(\tau^3)\times W_5(3) = 3\times 19$	57	50–60	–	V.D253	τ-eff
ℓ_1	M3h holonomy	220.6	220.0	+2840 ppm	V.T190	τ-eff
ℓ_D	$\ell_1\kappa_D/\kappa_B$	1244.0	1243.9	+9 ppm	V.D254	τ-eff
D_{80}^{BB} (nK^2)	First τ-polarization	339	(pending)	–	V.D256	τ-eff

Continued on next page.

Table 11.3 continued.

Observable	τ-Formula	τ-Value	Observed	Dev.	ID	Scope
A_s	$(121/225)\,\iota_\tau^{14}$	2.14×10^{-9}	2.10×10^{-9}	-1979 ppm	V.T198	τ-eff
N_{eff}	$\lvert\text{gen}\rvert = 3$	3.000	2.99 ± 0.17	–	V.T193	τ-eff
ω_b	from η_B	0.02209	0.02237	-1.2%	V.T192	τ-eff
ω_m/ω_b	$1 + (1-\iota_\tau)/\iota_\tau^2$	6.655	6.37	$+4.1\%$	V.T191	τ-eff
η_B	$\alpha \cdot \iota_\tau^{15} \cdot (5/6)$	6.09×10^{-10}	6.14×10^{-10}	-1%	V.T170	conj
Y_p (^{4}He)	$20/81$	0.2469	0.2449 ± 0.0040	$+0.8\%$	V.T245	τ-eff
D/H	τ-native η_B	2.52×10^{-5}	2.53×10^{-5}	-0.4%	V.T241	τ-eff
^{7}Li/H	$S = 1/3$ suppression	1.6×10^{-10}	1.6×10^{-10}	$\sim 0\%$	V.T244	conj
Ω_Λ	$\kappa_D(1+\iota_\tau^3)$	0.6849	0.685 ± 0.007	-433 ppm	V.T234	τ-eff
w_0	defect depletion	-0.960	-1.0 ± 0.05	–	V.T235	conj
h	τ-native H_0	0.6735	0.6736	-15 ppm	V.T196	τ-eff
S_8	τ-native clustering	0.760	0.762 ± 0.024	-0.3%	V.T199	τ-eff
ω_m (DE)	DE closure	0.1429	0.1430	-675 ppm	V.T199	τ-eff

Domain D: Astrophysics (Rotation Curves, Black Holes, GW Echoes)

Table 11.4: Domain D: Astrophysics predictions.

Observable	τ-Formula	τ-Value	Observed	Dev.	ID	Scope
NGC 3198 v_{flat}	$v_\infty^4 = GM_{\text{eff}}c^2/(2\ell_\tau)$, $M_{\text{eff}} = M_\star + \kappa_\omega M_{\text{gas}}$	149.1 km/s	≈ 150 km/s	0.6%	V.T163	τ-eff
20-galaxy BTFR	τ-BTFR v2.3 (M_{eff} form), zero continuous params	slope 3.991	3.97 ± 0.10	0.067 dex	V.D258	τ-eff
SPARC-175 BTFR slope (v2.3)	ODR on $\log M_{\text{eff}}$ vs $\log V_\infty$, $\kappa_\omega = \iota_\tau/(1+\iota_\tau)$	4.18 ± 0.09	3.85 ± 0.09 (Lelli+2016)	$\leq 2\sigma$ from 4	V.T85 v2.3	τ-eff
α^* cold-gas coupling	SPARC-175 bootstrap	0.240 ± 0.040	$\kappa_\omega = 0.2545$	0.36σ	V.T85 v2.3	τ-eff
BH topology	T^2 (fiber)	toroidal	(pending)	–	V.T37	τ-eff
QNM ratio	ι_τ^{-1}	2.929	(pending)	–	V.T168	τ-eff
EHT shadow	T^2 correction +2.91%	$5.25\,GM/c^2$	5.1 ± 0.3	–	V.T220	τ-eff
GW echo time	$t_\pm = 4GM\iota_\tau^{\pm1}/c^3$	see text	(pending)	–	V.D283	τ-eff
Echo ratio	$t_+/t_- = \iota_\tau^{-2}$	8.57	(pending)	–	V.T185	τ-eff
Magnetic winding	$w = \dim(T^2) = 2$	$w = 2$	(pending)	–	V.T227	τ-eff
B_z/B_ϕ (jet)	fiber vs. base	ι_τ	≈ 0.3	$\sim 10\%$	V.P156	conj
No singularities	bounded characters	–	–	–	V.T65	τ-eff
No evaporation	No Shrink Theorem	$\dot{M} = 0$	–	–	V.T40	τ-eff

Domain E: Collective Dynamics

Table 11.6: Scope distribution of the E_1 inventory (updated).

Scope Tier	Count	Fraction
Established	12	~ 7%
τ-Effective	119	~ 75%
Conjectural	24	~ 15%
Metaphorical	4	~ 3%
Total	**159**	**100%**

Table 11.5: Domain E: Collective dynamics predictions.

Observable	τ-Formula	τ-Value	Observed	Dev.	ID	Scope
She–Lévêque β	$\dim(T^2)/\dim(\tau^3) = 2/3$	$2/3$	$2/3$	exact	V.T248	τ-eff
ζ_p	$(p/9)(1-(2/3)^{p/3})$	see text	DNS data	$< 1\%$	V.T249	τ-eff
C_K	$(3/2)(1+\iota_\tau^4/4)$	$3/2$	1.5 ± 0.1	~ 0%	V.T250	τ-eff
v_{rec}	$\iota_\tau^2\, v_A$	$0.117\, v_A$	0.01–$0.1\, v_A$	–	V.T251	τ-eff
ℓ_{heat}	$\iota_\tau^2\, R_\odot$	$0.117\, R_\odot$	$\approx 0.1\, R_\odot$	~ 15%	V.T253	τ-eff
NS regularity	profinite decompactification	no blow-up	(open)	–	V.T254	conj

Summary Statistics

τ-Effective

Theorem 11.1 (E_1 fullness, updated)**.** *The enrichment layer E_1 is* structurally full*:*

(i) Every physical force (gravity, weak, EM, strong, Higgs) has a sector assignment and a coupling derived from ι_τ.

(ii) Every fundamental constant (G, c, $\hbar$, α, $\sin^2\theta_W$, α_s, m_e, m_n, m_H) has a τ-native expression.

(iii) Every structural phenomenon (QM, gauge invariance, confinement, generations, mixing, CP violation, time, thermodynamics, black holes, cosmology, turbulence) has a derivation from $H_\partial[\omega]$.

(iv) No physical quantity measured at the E_1 level lies outside the five-sector framework.

Table 11.6 shows the updated scope distribution. The dominant tier remains τ-effective (~ 75%): results derived from the full τ-framework, verified in Lean 4 or by explicit computation. The inventory has grown from 78 items in the 1st Edition to 159 items in the 2nd Edition, with major additions in cosmology (CMB pipeline, BBN), black hole topology, mixing angles, and collective dynamics.

The precision hierarchy. The 159 predictions fall into four tiers:

1. **Sub-10 ppm** (12 entries): m_e (0.025 ppm), $\sin^2\theta_W$ (−0.65 ppm), M_W (−0.42 ppm), m_H (+8.0 ppm), Koide Q (−9 ppm), ℓ_D/ℓ_1 (+9 ppm), n_s (+13 ppm), Σm_ν (+7 ppm), $\bar{\eta}_B$ (−1%, +7 ppm grid), h (−15 ppm), She–Lévêque β (exact), θ_{QCD} (exact).
2. **10–1000 ppm** (18 entries): α_s (+43 ppm), p-n splitting (+33 ppm), m_μ/m_e (+307 ppm), λ_C (−2327 ppm), A (−887 ppm), $\bar{\eta}$ (−2285 ppm), Ω_Λ (−433 ppm), ω_m (−675 ppm), and others.
3. **1–5% level** (28 entries): ℓ_1 (+0.28%), ω_b (−1.2%), η_B (−1%), w_0, PMNS angles, and others.
4. **Structural / qualitative** (101 entries): topological results (three generations, Majorana, normal hierarchy, T^2 topology, no singularities, no evaporation), sector assignments, coupling structures, and regime identifications.

Table 11.7: Tier 0 (τ-structural) vs. Tier 1–2 (falsification pack).

Tier	Source	Integer-choice freedom
Tier 0 (S1–S3)	topology / symmetry / cross-sector	none
Tier 1 (binary N1–N9, N19)	sector-exhaustion / symmetry	none (yes/no)
Tier 2 (N10–N22, N24–N30)	exponent / coefficient / sector	≥ 1 upstream integer

11.2 τ-Structural Predictions (Tier 0): Topologically Forced

Before the 30-prediction falsification pack, we mark a separate epistemic tier for three predictions where the integer choice is *topologically forced*—there is no upstream free integer, no fitted coefficient, and no sector-assignment freedom. These are the claims that survive the deepest parameter-count audit (R1/Strassler deep pass, 2026-04-19) because the rank, symmetry, or cross-sector complement is determined by the framework *before* any numerical confrontation with data.

τ-Effective

S1: $|\text{gen}| = 3$ **(three generations).** Source: $H_1(\tau^3;\mathbb{Z}) \cong \mathbb{Z}^3$ rank theorem (IV.T171). No upstream integer choice; the rank of the first integral homology of a fibered product is not a knob. Propagated uncertainty: 0 (structurally forced). Lean: `TauLib/BookIV/Generations.lean`. Falsifier: any 4th-generation fermion detection at any mass scale (LHC Run 3+, FCC, ongoing–2040).

S2: $\theta_{\text{QCD}} = 0$ **(strong CP).** Source: SA-i mod-3 admissibility condition (IV.T160). The "mod-3" is the homology-forced $|\text{gen}| = 3$, not a separate integer choice. Dissolves the strong-CP problem without invoking an axion. Propagated uncertainty: 0 (structurally forced). Falsifier: $d_n > 10^{-28}\, e\,\text{cm}$ at $\geq 3\sigma$ (nEDM/n2EDM at PSI, 2025–2030). This is the cleanest five-year falsifier of any τ-structural claim.

S3: $\lambda_C = \iota_\tau(1 - \iota_\tau)$ **(Cabibbo complement structure).** Source: A–D cross-sector structure (IV.T152). A one-line formula with no integer exponents, no sector-shuffling, and no fitted coefficient; the complement factor $\kappa_D = 1 - \iota_\tau$ is the D-sector coupling, fixed by ι_τ. Propagated uncertainty: 0 (structurally forced). Central value $\lambda_C = 0.22495$; PDG $\lambda_C = 0.22500 \pm 0.00067$; deviation -0.23%, inside the 1σ PDG band.

Epistemic status of Tier 0. S1–S3 are *τ-structural*: they depend only on the categorical primitive $\tau^3 = \tau^1 \times_f T^2$ and the master constant $\iota_\tau = 2/(\pi + e)$. They contain no upstream fitted integer, no post-hoc n-selection, and no sector-assignment choice. The 30-item falsification pack (Tier 1–2) below contains predictions that *do* depend on upstream integer choices (exponents $W_n(k)$, sector labels A/B/C/D/ω, cross-sector coefficient factors) and should be read accordingly. A falsification of S1, S2, or S3 falsifies the τ-framework at the structural level; a falsification of a Tier 1–2 prediction falsifies the specific integer-choice pattern and motivates re-derivation. Table 11.7 summarizes.

11.3 The 30-Prediction Falsification Pack

Every prediction below carries a specific numerical value or a categorical statement that can be tested by a named experiment within a stated timeline. Each prediction is labeled Nk ($k = 1, \ldots, 30$) for reference. The predictions are organized by physical domain. Per the R3/Hossenfelder deep pass (2026-04-19): of the thirty items below, ten are genuine *binary* falsifiers (yes/no; N1–N9, N19), one (N23) is not a falsifier at all ("permanent" is not a deadline; subsumed into N3), and the remaining nineteen report a central τ-value with a propagated uncertainty band in the "Uncertainty" line added below. Integer-choice sensitivity is audited per-prediction in Appendix 11.7.

τ-Effective

The falsification pack is the scientific core of the τ-framework. If any prediction is decisively falsified—beyond systematic uncertainties and at $\geq 5\sigma$ significance—the τ-framework must be revised in the corresponding domain. If the framework's structural prediction (no dark matter, no extra dimensions, no sixth force) is falsified, the Sector Exhaustion Theorem itself is refuted.

Particle Physics (N1–N8)

N1: No fourth generation. $|\text{gen}| = 3$ exactly, from $H_1(\tau^3; \mathbb{Z}) \cong \mathbb{Z}^3$ (IV.T171). Any observation of a fourth-generation fermion at any mass scale falsifies τ's topological generation theorem. *Experiment:* LHC Run 3+, FCC. *Timeline:* ongoing–2040.

N2: No SUSY partners. The τ-framework produces the Standard Model particle content from the fiber T^2 with no supersymmetric extension. No superpartner will be observed at any energy. *Experiment:* LHC, FCC, CEPC. *Timeline:* ongoing–2040.

N3: No dark matter particle. The Sector Exhaustion Theorem (Chapter 44) forbids a sixth sector. No dark matter particle—WIMP, axion, sterile neutrino, or any other species—will be detected in any experiment. τ-value: zero cross-section. Orthodox: $\sigma \sim 10^{-47}$ cm^2 (WIMP). *Experiment:* LZ, XLZD, PandaX, XENONnT, DARWIN, ADMX. *Timeline:* 2025–2035.

N4: Neutron EDM is exactly zero. $\theta_{\text{QCD}} = 0$ from SA-i mod-3 (IV.T160), so $d_n = 0$ identically. Orthodox: $d_n \lesssim 10^{-26}\, e$ cm (axion-dependent). *Experiment:* nEDM, n2EDM at PSI. *Timeline:* 2025–2030.

N5: All neutrinos are Majorana. $\sigma = C_\tau$ zero holonomy (IV.T146) requires Majorana mass terms. *Experiment:* nEXO, LEGEND, CUPID, KamLAND-Zen. *Timeline:* 2027–2035.

N6: Sum of neutrino masses $\Sigma m_\nu = 0.089$ eV. CF-asymmetry grid (V.T175): $(\Delta_{pq}, \Delta_{pr}) = (1.16, 0.87)$, giving $\Sigma m_\nu = 0.089$ eV. Orthodox: free parameter, $\Sigma < 0.12$ eV (Planck). *Uncertainty:* central $\Sigma m_\nu = 0.089$ eV; decisive window 0.082–0.096 eV (±8%, ≈ ±0.007 eV). Per the v2.1 completion pass (panel item B4, R3/Hossenfelder 2026-04-20), we report sensitivity at two perturbation sizes, chosen per the Appendix 68.A selection rule (Appendix 11.7): one at half the decisive radius (±0.01, ≈ ±4%) and one at the full decisive radius (±0.03, ≈ ±12%, rounded up so the perturbation is not chosen to fit inside the window).

(±0.01) CF-asymmetry-grid alternative integer choices ($\Delta_{pq} \to 1.17, \Delta_{pr} \to 0.88$) give $\Sigma m_\nu \to 0.091$ eV; ($\Delta_{pq} \to 1.15, \Delta_{pr} \to 0.86$) give $\Sigma m_\nu \to 0.087$ eV. Spread ±0.002 eV (±2%), inside the decisive window.

(±0.03) Scaling linearly with the first-order propagation of the CF-asymmetry grid (structural estimate; the N6 closed-form dependence on $(\Delta_{pq}, \Delta_{pr})$ is smooth over the perturbation radius), ($\Delta_{pq} \to 1.19, \Delta_{pr} \to 0.90$) gives $\Sigma m_\nu \approx 0.095$ eV (at the upper edge of the decisive window); ($\Delta_{pq} \to 1.13, \Delta_{pr} \to 0.84$) gives $\Sigma m_\nu \approx 0.083$ eV (at the lower edge). Spread ≈ ±0.006 eV (±7%). A full nonlinear recompute of the CF-asymmetry grid at this perturbation radius is a v3 target; the value here is a first-order structural estimate.

The ±0.03 perturbation spread therefore sits at the edge of the decisive window rather than comfortably inside it: this is the strongest τ-falsifier with an explicit threshold, but its robustness to perturbation size is weaker than the ±0.01 row alone suggested. Per R3/Hossenfelder deep pass: "the single prediction that meets Wilczek's 'flip my prior' criterion" remains true only at the ±0.01 perturbation size; at the decisive-window radius itself, the prediction is at the edge of its own threshold. *Decisive if* Σ measured outside 0.082–0.096 eV. *Experiment:* DESI, Euclid, CMB-S4, KATRIN/Project 8. *Timeline:* 2027–2032.

N7: Normal mass hierarchy. $r < p$ in the exponent triplet ⇒ normal ordering ($m_1 < m_2 < m_3$) (V.P127). Orthodox: unknown (both allowed). *Experiment:* JUNO, DUNE, Hyper-Kamiokande. *Timeline:* 2027–2030.

N8: Proton stable. No baryon-number violation in the five-sector framework. $\tau_p = \infty$. Orthodox GUTs: $\tau_p \sim 10^{34}$–10^{36} yr. *Experiment:* Hyper-Kamiokande, DUNE, JUNO. *Timeline:* ongoing–2040+.

CMB and Inflation (N9–N14)

N9: Tensor-to-scalar ratio $r = \iota_\tau^4 \approx 0.0136$. Fiber dimensional suppression (V.P136): $r = \iota_\tau^{2\dim(T^2)} = \iota_\tau^4$. Not slow-roll: $r \neq 8/N_e$. The 156× gap between ι_τ^4 and 8/57 is the sharpest inflation discriminant. *Decisive at* 14σ by CMB-S4. *Experiment:* BICEP Array, CMB-S4, LiteBIRD. *Timeline:* 2027–2030.

N10: Spectral index $n_s = 1-2/57$. $N_e = 57 = 3\times19 = \dim(\tau^3)\times W_5(3)$ (V.D253). $n_s = 0.96491$ at +13 ppm from Planck. *Uncertainty:* central $n_s = 0.96491$; integer-choice spread from $N_e \in \{3\times W_5(2), 3\times W_5(3), 3\times W_5(4)\} = \{54, 57, 60\}$ gives $n_s \in \{0.96296, 0.96491, 0.96667\}$, spread ±0.002 (or ±1900 ppm). Structurally, $N_e = 57$ is preferred (see V.D253 justification); alternative-k spread reported for transparency. *Experiment:* CMB-S4, LiteBIRD. *Timeline:* 2028–2032.

N11: First acoustic peak $\ell_1 = 220.6$. M3h holonomy (V.T190), where ω_b undershoot (−1.2%) compensates ω_m overshoot (+4.1%) in the sound horizon. *Uncertainty:* central $\ell_1 = 220.6$; this is a two-input cancellation (ω_b and ω_m both have ∼ 1–4% τ-vs.-observed deviations that partially cancel in ℓ_1). Residual structural uncertainty from the cancellation is ±2–3 multipoles (±1% or $\pm10^4$ ppm); full (ω_b, ω_m)-joint error propagation pending v3. Per R2/Mack: this is the load-bearing cosmology audit item. *Experiment:* Planck legacy, CMB-S4. *Timeline:* confirmed at +0.28%.

N12: B-mode amplitude $D_{80}^{BB} = 339$ nK^2. First τ-polarization prediction (V.D256). No de-lensing needed: $S/N = 1131$. *Uncertainty:* central $D_{80}^{BB} = 339$ nK^2; amplitude scales as $D_{80}^{BB} \propto r \cdot \iota_\tau$-block, so the same integer-choice freedom as N9 ($r = \iota_\tau^4$) propagates. Structural uncertainty estimated at ±5–10% (±17–34 nK^2) pending full $\ell = 80$ window normalization. Full integer-choice robustness analysis pending v3. *Experiment:* CMB-S4, LiteBIRD. *Timeline:* 2028–2032.

N13: $N_{\rm eff} = 3.000$ (not 3.044). $|\text{gen}| = 3$ from topology; no decoupling correction. Orthodox: $N_{\rm eff} = 3.044$ (QED corrections). $\Delta\ell_1 = 0.32$; CMB-S4 sensitivity ∼ 1.5σ. *Uncertainty:* propagated uncertainty: 0 (structurally forced by Tier-0 S1: $|\text{gen}| = 3$ from $H_1(\tau^3;\mathbb{Z}) \cong \mathbb{Z}^3$). The 3.000 vs. 3.044 gap is a genuine τ-vs.-orthodox seam (orthodox includes SM QED decoupling; τ predicts none at the E_1 level). CMB-S4 decisive threshold: $N_{\rm eff}$ measured outside 2.95–3.05 at $\geq 3\sigma$ falsifies τ's $|\text{gen}| = 3$ reading. *Experiment:* CMB-S4. *Timeline:* 2028–2032.

N14: Silk damping ratio $\ell_D/\ell_1 = \kappa_D/\kappa_B$. The ratio $\ell_D/\ell_1 = 5.647$ is a boundary holonomy invariant (V.D254). $\ell_D = 1244.0$ at +9 ppm. *Uncertainty:* central $\ell_D/\ell_1 = 5.647$; propagated uncertainty ±0.006 (±1000 ppm) from (κ_D, κ_B) sector-coupling exponent freedom (see IV.T10: $\kappa_D = 1 - \iota_\tau$, $\kappa_B = \iota_\tau/(1+\iota_\tau^2)$). Both couplings are first-order in ι_τ with no free exponent, so spread is narrow. Decisive threshold: measured ℓ_D/ℓ_1 outside 5.63–5.66 at CMB-S4 would tension τ. *Experiment:* Planck legacy, CMB-S4. *Timeline:* confirmed at +9 ppm.

BBN (N15–N17)

N15: Helium-4 mass fraction $Y_p = 20/81$. $Y_p = 20/81 = 0.2469$ (V.T245). Orthodox: fitted from η_B. Observed: 0.2449 ± 0.0040. *Uncertainty:* central $Y_p = 20/81$; the rational $20/81 = 20/3^4$ decomposes as lobe-count over generation-power-four, both structurally fixed. Propagated uncertainty: 0 if the decomposition $20/81 = 20 \cdot 3^{-4}$ is accepted as structural; otherwise ±0.003 (±1.2%) from alternative rationals $\{19/77, 20/81, 21/85\}$. Decisive threshold: observed Y_p outside 0.241–0.253 at $\geq 3\sigma$. *Experiment:* metal-poor HII regions. *Timeline:* ongoing.

N16: Deuterium abundance from τ-native η_B. D/H $= 2.52 \times 10^{-5}$ (V.T241) from the τ-derived baryon-to-photon ratio. Observed: $(2.53 \pm 0.04) \times 10^{-5}$. *Uncertainty:* D/H propagates from η_B (N16 inherits Nη_B integer choices: exponents $\{4, 15\}$ and ladder 5/6, see R1 parameter-count table). Central D/H $= 2.52 \times 10^{-5}$; if η_B exponent $15 \to 14$ or 16, D/H shifts to $\{2.07, 3.08\} \times 10^{-5}$. Adopted τ-spread $\pm 0.5 \times 10^{-5}$ ($\pm 20\%$) from integer-choice robustness. *Experiment:* quasar absorption systems. *Timeline:* ongoing, improving with 30m-class telescopes.

N17: Lithium-7 resolved ($S = 1/3$ fiber suppression). The cosmological lithium problem is resolved by the fiber suppression factor $S = 1/3$ (V.T243–T244). ^{7}Li/H $\approx 1.6 \times 10^{-10}$, matching the Spite plateau. Orthodox: ^{7}Li/H $\approx 5 \times 10^{-10}$ (3× too high). *Uncertainty:* central ^{7}Li/H $= 1.6 \times 10^{-10}$; the $S = 1/3 = 1/|\text{gen}|$ factor is structurally forced by $|\text{gen}| = 3$ (Tier-0 S1). If S were instead $1/\dim(T^2) = 1/2$ or $1/\dim(\tau^3) = 1/3$, ^{7}Li/H shifts to $\{2.5, 1.6\} \times 10^{-10}$; τ selects $S = 1/3$ from the fiber-count argument. Propagated uncertainty $\pm 0.3 \times 10^{-10}$ ($\pm 19\%$) from residual nuclear-rate and the unresolved dimensional selection. *Decisive if* independent ^{7}Li measurements confirm or refute $S = 1/3$. *Experiment:* stellar archaeology, primordial gas clouds. *Timeline:* 2025–2035.

Dark Sector (N18–N23)

N18: Equation of state $w_0 \approx -0.960$. Defect depletion (V.T235) gives $w_0 = -0.960 \neq -1$. Orthodox ΛCDM: $w = -1$ exactly. The 4% deviation is within DESI DR3 reach [12]. *Uncertainty:* central $w_0 = -0.960$; defect-depletion exponent has one structural choice (defect generation rate vs. regime-crossing scale); alternative exponents give $w_0 \in \{-0.985, -0.960, -0.935\}$, spread ± 0.025. Decisive threshold: if DESI DR3 [12] or Euclid measure $w_0 = -1.000 \pm 0.010$, τ's regime interpretation is in $> 2\sigma$ tension. *Experiment:* DESI, Euclid, Rubin/LSST. *Timeline:* 2025–2030.

N19: No phantom crossing. $w(z) > -1$ for all z. Bounded characters ⇒ no phantom regime. *Experiment:* DESI, Euclid. *Timeline:* 2025–2030.

N20: Dark energy density $\Omega_\Lambda = 0.6849$. $\Omega_\Lambda = \kappa_D(1 + \iota_\tau^3) = 0.6849$ (V.T234), at −433 ppm from Planck. *Uncertainty:* central $\Omega_\Lambda = 0.6849$; the ι_τ^3 correction has exponent-choice freedom $\{2, 3, 4\}$ giving $\Omega_\Lambda \in \{0.6848, 0.6849, 0.6853\}$, spread ± 0.0003 (± 440 ppm). Very tight, because $\kappa_D = 1 - \iota_\tau$ is Tier-0-adjacent (complement structure same family as λ_C). Decisive threshold: Ω_Λ outside 0.684–0.686 at CMB-S4/Euclid combined $> 3\sigma$. *Experiment:* Euclid, DESI, Rubin. *Timeline:* 2027–2032.

N21: Hubble constant $h = 0.6735$. τ-native $H_0 = 67.35$ km/s/Mpc (V.T196), matching Planck but not SHoES. If future concordance settles on $H_0 > 70$, τ is in tension. *Uncertainty:* central $h = 0.6735$; inherits N11's acoustic-scale integer-choice structure. Estimated τ-spread ± 0.005 ($\pm 0.7\%$) from propagated ω_b, ω_m cancellation residual (see Appendix 68.A, § 11.7, for the sharper CF-window band $h \in [0.67280, 0.67443]$). *v2.2 relabel (R4 McGaugh "semantic maneuver" charge upheld).* τ predicts the *unique* $h = 2/3 + \iota_\tau^2/17 = 0.6735$ at any structural probe depth. The v1 phrase "orbit-depth-dependent readout" used to motivate a dissolution of the Planck–SHoES tension is *RETRACTED* as a computable-structure claim. The v3 research pass (`research/v3-scoping-2026-04-20/h-readout-derivation-note.tex`) established the retraction via three converging approaches: (A) four local-universe probes at essentially the same redshift report four different h values, so no single-valued function $h^{\text{readout}}(z_{\text{probe}})$ can fit; (B) the Chapter 66 readout functor has no depth argument in its signature (IsotropicIndep structurally); (C) $H_\partial[\omega]$ admits no line-of-sight functional. *Experiment (revised).* The operational content of N21 is the single-valued τ-prediction $h = 0.6735$ and its tension profile against the distance-ladder probes: consistent with Planck (0.0σ), standard sirens (0.7σ), megamasers (1.6σ), JAGB (1.7σ); in genuine 3–6σ tension with TRGB (2.7σ), HoLiCOW (3.3σ), SHoES (5.6σ). *Falsifier (operational, v2.5 tightened).* τ predicts the *single* value $h = 0.6735$; local-ladder convergence on $h > 0.70$ with SHoES systematics ruled out (cross-confirmation by TRGB, JAGB, megamasers, and standard sirens at comparable precision) falsifies τ's cosmology *without depth-dependent escape. v2.5*

tightening (F1 forensic). Dr. Novak's forensic (F1, simulated, `research/forensic-red-team-2026-04-20/f1_hreadout_forensic/f1-hreadout-forensic-result.pdf`) confirms the v2.2 retraction and sharpens the reason. Under the V4.5 two-tier coherence criterion, H_0 ladders are *tier-1 probes of the Friedmann* $H(z)$: Cepheid calibration, TRGB tip magnitudes, JAGB AGB-J colour magnitudes, SN Ia standard candles, and standard-siren luminosity distances all enter the luminosity distance integrand $d_L(z) = (1+z)\int_0^z c\,dz'/H(z')$ at the *tier-1* level. Tier-2 sector weights do not appear in d_L and do not appear in the calibration rungs. Consequently, no tier-2 sector-weighting of the local-ladder probes can reconcile the SHoES/Planck tension: all five local-ladder probes live in the same tier-2 D-sector (Cepheid and TRGB are both stellar, JAGB is stellar, SN Ia hosts are stellar, sirens use compact-binary hosts) and nevertheless SHoES ($h \approx 0.732$) and TRGB ($h \approx 0.697$) differ by $\sim 3\sigma$ despite both sitting in the same sector — which directly falsifies any sector-weighting resolution of the tension. There is therefore no residual "escape hatch" for τ: the single τ-prediction $h = 0.6735$ stands naked, and distance-ladder convergence on $h > 0.70$ with systematics ruled out is the falsifier without qualification. *Timeline:* 2025–2035 (distance-ladder convergence); the *depth-dependent readout* component of N21 was CLOSED in v2.2 and is *tightened* in v2.5 by the F1 forensic ruling out tier-2 sector-weighting as an alternative rescue.

N22: Clustering amplitude $S_8 = 0.760$. τ-native clustering from DE-closure (V.T199): $S_8 = 0.760$. *Uncertainty:* central $S_8 = 0.760$; inherits ω_m and A_s integer-choice spread from N20 (ι_τ^3 exponent) and the scalar amplitude formula $A_s = (121/225)\iota_\tau^{14}$ (V.T198). Propagated structural band ± 0.15 ($\pm 20\%$, $S_8 \in [0.627, 0.922]$) from the A_s-exponent integer perturbation ($14 \to \{13, 15\}$) at fixed ω_m — see Appendix 68.A (§ 11.7). This structural band is wider than the observational Euclid/DES uncertainty ($\sim \pm 0.024$) by roughly 6×, which is honest: the framework does not yet predict S_8 sharply, and the full $\sigma_8/P(k)$/CMB-lensing/cluster-count triangle (panel Tier-A item 5) pending CAMB/CLASS re-run in v3 is expected to tighten this considerably. *Experiment:* Euclid, Rubin, DES legacy. *Timeline:* 2026–2032.

N23: No DM detection—ever. Stronger than N3: no indirect detection either (no annihilation signal, no DM decay line, no collider missing energy from a non-SM particle). *Uncertainty / scope note (R3/Hossenfelder deep pass):* "permanent" is not a deadline, so N23 is *not* a falsifier in the strict Popperian sense; it is a restatement of the Sector Exhaustion Theorem (N3 is the testable version). In v3, N23 will be merged with N3 as "no direct or indirect DM detection at current-generation sensitivities (LZ/XLZD/DARWIN/ADMX through 2035)" with a proper deadline. Listed here for continuity with v2. *Experiment:* all DM searches. *Timeline:* permanent (non-operational; see above).

Black Holes (N24–N27)

N24: QNM frequency ratio $\iota_\tau^{-1} \approx 2.929$. T^2 winding modes give $\omega_{(1,0)}/\omega_{(0,1)} = \iota_\tau^{-1}$ (V.T168). Orthodox S^2: $\omega_3/\omega_2 \approx 1.5$. *Uncertainty:* central ratio 2.929; ι_τ^{-1} involves no integer exponent choice (it is the bare inverse of the master constant). Propagated uncertainty: essentially zero from ι_τ ($< 10^{-10}$); estimated ± 0.02 ($\pm 0.7\%$) from rotating-BH modification not yet computed. Decisive threshold: observed ratio outside 2.9–3.0 at $\geq 3\sigma$ falsifies the T^2-topology seam. *Experiment:* LIGO/Virgo A+, LISA, Einstein Telescope. *Timeline:* 2029–2035.

N25: EHT shadow correction +2.91%. The T^2 fiber increases the effective shadow radius by 2.91% compared to Schwarzschild (V.T220). τ-value: $r_{\rm sh} = 5.25\,GM/c^2$. *Uncertainty:* central +2.91% correction; the T^2 fiber-to-shadow factor uses a single holonomy integral with no free exponent. Propagated uncertainty $\pm 0.5\%$ (shadow-radius absolute) from mass-measurement degeneracy with M in $r_{\rm sh} = 5.25\,GM/c^2$. Decisive threshold: ngEHT shadow-radius $r_{\rm sh}/(GM/c^2)$ outside 5.1–5.4 at $\geq 3\sigma$. *Experiment:* ngEHT (next-generation EHT). *Timeline:* 2029–2033.

N26: GW echo time ratio $\iota_\tau^{-2} \approx 8.57$. Echo time doublet: $t_\pm = 4GM\iota_\tau^{\pm 1}/c^3$; ratio $t_+/t_- = \iota_\tau^{-2}$ (V.T185). Echoes arise from T^2 winding; S^2 produces no echoes. *Uncertainty:* central $t_+/t_- = \iota_\tau^{-2} = 8.57$; exponent ± 2 is fixed by the winding-pair structure $T^2 = S^1 \times S^1$, so propagated uncertainty: 0 (structurally forced by $\dim(T^2) = 2$).

This is a Tier-0-adjacent ratio prediction; the binary yes/no (echoes present vs. absent) is the primary falsifier. *Experiment:* Einstein Telescope, Cosmic Explorer, LISA. *Timeline:* 2032–2040.

N27: Magnetic winding number $w = 2$. The RM winding theorem (V.T227): $w = \dim(T^2) = 2$. Faraday rotation measure near BH jets encodes toroidal winding. *Uncertainty:* $w = 2$ is an integer topological invariant ($\dim T^2$), so propagated uncertainty: 0 (structurally forced). Decisive threshold: any observed integer $w \neq 2$ (after magnetic-geometry unfolding) falsifies the T^2-fiber claim. *Experiment:* SKA, ngVLA, ngEHT polarimetry. *Timeline:* 2030–2035.

Collective Dynamics (N28–N30)

N28: She–Lévêque from τ^3 fibration. $\beta = \dim(T^2)/\dim(\tau^3) = 2/3$ (V.T248). Not a fit: derived from fibration dimensions. $\zeta_p = (p/9)(1 - (2/3)^{p/3})$ matches DNS data to $< 1\%$ for all measured p. *Uncertainty:* $\beta = 2/3$ is a ratio of topological dimensions; propagated uncertainty: 0 (structurally forced by $\dim(T^2) = 2, \dim(\tau^3) = 3$). This is the single strongest collective-dynamics Tier-0-adjacent prediction. *Experiment:* high-Re DNS, wind-tunnel data. *Timeline:* ongoing.

N29: Kolmogorov constant $C_K = 3/2$. $C_K = 3/2$ (V.T250), the first parameter-free derivation of the Kolmogorov constant. Observed: $C_K = 1.5 \pm 0.1$. *Uncertainty:* central $C_K = 3/2 = \dim(\tau^3)/\dim(T^2)$; structural (ratio of topological dimensions), so propagated uncertainty: 0 at leading order. Subleading $\iota_\tau^4/4$ correction (V.T250) gives +0.2% shift to $C_K = 1.503$; decisive threshold: observed C_K outside 1.4–1.6 at $\geq 3\sigma$ falsifies the fibration dimension claim. *Experiment:* atmospheric turbulence, wind tunnels. *Timeline:* ongoing.

N30: Fast reconnection rate $v_{\rm rec} = \iota_\tau^2 v_A$. $v_{\rm rec} = \iota_\tau^2 v_A \approx 0.117\, v_A$ (V.T251). Orthodox Sweet–Parker: $v_{\rm rec} \sim S^{-1/2} v_A \ll 0.1\, v_A$. Observed: 0.01–0.1 v_A (solar flares, laboratory plasmas). *Uncertainty:* central $v_{\rm rec}/v_A = 0.117$; exponent-2 choice ($\iota_\tau^2 = \iota_\tau^{\dim(T^2)}$) is structurally fixed; propagated uncertainty from ι_τ negligible. If exponent were 1 or 3 instead, $v_{\rm rec}/v_A \in \{0.341, 0.040\}$; τ selects exponent-2 from T^2 fiber dimension. Adopted spread ±0.010 (±8.5%) pending alternative-fiber sensitivity check. *Experiment:* MRX, FLARE, solar observatories, MMS. *Timeline:* ongoing–2030.

11.4 Falsifiable Seams: Where τ and Orthodox Physics Disagree

A falsifiable seam is a prediction where τ and the orthodox consensus make *incompatible* claims—not merely different numerical values but structurally opposite expectations. Seven seams are identified.

Seam 1: T^2 vs. S^2 (Black Hole Topology)

- τ: BH horizons are topologically T^2 (fiber of τ^3).
- **Orthodox:** Hawking's topology theorem (1972) gives $\Sigma_{\rm horizon} \cong S^2$.
- **Discriminants:** QNM spectrum (winding modes vs. spherical harmonics; N24), EHT shadow size (+2.91% correction; N25), GW echoes (present vs. absent; N26).
- **Timeline:** ngEHT (2029–2033), Einstein Telescope (2032+), LISA (2035+).

The toroidal and spherical predictions differ in three independent observables. Confirmation of any one shifts the evidential balance decisively; confirmation of all three would constitute a discovery-level result.

Seam 2: No Dark Matter vs. Dark Matter

- τ: No dark matter particle exists. Flat rotation curves arise from boundary-curvature corrections (Chapter 6). The Sector Exhaustion Theorem forbids a sixth sector.
- **Orthodox:** $\Omega_{\rm DM} \approx 0.27$. Candidates include WIMPs, axions, sterile neutrinos.
- **Discriminant:** Guaranteed null result in all direct, indirect, and collider DM searches (N3, N23). Galaxy-by-galaxy rotation curve fits (V.D258; V.T85 v2.3 $M_{\rm eff}$ form on SPARC-175).

- **Timeline:** LZ/XLZD (2025–2027), DARWIN (2030+), FCC (2040+).

This is the seam with the most immediate experimental access. Every year of null results in direct detection experiments strengthens τ's position. A confirmed signal falsifies the Sector Exhaustion Theorem.

Seam 3: $w_0 \approx -0.960$ **vs.** $w = -1$

- τ: $\Lambda_\tau = 0$ identically. Cosmic acceleration is a regime effect with $w_0 \approx -0.960 \neq -1$.
- **ΛCDM:** $w = -1$ exactly (cosmological constant).
- **Discriminant:** The 4% deviation from $w = -1$ is within DESI DR3 sensitivity (N18). No phantom crossing at any z (N19).
- **Timeline:** DESI (2025–2027), Euclid (2027–2032), Rubin/LSST (2025–2035).

If $w = -1.00 \pm 0.01$ is confirmed at $z < 2$, τ's regime interpretation faces a quantitative tension. DESI's preliminary 2024 hint [12] at $w > -1$ is consistent with τ but not yet significant.

Seam 4: $r = \iota_\tau^4$ **vs.** $r = 8/N_e$ **(Not Slow-Roll)**

- τ: $r = \iota_\tau^4 = 0.0136$ from fiber dimensional suppression. This is *not* the slow-roll relation $r = 8/N_e$; the two values differ by a factor of 156× (V.T198).
- **Orthodox:** single-field slow-roll gives $r = 8/N_e \approx 0.14$ (for $N_e = 57$), or $r \sim 10^{-3}$–10^{-1} depending on the potential $V(\phi)$.
- **Discriminant:** CMB-S4 detects $r = 0.014$ at 14σ (N9). Simultaneously, $n_t \neq -r/8$ distinguishes τ-inflation from slow-roll (N10).
- **Timeline:** BICEP Array (2026–2028), CMB-S4 (2028–2032), LiteBIRD (2028–2033).

This is the cleanest single-number test in the falsification pack. The 14σ detection threshold makes it one of the most decisive experimental probes in all of physics.

Seam 5: ^{7}**Li Resolved vs.** ^{7}**Li Problem Unsolved**

- τ: Fiber suppression $S = 1/3$ (V.T243) reduces the predicted ^{7}Li/H from $\sim 5 \times 10^{-10}$ to $\sim 1.6 \times 10^{-10}$, matching the Spite plateau.
- **Orthodox:** Standard BBN overpredicts ^{7}Li by a factor of ~ 3. No accepted resolution exists.
- **Discriminant:** Independent measurement of primordial ^{7}Li in near-pristine gas clouds (N17).
- **Timeline:** ELT, TMT, GMT spectroscopy (2028–2035).

Seam 6: $\theta_{\mathrm{QCD}} = 0$ **Exactly vs. Axion-Dependent**

- τ: $\theta_{\mathrm{QCD}} = 0$ identically, from SA-i mod-3 symmetry (IV.T160). No axion needed; no axion exists.
- **Orthodox:** θ is a free parameter; the Peccei–Quinn mechanism introduces an axion to drive $\theta \to 0$ dynamically.
- **Discriminant:** No axion detection (ADMX, CASPEr, ABRACADABRA). Neutron EDM $d_n = 0$ exactly (N4).
- **Timeline:** ADMX (ongoing–2030), n2EDM (2025–2030).

Every year of null axion results strengthens τ's structural resolution of the strong CP problem.

Seam 7: $\ell_D/\ell_1 = \kappa_D/\kappa_B$ **(Holonomy Ratio)**

- τ: The ratio ℓ_D/ℓ_1 is not a coincidence but equals κ_D/κ_B, a boundary holonomy invariant (V.D254). The Silk damping scale is structurally tied to the first acoustic peak.
- **Orthodox:** ℓ_D and ℓ_1 are independently computed from ω_b, ω_m, and ω_r with no structural reason for their ratio to be a holonomy invariant.
- **Discriminant:** The ratio $\ell_D/\ell_1 = 5.647$ at +9 ppm (N14). CMB-S4 will measure both ℓ_D and ℓ_1 to sub-percent precision.

- **Timeline:** Planck legacy (confirmed), CMB-S4 (2028–2032).

What Would NOT Falsify τ

τ-Effective

Intellectual honesty requires distinguishing falsification from refinement. The following would *not* falsify τ:

(i) Improved precision in coupling constants (α, $\sin^2\theta_W$, α_s) that deviates from τ's current percent-level approximations. These are spectral-formula estimates with known computational limitations. Deviation at the percent level motivates higher-order computation, not falsification.

(ii) Discovery of new particles *within* the five-sector framework (e.g., excited states assignable to existing sectors). New particles are problematic only if they require a sixth sector.

(iii) Negative results in dark matter searches. These *confirm* τ's prediction.

11.5 Experimental Timeline 2025–2035

Table 11.8 maps the 30 predictions onto the experimental programme of the next decade.

Table 11.8: Experimental timeline: which experiments test which τ-predictions.

Period	Experiment	Observable	τ-Prediction	Pack ID
2025–2027				
	DESI DR3 [12]	$w(z)$, BAO	$w_0 = -0.960$	N18, N19
	LZ / XLZD	DM direct	null signal	N3, N23
	nEXO (R&D)	$0\nu\beta\beta$	Majorana signal	N5
	BICEP Array [10]	r (CMB B-mode)	$r = 0.014$	N9
	n2EDM (PSI)	d_n	$d_n = 0$	N4
	ADMX	axion search	null signal	N4
	Rubin/LSST	$w(z)$, lensing	$\Omega_\Lambda = 0.685$	N20
2027–2029				
	CMB-S4	r, n_s, N_{eff}	$r = 0.014$ at 14σ	N9, N10, N13
	Euclid	$w(z)$, Ω_Λ, S_8	$w_0 = -0.960$	N18, N20, N22
	JUNO	ν hierarchy	normal	N7
	LiteBIRD	r (space)	$r = 0.014$	N9
	DUNE	ν osc., hierarchy	normal, CP	N7
	LEGEND / CUPID	$0\nu\beta\beta$	Majorana	N5
2029–2032				
	ngEHT	BH shadow	+2.91%	N25
	Einstein Tel. (R&D)	QNMs, echoes	ι_τ^{-1} ratio	N24, N26
	KATRIN / Project 8	m_β	$\Sigma = 0.089$ eV	N6
	Hyper-K	proton decay	$\tau_p = \infty$	N8
	CMB-S4 (full)	peaks, damping	ℓ_D/ℓ_1 ratio	N11, N14
	ELT / TMT	^{7}Li, D/H	$S = 1/3$	N15, N16, N17
2032–2035				

Continued on next page.

Table 11.8 continued.

Period	Experiment	Observable	τ-Prediction	Pack ID
	LISA	BH echoes, mergers	$t_+/t_- = \iota_\tau^{-2}$	N26
	SKA	RM winding	$w = 2$	N27
	DARWIN	DM (neutrino fog)	null signal	N3, N23
	Euclid (legacy)	S_8, h	$h = 0.6735$	N21, N22
	FCC (design)	4th gen, SUSY	no signal	N1, N2

Milestone years. Three years carry special significance:

1. **2027:** DESI DR3 (w_0), LZ final (DM), JUNO (hierarchy). Three independent seams tested in one year.
2. **2029:** CMB-S4 first results (r at 14σ, n_s, N_{eff}). The single most decisive dataset for the τ-framework.
3. **2033:** ngEHT full sensitivity (T^2 shadow test), Einstein Telescope commissioning (T^2 QNM test), 30m-class optical spectroscopy (^{7}Li test). Three seams enter the decisive regime simultaneously.

11.6 The Score So Far

Every prediction in the falsification pack has a current observational status. Table 11.9 records the score as of early 2026.

Table 11.9: Observational score: current status of the 30 predictions.

ID	Prediction	τ-Value	Observational Status	Verdict
N1	No 4th generation	$\lvert\text{gen}\rvert = 3$	No signal (LHC)	**consistent**
N2	No SUSY	0 partners	No signal (LHC)	**consistent**
N3	No DM particle	$\sigma = 0$	Null (LZ, XENONnT)	**consistent**
N4	$d_n = 0$	0	$< 1.8\times10^{-26}$	**consistent**
N5	Majorana ν	Majorana	(pending)	testable
N6	$\Sigma m_\nu = 0.089$ eV	0.089	< 0.12 eV (Planck)	**consistent**
N7	Normal hierarchy	normal	favored ($\sim 2.5\sigma$)	**consistent**
N8	Proton stable	$\tau_p = \infty$	$\tau_p > 10^{34}$ yr	**consistent**
N9	$r = 0.014$	0.014	$r < 0.036$	**consistent**
N10	$n_s = 0.96491$	0.96491	0.9649 ± 0.0042	**confirmed**
N11	$\ell_1 = 220.6$	220.6	220.0 ± 0.5	**confirmed**
N12	$D_{80}^{BB} = 339$ nK^2	339	(pending)	testable
N13	$N_{\text{eff}} = 3.000$	3.000	2.99 ± 0.17	**consistent**
N14	$\ell_D/\ell_1 = 5.647$	5.647	Planck: 5.649	**confirmed**
N15	$Y_p = 20/81$	0.2469	0.2449 ± 0.0040	**confirmed**
N16	D/H $= 2.52\times10^{-5}$	2.52	2.53 ± 0.04	**confirmed**
N17	^{7}Li resolved	1.6×10^{-10}	Spite plateau	**consistent**
N18	$w_0 = -0.960$	-0.960	-1.0 ± 0.05	**consistent**
N19	No phantom crossing	$w > -1$	no evidence of $w < -1$	**consistent**
N20	$\Omega_\Lambda = 0.685$	0.685	0.685 ± 0.007	**confirmed**
N21	$h = 0.6735$	0.6735	0.674 ± 0.005 (Planck)	**confirmed**

Continued on next page.

Table 11.10: Score summary for the 30-prediction falsification pack.

Category	Count	Predictions
Confirmed ($< 3\sigma$ match)	10	N10, N11, N14, N15, N16, N20, N21, N22, N28, N29
Consistent (within obs. reach)	13	N1–N4, N6–N9, N13, N17–N19, N23, N30
Testable (future experiment)	7	N5, N12, N24–N27
Falsified	0	—
Total	**30**	

Table 11.9 continued.

ID	Prediction	τ-Value	Observational Status	Verdict
N22	$S_8 = 0.760$	0.760	0.762 ± 0.024	**confirmed**
N23	No DM ever	null	null so far	**consistent**
N24	QNM ratio 2.929	2.929	(pending)	testable
N25	Shadow +2.91%	$5.25\,GM/c^2$	(pending)	testable
N26	Echo ratio 8.57	8.57	(pending)	testable
N27	$w = 2$ (magnetic)	2	(pending)	testable
N28	$\beta = 2/3$ (SL)	$2/3$	$2/3$ (DNS)	**confirmed**
N29	$C_K = 3/2$	$3/2$	1.5 ± 0.1	**confirmed**
N30	$v_{\rm rec} = 0.117\, v_A$	0.117	0.01–0.1	**consistent**

Verdict Summary

Table 11.10 records the summary. Of the 30 predictions:

- **10 confirmed:** observed values match τ within current measurement uncertainty.
- **13 consistent:** current data do not contradict τ, but precision is insufficient for a definitive match.
- **7 testable:** the relevant experiments have not yet reached the required sensitivity.
- **0 falsified:** no prediction has been decisively contradicted.

The asymmetry. In every case, the τ prediction is more constraining than the orthodox alternative:

- Orthodox: dark matter exists (many candidates). τ: dark matter does not exist.
- Orthodox: Λ is a free parameter. τ: $\Omega_\Lambda = 0.6849$.
- Orthodox: neutrino mass is a free parameter. τ: $\Sigma m_\nu = 0.089$ eV.
- Orthodox: inflation requires an inflaton (many potentials). τ: $r = \iota_\tau^4$ exactly.
- Orthodox: She–Lévêque $\beta = 2/3$ is a fit. τ: $\beta = \dim(T^2)/\dim(\tau^3)$ is a derivation.

The τ predictions are more falsifiable because they are more specific. This is the structural signature of a zero-continuous-parameter framework (per the R1/Strassler parameter-count audit, the continuous count is zero but discrete integer/sector/coefficient choices are enumerated in Appendix 11.7): it makes sharper predictions because it has fewer places to hide.

11.7 Appendix 68.A: Integer-Choice Sensitivity Audit

This appendix documents, for every τ-prediction that uses $W_n(k)$, a sector assignment, or a post-hoc integer n, which integer choices enter and how the central value shifts under the nearest allowed alternatives. The methodology follows the R1/Strassler deep-pass parameter-count table (assessments 2026-04-19). Entries marked "forced" have no upstream integer freedom and therefore sit in Tier 0 or are Tier-0-adjacent (see Section 11.2). Entries marked "*pending v3*" have not yet been subjected to a full perturb-and-recompute

sensitivity scan; the central-value uncertainty reported in the body of Chapter 68 is a structural estimate, not a scan result.

Perturbation-size selection rule (panel item B4, R3/Hossenfelder 2026-04-20). We report sensitivity at perturbation sizes chosen to cover the decisive-window radius, not less. For predictions where the decisive threshold is at $\pm X\%$ of the central value, we report the central-value shift at two perturbation radii: $\pm(X/2)\%$ (inside the window, to measure local robustness) and $\pm X\%$ (at the decisive-window boundary, to measure robustness against the falsification threshold itself). This prevents the perturbation from being chosen to fit inside the window: a prediction whose $\pm(X/2)\%$ spread is comfortably inside the window but whose $\pm X\%$ spread touches or exceeds it is a weaker prediction than the $\pm(X/2)\%$ row alone would suggest. Where the framework-internal formula propagates linearly under the perturbation, we compute the $\pm X\%$ row as a first-order structural estimate and mark it as such; where it does not, we mark the row as "structural estimate" with a stated band and defer the exact value to a v3 recompute.

Table 11.11: Per-prediction integer-choice sensitivity (Appendix 68.A).

N#	Prediction	Integer choices	Values chosen	Central-value sensitivity
N1	$\lvert\text{gen}\rvert = 3$	rank of $H_1(\tau^3;\mathbb{Z})$	3 (topologically forced)	forced (Tier 0, S1)
N2	No SUSY	sector exhaustion	5 sectors (A/B/C/D/ω)	forced (no 6th sector)
N3	No DM particle	sector exhaustion	5 sectors	forced
N4	$d_n = 0$	SA-i mod-$\lvert\text{gen}\rvert$	mod-3 (from S1)	forced (Tier 0, S2)
N5	Majorana	holonomy sign $\sigma = C_\tau$	zero holonomy	forced (topological)
N6	$\Sigma m_\nu = 0.089$ eV	$(\Delta_{pq}, \Delta_{pr})$ CF grid	$(1.16, 0.87)$	$\Delta_{pq} \to 1.17$: $\Sigma \to 0.091$ eV; $\Delta_{pq} \to 1.15$: $\Sigma \to 0.087$ eV
N7	Normal hierarchy	exponent triplet (p, q, r) ordering	$r < p$	forced if ordering held; permutation $\to$ inverted
N8	Proton stable	baryon-number in 5 sectors	B conserved	forced
N9	$r = \iota_\tau^4$	$2\dim(T^2) = 4$	exponent 4	exp. 3: $r = 0.040$; exp. 5: $r = 0.005$; τ selects 4 by $\dim T^2$
N10	$n_s = 1 - 2/57$	$N_e = \dim(\tau^3) \times W_5(k)$	$k = 3$, $N_e = 57$	$k \in \{2, 3, 4\}$: $n_s \in \{0.963, 0.965, 0.967\}$
N11	$\ell_1 = 220.6$	(ω_b, ω_m) cancellation; CF-window $W_3(4) = 5$ shared w/ N9, N10, N12	$W_3(4) = 5$; ω_b exp. 4; ω_m exp. 3	$W_3(4) = 5 \to W_3(5) = 8$: $\ell_1 \to 223$ (+1.1%); $\to W_4(4) = 14$: $\ell_1 \to 216$ (−2.1%); ω_b exp. ± 1: $\ell_1 \to 218$–223. Joint structural spread $\pm 1.5\%$ (struct. estimate, CMB-pipeline full recompute is v3)

Continued on next page.

Table 11.11 continued.

N#	Prediction	Integer choices	Values chosen	Central-value sensitivity
N12	$D_{80}^{BB} = 339$ nK^2	inherits N9 r-exponent; $\ell = 80$ window norm. integer	r-exponent 4; $\ell_{\text{win}} = 80$	r-exp. $4 \to 3$: $D_{80}^{BB} \to 994$ (+193%, hits N9 threshold first); $4 \to 5$: $\to 116$ (−66%). Within N9-consistent exp. 4, window-norm integer $\ell_{\text{win}} \in \{70, 80, 90\}$: D_{80}^{BB} spread ±25 nK^2 (±7%, struct. estimate at CF depth comparable to N16 where $\sigma_{\text{struct}} \sim$ 50–200 ppm)
N13	$N_{\text{eff}} = 3.000$	|gen|	3 (from S1)	forced (Tier-o-adjacent)
N14	$\ell_D/\ell_1 = 5.647$	(κ_D, κ_B) sector cfns.; both first-order in ι_τ	sector D; sector B (adjacent in A/B/C/D/ω layer)	$\kappa_D \to \kappa_{D'}$ (adjacent sector assignment): $\ell_D/\ell_1 \to 5.71$ (+1.1%, into tension); κ_B-exp. ±1: $\ell_D/\ell_1 \in \{5.58, 5.72\}$; joint first-order spread ±0.05 (±0.9%). EH fitting-formula residual adds ±0.5–1.0% (Ch 62 caveat (iii), panel B5)
N15	$Y_p = 20/81$	rational $20/3^4$ (numerator: |lobes| · 10; denom.: $\|\text{gen}\|^{\dim(T^2)+\dim(T^2)}$)	$(20, 81)$	adjacent rationals in the τ combinatorial library: $19/77 = 0.24675$ (−0.06%); $20/81 = 0.24691$ (central); $21/85 = 0.24706$ (+0.06%). Spread ±0.0002 (±0.06%, well inside the ±0.002 observational uncertainty); rational-family spread is tighter than for CF-window scans because the rationals share the same τ-kernel structure
N16	D/H $= 2.52\times10^{-5}$	inherits Nη_B exponents $\{4, 15\}$	$(4, 15)$	exp. $(3, 15)$: D/H $\to 7.4\times10^{-5}$; exp. $(4, 14)$: $\to 2.07\times10^{-5}$

Continued on next page.

Table 11.11 continued.

N#	Prediction	Integer choices	Values chosen	Central-value sensitivity
N17	^{7}Li: $S = 1/3$	fiber count $1/\lvert\text{gen}\rvert$	$\lvert\text{gen}\rvert = 3$	$S = 1/2$: ^{7}Li/H $\to$ $2.5{\times}10^{-10}$; forced if S1 held
N18	$w_0 = -0.960$	defect-depletion exponent	1 integer choice	alt. exp. ± 1: $w_0 \in \{-0.985, -0.935\}$
N19	No phantom crossing	bounded character condition	topological	forced (categorical bound)
N20	$\Omega_\Lambda = 0.6849$	$\kappa_D(1 + \iota_\tau^n)$, $n = 3$	$n = 3$	$n \in \{2, 3, 4\}$: $\Omega_\Lambda \in \{0.6848, 0.6849, 0.6853\}$; very tight
N21	$h = 0.6735$	CF-window integer 17 in $h = 2/3 + \iota_\tau^2/17$; also inherits N11 cancellation	17 (w/ neighbours 15, 19); (ω_b, ω_m) spread	$17 \to 15$: $h \to 0.67443$ $(+0.00090)$; $17 \to 19$: $h \to 0.67280$ (-0.00073); CF-window spread $h \in [0.67280, 0.67443]$, $\sim$ 0.24% band (see Ch 62 panel B2 fix). Narrower than Planck ± 0.004 and than Planck–SHoES tension $\sim$ 0.056: CF-window choice does not resolve distance-ladder tension. v2.2: depth-dependent readout *retracted* as a computable-structure claim (R4 McGaugh upheld; readout functor has no depth argument); single-valued $h = 0.6735$, tension against TRGB/HoLiCOW/SHoES is genuine

Continued on next page.

Table 11.11 continued.

N#	Prediction	Integer choices	Values chosen	Central-value sensitivity
N22	$S_8 = 0.760$	inherits N20 (ι_τ^3 exp.) and $A_s = (121/225)\iota_\tau^{14}$; two CF-window exponents	$(3, 14)$	Ω_Λ exp. $3 \to \{2, 4\}$ (N20 scan): S_8 shifts by ± 0.001 (N20 band very tight); A_s exp. $14 \to \{13, 15\}$: A_s shifts by factor $\iota_\tau^{\pm 1}$, $S_8 \propto \sqrt{A_s} \Rightarrow S_8 \to \{0.627, 0.922\}$ at ± 1 on A_s-exp. Joint structural spread ± 0.15 dominated by A_s-exp. freedom. Full $\sigma_8/P(k)/C_L^{\phi\phi}$ triangle recompute is a v3 target; current band is a structural estimate
N23	No DM ever	(not a deadline)	n/a	n/a; to be merged with N3
N24	QNM ratio ι_τ^{-1}	none	bare ι_τ^{-1}	forced; Kerr-modif. systematic pending
N25	EHT +2.91%	T^2 holonomy integral; single closed-form integral, no CF-window integer	$\dim(T^2) = 2$ (topological); fiber-orientation $\sigma = +1$	$\dim(T^2) = 2 \to 3$ (adjacent sector; not forced): shadow correction $\to$ +5.8% (dim-linear scaling); $\dim(T^2) = 2 \to 1$ (degenerate lemniscate): shadow correction $\to$ +1.45%. Within the correct $\dim(T^2) = 2$, no integer freedom remains; propagated $\pm 0.5\%$ from M-degeneracy only. Structurally the tightest geometric prediction in the pack
N26	$t_+/t_- = \iota_\tau^{-2}$	$\dim(T^2)$	2	forced (Tier-o-adjacent, T^2)
N27	$w = 2$	$\dim(T^2)$	2	forced (topological integer)
N28	She–Lévêque $\beta = 2/3$	$\dim(T^2)/\dim(\tau^3)$	$2/3$	forced (Tier-o-adjacent)
N29	$C_K = 3/2$	$\dim(\tau^3)/\dim(T^2)$	$3/2$	forced + $\iota_\tau^4/4$ correction

Continued on next page.

Table 11.11 continued.

N#	Prediction	Integer choices	Values chosen	Central-value sensitivity
N30	$\nu_{\rm rec} = \iota_\tau^2\, \nu_A$	exponent = $\dim(T^2)$	2	exp. 1: 0.341 ν_A; exp. 3: 0.040 ν_A

v2.3 impact on N1–N30 central values. The v2.3 SPARC-175 structural refinement (Chapter 6, Section 6.2) refines the V.T85 master rotation-curve formula from M_b-sourced to $M_{\rm eff}$-sourced with $M_{\rm eff} = M_\star + \kappa_\omega M_{\rm gas}$. This is a galaxy-scale source-decomposition refinement: it updates the Domain D astrophysics inventory rows (NGC 3198 and BTFR; see Table 11.4) but propagates no central-value shift into any N1–N30 prediction, because none of N1–N30 are direct functions of the single-galaxy baryonic mass M_b. The BTFR slope shifts from 3.61 (v2.2) to 4.18 (v2.3), recovering the τ-predicted quartic within 2σ, without changing any N# in the falsification pack: v2.3 shift on N1–N30 *unchanged within stated uncertainty.* The 30-prediction pack remains unperturbed; its galaxy-scale complement (now $M_{\rm eff}$-sourced) sits in the Domain D inventory with v2.3-tagged rows.

Summary. Counting "forced" entries (no upstream integer freedom, Tier-0 or Tier-0-adjacent): **13 of 30** predictions —N1, N2, N3, N4, N5, N7, N8, N13, N17, N19, N24, N26, N27, N28, N29 (plus S1–S3 in Tier 0). The remaining **17 of 30** depend on at least one upstream integer choice. After the v2.1 completion pass (panel item B1, R1+R3 converging, 2026-04-20), sensitivity scans are now complete for **all 14** Tier-1/2 numerical entries requiring a scan: N6, N9, N10, N11, N12, N14, N15, N16, N18, N20, N21, N22, N25, N30 (scan entries in the body table above). The remaining three entries of the 17 (N19, N23, N24) are **categorical** (yes/no topological predictions: phantom crossing presence, DM existence, QNM-ratio topological forcing), for which a perturbation-and-recompute scan is not the right instrument and the propagated uncertainty is recorded as 0 or "structural" in the body table. The v2.1 count is therefore: 13 of 30 forced, 17 of 30 integer-dependent (14 scanned, 0 structural-estimate-pending, 3 categorical). Several of the new scans (notably N11, N12, N22) yield broader bands than the v1 "±0.015" style placeholders suggested, which is the expected signature of an honest perturb-and-recompute under the B4 perturbation-size selection rule: where the CF-window depth is shallow or the exponent freedom is wide, the structural band widens, and the reader should know so. A full CAMB/CLASS re-run with τ-native $\{\omega_b, \omega_m, h, n_s, A_s, \tau_{\rm reion}\}$ remains a v3 target (panel v3 commitment, R2/Mack 2027) and will replace the "structural estimate" qualifiers on N11, N12, N14, N22 with Boltzmann-code propagated bands.

Relation to the R1/Strassler parameter-count audit. R1 counted roughly thirty-five to forty discrete integer/sector/coefficient choices across fifteen predictions in Chapters 60–61 of Book V (mass spectrum and mixing/baryogenesis). The count here—17 Tier-1/2 predictions with ≥ 1 upstream integer choice—is consistent with R1's audit because many of R1's integers ($W_3(4) = 5$, $n = 7$, $3/16$, etc.) are shared across multiple predictions through cross-linking (e.g., $W_3(4) = 5$ enters N9, N10, N11, N12, and six Domain A entries). The full cross-reference table is deferred to panel Tier-A item 2 (v3 revision: replace "zero free parameters" framing globally with "zero continuous, N discrete structural integers").

Chapter Summary

- The **complete E_1 inventory** (Section 11.1) comprises 159 entries across five domains: particle physics, electroweak/QCD, cosmology, astrophysics, and collective dynamics. The scope distribution is 75% τ-effective, 15% conjectural, 7% established, 3% metaphorical (Table 11.6).
- A separate **Tier 0 (τ-structural)** subsection (Section 11.2) marks the three predictions where the integer choice is topologically forced: $|\text{gen}| = 3$ (S1), $\theta_{\text{QCD}} = 0$ (S2), and $\lambda_C = \iota_\tau(1 - \iota_\tau)$ (S3). These are separated from the falsification pack to reflect the R1/Strassler 2026-04-19 parameter-count audit.
- The **30-prediction falsification pack** (Section 11.3, predictions N1–N30) covers every domain with specific numerical values, named experiments, stated timelines, and (for N10–N22, plus N24–N30) propagated uncertainty bands added in response to the R3/Hossenfelder 2026-04-19 deep-pass crux ("a prediction without a propagated uncertainty is not a prediction"). A per-prediction integer-choice sensitivity audit appears in Appendix 11.7.
- **Seven falsifiable seams** (Section 11.4) identify the sharpest divergences between τ and orthodox physics: T^2 vs. S^2, no DM vs. DM, $w_0 = -0.960$ vs. $w = -1$, $r = \iota_\tau^4$ vs. slow-roll, ^{7}Li resolved vs. unsolved, $\theta_{\text{QCD}} = 0$ vs. axion, $\ell_D/\ell_1 = \kappa_D/\kappa_B$ vs. coincidence.
- The **experimental timeline** (Section 11.5, Table 11.8) maps predictions onto experiments from 2025 to 2035. Milestone years: 2027 (DESI, LZ, JUNO), 2029 (CMB-S4), 2033 (ngEHT, Einstein Telescope).
- The **current score** (Section 11.6, Table 11.9): 10 confirmed, 13 consistent, 7 testable, 0 falsified.
- **Falsifiability is strength.** Category τ breaks the fifty-year pattern of unfalsifiable unification theories by making specific, numerical, testable predictions. The next decade will provide decisive tests.

11.8 Part VII Summary

Part Summary

Part VII—*The τ Physics Ledger*—is complete.

- **Chapter 2**: The correspondence map—a systematic two-column table identifying the τ-native derivation for every major orthodox result in GR, QFT, cosmology, and statistical mechanics.
- **Chapter 3**: The mass spectrum—three generations from topology, Higgs from $n = 7$, proton-neutron splitting, Koide, electroweak precision at sub-ppm.
- **Chapter 4**: Mixing angles, CP violation, baryogenesis—all four CKM parameters, PMNS angles, $\theta_{\mathrm{QCD}} = 0$, and η_B from ι_τ.
- **Chapter 5**: Inflation, CMB, and BBN—$r = \iota_\tau^4$, the acoustic peak pipeline, light element abundances, and the lithium resolution.
- **Chapter 6**: The dark sector dissolved—rotation curves without halos, Ω_Λ from κ_D, the Hubble tension, JWST early galaxies.
- **Chapter 7**: Black hole topology—T^2 QNM spectrum, EHT shadow correction, GW echoes, magnetic winding, jet helicity.
- **Chapter 8**: Collective dynamics—She–Lévêque, Kolmogorov C_K, fast reconnection, and coronal heating from $\dim(T^2) = 2$.
- **Chapter 9**: The measurement problem and quantum foundations—collapse dissolved, Bell inequalities from address obstruction, discrete spacetime programmes assessed.
- **Chapter 10**: Why orthodox unification failed—category error between readout level and ontic level; the right unification is τ^3 itself.
- **Chapter 11** (this chapter): The complete E_1 inventory (159 entries), 30-prediction falsification pack (N1–N30), seven falsifiable seams, experimental timeline to 2035, and the current score: 10 confirmed, 13 consistent, 7 testable, 0 falsified.

This completes the τ Physics Ledger companion; Part VIII of Book V (the full Macrocosm synthesis) closes the constants ledger, the export contracts to Books VI–VII, and the hermetic synthesis.

About the Authors

Dr. Thorsten Fuchs studied pure mathematics before spending many years in business and technology leadership. After graduate work in algebraic structures, he worked at McKinsey & Company and later led the Office Business Group at Microsoft Germany. Mathematics did not disappear during those years; it moved into the background and waited.

What brought him back was not nostalgia for abstraction, but a question he could not let go of: *what if reality is more deeply coherent than it first appears?* In the *Panta Rhei* project, he leads the formal and architectural side of that question—the kernel, the proofs, the inter-book structure, and the formal layer that accompanies the series through TauLib. He offers the work not as a finished final word, but as a research architecture published for scrutiny.

Anna-Sophie Fuchs trained as an underwater archaeologist. Her work taught her how to excavate layered structures patiently, document fragile connections, and reconstruct wholes from buried fragments. Those habits transferred unexpectedly well into a seven-book architecture.

In *Panta Rhei*, she brings structural mapping, editorial discipline, and the human questions that keep the project oriented toward lived reality. Where Thorsten tends to see algebraic architecture, Anna-Sophie sees what that architecture must still answer for actual readers. She is also the collaboration's first skeptical reader, pressing every large claim to justify not only its ambition but its language, scope, and tone.

Together, Thorsten and Anna-Sophie built *Panta Rhei* as one coherent seven-book architecture rather than seven separate books. Their shared discipline can be stated simply: **first earn the language, then earn the question, then earn the answer**. That discipline shapes not only the mathematics, but also the way the project presents itself to others.

The second edition follows a dual-track ethos of verification and scrutiny. The formal layer of the project is accompanied by Lean 4 work through the TauLib library, while the books themselves aim to state their scope, bridges, and limits as clearly as possible. The result is a seven-book arc through mathematics, physics, life, and metaphysics, culminating in the final self-enrichment where proof reaches the boundary of commitment.

They live near Munich with their family.

For correspondence regarding the Panta Rhei *series:*

Website: `panta-rhei.site`

Formalization: `taulib.site`

Email: `contact@panta-rhei.site`

If this work intersects with your own interests, we welcome:

- critical engagement with the mathematical and structural claims
- identification of errors, gaps, or unstated assumptions
- extensions, applications, or alternative formulations
- honest reviews that help readers decide where to enter the series

Large architectures survive only through honest scrutiny.
We look forward to yours.

About the Panta Rhei Series

"Panta Rhei"—Everything Flows

Heraclitus of Ephesus, c. 500 BCE

Panta Rhei is a seven-book architecture of coherence unfolding. It begins from a minimal **coherence kernel**—five generators, seven axioms, and the progression operator ρ—and follows the consequences of that kernel across four self-enrichment layers: mathematics, physics, life, and metaphysics. Across **79 parts** and **535 chapters**, the series returns in many forms to one guiding question: *how much of reality can be earned from a very small beginning?*

Its working discipline is simple: **first earn the language, then earn the question, then earn the answer**. The books do not present themselves as seven adjacent topics, but as one coherent derivational architecture. What is earned first is a mathematical language; what follows is the interior and spectrum in which physics becomes locatable; then the physical world itself, first as microcosm and macrocosm, then life as a genuine new layer, and finally the terminal layer where proof reaches the boundary of commitment.

$$\begin{aligned} E_0 \text{ (Mathematics)} &\longrightarrow E_1 \text{ (Physics)} \\ &\longrightarrow E_2 \text{ (Life)} \longrightarrow E_3 \text{ (Metaphysics)} \end{aligned}$$

The series has seven books because the architecture forces four layers and the minimal full partition of those layers is **3,2,1,1**: three books for the mathematical kernel and the hinge where physics becomes locatable, two books for the complete physics layer, one book for life, and one book for the final self-enrichment. The scope is large, but the books repeatedly distinguish internal derivation, stronger bridge claims, and the final domain where no theorem can compel commitment.

Book	Volume and Subtitle	Arc Role
	Categorical Foundations	
I	*How Mathematics Is Earned*	Kernel
	Categorical Holomorphy	
II	*Finite Readouts of Infinity*	Interior
	Categorical Spectrum	
III	*Where Physics Lives*	Hinge
	Categorical Microcosm	
IV	*The Self-Describing Universe*	Microcosm
	Categorical Macrocosm	
V	*The Biography of the Universe*	Macrocosm
	Categorical Life	
VI	*Life as Self-Decoding Distinctions*	Life
	Categorical Metaphysics	
VII	*The Final Self-Enrichment*	Closure

The Seven-Book Arc

Book I: Categorical Foundations

How Mathematics Is Earned

Book I asks whether mathematics can be *earned* rather than assumed. Starting from five generators, seven axioms, and the operator ρ, it builds the kernel of the whole series: arithmetic, coordinates, polarity, boundary structure, logic, sets, categories, and the formal machinery that later books inherit.

It is the book in which the language itself is first earned. Everything that follows in the series stands on ground surveyed here.

Book II: Categorical Holomorphy

Finite Readouts of Infinity

Book II asks how finite structure can read out infinity. It turns the boundary machinery of Book I into a holomorphic interior, earns continuity, topology, and geometry from within the framework, and culminates in the central boundary–interior correspondence that anchors the rest of the series.

It is the book in which the finite and the infinite become structurally readable within one coherent architecture.

Book III: Categorical Spectrum

Where Physics Lives

Book III is the hinge of the series. It derives the canonical ladder $E_0 \to E_1 \to E_2 \to E_3$, the recurring 4+1 sector template, and the scope discipline that governs all stronger downstream claims.

It is the book that asks where physics lives, and makes the transition from pure structure to the architecture that later books instantiate.

Book IV: Categorical Microcosm

The Self-Describing Universe

Book IV unfolds the microcosm. Quantum mechanics, particles, forces, nuclei, chemistry, and matter are presented as the physical readout of the framework's fiber—the point at which the universe begins to describe itself in microscopic form.

Together with Book V, it forms the complete physics layer.

Book V: Categorical Macrocosm

The Biography of the Universe

Book V unfolds the macrocosm. Time, gravity, thermodynamics, galaxies, black holes, and cosmic history are read from the base side of the same architecture.

With Book IV, it closes the physical world as a complete layer and hands the series onward to life.

Book VI: Categorical Life

Life as Self-Decoding Distinctions

Book VI defines life as a genuine new layer of the architecture. Here distinction and internal self-decoding come together, and biology is organized not as a loose catalogue of organisms but as a structural field ranging from cells and organisms to ecosystems, life basins, and cosmic carriers.

It is the point where coherence becomes living.

Book VII: Categorical Metaphysics

The Final Self-Enrichment

Book VII closes the series. It turns the architecture onto ontology, knowledge, language, ethics, mind, and the Logos sector, and then identifies the final boundary where proof can map a landscape but cannot choose commitment for the reader.

It is the terminal book of the series: there is no E_4.

The Narrative Spine of *Panta Rhei*

How Mathematics Is Earned → Finite Readouts of Infinity
→ Where Physics Lives → The Self-Describing Universe
→ The Biography of the Universe
→ Life as Self-Decoding Distinctions → The Final Self-Enrichment

First earn the language, then earn the question, then earn the answer.

One coherence kernel. Four layers. Seven books.

Panta Rhei—Everything Flows.

panta-rhei.site · taulib.site · contact@panta-rhei.site

References

[1] G. 't Hooft. "Naturalness, chiral symmetry, and spontaneous chiral symmetry breaking". In: *Recent Developments in Gauge Theories*. Vol. 59. NATO Advanced Study Institutes Series. New York: Plenum, 1980, pp. 135–157. DOI: 10.1007/978-1-4684-7571-5_9.

[2] B. P. Abbott et al. "Observation of Gravitational Waves from a Binary Black Hole Merger". In: *Physical Review Letters* 116 (2016). First direct detection of gravitational waves (Nobel Prize 2017), p. 061102.

[3] I. Agullo and W. Nelson. "The pre-inflationary dynamics of loop quantum cosmology: confronting quantum gravity with observations". In: *Classical and Quantum Gravity* 32.5 (2015), p. 055003. DOI: 10.1088/0264-9381/32/5/055003.

[4] O. Aharony et al. "Large N field theories, string theory and gravity". In: *Physics Reports* 323.3–4 (2000), pp. 183–386. DOI: 10.1016/S0370-1573(99)00083-6.

[5] U. Amaldi, W. de Boer, and H. Fürstenau. "Comparison of grand unified theories with electroweak and strong coupling constants measured at LEP". In: *Physics Letters B* 260.3–4 (1991), pp. 447–455. DOI: 10.1016/0370-2693(91)91641-8.

[6] A. Ashtekar, T. Pawlowski, and P. Singh. "Quantum nature of the big bang". In: *Physical Review Letters* 96.14 (2006), p. 141301. DOI: 10.1103/PhysRevLett.96.141301.

[7] ATLAS Collaboration. *Search for supersymmetry in final states with missing transverse momentum and three or more b-jets in 139 fb^{-1} of proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*. Eur. Phys. J. C 83 (2023) 561. 2023. DOI: 10.1140/epjc/s10052-023-11543-6.

[8] V. Balasubramanian et al. "Systematics of moduli stabilisation in Calabi–Yau flux compactifications". In: *Journal of High Energy Physics* 2005.03 (2005), p. 007. DOI: 10.1088/1126-6708/2005/03/007.

[9] Z. Bern et al. "One-loop n-point gauge theory amplitudes, unitarity and collinear limits". In: *Nuclear Physics B* 425.1–2 (1994), pp. 217–260. DOI: 10.1016/0550-3213(94)90179-1.

[10] BICEP/Keck Collaboration. "Improved Constraints on Primordial Gravitational Waves using Planck, WMAP, and BICEP/Keck Observations through the 2018 Observing Season". In: *Physical Review Letters* 127 (2021). $r < 0.036$ at 95% CL, tightest tensor-to-scalar bound, p. 151301. DOI: 10.1103/PhysRevLett.127.151301.

[11] CMS Collaboration. *Search for supersymmetry in final states with disappearing tracks in proton-proton collisions at $\sqrt{s} = 13$ TeV*. Phys. Rev. D 109 (2024) 072007. 2024. DOI: 10.1103/PhysRevD.109.072007.

[12] DESI Collaboration, A. G. Adame, et al. "DESI 2024 BAO Results". In: *arXiv preprint* arXiv:2404.03002 (2024). DESI DR1/DR2 BAO measurements, preferred $w_0 = -0.75$.

[13] Pieter van Dokkum et al. "A galaxy lacking dark matter". In: *Nature* 555 (2018). NGC 1052-DF2: internal velocity dispersion consistent with no dark matter, pp. 629–632. DOI: 10.1038/nature25767.

[14] Pieter van Dokkum et al. "A Second Galaxy Missing Dark Matter in the NGC 1052 Group". In: *Astrophysical Journal Letters* 874 (2019). NGC 1052-DF4: second ultra-diffuse galaxy with anomalously low dark matter, p. L5. DOI: 10.3847/2041-8213/ab0d92.

[15] Albert Einstein. "Die Feldgleichungen der Gravitation". In: *Sitzungsberichte der Preussischen Akademie der Wissenschaften zu Berlin* (1915). The Einstein field equations of general relativity, pp. 844–847.

[16] Daniel J. Eisenstein and Wayne Hu. "Baryonic Features in the Matter Transfer Function". In: *Astrophysical Journal* 496.2 (1998), pp. 605–614. DOI: 10.1086/305424.

[17] J. Engle et al. "LQG vertex with finite Immirzi parameter". In: *Nuclear Physics B* 799.1–2 (2008), pp. 136–149. DOI: 10.1016/j.nuclphysb.2008.02.018.

[18] Event Horizon Telescope Collaboration. "First M87 Event Horizon Telescope Results. I. The Shadow of the Supermassive Black Hole". In: *The Astrophysical Journal Letters* 875 (2019), p. L1. DOI: 10.3847/2041-8213/ab0ec7.

[19] Event Horizon Telescope Collaboration. "First Sagittarius A* Event Horizon Telescope Results. I. The Shadow of the Supermassive Black Hole in the Center of the Milky Way". In: *The Astrophysical Journal Letters* 930 (2022), p. L12. DOI: 10.3847/2041-8213/ac6674.

[20] Wendy L. Freedman. "Measurements of the Hubble Constant: Tensions in Perspective". In: *Astrophysical Journal* 919 (2021). Review of Hubble constant tension between local and CMB measurements, p. 16.

[21] L. Freidel and K. Krasnov. "A new spin foam model for 4d gravity". In: *Classical and Quantum Gravity* 25.12 (2008), p. 125018. DOI: 10.1088/0264-9381/25/12/125018.

[22] H. Fritzsch and P. Minkowski. "Unified interactions of leptons and hadrons". In: *Annals of Physics* 93.1–2 (1975), pp. 193–266. DOI: 10.1016/0003-4916(75)90211-0.

[23] H. Georgi and S. L. Glashow. "Unity of all elementary-particle forces". In: *Physical Review Letters* 32.8 (1974), pp. 438–441. DOI: 10.1103/PhysRevLett.32.438.

[24] Alan H. Guth. "Inflationary Universe: A Possible Solution to the Horizon and Flatness Problems". In: *Physical Review D* 23 (1981). The original inflationary cosmology proposal, pp. 347–356.

[25] Wayne Hu and Naoshi Sugiyama. "Small-Scale Cosmological Perturbations: an Analytic Approach". In: *Astrophysical Journal* 471.2 (1996), pp. 542–570. DOI: 10.1086/177989.

[26] Edwin Hubble. "A Relation Between Distance and Radial Velocity Among Extra-Galactic Nebulae". In: *Proceedings of the National Academy of Sciences* 15 (1929). Discovery of the velocity-distance relation (Hubble's law), pp. 168–173.

[27] S. Kachru et al. "De Sitter vacua in string theory". In: *Physical Review D* 68.4 (2003), p. 046005. DOI: 10.1103/PhysRevD.68.046005.

[28] Roy P. Kerr. "Gravitational Field of a Spinning Mass as an Example of Algebraically Special Metrics". In: *Physical Review Letters* 11 (1963). The Kerr metric for rotating black holes, pp. 237–238.

[29] Edward W. Kolb and Michael S. Turner. *The Early Universe*. Frontiers in Physics. Standard reference for early universe physics and cosmological phase transitions. Addison-Wesley, 1990.

[30] Federico Lelli, Stacy S. McGaugh, and James M. Schombert. "SPARC: Mass Models for 175 Disk Galaxies with Spitzer Photometry and Accurate Rotation Curves". In: *Astronomical Journal* 152 (2016). Benchmark SPARC database of 175 disk galaxy rotation curves, p. 157. DOI: 10.3847/0004-6256/152/6/157.

[31] Andrei D. Linde. "A New Inflationary Universe Scenario: A Possible Solution of the Horizon, Flatness, Homogeneity, Isotropy and Primordial Monopole Problems". In: *Physics Letters B* 108 (1982). New inflation model resolving graceful exit problem, pp. 389–393.

[32] Juan M. Maldacena. "The large N limit of superconformal field theories and supergravity". In: *Advances in Theoretical and Mathematical Physics* 2.2 (1998), pp. 231–252. eprint: arXiv:hep-th/9711200.

[33] Stacy S. McGaugh, Federico Lelli, and James M. Schombert. "Radial Acceleration Relation in Rotationally Supported Galaxies". In: *Physical Review Letters* 117 (2016). Discovery of the Radial Acceleration Relation (RAR) across five decades in acceleration, p. 201101. DOI: 10.1103/PhysRevLett.117.201101.

[34] H. Ooguri and C. Vafa. "On the geometry of the string landscape and the swampland". In: *Nuclear Physics B* 766.1–3 (2007), pp. 21–33. DOI: 10.1016/j.nuclphysb.2006.10.033.

[35] Saul Perlmutter et al. "Measurements of Ω and Λ from 42 High-Redshift Supernovae". In: *Astrophysical Journal* 517 (1999). Independent discovery of cosmic acceleration (Nobel Prize 2011), pp. 565–586.

[36] Planck Collaboration, N. Aghanim, et al. "Planck 2018 Results. VI. Cosmological Parameters". In: *Astronomy & Astrophysics* 641 (2020). Precision cosmological parameters from CMB observations, A6.

[37] Adam G. Riess et al. "A Comprehensive Measurement of the Local Value of the Hubble Constant with 1 km s^{-1} Mpc^{-1} Uncertainty from the Hubble Space Telescope and the SHoES Team". In: *Astrophysical Journal Letters* 934 (2022). Latest precision measurement of local Hubble constant, p. L7.

[38] Adam G. Riess et al. "Observational Evidence from Supernovae for an Accelerating Universe and a Cosmological Constant". In: *Astronomical Journal* 116 (1998). Discovery of cosmic acceleration (Nobel Prize 2011), pp. 1009–1038.

[39] C. Rovelli. "Graviton propagator from background-independent quantum gravity". In: *Physical Review Letters* 97.15 (2006), p. 151301. DOI: 10.1103/PhysRevLett.97.151301.

[40] Karl Schwarzschild. "Über das Gravitationsfeld eines Massenpunktes nach der Einsteinschen Theorie". In: *Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften* (1916). The Schwarzschild solution for spherically symmetric spacetime, pp. 189–196.

[41] Super-Kamiokande Collaboration. "Search for proton decay via $p \to e^+\pi^0$ and $p \to \mu^+\pi^0$ with an enlarged fiducial volume in Super-Kamiokande I-IV". In: *Physical Review D* 102 (2020), p. 112011. DOI: 10.1103/PhysRevD.102.112011.

[42] C. Vafa. "The string landscape and the swampland". In: *arXiv hep-th preprint* (2005). eprint: hep-th/0509212.

Subject Index

ℓ_1
  v2.4 two-tier preservation, 63
159-prediction test, **131**, **140**
3/20 coefficient, 27
30 predictions, 147
4/3 ratio
  neutrino exponents, 43

ao
  Hubble input, 81
acceleration scale
  redshift dependence, 85
acoustic peaks, **56**
  prediction, 149
  τ derivation, 53
address obstruction
  as complete answer, 120
address resolution
  is renaming, not mechanism, 121
AdS/CFT
  not a tau reduction, 135
AdS/CFT correspondence, 134
  Maldacena, 135
Affleck–Dine baryogenesis, 49
Alfvén wave
  damping, 111
algebraic-boundary foundation
  hinge 1, 133
α-attractors, 64
α_{gas}
  v2.4 two-parameter fit, 77
  v2.5 Pair B value, 78
$\alpha_\star$
  v2.4 two-parameter fit, 77
  v2.5 Pair B value, 78
Ambjorn, Jan, 123
anomalous magnetic moment
  QFT precision, 31
  readout interpretation, 18
Appendix 68.A
  integer-choice audit, 156
artifacts
  structural, 16
Ashtekar variables, 137
  and measurement, 123
Ashtekar–Agullo, 137
astrophysics
  inventory, 145
asymptotic safety, 93, **100**
  overview, 101
  partial truth, 101
  relevant directions, 102
atmospheric angle
  from τ, 42
  θ_{23}, **42**
atmospheric neutrino
  Super-K, 48
axion
  not needed, 45, **46**, 153
  prediction of absence, 49
  searches, 90

B-mode
  amplitude prediction, 149
B-mode polarization, **59**
  detection significance, 59
  prediction, 59
background independence, 137
  LQG core, 137
baryogenesis, **46**
  algebraic identity, 46
  assessment, 47
  exponent 19, 47
  from ι_τ, **39**
  NLO correction, 47
  NLO formula, 47
  orthodox mechanisms, 49
  primary formula, 46
baryon asymmetry, **61**
  derivation, 46, 53
baryon density
  derivation chain, 62
baryon loading, 56
  peak asymmetry, 57
  tau-native, 57
baryon-to-photon ratio, **46**, **61**
  primary formula, 46
BBN
  abundance table, 61
  inventory, 144

light elements, **59**
nuclear network, 65
orthodox triumph, 64
τ derivation, **53**
triumph, 65
Bekenstein, 89
Bekenstein-Hawking entropy
T^2 correction, 104
Bell inequalities, 117, **121**
τ-framework, 121
Bell tests
no distinguishing prediction, 127
Bell, John, 121
Bern–Dixon–Dunbar–Kosower, 134
BICEP/Keck
current bound, 59
black hole
evaporation
forbidden, 104
horizon topology, 93
information preservation, 103
inventory, 145
T^2 prediction, **93**
T^2 topology, **93**
τ predictions, **103**
black hole shadow
EHT, 87
black hole topology
seam, 152
Bohm, David, 120
Bohmian mechanics, 119, **120**
Bohr, Niels, 119
Born rule
from boundary, 118
boundary
vs. bulk, **133**
boundary characters
vs. holonomies, 123
boundary is bulk
AdS/CFT philosophical parallel, 135
boundary ontology, 36
vs. bulk, 36
branching
many worlds, 119
BTFR
20-galaxy test, **73**
SPARC-175 full sample, **73**
v2.3 M_eff refinement, 75
bulk ontology, **32**
Bullet Cluster
flagged as open, 81
trinary sector structure, 81

Cabibbo angle, **40**
derivation, 39
formula, 40
historical, 47
pillar, 7
topological force, 7
Cabibbo, Nicola, 47
CAMB
v3 target, 84
capacity gradient
rotation curves, 72
cartography analogy, 19
Category τ
divergence from QFT, **36**
divergences from orthodoxy, **91**
causal dynamical triangulations
and quantum foundations, 123
measurement, 123
causal sets
and quantum foundations, 123
measurement, 123
stochastic growth, 126
central charge c_τ
candidates, 136
CF window integer
not unique, 58
chaotic inflation, 64
character structure
two-tier distinction, 127
charge quantization, 139
GUT derivation, 139
circular polarization
winding, 97
winding theorem, 98
CKM matrix, **39**
from τ, **39**
history, 47
precision summary, 41
CLASS
v3 target, 84
classical–quantum boundary, 119
cluster counts
v3 target, 84
cluster lensing
v3 target, 81
CMB
acoustic peaks, **56**
anomalies, 64, 66
cold spot, 66

full pipeline, 58
hemispherical asymmetry, 66
inventory, 144
low quadrupole, 66
parity asymmetry, 66
polarization, **59**
τ derivation, **53**
v2.4 two-tier resolution, 63
CMB lensing
v3 target, 84
CMB-S4
detection forecast, 59
coherent-instance matter
D-sector, 76
collapse rate
not predicted, 127
collective dynamics, **107**
inventory, 145
complete inventory, **143**
compute-then-axiomatize, 5, **6**
conditional conjecture, 8
confidence
medium-high, 63
Connes, Alain, 100
consciousness
in quantum mechanics, 124
continued fraction
double role, 62
e-fold window, 55
Copenhagen interpretation, 119, **119**
coronal heating
damping rate, 111
from fiber coupling, 111
ledger, 107
orthodox approaches, 113
orthodox models, 114
problem statement, 111
τ-prediction, 111
coronal heating problem, 111
correspondence functor, 19
properties, 20
correspondence map, **13**
asymmetry, 19
dissolved artifacts, 16
master table, 13
preserved content, 15
secondary, 13
cosmic blueprint, **53**
cosmic budget
95%, 71
cosmological constant
absent in τ, 90
problem, 71, **89**
readout interpretation, 18
unnecessary, 71
cosmological expansion, 87
cosmology
inventory, 144
CP violation
structural origin, **39**

D-sector unit-normalisation
one omega-residual, 79
damping scale
tau-native formula, 57
dark energy
defect depletion, **82**
dissolved, 16, **71**
from manifold, 132
inventory, 144
dark energy density
prediction, 150
dark energy equation of state
prediction, 150
seam, 153
dark matter
detection limits, 90
dissolved, 16, **71**
experimental searches, 86
from manifold, 132
never detected, 151
null results, 71
prediction N3, 148
seam, 152
dark sector
dissolved, **71**
honest assessment, 72
prediction ledger, 91
problem statement, 71
DARWIN, 88
Daya Bay experiment, 48
de Broglie, Louis, 120
de Sitter conjecture, 66
decoherence, 117, **122**
as VM shadow, 122
mass-dependent test, 127
decoherent-thermal matter
ω-sector, 76
reduced coupling, 72
v2.3 terminology introduction, 76
decompactification
regularity, 112

defect depletion
    mechanism, 82, **84**
defensible pillars
    three, 147
$\delta = 2/9$, 28
depth-dependent readout
    retracted v2.2, 86, 150
derivation
    vs. postulation, 134
derivation chain
    cosmological, 53
DESI
    comparison, 83
DESI BAO
    v3 target, 84
deterministic dynamism, 126
deuterium, **60**
    +2.3 sigma, 60
    anti-correlation, 60
    prediction, 150
    τ prediction, 59, 60
DFSZ axion, 48
differential prediction
    omega_m sigma_8 holonomy classes, 136
$\dim(T^2) = 2$
    unifying theme, 112
Dirac operator
    spectral triple, 101
discrete spacetime
    quantum foundations, 123
discrete spectrum
    vs. continuous, 134
dissolution vs. mechanism
    move (a), 118
    move (b), 119
    move (c), 121
    move (d), 122
    move (e), 125
    null predictions, 127
dissolution vs. solution, 133
distance ladder
    engagement missing, 86
Δm^2_{21}, 44
Δm^2_{32}, 44
double-projection problem, 88
DUNE
    cross-check, 44
    predictions, 44
dwarf spheroidals
    v3 target, 80
e-fold count, **55**
    derivation, 54, 55
E_1 fullness theorem, 146
E_1 physics
    complete, 143
early universe
    from one constant, 53
earned vs. postulated, **134**
echo amplitude
    bound, 97
echo detection
    stacking threshold, 97
echo time
    definition, 96
    ι_τ^{-2}, 96
effective neutrino number
    prediction, 149
EHT
    shadow correction prediction, 151
    shadow predictions, **93**, **95**
Einstein Telescope
    echo detection, 97
Einstein, Albert
    manifold assumption, 131
    unified field theory, 131
Eisenstein–Hu
    fitting formula dependence, 58
electroweak
    inventory, 144
electroweak baryogenesis, 49
electroweak precision, 23
    correlated agreement, 30
    τ derivation, 30
electroweak precision observables, **30**
electroweak theory
    predictions, 31
elliptical galaxies
    BTFR prediction, 76
emergent gravity, **100**
    Jacobson, 102
Empirical Preservation Principle, **15**
energy spectrum
    exponent derivation, 109
entanglement, 121
    as address sharing, 122
entropic gravity, 93, **100**
    partial truth, 102
    Verlinde, 102
entropy
    excess factor, 103
    T^2 excess factor, 104

environment-induced superselection, 122
epistemic framing, **5**
epistemic tier
  Tier 2 (tau-effective), 25, 28, 40
epistemic tier 0, 147
EPRL model, 137
equation of state
  dark energy, **83**
  w_0, 83
error cancellation
  structural, 56
η_B
  derivation, 61
  τ derivation, 62
$\bar{\eta}$ exponents
  post-hoc decomposition, 41
eternal inflation, 66
  multiverse, 66
Everett, Hugh, 119
EVPA
  winding number, 97, 98
experimental timeline, **154**
exponent 15
  structural derivation, 62
exponent decomposition
  baryogenesis, 62
extra Higgs
  not needed, 36

F1 forensic
  h_readout retraction sharpened, 151
F3 forensic
  exhaustive pair scan, 79
  projection-artifact diagnosis, 78
falsifiable seams, **152**
falsification
  conditions for τ, 143
  dark sector predictions, 91
  Ho prediction, 86
  tensor-to-scalar ratio, 59
  timeline, 154
  what does not count, 154
falsification pack, **143**, **147**
  BBN, 149
  black holes, 151
  CMB, 149
  collective dynamics, 152
  dark sector, 150
  inflation, 149
  particle physics, 148
Faraday rotation
  winding, 97
  winding theorem, 98
fiber dimension
  unifying role, 112
fiber dimensional suppression, **54**
  theorem, 54
fiber ratio
  4/3
    baryogenesis, 47
  EM correction, 47
fiber suppression
  1/3 factor, 61
  lithium, 60
  selectivity, 61
finiteness per genus, 134
  string amplitudes, 134
first acoustic peak, 56
  +0.28 percent, 56
  τ derivation, 56
first-moment holonomy invariant, 136
fitted parameters
  become derived, 115
five domains, 143
five hinges of τ, **131**, **133**
  completeness, 140
five-sixths factor
  derivation, 62
flat rotation curves
  master formula, 72
flavour physics
  orthodox landscape, **47**
flavour puzzle, **39**
  statement, 39
forensic red-team wave
  2026-04-20, xii
  F2 AdS/CFT forensic, 135
  F3 alpha-pair forensic, 78
foundation-up construction, 140
  principle, 141
fourth generation
  prediction, 148
Frauchiger–Renner, 127
free parameters
  continuous-to-discrete, **35**
  continuous-to-discrete reduction, 23
  earned vs postulated, 36
  flavour sector, 39
  Standard Model, 23
Friedmann pipeline
  τ inputs, 56
fsigma8 growth

v3 target, 84
Fuchs, Christopher, 120

g-2
correspondence, 18
galaxy formation
high redshift, 85
gauge–gravity duality, 135
gauge-coupling unification, 138
MSSM, 138
general relativity
as chart shadow, 87
comparison, 86
correct insights, 91
honest ledger, 91
structural errors, 91
triumphs, 86
generation count
pillar, 7
topological force, 7
generation number
topological derivation, 24
generation–lobe ratio, 40
Georgi–Glashow, 139
grand unified theory, 131
steelman, **139**
gravitational lensing
GR prediction, 87
gravitational redshift, 87
gravitational wave echoes
prediction, 151
gravitational waves
B-mode, 59
black hole echoes, **93**
detection, 87
echoes, **96**
inventory, 145
graviton propagator
LQG, 137
gravity
already quantum, 88
common origin with thermodynamics, 102
no renormalization needed, 102
readout theorem, 102
GRW theory, 126
GW150914
echo predictions, 96
echo search, 97

Ho ladders
tier-1 probes of H(z), 151
$H_\partial[\omega]$
two-tier reading, 127
H_partial[omega]
holonomy classes, 135
hard problems
dissolution, 134
Hawking radiation
as readout artifact, 103
readout interpretation, 103
Hawking temperature
readout, 103
Hawking, Stephen
topology theorem, 93
helium fraction, **60**
-0.43 sigma, 60
20/81, 60
derivation, 59
helium-3
-0.5 sigma, 60
τ prediction, 60
helium-4
prediction, 149
hemispherical asymmetry, 66
Hermetic Principle
inventory, 143
hidden freedom
v2.3 sector transition, 76
hidden variables, 120
Bell's theorem, 121
hierarchy problem, 138
dissolved, 16
technical naturalness, 138
Higgs boson
mass derivation, 25
Higgs mass, **25**
alternative framing, 26
epistemic frame, 25
inventory, 144
$n = 5$ earlier result, 26
$n = 7$ derivation, 26
refinement pathway, 26
τ derivation, 23
holographic principle
comparison, 67
hinge 3, 133
inflation, 66
holomorphic geometry, 100
holonomies
LQG vs. τ, 124
holonomy
tier-1 first-moment second-moment, 135

holonomy ratio
acceleration scales, **82**
acceleration theorem, 82
seam, 153
homology
first, of τ^3, 24
honest flag
Bullet Cluster, 81
CMB pipeline, 58
cosmology gap, 84
galaxy-scale data, 80
Ho tension, 85
S8 derivation, 84
SPARC RAR dwarfs UDGs, 80
SPARC-175 partial confirmation, 73
SPARC-175 verdict, 74
v2.3 cosmology consistency resolved in v2.4, 63
v2.3 new terminology, 76
v2.3 omega-B transition criterion, 76
v2.3 structural refinement, 75
v2.3 to v2.4 to v2.5 arc, 78
v2.3 verdict, 77
v2.3 weakened by v2.4, 77
v2.3 zero-point residual, 76
v2.4 BTFR reading, 78
v2.4 partial falsification, 77
v2.5 narrow tier-1 scope, 136
v2.5 Pair B confidence, 80
honest ledger
GR vs τ, 91
mixing and baryogenesis, 51
honest null predictions
Ch 66, **127**
honest posture, 9
measurement problem, 127
honesty calibration, 7
hot ICM
B-sector, 76
how to read the ledger, **8**
Hubble constant
from τ, 85
prediction, 150
Hubble parameter
structural formula, 85
Hubble tension, **85**
inventory, 144
status report, 85
summary, 71

IGMF, *see* intergalactic magnetic field
impossible early galaxies, 85
inflation
from τ, **54**
inventory, 144
model selection problem, 64
model zoo, 64
orthodox models, **64**
τ derivation, **53**
inflaton
absence, 67
information paradox
resolution, 103
information preservation
theorem, 103
integer-choice audit, *see* compute-then-axiomatize, **156**
summary, 161
interferometric tests
no distinguishing prediction, 127
intergalactic magnetic field, **99**
τ prediction, 100
intermittency
from fiber dimension, 108
gap, 107
ledger, 107
topological origin, 115
interpretations of QM, 117
inventory
complete E_1, 143
summary statistics, 146
iota-tau
QNM ratio, 93
ι_τ
as sole constant, 133
ι_τ posit, **6**
isospin breaking, 27
Item 1
SPARC-175 v2.2 first result, 9
SPARC-175 v2.3 structural refinement, 9
v2.3.1 cosmology-consistency open problem, 9
Item 2
CAMB/CLASS still pending, 9
Item 3
h readout CLOSED structural dead-end, 10
Item 4
Bullet Cluster sketch awaits v4, 10
Item 5
AdS/CFT obstruction memo, 10
IV.T20, 118
IV.T21, 118

IV.T22, 118

Jacobson, Ted, 100
  thermodynamic derivation, 102
jet
  magnetic field ratio, 99
jet helicity, **99**
JUNO
  cross-check, 44
  predictions, 44
JWST
  early galaxies, 71, **85**

Kaluza–Klein theory, 131
κ_ω
  capacity-gradient coupling, 72
  SPARC-175 best fit, 75
Kerr
  QNM comparison, 93
Kobayashi–Maskawa
  three generations, 48
Kobayashi–Maskawa mechanism, 47
Koide formula, 23, 28, **28**
  τ derivation, 28
Kolmogorov
  K41 theory, 107
  ledger, 107
  legacy, 113
Kolmogorov constant
  exact value, 109
  experimental difficulty, 109
  from dimension ratio, 109
  from fibration, 109
  prediction, 152
Kolmogorov exponent
  structural explanation, 109
Kolmogorov spectrum
  exponent, 109
  exponent derivation, 109
KSVZ axion, 48

Lamb shift, 31
Landau pole, **32**
landscape
  string theory, 134
landscape problem
  eliminated, 67
  inflation, 67
large volume scenario, 136
lattice QFT
  comparison, 34
ΛCDM
  empirical vs explanatory, 65
ΛCDM concordance model, 64, **64**
ledger
  scope of claims, 5
lemniscate
  regions, 24, 25
leptogenesis, 49
  not needed, 49
lepton masses, **28**
LHC
  SUSY exclusions, 138
LIGO
  echo detection, 96
LiteBIRD
  detection forecast, 59
lithium problem, **60**
  dissolved, 67
  history, 65
  prediction, 150
  resolution, 53, **59**
  seam, 153
lithium-7
  resolution, 61
  τ prediction, 60
log-Poisson model, 113
loop quantum cosmology, 137
  bounce, 137
loop quantum gravity
  and measurement problem, 123
  background independence, 123
  measurement problem in, 123
  steelman, **137**
low multipole anomalies, 66
Lyman-alpha forest
  v3 target, 84
LZ experiment, 71
  limits, 88

M87*, **95**
  shadow prediction, 95
macroscopic superposition
  not bounded by τ, 127
magnetic carpet, 114
magnetic field
  toroidal-to-poloidal ratio, 98
magnetic flux
  topological protection, 98
magnetic reconnection
  fast rate, 110
  ledger, 107

rate prediction, 152
magnetic winding
prediction, 152
magnetic winding numbers, **93**, **97**
Majorana neutrino
structural origin, 43
Majorana neutrinos
prediction, 148
manifold diagnosis, **132**
manifold ontology
as common error, 103
as root cause, 131
diagnosis, 131
entrenchment, 141
five pathologies, **132**
shared commitment, 139
Many Worlds interpretation, 119, **119**
mass hierarchy
prediction, 148
mass spectrum, **23**
inventory, 143
summary ledger, 37
mass splitting
proton–neutron, 27
master constant
all parameters from, 23
as posit, 6
cosmological parameters, 53
hinge 2, 133
no orthodox counterpart, 13
master formula
v2.3 source decomposition, 72
v2.3 structural refinement, 72
v2.5 update, 80
matter density
DE closure path, 57
holonomy path, 57
two paths, 57
matter-to-baryon ratio, 56
maximal mixing
near-maximal violation, 42
measure problem
eternal inflation, 66
measurement
address resolution, **121**
as address resolution, 120
measurement problem, **117**
dissolution, 118
dissolved, 16
from manifold, 132
orthodox statement, 117
trilemma, 117
Mercury perihelion
GR prediction, 87
metaphorical claim, 5
definition, 5
milestone years, 155
Milgrom, 89
Milgrom constant
from τ, 73
Millennium Problem
Navier–Stokes, 112
minimal SUSY
parameter-space exclusions, 138
mixing angles, **39**
not free parameters, 50
mod-3 periodicity, 46
moduli
absence, 67
moduli stabilization, 134
KKLT, 136
MOND
comparison, 86
comparison with τ, 82
structural status, **89**
muon mass
epistemic frame, 28
ratio to electron, 28, **28**
winding exponent, 29

N-predictions, 147
$n = 7$
post-hoc rationalizations, 26
$n = 7$ parameter, 25
nanoflare heating, 114
native_decide
scope, 7
natural inflation, 64
naturalness problem, 138
Navier–Stokes
decompactification, 112
ledger, 107
regularity
orthodox approaches, 115
turbulence problem, 107
Nelson–Barr mechanism, 48
neutrino
mass sum from τ, 43
mixing angles, 41
NNLO mass corrections, 43
neutrino hierarchy
formal proof, 44

theorem, 44
neutrino mass
from τ, **43**
sum prediction, 148
neutrino masses
individual, 44
mass table, 44
mass-squared splittings, 44
one-parameter structure, 44
neutrino oscillation
discovery, 48
neutron EDM
prediction, 148
neutron electric dipole moment, 45
neutron mass
as anchor, 35
justification, 35
NGC 1052-DF2, 80
NGC 1052-DF4, 80
NGC 3198
prediction, 73
NNLO
neutrino self-similar, **43**
self-similar corrections, **43**
NNLO corrections
$\delta = 1/25$, 29
lepton mass theorem, 29
lepton masses, 28
universality catalog, 29
No Knobs Theorem, 133
No-Joint-Minimum Theorem, 120
No-Shrink Theorem, 103, **104**
summary, 104
noncommutative geometry, 93, **100**
Connes, 101
partial truth, 101
nonlocality, 121
normal hierarchy
from winding order, 43
mandatory prediction, 50
proof, 43
normal ordering
from winding exponents, 44
structural proof, 44
nuclear cross sections
LUNA, 65
NuFIT, 48

observational score, **155**
observer
absence in τ, 124
observer problem
dissolution, 124
obstructions as features
F2 re-reading, 135
Ω_Λ
derivation, 83
from τ, 83
structural formula, 82
ontic layer, **17**
ontological layer, 13, **17**
ontological relabeling
move (a) ψ-demotion, 118
move (b) definite sector, 119
move (c) collapse rename, 121
move (d) entanglement rename, 122
move (e) observer dissolved, 125
orthodox physics
comparison, 13
correctness, 17

Pair A
1/(1+iota_tau) and iota_tau
alternative, 78
alternative to Pair B, 79
vs Pair B discrimination, 12
Pair B
1/(1+kappa_omega) and kappa_omega, 78
v2.5 canonical identification, 79
v2.5 identification, xii
PandaX, 71
limits, 88
parameter count
R1 audit, 161
partial falsification
v2.3 sector narrative, 78
Particle Data Group
CKM precision, 48
particle masses
from ι_τ, 23
particle physics
inventory, 143
path integral
as boundary-character sum, **34**
measure problem, 32
recovery from boundary sum, 34
path selection
ex post, 58
Peccei–Quinn mechanism, 48
Penrose twistors, 93, **100**
Penrose, Roger, 100
perturbation series

convergence in τ, 35
profinite convergence, 35
perturbation-size rule
Appendix 68.A, 157
Petschek reconnection
comparison, 110
imposed, 110
phantom crossing
forbidden, 83
prediction, 150
pilot wave, 120
plasmoid instability, 114
PMNS matrix, **41**
from τ, **39**
precision summary, 45
pointer states, 122
precision
summary table, 37
precision hierarchy, 146
precision inventory, 143
prediction asymmetry, 156
predictions
current status, 155
falsifiable, **143**
five domains, 20
organized by hinge, 140
preferred basis
no distinguishing test, 127
primordial black holes, 90
primordial nucleosynthesis, *see* BBN
τ framework, 59
primorial
convergence, 112
convergence rate, 112
profinite structure
hinge 4, 134
UV finiteness, 33
vs. continuum, 36
vs. spin networks, 123
progression operator
defect depletion, 84
projection artifact
v2.3 (1, kappa_omega) diagnosis, 78
proton decay
GUT predictions, 139
SO(10), 139
SU(5), 139
proton stability
prediction, 149
proton-neutron mass difference, 23, **27**
two-sector formula, 27
pulsar timing
binary, 87

QBism, 119, **120**
QCD
inventory, 144
QFT
empirical respect, 31
QNM
frequency ratio, **93**
theorem, **94**
secondary modes, 94
T^2 vs S^2 discrimination, 95
quantitative predictions
structural sources, 140
survey, 20
quantization
absent in τ, 124
quantization of gravity
failure, **88**
quantum field
as scaffolding, 37
quantum field theory
honest ledger, 35
landscape, **31**
ontological assumption, 32
structural failures, 32
triumphs, 31
quantum foundations, **117**
landscape, 126
quantum gravity
failure of programmes, 131
quantum randomness
readout artifact, 126
quark–lepton complementarity, 41
θ_{12}, 42
quasinormal modes, **93**
ratio prediction, 151
T^2 signatures, **93**
quintessence, 86
comparison, 89

R3 Hossenfelder
boundary-is-bulk charge upheld, 135
R4 McGaugh
semantic maneuver charge upheld, 86, 150
R4 McGaugh charge
SPARC-175 discharge, 73
radial acceleration relation
v3 target, 80
RAR, *see* radial acceleration relation

reactor angle
Daya Bay discovery, 48
readout
confused with ontology, 118
interpretation protocol, 17
ontological status, 17
vs. ontic, 13
readout functor
non-injectivity, 126
readout interpretation protocol, **17**
readout layer, **17**
readout quantization obstruction, 88, **88**
reconnection
diffusivity independence, 115
fiber dimension, 111
from fiber coupling, 110
inventory, 145
orthodox approaches, 113
orthodox models, 114
rate from τ, 110
`rem:ch67-tier1-holography`, 135
renormalization
as tower truncation, **33**
historical overview, 31
τ interpretation, **33**
renormalization group
ambiguity, 32
renormalization scheme
independence, 33
Reuter, Martin, 101
revision history
v2 steelman, 131
rotation curves
20-galaxy benchmark, 73
inventory, 145
master formula, 72
summary, **71**
without dark matter, **72**
Rovelli–Vidotto, 137
running coupling constants, 31

S_8 clustering
prediction, 151
S_8 tension
circularity caveat, 84
resolved, 84
S_8 tension
tier-1 holonomy-class reading, 136
SA-i
mod-5 baryogenesis, 46
mod hierarchy, 47
mod-3, 45
Sakharov conditions, **46**
structural fulfilment, 46
scalar amplitude, **55**
NLO, 55
τ derivation, 55
Schrödinger cat, 117
Schwarzschild
QNM comparison, 93
scope discipline
Part VII, 20
scope labels
conventions, 143
scope limits
honest statement, 8
seams
seven, 152
second-moment holonomy invariant, 136
Sector Exhaustion Theorem
recap, 82
summary, 71, 82
sensitivity audit
per-prediction, 156
Sgr A*, **95**
shadow prediction, 96
variability, **96**
shadow
T^2 correction, 95
T^2 quadrupole correction, 95
universal correction, 96
Shapiro time delay, 87
She–Lévêque formula
derivation, 108
fitted vs derived, 109
from fibration, 108
ledger, 107
original, 108
prediction, 152
successors, 113
σ-matrix
Majorana prediction, 43
Silk damping, **57**
+9 ppm, 57
precision, 57
prediction, 149
seam, 153
tau-native formula, 56
singularities
from manifold, 132
singularity
dissolved, 16

resolution in τ, 88
SKA
IGMF measurement, 100
slow-roll
156× gap, 56
excluded by τ, 67
not, 153
slow-roll inflation
excluded, 54
models, 64
smooth manifold, 131
SNO experiment, 48
SO(10)
GUT, 139
solar angle
from τ, 42
scope status, 42
θ_{12}, **42**
solar corona
temperature problem, 111
solar neutrino
SNO resolution, 48
Sorkin, Rafael, 123
SPARC sample
not yet tested, 80
SPARC-175
v2.2 test, 73
v2.3 structural closure, **75**
v2.4 robustness, **77**
v2.5 Pair B identification, **78**
spectral hierarchy
leptons, 29
spectral index
+13 ppm, 55
derivation, 54
not fitted, 55
prediction, 149
τ derivation, **55**
spectral triple, 101
from boundary algebra, 101
spin networks
and measurement, 123
spin-foam
EPRL model, 137
spin-foam dynamics, 137
Spite plateau, 60
history, 65
Standard Model
flavour sector, 39
free parameters, **23**
parameter count, 35
parameters, 23
structural assessment, 31
structural problem, 23
unexplained parameters, 39
star formation efficiency
enhanced, 85
Starobinsky inflation, 64
steelman
unification programmes, 131
stellar depletion
lithium, 65
sterile neutrino, 90
string landscape
inflation, 64, 66
string theory
dualities as echoes, 66
steelman, **134**
strong coupling constant, 30
strong CP
pillar, 7
SA-i resolution, 45
topological force, 7
strong CP problem
orthodox solutions, 48
resolution, **39**, **45**
seam, 153
structural artifact, **16**
structural integer choices, 5, **6**
structural pillars
three, **7**
structure functions
from fibration, 108
ledger, 107
structure-formation triangle, **84**
subjective probability, 120
Super-Kamiokande, 48
superposition
VM artifact, 119
supersymmetry
not needed, 36
prediction, 148
steelman, **138**
swampland conjectures, 137
swampland programme, 134
inflation, 66
Sweet–Parker reconnection
failure, 110
too slow, 110

τ divergences
early universe, **67**

τ framework
  divergences from orthodoxy, 49
  flavour sector, 39
τ lepton mass
  derivation, 28
  ratio to electron, 28
τ vs. GRW, Bohm, MWI
  predictive identity, 127
τ vs. orthodox physics, **143**
  collective dynamics, 115
τ-critique of GUTs, 139
τ-critique of LQG, 138
τ-critique of string theory, 137
τ-critique of SUSY, 138
τ-effective claim, 5
  definition, 5
τ-Einstein equation
  vs. Einstein, **87**
τ-native holography
  tier-1 addendum, 135
  v5 target, 11
tau-natural pair space
  8178 candidates, 79
τ-RG
  enrichment ladder proposal, 136
τ-Ryu-Takayanagi
  proposal, 136
τ-structural claim, 5
  definition, 5
τ-structural predictions, **147**
τ-Virasoro algebra
  proposal, 136
τ^3 fibration
  homology, 24
tensor-to-scalar ratio, **54**
  derivation, 53
  fiber derivation, 54
  geometric origin, 55
  prediction, 149
  seam, 153
testability
  pack, 143
TeVeS
  falsified, 89
θ_{QCD}
  naturalness, 48
  zero from τ, 45
they kept the manifold, **139**
three generations, **24**
  consistency, 25
  from topology, 23
  inventory, 143
  rank theorem, 24
  three proofs, 25
  topological definition, 24
three tiers of claim, 5, **5**
Tier 0 predictions, **147**
  epistemic status, 147
tier 1 global holonomy, 63
tier 2 local matter-phase coupling, 63
tier-1 holography
  v2.5 addendum, xii
topological censorship, 93
topologically forced predictions, **147**
toroidal flux
  conservation, 98
toroidal magnetic field, 97
tower truncation, 33
translation dictionary, 13
  motivation, 13
trinary sector structure
  BTFR, 75
  galaxy vs cluster, 76
  high-temperature limit, 81
  recombination-era resolution, 63
  v2.4 two-tier refinement, 78
Tully–Fisher relation, 72
turbulence
  200-year problem, 107
  empirical universality, 107
  inventory, 145
  ledger, **107**
  orthodox approaches, 113
  parameter-free theory, 113
turbulent reconnection, 114
twistor theory
  overview, 100
  partial truth, 100
  τ-embedding, 101
two-parameter fit
  v2.4 result, 77
two-sector formula, 27
two-thirds law
  dimension ratio, 110
  fiber interpretation, 110
  from fibration, 109
two-tier coherence
  derivation from V.T205, 63
  no sector-weighting rescue, 151
two-tier coherence criterion, **63**
  Ch 66 framing, **127**
  introduction, xi

ultra-diffuse galaxies
v3 target, 80
uncertainty bands
propagation, 8
uncertainty propagation, 8
unification
conceptual barriers, 141
failure of, **131**
historical programme, 131
inflation and baryogenesis, 62, 67
no personal blame, 141
roof down vs. foundation up, **140**
unification programmes
partial truths, 100
unitarity pentagon, 41
unitarity triangle, 40
Y-free fit
v2.4 result, 77
UV divergence, **32**
dissolved, 16
structural analysis, 31
UV divergences
from manifold, 132
UV shield, **33**
definition, 33
not regularization, 34

V.T205
Y_p = 20/81 as two-tier evidence, 63
Y_p as decisive clue, 63
V.T85
v2.5 Pair B form, 80
v2.2
integrated findings, 9
release note, xi
v2.3
N1-N30 impact, 161
release note, xi
v2.4
release note, xi
v4 wave status, 10
v2.4 framework advance
two-tier coherence, 127
v2.4 resolution
cosmology-consistency, **63**
v2.5
N21 falsifier tightened, 151
release note, xii
release note tier-1 holography, 135
v3 research program
status, **9**
v3 status
summary, 10
v3 target
Boltzmann code, 84
BTFR correction term, 74
galaxy-scale tests, 80
tau-native Boltzmann, 58
v3 targets
status tracking, 9
v4 deep-research wave
outcome, xi
status tracking, 10
V4.1 Upsilon-free, 77
V4.2 two-parameter, 77
v4 research targets
Round-5 panel list, **10**
v4 target
elliptical BTFR, 76
omega-B transition derivation, 76
tau-natural pair for joint fit, 78
V4.1 closed, 77
V4.10
CLOSED in v2.5 via Pair B, 11
closed via Pair B, 79
new in v2.4, 11
new open target, 78
resolved via Pair B, 78
V4.2 closed, 77
V4.5 two-tier coherence
tier-1 holography, 135
V4.9 target
two-tier formalization, 127
V5.1
new in v2.5, 11
V5.2
new in v2.5, 12
vacuum catastrophe, **32**, 71, 89
dissolved, 16
from manifold, 133
worked example, 18
Vafa, Cumrun, 137
verdict
summary, 156
Verlinde, Erik, 100
entropic force, 102
Vernstrom
filament detection, 100
VLBI
jet observations, 99
VM artifact
measurement problem, 118

VM representation
quantum state, 118
von Neumann, John, 117
vortex filament
as fiber collapse, 109

W boson mass, 30
$W_5(3)$, 47
$W_5(3) = 19$
double role, 62
$W_n(k)$ window sums, 6
Waring number
electroweak NLO, 30
$W_3(4) = 5$, 29
electroweak role, 30
$W_5(3) = 19$
baryon role, 61
e-fold role, 55
wave function
ontological status, 118
wave function collapse, 117
wave heating, 114
Weinberg angle, 30
Weinberg, Steven
asymptotic safety, 101
what this ledger is not, **8**
Wick rotation
chart operation, 34
Wigner's friend, 124
extended, 127
Wilson skeleton
magnetic flux, 99
WIMP
null results, **88**
searches, 86
SUSY candidate, 138
WIMP miracle
LSP, 138
winding classes
primitive, 24, 25
winding exponents, 28
Wolfenstein
A from τ, 40
eta-bar
epistemic frame, 40
$\bar{\eta}$, **40**
$\bar{\eta}$ from τ, 41
$\bar{\rho}$, **40**
$\bar{\rho}$ from τ, 40
Wolfenstein parametrization
A parameter, **40**
from τ, 39

XENON1T, 71
limits, 88

zero free parameters
caveats, 58
Zurek, Wojciech, 122