

# Integration

Paolo Bettelini

## Contents

<b>1</b>	<b>Indefinite Integrals</b>	<b>2</b>
1.1	Definition . . . . .	2
1.2	Properties . . . . .	2
1.3	Substitution Rule . . . . .	2
1.4	Integration By Parts . . . . .	3
<b>2</b>	<b>Definite Integrals</b>	<b>4</b>
2.1	Area Problem . . . . .	4
2.2	Definition . . . . .	4
2.3	Properties . . . . .	4
2.4	Fundamental Theorem of Calculus . . . . .	5
<b>3</b>	<b>Average Function Value</b>	<b>5</b>
<b>4</b>	<b>Mean Value Theorem</b>	<b>5</b>
<b>5</b>	<b>Area Between Functions</b>	<b>5</b>

# 1 Indefinite Integrals

## 1.1 Definition

Given a function  $f(x)$ , an **anti-derivative** or **primitive** is any function  $F(x)$  such that

$$\frac{dF}{dx} = f(x)$$

The operator to find a primitive function is called the **indefinite integral**

$$\int f(x) dx = F(x) + C, \quad C \in \mathbb{R}$$

The function to integrate (integrand) is delimited by the integral symbol  $\int$  and a differential of the variable of integration  $dx$ .

A function has infinitely many primitives, hence the  $+C$  term. This essentially means that the derivative of a function is the same when the function is shifted up or down, the rate of change is the same. By reversing the process we don't know the up or down shift of the original function.

$$f(x) = \int \frac{df}{dx} dx + C$$

for some specific  $C$ .

## 1.2 Properties

If  $k$  is a constant

$$\int k f(x) dx = k \int f(x) dx$$

$$\int f(x) \pm g(x) dx = \int f(x) dx \pm \int g(x) dx$$

## 1.3 Substitution Rule

Given an integral in the form

$$\int f(g(x))g'(x) dx$$

Let

$$u = g(x)$$

The differential of  $u$  is then

$$du = g'(x)dx$$

meaning that we can rewrite the integral as

$$\int f(u) du = F(u) + C = F(g(x)) + C$$

## 1.4 Integration By Parts

Starting from the product rule

$$\frac{d}{dx}(f(x)g(x)) = f'(x)g(x) + f(x)g'(x)$$

if we integrate both parts we get


$$\begin{aligned} f(x)g(x) + C &= \int f'(x)g(x) dx + \int f(x)g'(x) dx \\ \int f(x)g'(x) dx &= f(x)g(x) + C - \int f'(x)g(x) dx \end{aligned}$$

Since the indefinite integral of  $f'(x)g(x)$  is equal to some function plus an arbitrary constant, we can ignore the  $+C$  term.

$$\int f(x)g'(x) dx = f(x)g(x) - \int f'(x)g(x) dx$$

## 2 Definite Integrals

### 2.1 Area Problem



We want to find the signed area between  $f(x)$  and the  $x$ -axis between the interval  $[a; b]$ .

One way to do it would be by dividing the area into  $n$  rectangles, each of width

$$\Delta x = \frac{b-a}{n}$$

The height of each rectangle is given by  $f(x_k^*)$ . The area under the curve is approximately

$$A \approx \sum_{k=0}^{n-1} f(x_k^*) \Delta x$$

Notice that the position of  $x_k^*$  within the base of each rectangle controls the type of the approximation of the area under the curve. By moving  $x_k^*$  within the base we may achieve an approximation

by abundance or defect. The type of approximation does not matter when we let  $n \rightarrow \infty$ . As the amount of rectangles approaches infinity, the approximation approaches the exact value of the area.

$$A = \lim_{n \rightarrow \infty} \sum_{k=0}^n f(x_k^*) \Delta x$$

### 2.2 Definition

Given a function  $f(x)$  continuous on the interval  $[a; b]$ , we divide the interval into  $n$  rectangles of width  $\Delta x = \frac{b-a}{n}$  and height  $f(x_k^*)$ . The definitive interval of  $f(x)$  from  $a$  to  $b$  is

$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{k=0}^n f(x_k^*) \Delta x$$

### 2.3 Properties

$$\int_a^b f(x) dx = - \int_b^a f(x) dx$$

If  $k$  is constant

$$\int_a^b c f(x) dx = c \int_a^b f(x) dx$$

$$\int_a^b f(x) \pm g(x) dx = \int_a^b f(x) dx \pm \int_a^b g(x) dx$$

$$\int_a^a f(x) dx = 0$$

## 2.4 Fundamental Theorem of Calculus

A primitive function  $F(x)$  (with  $C = 0$ , so passing through the origin) represents the area from 0 to  $x$  of  $f(x)$ .

Let  $f(x)$  be continuous on the interval  $I = [a; b]$  and  $F(x)$  any primitive of  $f(x)$ , then  $F(x)$  is also continuous on  $I$  and

$$\int_0^x f(x) dx = F(x)$$

and

$$\int_a^b f(x) dx = F(b) - F(a)$$

This also means that

$$f(x) = \int_0^x f'(t) dt$$

or

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

When the upper or lower limit is not constant,

$$\begin{aligned} \frac{d}{dx} \int_{v(x)}^{u(x)} f(t) dt &= \frac{d}{dx} [F(u(x)) - F(v(x))] \\ &= u'(x)f(u(x)) - v'(x)f(v(x)) \end{aligned}$$

## 3 Average Function Value

The average value of a continuous function  $f(x)$  over the interval  $[a; b]$  is given by


$$f_{\text{avg}} = \frac{1}{b-a} \int_a^b f(x) dx$$

## 4 Mean Value Theorem

If  $f(x)$  is a continuous function on  $I = [a; b]$ ,  $f(x)$  will at some point in  $I$  reach its average value, meaning

$$\exists c \mid f(c) = \frac{1}{b-a} \int_a^b f(x) dx$$

## 5 Area Between Functions



Given a function  $y = f(x)$  and  $y = g(x)$ , the area enclosed by the two functions in the interval  $I = [a; b]$  is given by


$$A = \int_a^b f(x) - g(x) dx$$

assuming that  $f(x) \geq g(x)$  when  $x \in I$ .

Note that  $A \geq 0$ .

If  $f(x) < g(x)$  for some  $x \in I$  this formula won't work. However, it is still possible to split the integral into multiple integrals at every point where  $f(x) - g(x)$  changes sign. To remove the sign constraint we could say

$$A = \int_a^b |f(x) - g(x)| dx$$



The same thing applies when we have functions in the form  $x = f(y)$  and  $x = g(y)$  and we want to find the areas enclosed by the functions when  $y \in [d; c]$

$$A = \int_d^c |f(y) - g(y)| dy$$