

Limits

Paolo Bettelini

Contents

1	Definition	2
2	Properties	2
3	Continuity	2
4	Intermediate value theorem	2
5	Bolzano's Theorem	3
6	Squeeze Theorem	3

1 Definition

A limit is usually used to describe the behavior of a function as its argument approaches a given value.

The limit towards a certain value c within a function can be approached both from the right and from the left.

The limit in a general sense exists if the value approached from both sides is the same and well-defined.

We define the limit of x approaching c from the left within the function $f(x)$ as

$$\lim_{x \rightarrow c^-} f(x)$$

We define the limit of x approaching c from the right within function $f(x)$ as

$$\lim_{x \rightarrow c^+} f(x)$$

We define the limit of x approaching c within function $f(x)$ as

$$\lim_{x \rightarrow c} f(x)$$

Formally, given a function $f : D \rightarrow \mathbb{R}$ the limit $L = \lim_{x \rightarrow c} f(x)$ exists if given an arbitrary small $\epsilon > 0$ there is another number $\delta > 0$ such that

$$|f(x) - L| < \epsilon, \quad \forall x \in D \text{ where } 0 < |x - c| < \delta$$

2 Properties

If the limit exists

$$\lim_{x \rightarrow c} f(g(x)) = f(\lim_{x \rightarrow c} g(x))$$

3 Continuity

A function f is continuous at a point c iff

$$\lim_{c_0 \rightarrow c^+} f(c_0) = \lim_{c_0 \rightarrow c^-} f(c_0) = f(c)$$

A function f is continuous on an interval $[a; b]$ iff it is continuous at each point $c \in [a; b]$

$$\forall c \in [a; b], \lim_{c_0 \rightarrow c^+} f(c_0) = \lim_{c_0 \rightarrow c^-} f(c_0) = f(c)$$

4 Intermediate value theorem

A function f continuous on an interval $[a; b]$ will take every value in the interval $[f(a); f(b)]$.

5 Bolzano's Theorem

If $f(x)$ is continuous on $[a; b]$ and $f(a) < f(b)$ then there is a root.

$$f(a) < f(b) \implies \exists c \in [a; b] \mid f(c) = 0$$

6 Squeeze Theorem

Let $h(x)$, $f(x)$ and $g(x)$ be three functions defined

TODO