

Taylor Series

Paolo Bettelini

Contents

1	Definition	2
2	Divergence	3

1 Definition

A function $f(x)$ can be approximated around a point a by a power series.

The polynomial must be centered around a , so the variable will be $x - a$ such as

$$\sum_{n=0}^{\infty} c_n (x - a)^n = c_0 + c_1(x - a) + c_2(x - a)^2 + c_3(x - a)^3 + \dots$$

Our goal is to find the coefficients c_n .

We notice that $f(a) = c_0$, which is the only coefficient that does not multiply a variable.

If we take the derivative, the coefficient c_1 will lose its variable

$$c_1 + 2c_2(x - a) + 3c_3(x - a)^2 + \dots$$

Now we have $f'(a) = c_1$.

We take the derivative of the polynomial again

$$2c_2 + 6c_3(x - a) + \dots$$

This time we have

$$\begin{aligned} f''(a) &= 2c_2 \\ c_2 &= \frac{f''(a)}{2} \end{aligned}$$

And again

$$\begin{aligned} f'''(a) &= 6c_3 \\ c_3 &= \frac{f'''(a)}{6} \end{aligned}$$

More generally, to extract the coefficient c_n we take the n -th derivative of the function.

By the power rule, we know that

$$\frac{d^k}{dx^k} (x^k) = k!$$

For example

$$\begin{aligned} \frac{d^4}{dx^4} (x^4) &= \frac{d^3}{dx^3} (4x^3) \\ &= \frac{d^2}{dx^2} (4 \cdot 3x^2) \\ &= \frac{d}{dx} (4 \cdot 3 \cdot 2x) \\ &= 4 \cdot 3 \cdot 2 \cdot 1 \\ &= 4! \end{aligned}$$

This brings us to

$$c_n = \frac{f^{(n)}(a)}{n!}$$

The Taylor series of $f(x)$ around the point a is defined as

$$\sum_{n=0}^{\infty} \frac{(x - a)^n f^{(n)}(a)}{n!}$$

2 Divergence