

## Sums of some infinite series

SGN-1156 Signal Processing Techniques

February 3, 2013

**Geometric series.** If  $a \in \mathfrak{R}$  and  $r \in \mathfrak{R}$ , a geometric series  $\{x_n\} = \{a, ar, ar^2, \dots, ar^n, \dots\}$  converges if and only if  $|r| < 1$  and then its sum is:

$$\sum_{n=0}^{\infty} x_n = \sum_{n=0}^{\infty} ar^n = \lim_{n \rightarrow \infty} a \frac{1 - r^{n+1}}{1 - r} = \frac{a}{1 - r} \quad (1)$$

**Telescopic series.** Given a series  $\{a_n\}$  we say that the series  $\{x_n\} = \{a_n - a_{n+1}\}$  is a telescopic series. Then,  $\{x_n\}$  is convergent if and only if  $\lim_{n \rightarrow \infty} a_n$  is finite. In that case, the sum of  $\{x_n\}$  is:

$$\sum_{n=1}^{\infty} x_n = \sum_{n=1}^{\infty} (a_n - a_{n+1}) = a_1 - \lim_{n \rightarrow \infty} a_n \quad (2)$$

**Arithmetic-geometric series.** Given  $a, b, r \in \mathfrak{R}$  we say that the series  $\{x_n\}$  is arithmetic-geometric if  $\{x_n\} = \{a, (a+b)r, (a+2b)r^2, \dots, (a+nb)r^n, \dots\}$ . Such series are convergent if and only if  $|r| < 1$  and, in that case, their sum is:

$$\sum_{n=0}^{\infty} x_n = \sum_{n=0}^{\infty} (a + nb)r^n = \frac{a(1 - r) + br}{(1 - r)^2} \quad (3)$$

**Hypergeometric series.** We say that a series  $\{x_n\}$  is hypergeometric if:

$$\frac{x_{n+1}}{x_n} = \frac{an + b}{an + c} \quad (4)$$

where  $a, b, c \in \mathfrak{R}$  and  $a \neq 0$ . This type of series converges if and only if  $\frac{(c-b)}{a} > 1$  and, in that case, its sum is:

$$\sum_{n=1}^{\infty} x_n = \frac{-x_1 c}{a + b - c} \quad (5)$$