

filtered_IDs.csv

List of precursors filtered at 1% FDR. Each row represents a single precursor (sequence, charge)

coeff	Coefficient estimated for the precursor
spec_id	MS2 spectrum ID where the precursor was found
Ms1_spec_id	Closest MS1 spectrum to the MS2 spectrum
seq	Precursor sequence
z	Precursor charge
window_mz	Center of the MS2 isolation window
rt	Retention time of the MS2 spectrum
num_lib	Number of library fragments matched
frac_lib_int	Fraction of the library intensity matched
frac_dia_int	Fraction of the observed intensity matched to the precursor
mz_error	Relative MS1 m/z error
rt_error	Retention time error (empirical RT - library RT)
frac_int_matched	Fraction of the total spectrum intensity with any match (any precursor)
frac_int_pred	Fraction of the total spectrum intensity predicted by all precursors
spec_r2	Pearson correlation for the predictions and observed spectrum
prec_r2	Pearson correlation for the predicted precursor and matched spectrum peaks
prec_r2_uniq	Pearson correlation for the predicted precursor and uniquely matched peaks
frac_int_uniq	Fraction of library intensity uniquely matched
frac_int_uniq_pred	Fraction of total spectrum intensity uniquely predicted
hyperscore	Sum of matched intensities times factorial(#b ions) times factorial(#y ions)
b_counts	Number of b ions matched
y_counts	Number of y ions matched

longest_y_ions	Index of largest y ion matched
scribe_scores	Scribe score as described here: https://pubs.acs.org/doi/abs/10.1021/acs.jproteome.2c00672
max_unmatched_residuals	Maximum residual for unmatched peaks, normalized and log-transformed.
max_matched_residuals	Maximum residual for matched peaks, normalized and log-transformed.
gof_stats	Goodness-of-fit score for each precursor (log2 of residuals/fitted).
manhattan_distances	Manhattan distances between precursor peaks and observed peaks. (normalized and log-transformed)
fitted_spectral_contrasts	Spectral contrast angles for each precursor
frac_int_matched_pred	Fraction of the matched spectrum intensity predicted by all precursors
frac_int_matched_pred_sigcoeff	Fraction of spectrum intensity predicted by precursors with coeff > 1
cosine	Similarity between predicted spectra and observed intensities
mz	Precursor m/z
frag_names	Names of fragments matched
frag_errors	m/z errors of fragments matched
frag_mz	Theoretical m/z of fragments matched
frag_int	Library intensity of fragments matched
obs_int	Observed intensity of fragments matched
unique_frag_mz	Fragments that uniquely matched a peak
unique_obs_int	Intensities for fragments that uniquely matched a peak
file_name	Name of the file searched
protein	Protein associated with the precursor
n_scans	Number of scans where that precursor was identified.
debug_window_value	
manhattan_distances_nearby_max	Maximum manhattan_distances of nearby scans
max_matched_residuals_nearby	Minimum max_matched_residuals of nearby scans

_min	
gof_stats_nearby_min	Minimum gof_stats of nearby scans
scribe_scores_nearby_min	Minimum scribe_scores of nearby scans
smoothness	The smoothness of coeff for precursor for that precursor
stripped_seq	Amino acid sequence without tags or mods.
pep_len	Peptide sequence length
sq_rt_error	Square root of the rt_error
sq_mz_error	Square root of the mz_error
untag_seq	Peptide sequence without the tag (contains mods)
untag_prec	untag_seq and charge combined
channels_matched	Number of channels from the plex set matched
channel	Channel name of the precursor
med_frag_error	Median absolute mz error of the matched fragments
decoy	If the precursor is a decoy
PredVal	Raw score from the target-decoy classifier (1 is best)
Qvalue	Q-value of the precursor
BestChannel_Qvalue	Best q-value of the precursor set
plexfitMS1	Fitted MS1 coefficient at max
plexfitMS1_p	Correlation of theoretical MS1 isotope distribution with observed peaks
plexfittrace	Fitted MS1 coefficients of elution trace
plexfit_ps	Correlation of theoretical MS1 isotope distribution with observed peaks for each fit in elution trace
plexfittrace_spec_all	All MS1 spectra used for MS1 fitting
plexfittrace_all	All fitted MS1 coefficients
plexfittrace_ps_all	Correlation of theoretical MS1 isotope distribution with observed peaks for each fit
plex_Area	Area under fitted MS1 coefficients
ms1_cor	Correlation between MS2 coefficients and monoisotopic trace

traceproduct	Product of MS1xMS1 correlations
iso_cor	Correlation between theoretical and observed MS1 isotopes
MS1_Int	Observed intensity of MS1 monoisotopic peak at max
all_ms1_specs	MS1 spectra searched for monoisotopic peak
MS1_Area	Area under elution trace of MS1 monoisotopic trace
all_ms1_iso0vals	Observed MS1 intensities for the monoisotopic peak
all_ms1_iso1vals	Observed MS1 intensities for the first isotopic peak
all_ms1_iso2vals	Observed MS1 intensities for the second isotopic peak
all_ms1_iso3vals	Observed MS1 intensities for the third isotopic peak
all_ms1_iso4vals	Observed MS1 intensities for the fourth isotopic peak
all_ms1_iso5vals	Observed MS1 intensities for the fifth isotopic peak
last_aa	Last amino acid in the peptide sequence
seq_len	Length of the amino acid sequence
run_chan	Sample run and channel concatenated
Protein_Qvalue	Protein q-value