Statistics
Parameter Estimation, Hypothesis Testing, Regression

D Manjunath

CMInDS, EE
IIT Bombay

October 21, 2026

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

Lecture 5

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

Parameter Estimation

- Many times, we have a model for the CDF of pdf of population feature of interest.

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

Parameter Estimation

- Many times, we have a model for the CDF of pdf of population feature of interest.
- The model CDF will have parameters.

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

Parameter Estimation

- Many times, we have a model for the CDF of pdf of population feature of interest.
- The model CDF will have parameters.
- We will now discuss how to determine the **best estimate** of the paramter(s).

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

Parameter Estimation

- Many times, we have a model for the CDF of pdf of population feature of interest.
- The model CDF will have parameters.
- We will now discuss how to determine the **best estimate** of the paramter(s).
- As before we will obtain n independent sample realizations of the parameter of interest.

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

What is being estimated?



This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

What is being estimated?



And what is the method?

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

Likelihood Function

- $x_1, \ldots x_K$ is the set of measurements.

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

Likelihood Function

- $x_1, \ldots x_K$ is the set of measurements.
- Assume that the parameter set is θ.

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

Likelihood Function

- $x_1, \ldots x_K$ is the set of measurements.
- Assume that the parameter set is θ.
- Question: What is the *likelihood* of observing of $x_1, \ldots x_K$ if the parameter value was θ?

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

Likelihood Function

- $x_1, \ldots x_K$ is the set of measurements.
- Assume that the parameter set is θ.
- Question: What is the *likelihood* of observing of $x_1, \ldots x_K$ if the parameter value was θ?
  - Assume the population is Bernoulli with parameter p.

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

Likelihood Function

- $x_1, \ldots x_K$ is the set of measurements.
- Assume that the parameter set is θ.
- Question: What is the *likelihood* of observing of $x_1, \ldots x_K$ if the parameter value was θ?
  - Assume the population is Bernoulli with parameter p.
  - Assume the population is Poisson with parameter λ.

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

Likelihood Function

- $x_1, \ldots x_K$ is the set of measurements.
- Assume that the parameter set is θ.
- Question: What is the *likelihood* of observing of $x_1, \ldots x_K$ if the parameter value was θ?
  - Assume the population is Bernoulli with parameter p.
  - Assume the population is Poisson with parameter λ.
  - Assume the population is uniform in the interval $(0, a)$.

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

Likelihood Function

- $x_1, \ldots x_K$ is the set of measurements.
- Assume that the parameter set is θ.
- Question: What is the *likelihood* of observing of $x_1, \ldots x_K$ if the parameter value was θ?
  - Assume the population is Bernoulli with parameter p.
  - Assume the population is Poisson with parameter λ.
  - Assume the population is uniform in the interval $(0, a)$.
  - Assume the population is Gaussian with unit variance and mean μ.

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

Likelihood Function

- $x_1, \ldots x_K$ is the set of measurements.
- Assume that the parameter set is θ.
- Question: What is the *likelihood* of observing of $x_1, \ldots x_K$ if the parameter value was θ?
  - Assume the population is Bernoulli with parameter p.
  - Assume the population is Poisson with parameter λ.
  - Assume the population is uniform in the interval $(0, a)$.
  - Assume the population is Gaussian with unit variance and mean μ.
  - Assume the population is exponential with parameter μ.

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

Maximum Likelihood Estimate

- Find the θ that maximises the likelihood of the observations.

$$\hat{\theta} = \arg\max_{\theta} L(x_1, \ldots, x_K|\theta)$$

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

Maximum Likelihood Estimate

- Find the θ that maximises the likelihood of the observations.

$$\hat{\theta} = \arg\max_{\theta} L(x_1, \ldots, x_K|\theta)$$

- Can also maximise the Log Likelihood function.

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

Maximum Likelihood Estimate

- Find the θ that maximises the likelihood of the observations.

$$\hat{\theta} = \arg\max_{\theta} L(x_1, \ldots, x_K|\theta)$$

- Can also maximise the Log Likelihood function.
- Likelihood and log likelihood functions and MLEs

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

Maximum Likelihood Estimate

- Find the θ that maximises the likelihood of the observations.

$$\hat{\theta} = \arg\max_{\theta} L(x_1, \ldots, x_K|\theta)$$

- Can also maximise the Log Likelihood function.
- Likelihood and log likelihood functions and MLEs
- Bernouilli

$$\begin{aligned} L(\underline{x}; p) &= \prod_{i=1}^{K} p^{x_i}(1-p)^{1-x_i} = p^n(1-p)^{K-n} \\ \hat{p} &= \frac{\sum_{i=1}^{K} x_i}{K} \end{aligned}$$

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

Maximum Likelihood Estimate

- Poisson

$$L(\underline{x}; \lambda) \quad =$$

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

Maximum Likelihood Estimate

- Poisson

$$L(\underline{x}; \lambda) \quad =$$

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

Maximum Likelihood Estimate

- Poisson

$$L(\underline{x};\lambda) \quad = \quad \prod_{i=1}^{K} e^{-\lambda} \frac{\lambda x_i}{x_i!} \quad =$$

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

Maximum Likelihood Estimate

- Poisson

$$L(\underline{x};\lambda) \quad = \quad \prod_{i=1}^{K} e^{-\lambda}\frac{\lambda x_i}{x_i!} \quad = \quad e^{-K\lambda}\frac{\lambda^{\sum_{i=1}^{K} x_i}}{\prod_{i=1}^{K} x_i}$$

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

Maximum Likelihood Estimate

- Poisson

$$
\begin{aligned}
L(\underline{x};\lambda) &= \prod_{i=1}^{K} e^{-\lambda}\frac{\lambda x_i}{x_i!} = e^{-K\lambda}\frac{\lambda^{\sum_{i=1}^{K} x_i}}{\prod_{i=1}^{K} x_i} \\
\log L(\underline{x};\lambda) &= -K\lambda + \sum_{i=1}^{K} x_i \log\lambda - \log(\prod_{i=1}^{K} x_i)
\end{aligned}
$$

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

Maximum Likelihood Estimate

- Poisson

$$
\begin{aligned}
L(\underline{x};\lambda) &= \prod_{i=1}^{K} e^{-\lambda}\frac{\lambda x_i}{x_i!} = e^{-K\lambda}\frac{\lambda^{\sum_{i=1}^{K} x_i}}{\prod_{i=1}^{K} x_i} \\
\log L(\underline{x};\lambda) &= -K\lambda + \sum_{i=1}^{K} x_i \log\lambda - \log(\prod_{i=1}^{K} x_i) \\
\hat{\lambda} &= \frac{\sum_{i=1}^{K} x_i}{K}
\end{aligned}
$$

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

Maximum Likelihood Estimate

- Gaussian with known σ and unknown μ.

$$L(\underline{x};\lambda) = \prod_{i=1}^{K} \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x_i-\mu)^2}{2\sigma^2}} = \frac{1}{(\sqrt{2\pi}\sigma)^K} e^{-\sum_{i=1}^{K}\frac{(x_i-\mu)^2}{2\sigma^2}}$$

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

Maximum Likelihood Estimate

- Gaussian with known σ and unknown μ.

$$L(\underline{x};\lambda) = \prod_{i=1}^{K} \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x_i-\mu)^2}{2\sigma^2}} = \frac{1}{(\sqrt{2\pi}\sigma)^K} e^{-\sum_{i=1}^{K}\frac{(x_i-\mu)^2}{2\sigma^2}}$$

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

Maximum Likelihood Estimate

- Gaussian with known σ and unknown μ.

$$L(\underline{x};\lambda) = \prod_{i=1}^{K} \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x_i-\mu)^2}{2\sigma^2}} = \frac{1}{(\sqrt{2\pi}\sigma)^K} e^{-\sum_{i=1}^{K}\frac{(x_i-\mu)^2}{2\sigma^2}}$$

$$\log L(\underline{x};\lambda) = -K\log(\sqrt{2\pi}\sigma) - \sum_{i=1}^{K}\frac{(x_i-\mu)^2}{2\sigma^2}$$

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

Maximum Likelihood Estimate

- Gaussian with known σ and unknown μ.

$$
\begin{aligned}
L(\underline{x};\lambda) &= \prod_{i=1}^{K} \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x_i-\mu)^2}{2\sigma^2}} = \frac{1}{(\sqrt{2\pi}\sigma)^K} e^{-\sum_{i=1}^{K}\frac{(x_i-\mu)^2}{2\sigma^2}} \\
\log L(\underline{x};\lambda) &= -K\log(\sqrt{2\pi}\sigma) - \sum_{i=1}^{K}\frac{(x_i-\mu)^2}{2\sigma^2} \\
\hat{\mu} &= \frac{\sum_{i=1}^{K} x_i}{K}
\end{aligned}
$$

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

Maximum Likelihood Estimate

- MLE for Uniform in $(0, a)$

$$L(\underline{x}; \lambda) = \frac{1}{a^K}$$

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

Maximum Likelihood Estimate

- MLE for Uniform in $(0, a)$

$$L(\underline{x}; \lambda) = \frac{1}{a^K}$$

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

Maximum Likelihood Estimate

- MLE for Uniform in $(0, a)$

$$\begin{aligned} L(\underline{x}; \lambda) &= \frac{1}{a^K} \\ \hat{a} &= \max\{x_1, x_2, \ldots, x_K\} \end{aligned}$$

- Expectation of Estimate is the **bias** and variance is the mean squared error.

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

More on estimation methods

- Best guess of a random variable before measurement

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

More on estimation methods

- Best guess of a random variable before measurement
- What do I mean by best?

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

More on estimation methods

- Best guess of a random variable before measurement
- What do I mean by best?

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

More on estimation methods

- Best guess of a random variable before measurement
- What do I mean by best? Two possibilities: Most likely value and the one that makes least error.

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

More on estimation methods

- Best guess of a random variable before measurement
- What do I mean by best? Two possibilities: Most likely value and the one that makes least error. latter is further characterized as the one that minimizes mean squared error.
- 0-1 Bernoulli,

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

More on estimation methods

- Best guess of a random variable before measurement
- What do I mean by best? Two possibilities: Most likely value and the one that makes least error. latter is further characterized as the one that minimizes mean squared error.
- 0-1 Bernoulli,

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

More on estimation methods

- Best guess of a random variable before measurement
- What do I mean by best? Two possibilities: Most likely value and the one that makes least error. latter is further characterized as the one that minimizes mean squared error.
- 0-1 Bernoulli, Loaded die with $p_1 = 0.2$ and all other values having probability 0.16,

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

More on estimation methods

- Best guess of a random variable before measurement
- What do I mean by best? Two possibilities: Most likely value and the one that makes least error. latter is further characterized as the one that minimizes mean squared error.
- 0-1 Bernoulli, Loaded die with $p_1 = 0.2$ and all other values having probability 0.16, Exponential with mean 1,

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

More on estimation methods

- Best guess of a random variable before measurement
- What do I mean by best? Two possibilities: Most likely value and the one that makes least error. latter is further characterized as the one that minimizes mean squared error.
- 0-1 Bernoulli, Loaded die with $p_1 = 0.2$ and all other values having probability 0.16, Exponential with mean 1, Gaussian with mean 2 and variance 1.

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

More on estimation methods

- Best guess of a random variable before measurement
- What do I mean by best? Two possibilities: Most likely value and the one that makes least error. latter is further characterized as the one that minimizes mean squared error.
- 0-1 Bernoulli, Loaded die with $p_1 = 0.2$ and all other values having probability 0.16, Exponential with mean 1, Gaussian with mean 2 and variance 1.
- Typically, we have a belief about the parameter, expressed as a CDF (or PDF) for the parameter, say $F(\theta)$.

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

More on estimation methods

- Best guess of a random variable before measurement
- What do I mean by best? Two possibilities: Most likely value and the one that makes least error. latter is further characterized as the one that minimizes mean squared error.
- 0-1 Bernoulli, Loaded die with $p_1 = 0.2$ and all other values having probability 0.16, Exponential with mean 1, Gaussian with mean 2 and variance 1.
- Typically, we have a belief about the parameter, expressed as a CDF (or PDF) for the parameter, say $F(\theta)$.
- After measurement, you have additional information. You can now change your belief based on the measurement using Baye's theorem.

$$F(\theta \mid x_1, x_2, \ldots, x_n) = \frac{F(x_1, x_2, \ldots, x_n \mid \theta) F(\theta)}{\int_\theta F(x_1, x_2, \ldots, x_n \mid \theta) F(\theta)\, d\theta}$$

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

More on estimation methods

- Best guess of a random variable before measurement
- What do I mean by best? Two possibilities: Most likely value and the one that makes least error. latter is further characterized as the one that minimizes mean squared error.
- 0-1 Bernoulli, Loaded die with $p_1 = 0.2$ and all other values having probability 0.16, Exponential with mean 1, Gaussian with mean 2 and variance 1.
- Typically, we have a belief about the parameter, expressed as a CDF (or PDF) for the parameter, say $F(\theta)$.
- After measurement, you have additional information. You can now change your belief based on the measurement using Baye's theorem.

$$F(\theta \mid x_1, x_2, \ldots, x_n) = \frac{F(x_1, x_2, \ldots, x_n \mid \theta) F(\theta)}{\int_\theta F(x_1, x_2, \ldots, x_n \mid \theta) F(\theta)\, d\theta}$$

- THis is called the Maximum a Posteriori (MAP) estimate

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

MAP Estimate

Bernoulli with priors $\mathsf{Prob}(p = 0.3) = 0.5$ $\mathsf{Prob}(p = 0.5) = 0.2$ $\mathsf{Prob}(p = 0.7) = 0.3$. Determine the expression for the posterior based on five samples.

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.

MAP Estimate

Bernoulli with priors $\mathsf{Prob}(p = 0.3) = 0.5$ $\mathsf{Prob}(p = 0.5) = 0.2$ $\mathsf{Prob}(p = 0.7) = 0.3$. Determine the expression for the posterior based on five samples.

This file is meant for personal use by parikshithrnayak@gmail.com only.
Sharing or publishing the contents in part or full is liable for legal action.