

OpenDyslexic

# F61 - Nuclear Magnetic Resonance

Paris J. Huth - Q'inich Figueroa

May 2024

## 1 Introduction

In Protokoll we will examine the usage nuclear magnetic resonance to identify probes and reveal the structure of objects.

## 2 Basics

Any nuclei with an existent spin  $S$  has a magnetic dipole moment:

$$\vec{\mu} = \hbar\gamma\vec{S}$$

where  $\gamma$  is the gyromagnetic ration.

This magnetic dipole  $\vec{\mu}$  interacts with an external magnetic field  $\vec{B}_0$  and is associated with an interaction energy  $\Delta E$ :

$$\Delta E = -\vec{\mu} \cdot \vec{B}_0$$

This interaction yields both a parallel  $\mu_+$  and antiparallel  $\mu_-$  orientation of the protons magnetic dipole in the external field. For a macroscopic sample of  $N$  protons, the number of occupied states  $N_+$  and  $N_-$ , the sum of which comprises  $N$ , can be approximated by a Boltzmann distribution:

$$N_{\pm} = N_0 e^{-\frac{E_0 \pm \Delta E}{kT}}$$

with a normalization factor  $N_0$ .

## 3 Measurements

## 4 Analysis

## 5 Critical Discussion