Food Customers: Exploratory Data Analysis

MAS531 Group Project "Fo"

Fanourios Loukas Orthodoxou
fortho01@ucy.ac.cy

Olga Kappai
okappa01@ucy.ac.cy

Davide Vicentini
dvicen01@ucy.ac.cy

Panagiotis Parpas
pparpa01@ucy.ac.cy

MSc Data Science, University of Cyprus

November 2026

Abstract

We analyse the marketing data of a food company (2,205 customers, 27 variables) to describe its customers and the relationships between their characteristics and their purchasing behaviour. After removing 203 duplicate rows and 5 inconsistent records, 1,997 customers remain. None of the numerical variables is normally distributed, so non-parametric methods are used throughout. Income is by far the strongest driver of spending (Spearman $\rho = 0.87$). Customers with children spend much less (median 182 vs 1,117) and rely far more on discounts, and this effect remains at the same income level. Education and marital status have little or no effect once income is taken into account. A k-means clustering identifies three customer segments: *budget families*, *affluent big spenders* and *established families*.

Contents

1 Introduction **3**

2 Data **3**

3 Data Cleaning and Preparation **4**
3.1 Missing values and duplicates 4
3.2 Logic checks 4
3.3 New variables 4

4 Univariate Analysis **5**
4.1 Numerical variables 5
4.2 Categorical variables 6
4.3 Normality 6

5 Bivariate Analysis **7**
5.1 Correlations 7
5.2 Income and spending 8
5.3 Children, education and marital status 9
5.4 Purchase channels 9
5.5 Discounts 10
5.6 Website visits 10
5.7 Recency and length of relationship 11

6 Hypothesis Testing **11**

7 Customer Segments and Regression **13**
7.1 Principal Component Analysis 13
7.2 Clustering with k-means 13
7.3 Regression: what explains spending? 16

8 Conclusions and Recommendations **17**
8.1 Main findings 17
8.2 Recommendations for the company 17

9 Limitations **17**

A Additional Figures and Tables **18**
A.1 Histograms of all numerical variables 18
A.2 Categorical variables 19
A.3 Outliers 19
A.4 Pairwise comparisons 20

B R Code **20**

1 Introduction

The company wants a full exploratory analysis of its customer data in order to understand the profiles of its customers and how the variables vary together. Our aims are to:

1. clean the data and check it for errors and inconsistencies;
2. describe each variable (summary statistics, plots, distribution shape);
3. study the relationships between variables, especially with total spending;
4. test the most important patterns formally with hypothesis tests;
5. group customers into segments and measure the effect of each characteristic on spending.

All analysis was done in R. The code is submitted with this report and is commented section by section.

2 Data

The dataset contains 2,205 customers and 27 variables, described in Table 1. Spending amounts are treated as continuous. Counts (children, purchases, visits) and days are discrete. The marital status and education columns are 0/1 indicators, and each group is combined into one categorical variable during cleaning (Section 3.3).

Variable	**Type**	**Description**
Customer profile		
`Income`	Numerical, continuous	Annual household income ($)
`Age`	Numerical, discrete	Age of the customer (years)
`Kidhome`	Numerical, discrete	Number of small children at home (0–2)
`Teenhome`	Numerical, discrete	Number of teenagers at home (0–2)
`Divorced, Married, Single, Living_together, Widow`	Binary (0/1)	Marital status, one indicator per status (5 columns)
`Up_to_high_school, Up_to_undergraduate_degree, Up_to_masters_degree, PhD`	Binary (0/1)	Highest education level, one indicator per level (4 columns)
Spending in the last 2 years		
`MntWines, MntFruits, MntMeatProducts, MntFishProducts, MntSweetProducts`	Numerical, continuous	Amount spent on each product category (5 columns)
`Total_amount_spent`	Numerical, continuous	Total amount spent (sum of the 5 columns above)
Purchases		
`NumWebPurchases`	Numerical, discrete	Purchases made through the website
`NumCatalogPurchases`	Numerical, discrete	Purchases made through the catalogue
`NumStorePurchases`	Numerical, discrete	Purchases made in stores
`NumDealsPurchases`	Numerical, discrete	Purchases made with a discount
`NumWebVisitsMonth`	Numerical, discrete	Visits to the website in the last month
Relationship with the company		
`Customer_Days`	Numerical, discrete	Number of days as a customer
`Recency`	Numerical, discrete	Days since the last purchase
`Complain`	Binary (0/1)	1 if the customer complained in the last 2 years

Table 1: The 27 variables of the original dataset.

3 Data Cleaning and Preparation

3.1 Missing values and duplicates

There are no missing values. However, 203 rows are identical to another row in all 27 columns, including the exact income and the exact number of days as a customer. Two different customers matching on every column is very unlikely, so these rows were treated as repeated entries and removed (2,205 → 2,002 customers).

3.2 Logic checks

We ran 13 checks to make sure the values are consistent with each other (Table 2). Two of them failed for a few customers:

- 4 customers made 0 purchases but spent 4–8, which is impossible. They were removed.
- 1 customer made more discount purchases than purchases in total. This customer was removed.

After cleaning, **1,997 customers** remain. Check 1 also shows that `Total_amount_spent` is exactly the sum of the five `Mnt*` columns, so it is never used in the same test or model as those columns. Check 13 found 4 customers who were 16–18 years old when they joined. This is possible, so they were kept.

#	Check	Result
1	Total spent equals the sum of the `Mnt*` columns	Pass
2	No customer with 0 purchases has spent money	4 removed
3	Discount purchases do not exceed total purchases	1 removed
4	Each customer has exactly one marital status	Pass
5	Each customer has exactly one education level	Pass
6	Indicator columns contain only 0 or 1	Pass
7	No negative values	Pass
8	All counts and amounts are whole numbers	Pass
9	`Kidhome` and `Teenhome` only take values 0–2	Pass
10	Income is positive and age is realistic (24–80)	Pass
11	Last purchase is not before the customer joined	Pass
12	No customer with purchases has spent nothing	Pass
13	Age when joining (4 customers were 16–18)	Kept

Table 2: Logic checks on the data.

3.3 New variables

The indicator columns were combined into two categorical variables:

- `Marital_State`: Divorced, Living together, Married, Single or Widow.
- `schoolinglevel`: an ordered factor, high school < bachelor's < master's < PhD.

We also created:

- `Children` = `Kidhome` + `Teenhome` (0–3), and `HasChildren` (yes/no);
- `TotalPurchases` = web + catalogue + store purchases;
- `AgeGroup`: under 30, 30–44, 45–59, 60+;
- `IncomeGroup`: income quartiles (boundaries 35,641, 51,529 and 68,352);

- `WebPercent`, `CataloguePercent`, `StorePercent`: the share of each customer's purchases made through each channel (%);
- `DealPercent`: the share of each customer's purchases made with a discount (%).

4 Univariate Analysis

4.1 Numerical variables


Table 3 summarises the 15 numerical variables. All values are in a sensible range.

Variable	Min	Median	Mean	SD	Max	Skewness
Income	2,447	51,529	51,803.5	20,618.9	113,734	0.01
Age	24	50	51.2	11.7	80	0.09
Recency	0	49	49.0	29.0	99	−0.01
Customer_Days	2,159	2,510	2,511.3	202.0	2,858	0.00
MntWines	0	179	306.6	337.0	1,493	1.16
MntFruits	0	8	26.5	39.9	199	2.09
MntMeatProducts	0	69	166.5	219.5	1,725	1.82
MntFishProducts	0	12	37.8	55.0	259	1.91
MntSweetProducts	0	8	27.4	41.6	262	2.10
Total_amount_spent	6	345	564.9	576.1	2,491	0.91
NumDealsPurchases	0	2	2.3	1.8	15	2.13
NumWebPurchases	0	4	4.1	2.8	27	1.24
NumCatalogPurchases	0	2	2.7	2.8	28	1.40
NumStorePurchases	0	5	5.8	3.2	13	0.72
NumWebVisitsMonth	0	6	5.3	2.4	20	−0.01

Table 3: Summary statistics of the numerical variables (1,997 customers).

The main observations are:

- **Spending is strongly right-skewed.** For every `Mnt*` column the mean is far above the median (e.g. fruit: median 8, mean 26.5). Most customers spend little, and a few spend a lot (Figure 1).
- **Purchase counts are also right-skewed**, especially discount, web and catalogue purchases.
- **Recency and Customer_Days are roughly flat (uniform)**: customers are spread evenly over the range.
- **Income and Age are roughly symmetric** with one main peak.



(a) Total amount spent

(b) Income

Figure 1: Histograms of total spending (right-skewed) and income (roughly symmetric).

Outliers. Using the $1.5 \times$ IQR rule, Income, Age, Recency, Customer_Days and store purchases have no outliers. The spending columns have many (e.g. 220 for fruit, 166 for meat), but these come from the strong right skew rather than from errors: some customers simply spend much more than others. They are real customers, so all of them were kept. The full outlier counts are in Table 8 (Appendix A), and the histograms of all 15 variables in Figure 14.

4.2 Categorical variables

The bar charts of these variables are in Figure 15 (Appendix A).

- **Marital status:** married 777 (38.9%), living together 505 (25.3%), single 437 (21.9%), divorced 209 (10.5%) and widow 69 (3.5%).
- **Education:** bachelor's 1,006 (50.4%), master's 512 (25.6%), PhD 430 (21.5%) and high school only 49 (2.5%).
- **Children:** 561 customers (28.1%) have no children, 1,017 (50.9%) have one, 374 (18.7%) have two and 45 (2.3%) have three.
- **Complaints:** only 19 customers (0.95%) complained in the last 2 years.

Some groups (high school, widows, complainers, 3 children) are small, so tests involving them have low power.

4.3 Normality


A normality test only makes sense when the histogram looks roughly bell-shaped. From the histograms, 13 of the 15 variables are clearly not normal: spending and purchase counts are strongly right-skewed, and Recency and Customer_Days are flat. These were not tested. Only Income and Age look roughly symmetric, so only these two were checked, with a normal Q-Q plot, the Kolmogorov–Smirnov (K-S) test and the Lilliefors test (Figure 2).

- **Income:** $D = 0.047$ (K-S $p = 0.0003$, Lilliefors $p < 0.001$).
- **Age:** $D = 0.058$ (K-S $p < 0.001$, Lilliefors $p < 0.001$).

Both tests reject normality for both variables. The Lilliefors test is the correct version here, because the mean and standard deviation are estimated from the same data, which makes the

plain K-S test too lenient. The departures from normality are small ($D < 0.06$), but with about 2,000 customers even small differences are significant. The Q-Q plots show that the points bend away from the line at both ends: the distributions have a flatter top and thinner tails than a normal curve.

Conclusion: none of the numerical variables is normally distributed, so the rest of the analysis uses non-parametric methods (Spearman correlation, Wilcoxon and Kruskal–Wallis tests).



(a) Income

(b) Age

Figure 2: Normal Q-Q plots of Income and Age.

5 Bivariate Analysis

5.1 Correlations

Figure 3 shows the Spearman correlations between the numerical variables. Spearman is used because the variables are skewed and it measures any monotonic relationship, not only a straight line. The main patterns are:

- **Income drives almost everything.** It has a very strong correlation with total spending ($\rho = 0.87$), with wine (0.84) and meat (0.82) spending, and with catalogue (0.80) and store (0.75) purchases. Its correlation with web purchases is weaker (0.58).
- **Customers who buy one product buy them all.** The five spending categories are all positively correlated with each other ($\rho = 0.50$ to 0.73). Customers differ mainly in *how much* they buy, not in *what* they buy. Wine and meat each have $\rho = 0.94$ with total spending, but this is mostly built in, since they are the two largest parts of the total.
- **Catalogue purchases go with high spending.** Of the three channels, catalogue purchases have the strongest correlation with total spending (0.89), followed by store (0.82) and web (0.73).
- **Website visits are the clear exception.** Web visits are strongly negatively correlated with income ($\rho = -0.63$, the strongest negative value in the matrix), and also with every spending category (-0.39 to -0.49) and with catalogue (-0.54) and store (-0.46) purchases. Lower-income customers visit the website more but buy less. Web visits are almost unrelated to web purchases (-0.09, Section 5.6), but they are positively related to discount purchases (0.39). This suggests that frequent visitors are looking for offers.

- **Discount purchases are not related to spending** ($\rho = -0.02$). They are weakly negative with income (-0.20), and positive with web visits (0.39) and days as a customer (0.22).
- **Age and length of relationship matter little.** Age has only weak correlations with all the other variables (at most 0.24, with wine). Days as a customer is weakly related to spending (0.18) and to web visits (0.31).
- **Recency is unrelated to everything** (all $|\rho| \leq 0.03$).





Figure 3: Spearman correlation heatmap (red = positive, blue = negative).

5.2 Income and spending

Spending rises clearly with income, but not in a straight line (Figure 4a): it stays low up to an income of about 40,000 and then rises steeply. Higher-income customers also buy more often: the average number of purchases rises from 5.5 in the lowest income quartile to 19.4 in the highest (Figure 4b).




(a) Total spending, with a smooth trend line (red) (b) Average number of purchases by income group

Figure 4: Spending and purchases by income.

5.3 Children, education and marital status

- **Children:** median spending falls with every extra child, from 1,117 with no children to 264 (1 child), 85 (2) and 62 (3) (Figure 5a).
- **Education:** high-school customers spend much less (median 40) than the others. PhD customers have the highest median (448.5), but spending does not rise steadily with education: bachelor's (353) is above master's (297) (Figure 5b).
- **Marital status:** widows have the highest median spending (597), and the other groups are similar (303–350) (Figure 5c). The widow group is small (69) and older, so this may be an age effect (tested in Section 6).




(a) By number of children (b) By education level (c) By marital status

Figure 5: Total spending by number of children, education level and marital status.

5.4 Purchase channels

On average, half of a customer's purchases are made in store (50.5%), a third on the website (33.2%) and a sixth through the catalogue (16.4%). The store is the main channel in every age group (Figure 6).



(a) All customers


(b) By age group

Figure 6: Average share of purchases made through each channel.

5.5 Discounts

Families and lower-income customers are clearly more price-sensitive (Figure 7):

- The average share of purchases made with a discount rises from 8.8% for customers without children to 24.9% with one child and about 43% with two or three.
- It falls steadily with income: 36.4% in the lowest quartile, 32.6%, 19.7% and only 7.6% in the highest ($\rho = -0.73$).



(a) By number of children

(b) By income group

Figure 7: Share of purchases made with a discount.

5.6 Website visits

Website visits do not turn into website purchases (Figure 8): the correlation is almost zero ($\rho = -0.09$), and customers make about 4 web purchases on average whatever their number of visits. The most frequent visitors are the lower-income, lower-spending customers: median income falls from 77,382 (0–2 visits a month) to 31,814 (9 or more), and median spending from 1,112 to 118. A possible explanation is that these customers browse the website for offers rather than to buy. Note that visits cover only the last month, while purchases cover a longer period.






Figure 8: Website visits in the last month vs web purchases, with a smooth trend line (red). The points are moved slightly at random ("jittered") so that customers with the same values do not hide behind each other.

5.7 Recency and length of relationship

How recently a customer last bought is not related to how much they spend ($\rho = 0.02$): the cloud of points in Figure 9a is flat. The length of the relationship has a weak positive relationship with spending ($\rho = 0.18$, Figure 9b): median spending rises from 150.5 for the newest quarter of customers to 523 for the longest-standing quarter. Spending covers the last 2 years only, and every customer has been with the company for at least 5.9 years, so all customers are compared over the same period.



(a) Days since the last purchase



(b) Days as a customer

Figure 9: Total spending vs recency and length of relationship, with smooth trend lines (red).

6 Hypothesis Testing

Since no numerical variable is normal, we use non-parametric tests:

- **Wilcoxon rank-sum test** to compare a numerical variable between two groups;
- **Kruskal–Wallis test** for three or more groups, followed by pairwise Wilcoxon tests (with the Benjamini–Hochberg correction) to find which groups differ;
- **Spearman correlation test** for two numerical variables;

- **Chi-square test of independence** for two categorical variables.

The significance level is 0.05. Table 4 summarises the results.

#	Question	Test	Result	Decision
1	Spending: children vs no children	Wilcoxon	$p < 0.001$	Reject H_0
2	Spending by number of children	Kruskal–Wallis	$p < 0.001$	Reject H_0
3	Spending by education	Kruskal–Wallis	$p < 0.001$	Reject H_0
4	Spending by marital status	Kruskal–Wallis	$p = 0.24$	Do not reject
5	Income vs spending	Spearman	$\rho = 0.87$, $p < 0.001$	Reject H_0
6	Marital status vs education	Chi-square	$p = 0.26$	Do not reject
7a	Income: complained vs not	Wilcoxon	$p = 0.22$	Do not reject
7b	Spending: complained vs not	Wilcoxon	$p = 0.13$	Do not reject
8	Discount share by children	Kruskal–Wallis	$p < 0.001$	Reject H_0
9	Discount share vs income	Spearman	$\rho = -0.73$, $p < 0.001$	Reject H_0
10	Web visits vs web purchases	Spearman	$\rho = -0.09$, $p < 0.001$	Reject H_0
11	Days as customer vs spending	Spearman	$\rho = 0.18$, $p < 0.001$	Reject H_0
12	Recency vs spending	Spearman	$\rho = 0.02$, $p = 0.43$	Do not reject

Table 4: Summary of the hypothesis tests. In each test, H_0 states that there is no difference between the groups (or no monotonic relationship).

Children (tests 1, 2, 8). Customers without children spend far more (median 1,117 vs 182). The pairwise tests show that spending falls with each extra child from 0 to 2, while the groups with 2 and 3 children do not differ significantly ($p = 0.15$). The discount share shows the same pattern in reverse: it rises from 0 to 2 children (medians 5.9%, 25.0% and 43.3%, all pairs $p < 0.001$), and 2 and 3 children do not differ ($p = 0.9$). All the pairwise p-values are in Table 9 (Appendix A).

Education (test 3). High-school customers spend much less than every other level (all $p < 0.001$). PhD customers spend more than master's customers ($p = 0.004$), but bachelor's does not differ significantly from master's ($p = 0.10$) or PhD ($p = 0.07$). The clear finding is therefore high school versus the rest. The high-school group is small (49 customers).

Marital status (test 4). There is no evidence that spending differs by marital status. Widows have the highest median, but they are also older (median age 63, vs 48–53 for the other groups), and older customers spend more ($\rho = 0.18$, $p < 0.001$). Among customers aged 60 and over, widows (median 606) spend about the same as non-widows (median 546; Wilcoxon $p = 0.67$). The higher spending of widows is therefore explained by their age, not by their marital status.

Complaints (test 7). Customers who complained have lower medians (income 39,684 vs 51,566; spending 89 vs 345), but the differences are not significant. With only 19 complainers, the tests have very little power, so we cannot conclude that there is no difference either.

Significant but weak (tests 10 and 11). With about 2,000 customers, even very small correlations are statistically significant. The web visits correlation ($\rho = -0.09$) is significant but negligible in practice. A p-value tells us whether a relationship exists, while ρ tells us how strong it is.

Multiple testing. Running many tests at the 0.05 level increases the chance that some are significant by chance. After adjusting all p-values with the Benjamini–Hochberg correction, the conclusions do not change. Most significant p-values are below 0.001, so even the stricter Bonferroni threshold ($0.05/13 \approx 0.0038$) would give the same conclusions.

7 Customer Segments and Regression

7.1 Principal Component Analysis

Principal Component Analysis (PCA) summarises many variables with a few new axes, called principal components, that keep as much of the variation as possible. We applied it to 10 variables describing customer behaviour: income, the five spending columns, the three purchase channels and website visits. The variables were standardised first, so that income does not dominate.

The first component (PC1) explains 56% of the variation and the second (PC2) another 12%, so together they keep 68% of the information in the 10 variables (Table 5).

- **PC1: overall buying power.** Income, all the spending columns and all the purchase channels have similar loadings, and website visits has the opposite sign. The sign of a component is arbitrary. Here a *low* PC1 score means high income and high spending.
- **PC2: online wine buyers.** It is highest for web purchases, web visits and wine, and negative for fish, fruit and sweets.

Most of the differences between customers therefore come down to one thing: how much they earn and spend. The second, much smaller difference is whether they buy online and buy wine, or buy more fresh and sweet products.

Variable	PC1	PC2
Income	−0.38	0.07
MntWines	−0.32	0.40
MntFruits	−0.30	−0.24
MntMeatProducts	−0.35	−0.14
MntFishProducts	−0.31	−0.26
MntSweetProducts	−0.30	−0.21
NumWebPurchases	−0.23	0.63
NumCatalogPurchases	−0.35	0.01
NumStorePurchases	−0.32	0.24
NumWebVisitsMonth	0.27	0.44
Variance explained	56%	12%

Table 5: Loadings of the first two principal components.


7.2 Clustering with k-means

To find groups of customers with similar behaviour, we used k-means clustering on the same 10 variables: income, the five spending columns, the three purchase channels and website visits. Total spending was left out because it is the sum of the spending columns. All variables were standardised first (mean 0, SD 1), so that income, measured in tens of thousands, does not dominate the purchase counts.

Choosing the number of clusters. Two methods were used (Figure 10):

- **Elbow method:** the within-cluster sum of squares drops sharply from $k = 1$ to 2 and from 2 to 3, and then flattens, so the elbow is at $k = 2$–3.
- **Silhouette method:** the average silhouette width is highest for $k = 2$ (0.43), followed by $k = 3$ (0.35).

We chose $k = 3$. With $k = 2$ the customers are only split into high and low spenders, while $k = 3$ also separates a distinct middle group, which is more useful for describing customer profiles. Its silhouette (0.35) is still reasonable, and it sits at the elbow.



(a) Elbow method

(b) Silhouette method

Figure 10: Choosing the number of clusters.

The three segments. Table 6 describes each cluster using the medians of the original variables. Age, children and the purchase shares were not used to build the clusters, but they help describe them.

	1. Budget families	**2. Affluent big spenders**	**3. Established families**
Customers	960	491	546
Income	35,081	76,412	59,987
Total spending	54	1,290	652
Wine spending	23	548	434
Meat spending	15	435	127
Age	48	51	54
Have children	89%	25%	84%
Discount share	33%	6%	15%
Web share	33%	26%	36%
Catalogue share	0%	30%	16%
Store share	57%	43%	46%
Web visits per month	7	2	6

Table 6: Cluster profiles (medians, except "Have children").

- **Budget families** (960 customers, the largest group): the lowest income and very low spending. Almost all have children. A third of their purchases are made with a discount, mostly in store, and they almost never use the catalogue. They visit the website often but buy little there.
- **Affluent big spenders** (491): the highest income and spending, especially on wine and meat. Most have no children. They rarely use discounts and use the catalogue the most.
- **Established families** (546): mid-to-high income and spending, mostly on wine. Most have children, and they are slightly older. They use some discounts and buy online the most.

Figure 12 shows that budget families are clearly separated from the other two groups by income and spending. Clusters 2 and 3 overlap at mid-to-high incomes: they differ less in income and more in *how* they buy (children, catalogue vs web, wine vs meat), which a two-variable plot cannot show. Five of the ten clustering variables are spending columns, so spending has a large weight in the clustering, which is why the clusters follow spending level closely.

The PCA plot separates the clusters much better (Figure 11). Along PC1 they are ordered by buying power: affluent big spenders (median PC1 score -3.27), established families (-0.59) and budget families (2.20). Along PC2, established families score highest (median 1.06), because they buy online and buy wine the most. So clusters 2 and 3, which overlap in income, are told apart by *how* they buy.




Figure 11: Customers on the first two principal components, coloured by cluster.




(a) Total spending by cluster

(b) Income vs spending, coloured by cluster

Figure 12: The three customer segments.

7.3 Regression: what explains spending?

The tests in Section 6 look at one variable at a time. Income, children, age and education are related to each other, so we fitted a linear regression to measure the effect of each one *while holding the others fixed*. Spending is strongly right-skewed, so the response is log(Total_amount_spent), which is much more symmetric (Figure 13; the minimum spending is 6, so there are no zeros).



(a) Total amount spent

(b) log(Total amount spent)

Figure 13: Spending before and after the log transformation.

The model is:

$$\log(\text{Spending}) = \beta_0 + \beta_1\,\text{Income} + \beta_2\,\text{Children} + \beta_3\,\text{Age} + \text{Education} + \text{Marital status} + \varepsilon.$$

With a log response, a coefficient β means that spending changes by about $(e^{\beta} - 1) \times 100\%$ for a one-unit increase in that predictor. High school and divorced are the reference groups.

Predictor	Coefficient	p-value	Effect on spending
Income (per 10,000)	0.577	< 0.001	+78%
Children (per child)	−0.344	< 0.001	−29%
Age (per year)	0.0045	0.006	+0.45%
Education (vs high school)	0.01 to 0.05	all > 0.6	not significant
Marital status (vs divorced)	−0.01 to 0.08	all > 0.4	not significant

Table 7: Regression of log(spending). $R^2 = 0.73$.

- **Income** is the strongest predictor: 10,000 more income goes with about 78% higher spending.
- **Each extra child** goes with about 29% lower spending *at the same income*. Children therefore reduce spending on their own, not only because families with children earn less.
- **Age** has a small positive effect (about 0.45% per year).
- **Education is not significant** once income is in the model. The low spending of high-school customers (test 3) is therefore mostly explained by their lower income.
- **Marital status is not significant**, which agrees with test 4.
- The model explains 73% of the variation in log(spending) ($R^2 = 0.73$), mostly through income and children.

8 Conclusions and Recommendations

8.1 Main findings

1. **Income is the main driver of spending** ($\rho = 0.87$; +78% spending per 10,000 of income).
2. **Children reduce spending strongly**, even at the same income (about −29% per child), and families rely much more on discounts.
3. **Lower-income customers are more price-sensitive**: their share of discounted purchases is much higher ($\rho = -0.73$).
4. **Education and marital status matter little.** High-school customers spend less, but this is explained by their income. The higher spending of widows is explained by their age.
5. **Website visits do not turn into website purchases.** The most frequent visitors are low-income, low-spending customers.
6. **The store is the main channel** (about half of all purchases) in every age group.
7. **There are three customer segments**: budget families, affluent big spenders and established families.

8.2 Recommendations for the company

- **Budget families:** target them with discounts and family offers, mainly in store. Since they visit the website often but rarely buy there, online offers could help turn their visits into purchases.
- **Affluent big spenders:** focus on premium wine and meat products, through the catalogue. Discounts are not needed to keep them.
- **Established families:** this group buys online the most and spends a lot on wine, so it is a good target for online wine promotions.
- **Website:** investigate why visits do not lead to purchases, for example the checkout process or the offers shown to frequent visitors.

9 Limitations

- **Small groups:** high school (49), widows (69), 3 children (45) and complainers (19). Tests on these groups have low power.
- **Different time periods:** website visits cover only the last month, while spending and purchases cover the last 2 years.
- **Correlation is not causation:** the analysis shows associations, not causes. For example, children are related to lower spending, but other factors linked to having children may also play a role.
- **Regression:** the straight-line model does not fully capture the curved effect of income.
- **Clustering:** the choice of $k = 3$ is partly a judgement call ($k = 2$ had a higher silhouette), and the result depends on which variables are used.
- **Duplicates:** we assumed that the 203 identical rows are repeated entries. If some were real customers, a small amount of data was lost.

A Additional Figures and Tables

A.1 Histograms of all numerical variables

Figure 14 shows the distribution of each of the 15 numerical variables. Spending and purchase counts are right-skewed, Recency and Customer_Days are roughly flat, and only Income and Age look roughly symmetric (Section 4.3).






Figure 14: Histograms of the 15 numerical variables.


A.2 Categorical variables




(a) Marital status



(b) Education level



(c) Small children at home



(d) Teenagers at home

Figure 15: Number of customers in each group of the categorical variables.

A.3 Outliers

Table 8 gives the number of outliers of each numerical variable by the $1.5 \times$ IQR rule: values below $Q_1 - 1.5\,\text{IQR}$ or above $Q_3 + 1.5\,\text{IQR}$. For every variable except Age the lower limit is negative, so all the outliers are unusually *high* values. They come from the right skew of the variables, not from errors, so all of them were kept.

Variable	Lower limit	Upper limit	Outliers	% of customers
Income	−13,426	117,419	0	0.0
Age	16	88	0	0.0
Recency	−51	149	0	0.0
Customer_Days	1,812	3,212	0	0.0
MntWines	−701	1,232	32	1.6
MntFruits	−45	80	220	11.0
MntMeatProducts	−305	551	166	8.3
MntFishProducts	−68	121	201	10.1
MntSweetProducts	−49	84	217	10.9
Total_amount_spent	−1,306	2,326	2	0.1
NumDealsPurchases	−2	6	70	3.5
NumWebPurchases	−4	12	3	0.2
NumCatalogPurchases	−6	10	19	1.0
NumStorePurchases	−4.5	15.5	0	0.0
NumWebVisitsMonth	−3	13	3	0.2

Table 8: Outliers by the $1.5 \times$ IQR rule (1,997 customers).

A.4 Pairwise comparisons

Table 9 gives the p-values of the pairwise Wilcoxon tests that follow the significant Kruskal–Wallis tests 2, 3 and 8, corrected with the Benjamini–Hochberg method.

Spending by children

Pair	p
0 vs 1	< 0.001
0 vs 2	< 0.001
0 vs 3	< 0.001
1 vs 2	< 0.001
1 vs 3	< 0.001
2 vs 3	0.148

Spending by education

Pair	p
HS vs Bach.	< 0.001
HS vs Mast.	< 0.001
HS vs PhD	< 0.001
Bach. vs Mast.	0.102
Bach. vs PhD	0.072
Mast. vs PhD	0.004

Discount share by children

Pair	p
0 vs 1	< 0.001
0 vs 2	< 0.001
0 vs 3	< 0.001
1 vs 2	< 0.001
1 vs 3	< 0.001
2 vs 3	0.904

Table 9: Pairwise Wilcoxon tests (BH-corrected p-values). HS = high school, Bach. = bachelor's, Mast. = master's.

B R Code

The analysis is submitted as a commented R file. It is organised in the following parts:

1. Setup: file paths, the function that saves figures and the lists of column names.
2. Cleaning: import, logic checks, cleaning and new variables.
3. Descriptive analysis: summary statistics, outliers, categorical variables and normality.
4. Bivariate analysis: correlations, group comparisons and scatterplots.
5. Hypothesis tests.
6. PCA, clustering and regression.

The R packages used are `e1071` (skewness and kurtosis), `nortest` (Lilliefors test), `ggplot2` (correlation heatmap and violin plot) and `cluster` (silhouette).