



ITMO UNIVERSITY

How to Win Coding Competitions: Secrets of Champions


Week 6: Algorithms on Graphs 2

Lecture 1: Eulerian paths and Eulerian tours

Maxim Buzdalov
Saint Petersburg 2016


An **Eulerian path** is a path in a graph that contains each edge of the graph exactly once

FBCEDFGCADBAE




An **Eulerian path** is a path in a graph that contains each edge of the graph exactly once

FBCEDFGCADBAE




An **Eulerian path** is a path in a graph that contains each edge of the graph exactly once

FBCEDFGCADBAE




An **Eulerian path** is a path in a graph that contains each edge of the graph exactly once

FBCEDFGCADBAE




An **Eulerian path** is a path in a graph that contains each edge of the graph exactly once

FBCEDFGCADBAE




An **Eulerian path** is a path in a graph that contains each edge of the graph exactly once

FBCE**DFGCADBAE**




An **Eulerian path** is a path in a graph that contains each edge of the graph exactly once

FBCEDFGCADBAE




An **Eulerian path** is a path in a graph that contains each edge of the graph exactly once

FBCEDFGCADBAE




An **Eulerian path** is a path in a graph that contains each edge of the graph exactly once

FBCEDFG**CADBAE**




An **Eulerian path** is a path in a graph that contains each edge of the graph exactly once

FBCEDFGCADBAE




An **Eulerian path** is a path in a graph that contains each edge of the graph exactly once

FBCEDFGCADA**EB**




An **Eulerian path** is a path in a graph that contains each edge of the graph exactly once

FBCEDFGCADBAE




An **Eulerian path** is a path in a graph that contains each edge of the graph exactly once

FBCEDFGCADBAE




An **Eulerian path** is a path in a graph that contains each edge of the graph exactly once

FBCEDFGCADBAE




An **Eulerian path** is a path in a graph that contains each edge of the graph exactly once

FBCEDFGCADBAE




An **Eulerian tour** is a Eulerian path which starts and ends on the same vertex

FEABGCEDBCADF




An **Eulerian path** is a path in a graph that contains each edge of the graph exactly once

FBCEDFGCADBAE




An **Eulerian tour** is a Eulerian path which starts and ends on the same vertex

FEABGCEDBCADF




An **Eulerian path** is a path in a graph that contains each edge of the graph exactly once

FBCEDFGCADBAE




An **Eulerian tour** is a Eulerian path which starts and ends on the same vertex

FEABGCEDBCADF




An **Eulerian path** is a path in a graph that contains each edge of the graph exactly once

FBCEDFGCADBAE




An **Eulerian tour** is a Eulerian path which starts and ends on the same vertex

FEABGCEDBCADF




An **Eulerian path** is a path in a graph that contains each edge of the graph exactly once

FBCEDFGCADBAE




An **Eulerian tour** is a Eulerian path which starts and ends on the same vertex

FEABGCEDBCADF




An **Eulerian path** is a path in a graph that contains each edge of the graph exactly once

FBCEDFGCADBAE




An **Eulerian tour** is a Eulerian path which starts and ends on the same vertex

FEABGCEDBCADF




An **Eulerian path** is a path in a graph that contains each edge of the graph exactly once

FBCEDFGCADBAE




An **Eulerian tour** is a Eulerian path which starts and ends on the same vertex

FEABGCEDBCADF




An **Eulerian path** is a path in a graph that contains each edge of the graph exactly once

FBCEDFGCADBAE




An **Eulerian tour** is a Eulerian path which starts and ends on the same vertex

FEABGCEDBCADF




An **Eulerian path** is a path in a graph that contains each edge of the graph exactly once

FBCEDFGCADBAE




An **Eulerian tour** is a Eulerian path which starts and ends on the same vertex

FEABGCEDBCADF




An **Eulerian path** is a path in a graph that contains each edge of the graph exactly once

FBCEDFGCADBAE




An **Eulerian tour** is a Eulerian path which starts and ends on the same vertex

FEABGCEDBCADF




An **Eulerian path** is a path in a graph that contains each edge of the graph exactly once

FBCEDFGCADBAE




An **Eulerian tour** is a Eulerian path which starts and ends on the same vertex

FEABGCEDBCADF




An **Eulerian path** is a path in a graph that contains each edge of the graph exactly once

FBCEDFGCADBAE




An **Eulerian tour** is a Eulerian path which starts and ends on the same vertex

FEABGCEDBCADF




An **Eulerian path** is a path in a graph that contains each edge of the graph exactly once

FBCEDFGCADBAE




An **Eulerian tour** is a Eulerian path which starts and ends on the same vertex

FEABGCEDBCAD~~DF~~




An **Eulerian path** is a path in a graph that contains each edge of the graph exactly once

FBCEDFGCADBAE



An **Eulerian tour** is a Eulerian path which starts and ends on the same vertex

FEABGCEDBCADF



When does Eulerian path exist?

When does Eulerian path exist?

- ▶ Undirected graph:
 - ▶ The graph is connected
 - ▶ There are at most two vertices with odd degree

When does Eulerian path exist?

- ▶ Undirected graph:
 - ▶ The graph is connected
 - ▶ There are at most two vertices with odd degree
- ▶ Directed graph:
 - ▶ The graph is connected (when directions are removed)
 - ▶ At most one vertex u has $\deg^+(u) - \deg^-(u) = +1$
 - ▶ At most one vertex v has $\deg^+(v) - \deg^-(v) = -1$
 - ▶ All other vertices have $\deg^+(x) = \deg^-(x)$

When does Eulerian path exist?

- ▶ Undirected graph:
 - ▶ The graph is connected
 - ▶ There are at most two vertices with odd degree
- ▶ Directed graph:
 - ▶ The graph is connected (when directions are removed)
 - ▶ At most one vertex u has $\deg^+(u) - \deg^-(u) = +1$
 - ▶ At most one vertex v has $\deg^+(v) - \deg^-(v) = -1$
 - ▶ All other vertices have $\deg^+(x) = \deg^-(x)$

When does Eulerian tour exist?

When does Eulerian path exist?

- ▶ Undirected graph:
 - ▶ The graph is connected
 - ▶ There are at most two vertices with odd degree
- ▶ Directed graph:
 - ▶ The graph is connected (when directions are removed)
 - ▶ At most one vertex u has $\deg^+(u) - \deg^-(u) = +1$
 - ▶ At most one vertex v has $\deg^+(v) - \deg^-(v) = -1$
 - ▶ All other vertices have $\deg^+(x) = \deg^-(x)$

When does Eulerian tour exist?

- ▶ Undirected graph:
 - ▶ The graph is connected
 - ▶ All vertex degrees are even

When does Eulerian path exist?

- ▶ Undirected graph:
 - ▶ The graph is connected
 - ▶ There are at most two vertices with odd degree
- ▶ Directed graph:
 - ▶ The graph is connected (when directions are removed)
 - ▶ At most one vertex u has $\deg^+(u) - \deg^-(u) = +1$
 - ▶ At most one vertex v has $\deg^+(v) - \deg^-(v) = -1$
 - ▶ All other vertices have $\deg^+(x) = \deg^-(x)$

When does Eulerian tour exist?

- ▶ Undirected graph:
 - ▶ The graph is connected
 - ▶ All vertex degrees are even
- ▶ Directed graph:
 - ▶ The graph is strongly connected
 - ▶ All vertices have $\deg^+(x) = \deg^-(x)$

- ▶ Induction: assume that your graph has $|E|$ edges,
and the theorem was proven for all $e < |E|$ numbers of edges

- ▶ Induction: assume that your graph has $|E|$ edges,
and the theorem was proven for all $e < |E|$ numbers of edges
- ▶ Pick a vertex
 - ▶ Random when you don't have "special" vertices
 - ▶ A "special" vertex if you have one
 - ▶ The one with $\deg^+(v) > \deg^-(v)$ if the graph is directed

- ▶ Induction: assume that your graph has $|E|$ edges,
and the theorem was proven for all $e < |E|$ numbers of edges
- ▶ Pick a vertex
 - ▶ Random when you don't have "special" vertices
 - ▶ A "special" vertex if you have one
 - ▶ The one with $\deg^+(v) > \deg^-(v)$ if the graph is directed
- ▶ Traverse the graph and remove the traversed edges

- ▶ Induction: assume that your graph has $|E|$ edges, and the theorem was proven for all $e < |E|$ numbers of edges
- ▶ Pick a vertex
 - ▶ Random when you don't have "special" vertices
 - ▶ A "special" vertex if you have one
 - ▶ The one with $\deg^+(v) > \deg^-(v)$ if the graph is directed
- ▶ Traverse the graph and remove the traversed edges
- ▶ If you cannot do this anymore, what happens?
 - ▶ Either there are no more edges \rightarrow path/tour is found
 - ▶ Some edges remain
 - ▶ There are (maybe several) connected subgraphs
 - ▶ Find the paths/tours in them (can do this **by induction**)
 - ▶ Connect them with the path/tour constructed from removed edges

- ▶ Induction: assume that your graph has $|E|$ edges, and the theorem was proven for all $e < |E|$ numbers of edges
- ▶ Pick a vertex
 - ▶ Random when you don't have "special" vertices
 - ▶ A "special" vertex if you have one
 - ▶ The one with $\deg^+(v) > \deg^-(v)$ if the graph is directed
- ▶ Traverse the graph and remove the traversed edges
- ▶ If you cannot do this anymore, what happens?
 - ▶ Either there are no more edges \rightarrow path/tour is found
 - ▶ Some edges remain
 - ▶ There are (maybe several) connected subgraphs
 - ▶ Find the paths/tours in them (can do this **by induction**)
 - ▶ Connect them with the path/tour constructed from removed edges
- ▶ That's what **Depth First Search** can do!

```
 $G = \langle V, E \rangle$   
 $A(v) = \{u \mid (v, u) \in E\}$   
 $R \leftarrow []$   
procedure DFS( $v$ )  
  for  $u \in A(v)$  do  
    Remove  $u$  from  $A(v)$   
    if graph is undirected then  
      Remove  $v$  from  $A(u)$   
    end if  
    DFS( $u$ )  
  end for  
   $R \leftarrow [v] + R$   
end procedure
```

$G = \langle V, E \rangle$ $A(v) = \{u \mid (v, u) \in E\}$ ▷ No U set is used: can visit a vertex more than once! $R \leftarrow []$ **procedure** DFS(v) **for** $u \in A(v)$ **do** Remove u from $A(v)$ **if** graph is undirected **then** Remove v from $A(u)$ **end if** DFS(u) **end for** $R \leftarrow [v] + R$ **end procedure**

$G = \langle V, E \rangle$ $A(v) = \{u \mid (v, u) \in E\}$ ▷ No U set is used: can visit a vertex more than once! $R \leftarrow []$

▷ The result storage

procedure DFS(v) **for** $u \in A(v)$ **do** Remove u from $A(v)$ **if** graph is undirected **then** Remove v from $A(u)$ **end if** DFS(u) **end for** $R \leftarrow [v] + R$ **end procedure**

$G = \langle V, E \rangle$
 $A(v) = \{u \mid (v, u) \in E\}$

▷ No U set is used: can visit a vertex more than once!

 $R \leftarrow []$

▷ The result storage

procedure DFS(v)

▷ Should be called on a **non-regular vertex**, if any

for $u \in A(v)$ **do**

 Remove u from $A(v)$

if graph is undirected **then**

 Remove v from $A(u)$

end if

 DFS(u)

end for

$R \leftarrow [v] + R$

end procedure

$G = \langle V, E \rangle$
 $A(v) = \{u \mid (v, u) \in E\}$

▷ No U set is used: can visit a vertex more than once!

 $R \leftarrow []$

▷ The result storage

procedure DFS(v)

▷ Should be called on a non-regular vertex, if any

for $u \in A(v)$ **do**

 Remove u from $A(v)$

▷ Will never follow the (v, u) edge anymore

if graph is undirected **then**

 Remove v from $A(u)$
end if

 DFS(u)

end for
 $R \leftarrow [v] + R$
end procedure

$G = \langle V, E \rangle$
 $A(v) = \{u \mid (v, u) \in E\}$

▷ No U set is used: can visit a vertex more than once!

 $R \leftarrow []$

▷ The result storage

procedure DFS(v)

▷ Should be called on a non-regular vertex, if any

for $u \in A(v)$ **do**

 Remove u from $A(v)$

▷ Will never follow the (v, u) edge anymore

if graph is undirected **then**

 Remove v from $A(u)$

▷ If undirected, the anti-edge must be removed

end if

 DFS(u)

end for

$R \leftarrow [v] + R$

end procedure

$G = \langle V, E \rangle$
 $A(v) = \{u \mid (v, u) \in E\}$

▷ No U set is used: can visit a vertex more than once!

 $R \leftarrow []$

▷ The result storage

procedure DFS(v)

▷ Should be called on a non-regular vertex, if any

for $u \in A(v)$ **do**

 Remove u from $A(v)$

▷ Will never follow the (v, u) edge anymore

if graph is undirected **then**

 Remove v from $A(u)$

▷ If undirected, the anti-edge must be removed


end if


 DFS(u)


end for
 $R \leftarrow [v] + R$


▷ Prepend v to the answer


end procedure





























E




E




E




E




E




E




CE




CE




BCE




BCE




DBCE




DBCE





ADBCE





ADBCE




CADBCE

CADBCE


GCADBCE


GCADBCE

FGCADBCE




FGCADBCE




DFGCADBCE


DFGCADBCE




EDFGCADBCE




EDFGCADBCE




AEDFGCADBCE




AEDFGCADBCE




BAEDFGCADBCE



BAEDFGCADBCE



FBAEDFGCADBCE



FBAEDFGCADBCE

