## Medication Overview:

- 1. Dalo (1 Morning, 1 Dinner for 10 days):
  - Dosage Instructions: Administer 1 tablet during the morning and 1 tablet during dinner.
  - Special Instructions: Do not crush or chew tablets. Report any adverse reactions promptly.

## 2. Anvil (1 Lunch for 5 days):

- Prescribed for: [Indication/Condition]
- Dosage Instructions: Take 1 tablet during the lunchtime.
- Special Instructions: Inform about any history of gastric issues. Avoid alcohol during the medication period.

## 3. Citrizine (1 Morning, 1 Lunch for 2 days):

- Dosage Instructions: Consume 1 tablet in the morning and 1 tablet during lunch.
- Special Instructions: May cause drowsiness; refrain from operating machinery. Seek medical attention for severe symptoms.

## Additional Guidance:

- Medications are tailored to the patient's specific health needs; modifications should be made only under professional consultation.
- Missed doses should be taken as soon as remembered; skip if it's almost time for the next dose.
- Scheduled follow-up appointments are crucial for a comprehensive health assessment.
- Immediate reporting of any health changes, new symptoms, or concerns is encouraged.

Disclaimer: This document is intended for reference by healthcare professionals and should not replace di rect consultation. For any queries or modifications to the prescribed medications, please contact the prescribing physician directly.