

CHAPTER-7
COORDINATE GEOMETRY

Exercise 7.4

Q4. The two opposite vertices of a square are $(-1, 2)$ and $(3, 2)$. Find the coordinates of the other two vertices.

Solution :

Given points $A = \begin{pmatrix} -1 \\ 2 \end{pmatrix}$, $C = \begin{pmatrix} 3 \\ 2 \end{pmatrix}$

Let $B = \begin{pmatrix} x_1 \\ y_1 \end{pmatrix}$ and $D = \begin{pmatrix} x_2 \\ y_2 \end{pmatrix}$




Figure 1:

Shifting point A to origin $A' = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$, $B' = B - A = \begin{pmatrix} x_1 + 1 \\ y_1 - 2 \end{pmatrix}$
 $C' = C - A = \begin{pmatrix} 4 \\ 0 \end{pmatrix}$ and $D' = D - A = \begin{pmatrix} x_2 + 1 \\ y_2 - 2 \end{pmatrix}$




Figure 2:

Now calculating the angle the **AB** makes with the x-axis.

Let **AB** makes an angle θ with the x-axis and we know $\angle C'A'B' = 45^\circ$

Angle made by **AC** = $\theta + 45$ on the x-axis

$$\tan(\theta + 45) = \frac{0 - 0}{4 - 0}$$

$$\theta = -45^\circ$$

We know the rotation matrix is given as $P = \begin{pmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{pmatrix}$

Now the transformed coordinates are

$$C'' = P(C - A) = \begin{pmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix} \begin{pmatrix} 4 \\ 0 \end{pmatrix} = \begin{pmatrix} \frac{4}{\sqrt{2}} \\ -\frac{4}{\sqrt{2}} \end{pmatrix}$$

$$B'' = \begin{pmatrix} 0 \\ -\frac{4}{\sqrt{2}} \end{pmatrix}, D'' = \begin{pmatrix} \frac{4}{\sqrt{2}} \\ 0 \end{pmatrix} \text{ and } A'' = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$




Figure 3:

Again transforming the coordinates back to the original axis.
We know for anti-clockwise direction the rotation matrix is given as

$$P = \begin{pmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{pmatrix}$$

So now the transformed coordinates B and D will be

$$B' = PB'' = \begin{pmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix} \begin{pmatrix} 0 \\ -\frac{4}{\sqrt{2}} \end{pmatrix} = \begin{pmatrix} 2 \\ -2 \end{pmatrix}$$

$$D' = PD'' = \begin{pmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix} \begin{pmatrix} \frac{4}{\sqrt{2}} \\ 0 \end{pmatrix} = \begin{pmatrix} 2 \\ 2 \end{pmatrix}$$




Figure 4:

Again transforming the axis back to the original

$$B = B' + A = \begin{pmatrix} 2 \\ -2 \end{pmatrix} + \begin{pmatrix} -1 \\ 2 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$D = D' + A = \begin{pmatrix} 2 \\ 2 \end{pmatrix} + \begin{pmatrix} -1 \\ 2 \end{pmatrix} = \begin{pmatrix} 1 \\ 4 \end{pmatrix}$$

Hence, the other two vertices are $B(1,0)$ and $D(1,4)$




Figure 5: