CHAPTER-7 COORDINATE GEOMETRY

Excercise 7.4

Q4. The two opposite vertices of a square are (-1,2) and (3,2). Find the coordinates of the other two vertices.

Solution:

Given points

$$\mathbf{A} = \begin{pmatrix} -1\\2 \end{pmatrix}, \mathbf{C} = \begin{pmatrix} 3\\2 \end{pmatrix} \tag{1}$$




Figure 1:

Shifting point A to origin with refrence to Figure 2 $\,$

$$\mathbf{A}' = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \mathbf{C}' = \mathbf{C} - \mathbf{A} = \begin{pmatrix} 4 \\ 0 \end{pmatrix} \tag{2}$$




Figure 2:

For a generalised case, the angle made by AC on the x-axis will be $\theta+45^{\circ}$, where θ is the angle made by AB on the x-axis. Hence, we can calculate the angle made by AB on the x-axis as.

dierection vector is given as

$$\mathbf{C} - \mathbf{A} = \begin{pmatrix} 4 \\ 0 \end{pmatrix} \tag{3}$$

However in this particular case, angle made by AC on the x-axis

$$tan\beta = \frac{0}{4} \implies sin\beta = \frac{0}{4} \tag{4}$$

$$\beta = 0^{\circ} \tag{5}$$

Therefore, here $\theta = -45^{\circ}$

We know the rotation matrix for CW rotation is given as

$$\mathbf{P}^{\top} = \begin{pmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{pmatrix} \tag{6}$$

Since the angle here is negative so ultimately the coordinates will be rotated in ACW direction. Now the transformed (rotated) coordinates are with refrence to Figure 3.

$$\mathbf{C}'' = \mathbf{P}(\mathbf{C} - \mathbf{A}) = \begin{pmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix} \begin{pmatrix} 4 \\ 0 \end{pmatrix} = \begin{pmatrix} \frac{4}{\sqrt{2}} \\ \frac{4}{\sqrt{2}} \end{pmatrix}$$
(7)

$$\mathbf{B}'' = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \mathbf{C}'' = \begin{pmatrix} \frac{4}{\sqrt{2}} \\ 0 \end{pmatrix}, \mathbf{D}'' = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \mathbf{C}'' = \begin{pmatrix} 0 \\ \frac{4}{\sqrt{2}} \end{pmatrix} \text{ and } \mathbf{A}'' = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$
(8)




Figure 3:

Again tranforming(rotating) the coordinates back to the original axis. We know for anti-clockwise direction the rotation matrix is given as

$$\mathbf{P} = \begin{pmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{pmatrix} \tag{9}$$

Again we know that the angle is negative so the rotation will be in clockwise direction. So now the transformed(rotated) coordinates ${\bf B}$ and ${\bf D}$ are with refrence to Figure 4

$$\mathbf{B}' = \mathbf{P}\mathbf{B}' = \begin{pmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix} \begin{pmatrix} \frac{4}{\sqrt{2}} \\ 0 \end{pmatrix} = \begin{pmatrix} 2 \\ -2 \end{pmatrix}$$
 (10)

$$\mathbf{D}' = \mathbf{P}\mathbf{D}'' = \begin{pmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix} \begin{pmatrix} 0 \\ \frac{4}{\sqrt{2}} \end{pmatrix} = \begin{pmatrix} 2 \\ 2 \end{pmatrix}$$
 (11)




Figure 4:

Again transforming (shifting) the axis back to the original with refrence to Figure $5\,$

$$\mathbf{B} = \mathbf{B}' + \mathbf{A} = \begin{pmatrix} 2 \\ -2 \end{pmatrix} + \begin{pmatrix} -1 \\ 2 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \tag{12}$$

$$\mathbf{D} = \mathbf{D}' + \mathbf{A} = \begin{pmatrix} 2\\2 \end{pmatrix} + \begin{pmatrix} -1\\2 \end{pmatrix} = \begin{pmatrix} 1\\4 \end{pmatrix} \tag{13}$$

Hence, the other two vertices are $\mathbf{B}(1,0)$ and $\mathbf{D}(1,4)$ The direct formula for calculation of the vertices is:

$$\mathbf{B} = \mathbf{A} + \mathbf{P} \begin{pmatrix} \mathbf{e} \mathbf{1} & \mathbf{0} \end{pmatrix} \mathbf{P}^{\top} (\mathbf{C} - \mathbf{A})$$
 (14)

$$\mathbf{D} = \mathbf{A} + \mathbf{P} \begin{pmatrix} \mathbf{0} & \mathbf{e2} \end{pmatrix} \mathbf{P}^{\top} (\mathbf{C} - \mathbf{A})$$
 (15)

(16)

where ${\bf P}$ is the rotation matrix and ${\bf A}$ and ${\bf C}$ are the position vectors of opposite vertices.




Figure 5: