OPTIMIZATION

Excercise 6.6

Q4. Find the equation of normal to the curve $x^2 = 4y$ which passes through the point (4,-2) **Solution:** The given equation of the curve can be written as

$$g(\mathbf{x}) = \mathbf{x}^{\mathsf{T}} \mathbf{V} \mathbf{x} + 2\mathbf{u}^{\mathsf{T}} \mathbf{x} + f = 0 \tag{1}$$

where

$$\mathbf{V} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \tag{2}$$

$$\mathbf{u} = \begin{pmatrix} 0 \\ -2 \end{pmatrix} \tag{3}$$

$$f = 0 (4)$$

We are given that

$$\mathbf{h} = \begin{pmatrix} 4 \\ -2 \end{pmatrix} \tag{5}$$

This can be formulated as optimization problem as follows:

$$\min_{\mathbf{x}} \quad f(\mathbf{x}) = \|\mathbf{x} - \mathbf{h}\|^2 \tag{6}$$

s.t.
$$g(\mathbf{x}) = \mathbf{x}^{\mathsf{T}} \mathbf{V} \mathbf{x} + 2 \mathbf{u}^{\mathsf{T}} \mathbf{x} + f = 0$$
 (7)

It is already proved that the optimization problem is non-convex. We use Gradient Descent to find the optimum value. We define

$$\mathbf{x}_{n+1} = \mathbf{x}_n - \alpha \nabla g \left(\mathbf{x}_n \right) \tag{8}$$

And the condition is given as

$$(\mathbf{x} - \mathbf{h})^{\top} \nabla g (\mathbf{x}_n) \neq 0 \tag{9}$$

Now we choose the parameters as

- 1. $\alpha = 0.001$
- 2. precision = 0.001
- 3. n = 10000
- 4. $\mathbf{x}_0 = 4$

We get the minimum value of \mathbf{x} as

$$\mathbf{x}_{min} = \begin{pmatrix} 1.695\\ 0.718 \end{pmatrix} \tag{10}$$

See figure 1




Figure 1: