

All Pairs Shortest Paths

Book: Cormen; Chapter: 25

Floyd-Warshall : Section: 25.2

All-Pairs Shortest Paths Problem

- Problem:

Given a directed graph $G=(V, E)$, and a weight function $w: E \rightarrow \mathbb{R}$, for each pair of vertices u, v , compute the shortest path weight $\delta(u, v)$, and a shortest path if exists.

- Output:

- A $V \times V$ matrix $D = (d_{ij})$, where, d_{ij} contains the shortest path weight from vertex i to vertex j .
- A $V \times V$ matrix $\Pi = (\pi_{ij})$, where, π_{ij} is NIL if either $i=j$ or there is no path from i to j , otherwise π_{ij} is the predecessor of j on some shortest path from i .

Adjacency Matrix: $W = (w_{ij})$

$$w_{ij} = \begin{cases} 0, & \text{if } i = j \\ \text{the weight of edge } (i, j) & \text{if } i \neq j \text{ and } (i, j) \in E \\ \infty & \text{if } i \neq j \text{ and } (i, j) \notin E \end{cases}$$

Methods

- ❑ Single-source shortest-path algorithms
- ❑ Direct methods
 - Matrix multiplication
 - **Floyd-Warshall Algorithm**
 - Johnson's Algorithm for sparse graphs
- ❑ Transitive closure (Floyd-Warshall Algorithm)

Floyd-Warshall Algorithm

Structure of a Shortest Path





Figure 25.3 Path p is a shortest path from vertex i to vertex j , and k is the highest-numbered intermediate vertex of p . Path p_1 , the portion of path p from vertex i to vertex k , has all intermediate vertices in the set $\{1, 2, \dots, k-1\}$. The same holds for path p_2 from vertex k to vertex j .

Floyd-Warshall Algorithm

Structure of a Shortest Path (Explanation)



Floyd-Warshall Algorithm

Recursive Solution

- $d_{ij}^{(k)}$:
shortest path weight from i to j
with intermediate vertices **(excluding i, j)**
from the set $\{1, 2, \dots, k\}$
- $d_{ij}^{(0)} = w_{ij}$;
- $d_{ij}^{(k)} = \min(d_{ij}^{(k-1)}, d_{ik}^{(k-1)} + d_{kj}^{(k-1)})$; if $k \geq 1$

Result: $D^{(n)} = (d_{ij}^{(n)})$

Constructing a Shortest Path

- For $k = 0$

$$\pi_{ij}^{(0)} = \begin{cases} \text{NIL} & \text{if } i = j \text{ or } w_{ij} = \infty \\ i & \text{if } i \neq j \text{ and } w_{ij} < \infty \end{cases}$$

- For $k \geq 1$

$$\pi_{ij}^{(k)} = \begin{cases} \pi_{ij}^{(k-1)} & \text{if } d_{ij}^{(k-1)} \leq d_{ik}^{(k-1)} + d_{kj}^{(k-1)} \\ \pi_{kj}^{(k-1)} & \text{if } d_{ij}^{(k-1)} > d_{ik}^{(k-1)} + d_{kj}^{(k-1)} \end{cases}$$

Compute Weights and Construct Paths

□ Solve the problem stage by stage:

➤ $D(o), \Pi(o)$

➤ $D^{(1)}, \Pi^{(1)}$


➤ $D^{(2)}, \Pi^{(2)}$

➤ \dots, \dots

➤ $D^{(n)}, \Pi^{(n)}$

□ where $D^{(k)}$ contains the shortest path weight with all the **intermediate vertices** from set $\{1, 2, \dots, k\}$.

Example (Page - 690)



Example (Page - 696)

$$D^{(0)} = \begin{pmatrix} 0 & 3 & 8 & \infty & -4 \\ \infty & 0 & \infty & 1 & 7 \\ \infty & 4 & 0 & \infty & \infty \\ 2 & \infty & -5 & 0 & \infty \\ \infty & \infty & \infty & 6 & 0 \end{pmatrix}$$

$$\Pi^{(0)} = \begin{pmatrix} \text{NIL} & 1 & 1 & \text{NIL} & 1 \\ \text{NIL} & \text{NIL} & \text{NIL} & 2 & 2 \\ \text{NIL} & 3 & \text{NIL} & \text{NIL} & \text{NIL} \\ 4 & \text{NIL} & 4 & \text{NIL} & \text{NIL} \\ \text{NIL} & \text{NIL} & \text{NIL} & 5 & \text{NIL} \end{pmatrix}$$

$$D^{(1)} = \begin{pmatrix} 0 & 3 & 8 & \infty & -4 \\ \infty & 0 & \infty & 1 & 7 \\ \infty & 4 & 0 & \infty & \infty \\ 2 & 5 & -5 & 0 & -2 \\ \infty & \infty & \infty & 6 & 0 \end{pmatrix}$$

$$\Pi^{(1)} = \begin{pmatrix} \text{NIL} & 1 & 1 & \text{NIL} & 1 \\ \text{NIL} & \text{NIL} & \text{NIL} & 2 & 2 \\ \text{NIL} & 3 & \text{NIL} & \text{NIL} & \text{NIL} \\ 4 & 1 & 4 & \text{NIL} & 1 \\ \text{NIL} & \text{NIL} & \text{NIL} & 5 & \text{NIL} \end{pmatrix}$$

Example (Page - 696)

$$D^{(2)} = \begin{pmatrix} 0 & 3 & 8 & 4 & -4 \\ \infty & 0 & \infty & 1 & 7 \\ \infty & 4 & 0 & 5 & 11 \\ 2 & 5 & -5 & 0 & -2 \\ \infty & \infty & \infty & 6 & 0 \end{pmatrix}$$

$$\Pi^{(2)} = \begin{pmatrix} NIL & 1 & 1 & 2 & 1 \\ NIL & NIL & NIL & 2 & 2 \\ NIL & 3 & NIL & 2 & 2 \\ 4 & 1 & 4 & NIL & 1 \\ NIL & NIL & NIL & 5 & NIL \end{pmatrix}$$

$$D^{(3)} = \begin{pmatrix} 0 & 3 & 8 & 4 & -4 \\ \infty & 0 & \infty & 1 & 7 \\ \infty & 4 & 0 & 5 & 11 \\ 2 & -1 & -5 & 0 & -2 \\ \infty & \infty & \infty & 6 & 0 \end{pmatrix}$$

$$\Pi^{(3)} = \begin{pmatrix} NIL & 1 & 1 & 2 & 1 \\ NIL & NIL & NIL & 2 & 2 \\ NIL & 3 & NIL & 2 & 2 \\ 4 & 3 & 4 & NIL & 1 \\ NIL & NIL & NIL & 5 & NIL \end{pmatrix}$$

Example (Page - 696)


$$D^{(4)} = \begin{pmatrix} 0 & 3 & -1 & 4 & -4 \\ 3 & 0 & -4 & 1 & -1 \\ 7 & 4 & 0 & 5 & 3 \\ 2 & -1 & -5 & 0 & -2 \\ 8 & 5 & 1 & 6 & 0 \end{pmatrix}$$

$$\Pi^{(4)} = \begin{pmatrix} NIL & 1 & 4 & 2 & 1 \\ 4 & NIL & 4 & 2 & 1 \\ 4 & 3 & NIL & 2 & 1 \\ 4 & 3 & 4 & NIL & 1 \\ 4 & 3 & 4 & 5 & NIL \end{pmatrix}$$

$$D^{(5)} = \begin{pmatrix} 0 & 1 & -3 & 2 & -4 \\ 3 & 0 & -4 & 1 & -1 \\ 7 & 4 & 0 & 5 & 3 \\ 2 & -1 & -5 & 0 & -2 \\ 8 & 5 & 1 & 6 & 0 \end{pmatrix}$$

$$\Pi^{(5)} = \begin{pmatrix} NIL & 3 & 4 & 5 & 1 \\ 4 & NIL & 4 & 2 & 1 \\ 4 & 3 & NIL & 2 & 1 \\ 4 & 3 & 4 & NIL & 1 \\ 4 & 3 & 4 & 5 & NIL \end{pmatrix}$$

Shortest Path from 1 to 2



Algorithm (Page - 695)

FLOYD-WARSHALL(W)

1 $n = W.rows$

2 $D^{(0)} = W$

3 **for** $k = 1$ **to** n

4 let $D^{(k)} = (d_{ij}^{(k)})$ be a new $n \times n$ matrix

5 **for** $i = 1$ **to** n

6 **for** $j = 1$ **to** n

7 $d_{ij}^{(k)} = \min(d_{ij}^{(k-1)}, d_{ik}^{(k-1)} + d_{kj}^{(k-1)})$

8 **return** $D^{(n)}$

Print Shortest Paths (Page - 685)

PRINT-ALL-PAIRS-SHORTEST-PATH(Π, i, j)

```
1  if  $i = j$ 
2      then print  $i$ 
3      else if  $\pi_{ij} = \text{NIL}$ 
4          then print "no path from"  $i$  "to"  $j$  "exists"
5          else PRINT-ALL-PAIRS-SHORTEST-PATH( $\Pi, i, \pi_{ij}$ )
6              print  $j$ 
```


Thank You

Stay Safe