

Lec 10 多元函数微分法习题课 (1)

10.1 例题

例 10.1 设方程: $u^3 - 3(x+y)u^2 + z^3 = 0$ 确定了隐函数 $u = f(x, y, z)$, 求 du .

解 解法一:

原方程两边取全微分 d 得

$$\begin{aligned}d(u^3 - 3(x+y)u^2 + z^3) &= d(0) = 0 \\ \Rightarrow d(u^3) - 3d((x+y)u^2) + d(z^3) &= 0 \\ \Rightarrow 3u^2 du - 3u^2(dx + dy) - 3(x+y)2u du + 3z^2 dz &= 0\end{aligned}$$

整理得

$$du = \frac{3u^2(dx + dy) + 3z^2 dz}{3u^2 + 6(x+y)u} = \frac{u^2 dx + u^2 dy - z^2 dz}{u^2 - 2(x+y)u}$$

解 解法二:

令 $F(x, y, z, u) = u^3 - 3(x+y)u^2 + z^3$, 其中 x, y, z, u 地位相同, 则 $F'_u = 3u^2 - 6(x+y)u, F'_x = -3u^2, F'_y = -3u^2, F'_z = 3z^2$. 从而

$$\begin{aligned}\frac{\partial u}{\partial x} &= -\frac{F'_x}{F'_u} = \frac{u^2}{-2(x+y)u + u^2}; \\ \frac{\partial u}{\partial y} &= -\frac{F'_y}{F'_u} = \frac{u^2}{-2(x+y)u + u^2}; \\ \frac{\partial u}{\partial z} &= \frac{F'_z}{F'_u} = \frac{z^2}{-2(x+y)u + u^2}.\end{aligned}$$

因此

$$du = \frac{\partial u}{\partial x} dx + \frac{\partial u}{\partial y} dy + \frac{\partial u}{\partial z} dz = \frac{u^2 dx + u^2 dy - z^2 dz}{u^2 - 2(x+y)u}.$$

解 解法三:

原方程两边分别对 x, y, z 求偏导数, 解出 $\frac{\partial u}{\partial x}, \frac{\partial u}{\partial y}, \frac{\partial u}{\partial z}$ 再代入 $du = \frac{\partial u}{\partial x} dx + \frac{\partial u}{\partial y} dy + \frac{\partial u}{\partial z} dz$ 即可.

例 10.2 试证明方程

$$\frac{\partial^2 u}{\partial x^2} + 2\frac{\partial^2 u}{\partial x \partial y} - 3\frac{\partial^2 u}{\partial y^2} + 2\frac{\partial u}{\partial x} + 6\frac{\partial u}{\partial y} = 0$$

在线性变换

$$\begin{cases} \xi = x + y, \\ \eta = 3x - y \end{cases}$$

下可以化简为

$$\frac{\partial^2 u}{\partial \xi \partial \eta} + \frac{1}{2} \frac{\partial u}{\partial \xi} = 0$$

解 证法一:

从线性变换 $\begin{cases} \xi = x + y, \\ \eta = 3x - y \end{cases}$ 可得 $x = \frac{1}{4}(\xi + \eta), y = \frac{1}{4}(3\xi - \eta)$, 因此 $\frac{\partial x}{\partial \xi} = \frac{1}{4}, \frac{\partial x}{\partial \eta} = \frac{1}{4}, \frac{\partial y}{\partial \xi} = \frac{3}{4}, \frac{\partial y}{\partial \eta} = -\frac{1}{4}$. 由此得

$$\begin{aligned} \frac{\partial u}{\partial \xi} &= \frac{\partial u}{\partial x} \frac{\partial x}{\partial \xi} + \frac{\partial u}{\partial y} \frac{\partial y}{\partial \xi} = \frac{1}{4} \frac{\partial u}{\partial x} + \frac{3}{4} \frac{\partial u}{\partial y} \\ \Rightarrow \frac{\partial^2 u}{\partial \eta \partial \xi} &= \left(\frac{\partial u}{\partial \xi} \right)'_{\eta} = \left(\frac{1}{4} \frac{\partial u}{\partial x} \right)'_{\eta} + \left(\frac{3}{4} \frac{\partial u}{\partial y} \right)'_{\eta} \\ &= \frac{1}{4} \left(\frac{\partial^2 u}{\partial x^2} \frac{\partial x}{\partial \eta} + \frac{\partial^2 u}{\partial x \partial y} \frac{\partial y}{\partial \eta} \right) + \frac{3}{4} \left(\frac{\partial^2 u}{\partial x \partial y} \frac{\partial x}{\partial \eta} + \frac{\partial^2 u}{\partial y^2} \frac{\partial y}{\partial \eta} \right) \\ &= \frac{1}{4} \left(\frac{\partial^2 u}{\partial x^2} \frac{1}{4} + \frac{\partial^2 u}{\partial x \partial y} \left(-\frac{1}{4} \right) \right) + \frac{3}{4} \left(\frac{\partial^2 u}{\partial x \partial y} \frac{1}{4} + \frac{\partial^2 u}{\partial y^2} \left(-\frac{1}{4} \right) \right) \\ &= \frac{1}{16} \left(\frac{\partial^2 u}{\partial x^2} + 2 \frac{\partial^2 u}{\partial x \partial y} - 3 \frac{\partial^2 u}{\partial y^2} \right) \end{aligned}$$

即有

$$\begin{cases} \frac{\partial^2 u}{\partial x^2} + 2 \frac{\partial^2 u}{\partial x \partial y} - 3 \frac{\partial^2 u}{\partial y^2} = 16 \frac{\partial^2 u}{\partial \eta \partial \xi} \\ \frac{\partial u}{\partial x} + 3 \frac{\partial u}{\partial y} = 4 \frac{\partial u}{\partial \xi} \end{cases}$$

故原偏微分方程化简为

$$\begin{aligned} 16 \frac{\partial^2 u}{\partial \eta \partial \xi} + 2 \left(4 \frac{\partial u}{\partial \xi} \right) &= 0 \\ \Rightarrow \frac{\partial^2 u}{\partial \eta \partial \xi} + \frac{1}{2} \frac{\partial u}{\partial \xi} &= 0 \end{aligned}$$

解 解法二:

从线性变换 $\begin{cases} \xi = x + y, \\ \eta = 3x - y \end{cases} \Rightarrow \frac{\partial \xi}{\partial x} = 1, \frac{\partial \xi}{\partial y} = 1, \frac{\partial \eta}{\partial x} = 3, \frac{\partial \eta}{\partial y} = -1$. 而 $u(x, y)$ 通过中间变量可视为 ξ, η 的函数, 从而

$$\begin{cases} \frac{\partial u}{\partial x} = \frac{\partial u}{\partial \xi} \frac{\partial \xi}{\partial x} + \frac{\partial u}{\partial \eta} \frac{\partial \eta}{\partial x} = \frac{\partial u}{\partial \xi} + 3 \frac{\partial u}{\partial \eta}, \\ \frac{\partial u}{\partial y} = \frac{\partial u}{\partial \xi} \frac{\partial \xi}{\partial y} + \frac{\partial u}{\partial \eta} \frac{\partial \eta}{\partial y} = \frac{\partial u}{\partial \xi} - \frac{\partial u}{\partial \eta}. \end{cases},$$

从而

$$\begin{cases} \frac{\partial^2 u}{\partial x^2} = \left(\frac{\partial u}{\partial \xi} \right)'_x + 3 \left(\frac{\partial u}{\partial \eta} \right)'_x = \left(\frac{\partial^2 u}{\partial \xi^2} + 3 \frac{\partial^2 u}{\partial \xi \partial \eta} \right) + 3 \left(\frac{\partial^2 u}{\partial \xi \partial \eta} + 3 \frac{\partial^2 u}{\partial \eta^2} \right), \\ \frac{\partial^2 u}{\partial x \partial y} = \left(\frac{\partial u}{\partial \xi} - \frac{\partial u}{\partial \eta} \right)'_x = \left(\frac{\partial^2 u}{\partial \xi^2} + 3 \frac{\partial^2 u}{\partial \xi \partial \eta} \right) - \left(\frac{\partial^2 u}{\partial \xi \partial \eta} + 3 \frac{\partial^2 u}{\partial \eta^2} \right), \\ \frac{\partial^2 u}{\partial y^2} = \left(\frac{\partial u}{\partial \xi} - \frac{\partial u}{\partial \eta} \right)'_y = \left(\frac{\partial^2 u}{\partial \xi^2} - \frac{\partial^2 u}{\partial \xi \partial \eta} \right) - \left(\frac{\partial^2 u}{\partial \xi \partial \eta} - \frac{\partial^2 u}{\partial \eta^2} \right). \end{cases}$$

即

$$\begin{cases} \frac{\partial^2 u}{\partial x^2} = \frac{\partial^2 u}{\partial \xi^2} + 6 \frac{\partial^2 u}{\partial \xi \partial \eta} + 9 \frac{\partial^2 u}{\partial \eta^2}, \\ \frac{\partial^2 u}{\partial x \partial y} = \frac{\partial^2 u}{\partial \xi^2} + 2 \frac{\partial^2 u}{\partial \xi \partial \eta} - 3 \frac{\partial^2 u}{\partial \eta^2}, \\ \frac{\partial^2 u}{\partial y^2} = \frac{\partial^2 u}{\partial \xi^2} - 2 \frac{\partial^2 u}{\partial \xi \partial \eta} + \frac{\partial^2 u}{\partial \eta^2}. \end{cases}$$

且

$$2 \frac{\partial u}{\partial x} + 6 \frac{\partial u}{\partial y} = 2 \left(\frac{\partial u}{\partial \xi} + 3 \frac{\partial u}{\partial \eta} \right) + 6 \left(\frac{\partial u}{\partial \xi} - \frac{\partial u}{\partial \eta} \right) = 8 \frac{\partial u}{\partial \xi},$$

从而原方程化为

$$\frac{\partial^2 u}{\partial \xi^2} (1 + 2 - 3) + \frac{\partial^2 u}{\partial \xi \partial \eta} (6 + 4 + 6) + \frac{\partial^2 u}{\partial \eta^2} (9 - 6 - 3) + 8 \frac{\partial u}{\partial \xi} = 0,$$

即

$$\frac{\partial^2 u}{\partial \xi \partial \eta} + \frac{1}{2} \frac{\partial u}{\partial \xi} = 0.$$

例 10.3 设 $u = f(x, y, z)$, $\varphi(x^2, e^y, z) = 0$, $y = \sin x$, 且 $f, \varphi \in C^1$, $\frac{\partial \varphi}{\partial z} \neq 0$, 求 $\frac{du}{dx}$.

解 从 $\varphi(x^2, e^{\sin x}, z) = 0$ 及 $\varphi'_z \neq 0$ 可知, 由方程 $\varphi(x^2, e^{\sin x}, z) = 0$ 可确定 z 是 x, y 的隐函数, 从而 z 是 x 的复合函数. 故从 $u = f(x, y, z)$ 知, u 是 x 的一元函数.

注 助教注: 这个地方可以理解为由隐函数定理 $F(x, y, z) = \varphi(x^2, e^y, z) = 0$, 确定了隐函数 $z = z(x, y)$, 从而 $u = f(x, y, z) = f(x, y, z(x, y)) = f(x, \sin x, z(x, \sin x))$ 确定了 u 是 x 的函数.

$$\frac{du}{dx} = f'_1 \cdot 1 + f'_2 \cdot y'_x + f'_3 \cdot z'_x = f'_1 + f'_2 \cdot \cos x \cdot 2x + f'_3 \cdot z'_x.$$

令 $F(x, y, z) = \varphi(x^2, e^y, z)$, 则

$$\begin{cases} F'_x(x, y, z) = \varphi'_1 \cdot 2x + \varphi'_2 e^{\sin x} \cos x, \\ F'_y(x, y, z) = \varphi'_3 \cdot 1 = \varphi'_3. \end{cases}$$

故 $z'_x = -\frac{F'_x}{F'_y} = -\frac{\varphi'_1 \cdot 2x + \varphi'_2 e^{\sin x} \cos x}{\varphi'_3}$. 代入 $\frac{du}{dx}$ 即有

$$\frac{du}{dx} = f'_1 + f'_2 \cdot y'_x + f'_3 \cdot \left(-\frac{\varphi'_1 \cdot 2x + \varphi'_2 e^{\sin x} \cos x}{\varphi'_3} \right).$$

例 10.4 证明: 全微分也具有一阶微分形式不变性, 即, 若 $f(x, y)$ 可微, 则不论 x, y 是自变量还是中间变量, 则 $z = f(x, y)$, 总有

$$dz = df(x, y) = \frac{\partial z}{\partial x} dx + \frac{\partial z}{\partial y} dy = f'_x dx + f'_y dy.$$

证明

1. 当 x, y 是自变量时, 显然有 $dz = f'_x dx + f'_y dy$.

2. 当 x, y 是中间变量时, 设 $\begin{cases} x = g(s, t), \\ y = h(s, t), \end{cases}$ 可微, 且 $f(g(s, t), h(s, t))$ 有意义时, z 通过中间

变量 x, y 成为 s, t 的复合函数, 且有求偏导数的链式法则如下:

$$\begin{aligned} dx &= \frac{\partial x}{\partial s} ds + \frac{\partial x}{\partial t} dt, \\ dy &= \frac{\partial y}{\partial s} ds + \frac{\partial y}{\partial t} dt. \end{aligned}$$

且

$$\begin{aligned} dz &= \frac{\partial z}{\partial s} ds + \frac{\partial z}{\partial t} dt = \left(\frac{\partial z}{\partial x} \cdot \frac{\partial x}{\partial s} + \frac{\partial z}{\partial y} \cdot \frac{\partial y}{\partial s} \right) ds + \left(\frac{\partial z}{\partial x} \cdot \frac{\partial x}{\partial t} + \frac{\partial z}{\partial y} \cdot \frac{\partial y}{\partial t} \right) dt \\ &= \frac{\partial z}{\partial x} \left(\frac{\partial x}{\partial s} ds + \frac{\partial x}{\partial t} dt \right) + \frac{\partial z}{\partial y} \left(\frac{\partial y}{\partial s} ds + \frac{\partial y}{\partial t} dt \right) \\ &= \frac{\partial z}{\partial x} dx + \frac{\partial z}{\partial y} dy. \end{aligned}$$

即 x, y 是中间变量时, 也有 $dz = f'_x dx + f'_y dy$.

注 利用全微分的一阶微分形式不变性, 可导出多元可微函数的如下的微分四则运算法则:

1. $d(u \pm v) = du \pm dv$;
2. $d(uv) = u dv + v du$;
3. $d\left(\frac{u}{v}\right) = \frac{v du - u dv}{v^2}$, 其中 u, v 均可微, 且 $v \neq 0$;

证明

1. 令 $f(u, v) = u + v$, 则 $f(u, v) \in C^1$, 从而 $f(u, v)$ 可微, 无论 u, v 是自变量还是中间变量, 总有

$$d(u + v) = df(u, v) = \frac{\partial f}{\partial u} du + \frac{\partial f}{\partial v} dv = du + dv$$

从而有 $d(u \pm v) = du \pm dv$. 这里 d 是全微分.

2. 令 $f(u, v) = uv$, 则 $f(u, v) \in C^1$, 从而 $f(u, v)$ 可微, 无论 u, v 是自变量还是中间变量, 总有

$$d(uv) = df(u, v) = \frac{\partial f}{\partial u} du + \frac{\partial f}{\partial v} dv = v du + u dv$$

从而有 $d(uv) = u dv + v du$.

3. 令 $f(u, v) = \frac{u}{v}$, 则 $f(u, v) \in C^1$, 从而 $f(u, v)$ 可微, 无论 u, v 是自变量还是中间变量, 总有

$$d\left(\frac{u}{v}\right) = df(u, v) = \frac{\partial f}{\partial u} du + \frac{\partial f}{\partial v} dv = \frac{1}{v} du + \left(-\frac{u}{v^2}\right) dv = \frac{v du - u dv}{v^2}$$

注 二阶及以上的微分通常没有形式不变性, 具体而言, 设 $f(x, y) \in C^2$, 则 $z = f(x, y) \Rightarrow dz = \frac{\partial z}{\partial x} dx + \frac{\partial z}{\partial y} dy$.

$$\begin{aligned} d(dz) &:= d^2z = d\left(\frac{\partial z}{\partial x} dx + \frac{\partial z}{\partial y} dy\right) \\ &= \left(\frac{\partial z}{\partial x} dx + \frac{\partial z}{\partial y} dy\right)'_x dx + \left(\frac{\partial z}{\partial x} dx + \frac{\partial z}{\partial y} dy\right)'_y dy \\ &= \left(\frac{\partial^2 z}{\partial x^2} dx + \frac{\partial^2 z}{\partial x \partial y} dy\right) dx + \left(\frac{\partial^2 z}{\partial x \partial y} dx + \frac{\partial^2 z}{\partial y^2} dy\right) dy \\ &= \frac{\partial^2 z}{\partial x^2} (dx)^2 + 2 \frac{\partial^2 z}{\partial x \partial y} dx dy + \frac{\partial^2 z}{\partial y^2} (dy)^2. \end{aligned}$$

d^2z 是 x, y 是自变量时的 $z = f(x, y)$ 的二阶微分, 而 d^2z 是 x, y 是中间变量时的 $z = f(x, y)$ 的二阶微分, 二者通常不相等.

例 10.5 设 $u = u(x, y), v = v(x, y)$ 是由方程组

$$\begin{cases} u = f(ux, v + y), \\ v = g(u - x, v^2y) \end{cases}$$

所确定的隐函数组, 求变换 $\begin{cases} u = u(x, y), \\ v = v(x, y) \end{cases}$ 的 Jacobi 行列式:

$$\begin{vmatrix} u'_x & u'_y \\ v'_x & v'_y \end{vmatrix} := \frac{\partial(u, v)}{\partial(x, y)} \quad f, g \in C^1.$$

解 令 $A = ux, B = v + y, E = u - x, F = v^2y$, 则方程组可化为

$$\begin{cases} u = f(A, B), \\ v = g(E, F). \end{cases}$$

方程组两边关于 x 求偏导

$$\begin{cases} u'_x = f'_1 \cdot (u + xu'_x) + f'_2 \cdot (v'_x + 0), \\ v'_x = g'_1 \cdot (u'_x - 1) + g'_2 \cdot 2vv'_xy. \end{cases}$$

标准化为

$$\begin{cases} (xf'_1 - 1)u'_x + f'_2v'_x = -f'_1u, \\ g'_1u'_x + (2vg'_2y - 1)v'_x = g'_1. \end{cases}$$

令 $D = \begin{vmatrix} xf'_1 - 1 & f'_2 \\ g'_1 & 2vg'_2y - 1 \end{vmatrix}$, 则 $D \neq 0$, 再令

$$D_1 = \begin{vmatrix} -f'_1u & f'_2 \\ g'_1 & 2vg'_2y - 1 \end{vmatrix}, \quad D_2 = \begin{vmatrix} xf'_1 - 1 & -f'_1u \\ g'_1 & g'_1 \end{vmatrix},$$

由克莱姆法则可得

$$u'_x = \frac{D_1}{D}, \quad v'_x = \frac{D_2}{D}.$$

方程组 $\begin{cases} u = f(A, B), \\ v = g(E, F) \end{cases}$ 两边同时对 y 求偏导, 可得

$$\begin{cases} u'_y = f'_1 \cdot u'_y \cdot x + f'_2 \cdot (v'_y + 1), \\ v'_y = g'_1 \cdot u'_y + g'_2 \cdot (2vv'_yy + 2v^2). \end{cases}$$

标准化为

$$\begin{cases} (xf'_1 - 1)u'_y + f'_2v'_y = -f'_2, \\ g'_1u'_y + (2vg'_2y - 1)v'_y = 2vg'_2. \end{cases}$$

令 $D = \begin{vmatrix} xf'_1 - 1 & f'_2 \\ g'_1 & 2vg'_2y - 1 \end{vmatrix}$, 则 $D \neq 0$, 再令

$$\tilde{D}_1 = \begin{vmatrix} -f'_2 & f'_2 \\ v^2g'_2 & 2vg'_2y - 1 \end{vmatrix}, \quad \tilde{D}_2 = \begin{vmatrix} xf'_1 - 1 & -f'_2 \\ g'_1 & g'_1v^2 \end{vmatrix},$$

由克莱姆法则可得

$$u'_y = \frac{\tilde{D}_1}{D}, \quad v'_y = \frac{\tilde{D}_2}{D}.$$

从而

$$\frac{\partial(u, v)}{\partial(x, y)} = \begin{vmatrix} u'_x & u'_y \\ v'_x & v'_y \end{vmatrix} = \frac{\begin{vmatrix} D_1 & \tilde{D}_1 \\ D_2 & \tilde{D}_2 \end{vmatrix}}{D}$$

例 10.6 设 $\begin{cases} u = u(x, y), \\ v = v(x, y) \end{cases}$ 是由方程组

$$\begin{cases} F(x, y, u, v) = 0, \\ G(x, y, u, v) = 0 \end{cases}$$

确定的隐函数组, $F, G \in C^1$, 且 $\frac{\partial(F, G)}{\partial(u, v)} \neq 0$, 求 du, dv .

解 解法一:

$$du = \frac{\partial u}{\partial x} dx + \frac{\partial u}{\partial y} dy, \quad dv = \frac{\partial v}{\partial x} dx + \frac{\partial v}{\partial y} dy, \quad \text{对 } \begin{cases} F(x, y, u, v) = 0, \\ G(x, y, u, v) = 0 \end{cases} \quad \text{两边关于 } x \text{ 求偏导, 可}$$

得

$$\begin{cases} F'_x \cdot 1 + F'_u \cdot \frac{\partial u}{\partial x} + F'_v \cdot \frac{\partial v}{\partial x} = 0, \\ G'_x \cdot 1 + G'_u \cdot \frac{\partial u}{\partial x} + G'_v \cdot \frac{\partial v}{\partial x} = 0. \end{cases}$$

标准化为

$$\begin{cases} F'_u \frac{\partial u}{\partial x} + F'_v \frac{\partial v}{\partial x} = -F'_x, \\ G'_u \frac{\partial u}{\partial x} + G'_v \frac{\partial v}{\partial x} = -G'_x. \end{cases}$$

令 $D = \begin{vmatrix} F'_u & F'_v \\ G'_u & G'_v \end{vmatrix}$, 则 $D \neq 0$, 再令

$$D_1 = \begin{vmatrix} -F'_x & F'_v \\ -G'_x & G'_v \end{vmatrix}, \quad D_2 = \begin{vmatrix} F'_u & -F'_x \\ G'_u & -G'_x \end{vmatrix},$$

此时注意到 $D_1 = \begin{vmatrix} F'_v & F'_x \\ G'_v & G'_x \end{vmatrix} = \frac{\partial(F, G)}{\partial(v, x)}$, $D_2 = \begin{vmatrix} F'_u & F'_v \\ G'_u & G'_v \end{vmatrix} = \frac{\partial(F, G)}{\partial(u, v)}$, 由克莱姆法则可得

$$\begin{aligned}\frac{\partial u}{\partial x} &= \frac{D_1}{D} = \frac{\frac{\partial(F, G)}{\partial(v, x)}}{\frac{\partial(F, G)}{\partial(u, v)}}, \\ \frac{\partial v}{\partial x} &= \frac{D_2}{D} = \frac{\frac{\partial(F, G)}{\partial(u, x)}}{\frac{\partial(F, G)}{\partial(u, v)}}.\end{aligned}$$

对原方程组两边对 y 求偏导, 同样可得

$$\begin{aligned}\frac{\partial u}{\partial y} &= \frac{\frac{\partial(F, G)}{\partial(v, y)}}{\frac{\partial(F, G)}{\partial(u, v)}}, \\ \frac{\partial v}{\partial y} &= \frac{\frac{\partial(F, G)}{\partial(u, y)}}{\frac{\partial(F, G)}{\partial(u, v)}}.\end{aligned}$$

从而

$$\begin{aligned}du &= \frac{\partial u}{\partial x} dx + \frac{\partial u}{\partial y} dy = \frac{\frac{\partial(F, G)}{\partial(v, x)} dx + \frac{\partial(F, G)}{\partial(v, y)} dy}{\frac{\partial(F, G)}{\partial(u, v)}}, \\ dv &= \frac{\partial v}{\partial x} dx + \frac{\partial v}{\partial y} dy = \frac{\frac{\partial(F, G)}{\partial(u, x)} dx + \frac{\partial(F, G)}{\partial(u, y)} dy}{\frac{\partial(F, G)}{\partial(u, v)}}.\end{aligned}$$

解 解法二:

对原方程两边同时取全微分, 可得


$$\begin{cases} F'_x dx + F'_y dy + F'_u du + F'_v dv = 0, \\ G'_x dx + G'_y dy + G'_u du + G'_v dv = 0. \end{cases}$$

以 du, dv 为变量. 依 cramer 法则, 解得

$$\begin{aligned}du &= \frac{D_1}{D} = \frac{\partial u}{\partial x} dx + \frac{\partial u}{\partial y} dy = \frac{\frac{\partial(F, G)}{\partial(v, x)} dx + \frac{\partial(F, G)}{\partial(v, y)} dy}{\frac{\partial(F, G)}{\partial(u, v)}}, \\ dv &= \frac{D_2}{D} = \frac{\partial v}{\partial x} dx + \frac{\partial v}{\partial y} dy = \frac{\frac{\partial(F, G)}{\partial(u, x)} dx + \frac{\partial(F, G)}{\partial(u, y)} dy}{\frac{\partial(F, G)}{\partial(u, v)}}.\end{aligned}$$

其中,

$$D_1 = \begin{vmatrix} -(F'_x dx + F'_y dy) & F'_v \\ -(G'_x dx + G'_y dy) & G'_v \end{vmatrix}, \quad D_2 = \begin{vmatrix} F'_u & -(F'_x dx + F'_y dy) \\ G'_u & -(G'_x dx + G'_y dy) \end{vmatrix}, \quad D = \begin{vmatrix} F'_u & F'_v \\ G'_u & G'_v \end{vmatrix} = \frac{\partial(F, G)}{\partial(u, v)}.$$

 **作业** ex9.2:31;ex9.3:6,7,8,10,11(1),14.