Convergence of Allen-Cahn equations to multi-phase mean curvature flow

Pascal Maurice Steinke

Born 22.05.1999 in Gießen, Germany 09.09.2022

Master's Thesis Mathematics

Advisor: Prof. Dr. Tim Laux

Second Advisor: Prof. Dr. X Y

Institute for Applied Mathematics

Mathematisch-Naturwissenschaftliche Fakultät der Rheinischen Friedrich-Wilhelms-Universität Bonn

Contents

1 The Allen–Cahn equation

1

1 The Allen-Cahn equation

Let $\Lambda>0$ and define the flat torus $\mathbb{T}=[0,\Lambda)^d\subset\mathbb{R}^d$, where we as usual assume periodic boundary conditions. Then for $u\colon [0,\infty)\times\mathbb{T}\to\mathbb{R}^N$ and some potential $W\colon\mathbb{R}^N\to[0,\infty)$, the Allen–Cahn equation with parameter $\varepsilon>0$ is given by

$$\partial_t u = \Delta u - \frac{1}{\varepsilon^2} \partial_u W(u). \tag{1.1}$$

To understand this equation better, we consider the ${\it Cahn-Hilliard\ energy}$ assigns to u for a fixed time the real number

$$E_{\varepsilon}(u) := \int_{\mathbb{T}} \frac{1}{\varepsilon} W(u) + \frac{\varepsilon}{2} \|\nabla u\|^2 dx.$$
 (1.2)

If everything is nice and smooth, we can compute that under the assumption that u satisfies equation 1.1, we have that

$$\frac{\mathrm{d}}{\mathrm{d}t} \, \mathrm{E}_{\varepsilon}(u) = \int stuff$$

Bibliography

[LS16] Tim Laux and Thilo Simon. "Convergence of the Allen-Cahn Equation to Multiphase Mean Curvature Flow". In: Communications on Pure and Applied Mathematics 71 (June 2016). DOI: 10.1002/cpa.21747.