

# Causal Data Science: A general framework for data fusion and causal inference

Elias Bareinboim

Columbia University

Twitter: @eliasbareinboim

Machine Learning Research School  
Bangkok, August, 2019

# Logistics

---

# Logistics

---

- Website
  - <https://causalai.net/mlss19/> (soon)
  - slides – l: mlss19, p: mlss19

# Logistics

---

- Website
  - <https://causalai.net/mlss19/> (soon)
  - slides – l: mlss19, p: mlss19
- Instructor

# Logistics

---

- Website
  - <https://causalai.net/mlss19/> (soon)
  - slides – l: mlss19, p: mlss19
- Instructor
- Census

# Logistics

---

- Website
  - <https://causalai.net/mlss19/> (soon)
  - slides – l: mlss19, p: mlss19
- Instructor
  - 1. prob. & statistics
  - 2. graphical models
  - 3. d-separation
  - 4. experimental design
  - 5. causal inference
- Census

# Logistics

---

- Website
  - <https://causalai.net/mlss19/> (soon)
  - slides – l: mlss19, p: mlss19
- Instructor
  - 1. prob. & statistics
  - 2. graphical models
  - 3. d-separation
  - 4. experimental design
  - 5. causal inference
- Census
- Background

# Logistics

---

- Website
  - <https://causalai.net/mlss19/> (soon)
  - slides – l: mlss19, p: mlss19
- Instructor
- Census
- Background
  - 1. prob. & statistics
  - 2. graphical models
  - 3. d-separation
  - 4. experimental design
  - 5. causal inference
- Software (email for access)

# Logistics

---

- Website
    - <https://causalai.net/mlss19/> (soon)
    - slides – l: mlss19, p: mlss19
  - Instructor
  - Census
  - Background
  - Software (email for access)
  - Bibliography
1. prob. & statistics
  2. graphical models
  3. d-separation
  4. experimental design
  5. causal inference

# References

---

# References

---

- Causal Inference and the Data-fusion Problem
  - E. Bareinboim and J. Pearl
  - Proc. National Academy of Sciences, 2016

# References

---

- Causal Inference and the Data-fusion Problem
  - E. Bareinboim and J. Pearl
  - Proc. National Academy of Sciences, 2016

The screenshot shows the homepage of the Proceedings of the National Academy of Sciences (PNAS) website. The header features the PNAS logo and the full title "Proceedings of the National Academy of Sciences of the United States of America". A search bar is located in the top right corner. Below the header, a navigation menu includes "Home", "Articles" (which is the active tab), "Front Matter", "News", "Podcasts", and "Authors". A dropdown menu under "NEW RESEARCH IN" shows "Physical Sciences" and "Social Sciences". The main content area displays the article "Causal inference and the data-fusion problem" by Elias Bareinboim and Judea Pearl, published on July 5, 2016. The article summary, authors, and publication details are visible. A "Check for updates" button is in the bottom right corner of the article box.

PNAS  
Proceedings of the  
National Academy of Sciences  
of the United States of America

Keyword, Author,

Home Articles Front Matter News Podcasts Authors

NEW RESEARCH IN Physical Sciences ▾ Social Sciences

**Causal inference and the data-fusion problem**

Elias Bareinboim and Judea Pearl  
PNAS July 5, 2016 113 (27) 7345-7352; published ahead of print July 5, 2016 <https://doi.org/10.1073/pnas.1510507113>  
Edited by Richard M. Shiffrin, Indiana University, Bloomington, IN, and approved March 15, 2016 (received for review June 29, 2015)

Check for updates

# References

---

- Causal Inference and the Data-fusion Problem
  - E. Bareinboim and J. Pearl
  - Proc. National Academy of Sciences, 2016

# References

---

- Causal Inference and the Data-fusion Problem
  - E. Bareinboim and J. Pearl
  - Proc. National Academy of Sciences, 2016

# References

---

- Causal Inference and the Data-fusion Problem
  - E. Bareinboim and J. Pearl
  - Proc. National Academy of Sciences, 2016
- The Book of Why: The New Science of Cause and Effect
  - J. Pearl and D. Mackenzie
  - Basic Books, 2018

# References

---

- Causal Inference and the Data-fusion Problem
  - E. Bareinboim and J. Pearl
  - Proc. National Academy of Sciences, 2016
- The Book of Why: The New Science of Cause and Effect
  - J. Pearl and D. Mackenzie
  - Basic Books, 2018
- Causal Inference in Statistics: A primer
  - J. Pearl, M. Glymour, N. Jewel
  - Cambridge Press, 2016

# References (cont'd)

---

# References (cont'd)

---

- Causality: Models, Reasoning, and Inference
  - J. Pearl
  - Cambridge Press, 2000

# References (cont'd)

---

- Causality: Models, Reasoning, and Inference
  - J. Pearl
  - Cambridge Press, 2000
- General Identifiability with Arbitrary Surrogate Experiments
  - S. Lee, J. Correa, and E. Bareinboim
  - Proc. of 35th UAI conference

# References (cont'd)

---

- Causality: Models, Reasoning, and Inference
  - J. Pearl
  - Cambridge Press, 2000
- General Identifiability with Arbitrary Surrogate Experiments
  - S. Lee, J. Correa, and E. Bareinboim
  - Proc. of 35th UAI conference
- Probabilistic Graphical Models
  - D. Koller and N. Friedman
  - MIT Press, 2009

# Research in CausalAI - Big Picture

# Research in CausalAI - Big Picture

## Structural Causal Models



# Research in CausalAI - Big Picture

## Structural Causal Models

### 1. Explainability

(Effect identification and decomposition, Bias Analysis and Fairness, Robustness and Generalizability)




# Research in CausalAI - Big Picture

## Structural Causal Models

### 1. Explainability

(Effect identification and decomposition, Bias Analysis and Fairness, Robustness and Generalizability)



### 2. Decision-Making


(Reinforcement Learning, Randomized Clinical Trials, Personalized Decision-Making)

# Research in CausalAI - Big Picture

## Structural Causal Models

### 1. Explainability

(Effect identification and decomposition, Bias Analysis and Fairness, Robustness and Generalizability)



### 2. Decision-Making

(Reinforcement Learning, Randomized Clinical Trials, Personalized Decision-Making)

# Research in CausalAI - Big Picture


## Structural Causal Models

### 1. Explainability

(Effect identification and decomposition, Bias Analysis and Fairness, Robustness and Generalizability)

### 2. Decision-Making

(Reinforcement Learning, Randomized Clinical Trials, Personalized Decision-Making)



Applications

# Research in CausalAI - Big Picture


## Structural Causal Models

### 1. Explainability

(Effect identification and decomposition, Bias Analysis and Fairness, Robustness and Generalizability)

### 2. Decision-Making

(Reinforcement Learning, Randomized Clinical Trials, Personalized Decision-Making)



### AI-ML:

Principles and tools for designing robust and adaptable learning systems.

# Research in CausalAI - Big Picture


## Structural Causal Models

### 1. Explainability

(Effect identification and decomposition, Bias Analysis and Fairness, Robustness and Generalizability)

### 2. Decision-Making

(Reinforcement Learning, Randomized Clinical Trials, Personalized Decision-Making)



### Data Science:

Principled (“scientific”) inferences from large data collections.

### AI-ML:

Principles and tools for designing robust and adaptable learning systems.

# Research in CausalAI - Big Picture


## Structural Causal Models

### 1. Explainability

(Effect identification and decomposition, Bias Analysis and Fairness, Robustness and Generalizability)

### 2. Decision-Making

(Reinforcement Learning, Randomized Clinical Trials, Personalized Decision-Making)



### Data Science:

Principled (“scientific”) inferences from large data collections.

### AI-ML:

Principles and tools for designing robust and adaptable learning systems.

# Research in CausalAI - Big Picture


## Structural Causal Models

### 1. Explainability

(Effect identification and decomposition, Bias Analysis and Fairness, Robustness and Generalizability)

### 2. Decision-Making

(Reinforcement Learning, Randomized Clinical Trials, Personalized Decision-Making)



## Applications

### Data Science:

Principled (“scientific”) inferences from large data collections.

### AI-ML:

Principles and tools for designing robust and adaptable learning systems.

# MOTIVATION

# The aspirations & challenges of modern Data Science (circa 2019)

---

# The aspirations & challenges of modern Data Science (circa 2019)

---

- *“The ability to take data — to be able to understand it, to extract value from it, to visualize it, to communicate it - that’s going to be hugely important in the next decades”*

Hal Varian, chief economist at Google and UC Berkeley  
Professor of Information Sciences, Business, and Economics.

# The aspirations & challenges of modern Data Science (circa 2019)

---

- *“The ability to take data — to be able to understand it, to extract value from it, to visualize it, to communicate it - that’s going to be hugely important in the next decades”*  
Hal Varian, chief economist at Google and UC Berkeley  
Professor of Information Sciences, Business, and Economics.
- *“Big data is not about the data!”*  
Gary King, Political Scientist, University Professor, Harvard University.

# The aspirations & challenges of modern Data Science (circa 2019)

---

- “*The ability to take data — to be able to understand it, to extract value from it, to visualize it, to communicate it - that's going to be hugely important in the next decades*”

Hal Varian, chief economist at Google and UC Berkeley  
Professor of Information Sciences, Business, and Economics.

- “*Big data is not about the data!*”

Gary King, Political Scientist, University Professor, Harvard University.

- “*Data Science is only as much of a science as it facilitates the interpretation of data - a two body problem, connecting data to reality*”.

Judea Pearl, Professor of Computer Science & Statistics, UCLA.

# CURRENT STATE OF AFFAIRS (REPORT FROM THE TRENCHES)

Edition: U.S. | International | MÉXICO | ARABIC  
TV: CNN | CNNI | CNN en Español | HLN

# CNN Health

SEARCH POWERED BY Google

Home TV & Video CNN Trends U.S. World Politics Justice Entertainment Tech Health Living Travel Opinion iReport Money Sports

## the chart

Dr. Sanjay Gupta Children's Health Expert Doctor Q&A Sleep Sex and You

2.3k 102 31 20 Share Tweet g+1 More sharing

CONTINUE Privacy Policy Close

### voiceFIVE

Are you aware of any of the following?

- UnitedHealthcare
- Aetna
- Humana
- Blue Cross

1 of 5 Only 5 questions? Easy. Too easy

ADVERTISEMENT

August 15th, 2013 08:00 PM ET

#### Study: Heavy coffee drinking in people under 55 linked to early death

Dr. Sanjay Gupta

<>

Edition: U.S. | International | MÉXICO | ARABIC  
TV: CNN | CNNI | CNN en Español | HLN

# CNN Health

SEARCH POWERED BY Google

Home TV & Video CNN Trends U.S. World Politics Justice Entertainment Tech Health Living Travel Opinion iReport Money Sports

## the chart

Dr. Sanjay Gupta Children's Health Expert Doctor Q&A Sleep Sex and You

2.3k 102 31 20 Share Tweet g+1 More sharing

CONTINUE Privacy Policy Close

### voiceFIVE

Are you aware of any of the following?

- UnitedHealthcare
- Aetna
- Humana
- Blue Cross

1 of 5 Only 5 questions? Easy. Too easy

ADVERTISEMENT

August 15th, 2013 08:00 PM ET

Study: Heavy coffee drinking in people under 55 linked to early death

Dr. Sanjay Gupta

<>

ACP AMERICAN COLLEGE OF PHYSICIANS  
INTERNAL MEDICINE | Doctors for Adults®

UCLA Biomedical Lib/Serial

# Annals of Internal Medicine

ESTABLISHED IN 1927 BY THE AMERICAN COLLEGE OF PHYSICIANS

Home Current Issue All Issues Online First Collections In the Clinic Journal Club CME An

17 June 2008, Vol 148, No. 12>

< Previous in this Issue Next in this Issue >

Email Share Get Citation Figures Slideset (.ppt) PDF

Articles 17 June 2008

Dr. Sanjay Gupta

The Relationship of Coffee Consumption with Mortality

Esther Lopez-Garcia, PhD; Rob M. van Dam, PhD; Tricia Y. Li, MD; Fernando Rodriguez-Artalejo, MD, PhD; and Frank B. Hu, MD, PhD

[+] Article and Author Information

Ann Intern Med. 2008;148(12):904-914. doi:10.7326/0003-4819-148-12-200806170-00003 Text Size: A A A

Article Figures Tables References Audio/Video Summary for Patients Comments (2)

## Abstract

Abstract | Context | Contribution | Caution | Methods | Results | Discussion | References ▾

**Background:** Coffee consumption has been linked to various beneficial and detrimental health effects, but data on its relation with mortality are sparse.

**Study:** Heavy coffee consumption linked to early death

August 15th, 2013  
08:00 PM ET

**CBSNEWS**

EDITION:  
TV: CNN

Video | US | World | Politics | Entertainment | Health

RELAX AND  
ENJOY THE LUXURY.  
A full-size sedan designed with you in mind.

February 15, 2013, 3:36 PM

By MICHELLE CASTILLO CBS NEWS

# Alcohol causes 20,000 cancer deaths in the U.S. annually



In Texas it is illegal to take more than three sips of beer at a time while standing. / AP / FILE

41 Comments / [Shares](#) / [Tweets](#) / [Stumble](#) / [Email](#)

More ▾

UCLA Biomedical Lib/Serial

e

the Clinic Journal Club CME

PDF

## Mortality

io, MD, PhD; and Frank

Size: A A A

ts (2)

cial and detrimental health effects,

## Science News

from universities, journals, and other research organizations

# One Drink Of Red Wine Or Alcohol Is Relaxing To Circulation, But Two Drinks Are Stressful

Feb. 13, 2008 — One drink of either red wine or alcohol slightly benefits the heart and blood vessels, but the positive effects on specific biological markers disappear with two drinks, say researchers at the Peter Munk Cardiac Centre of the Toronto General Hospital.

### Related Topics

#### Health & Medicine

- ▶ Heart Disease
- ▶ Hypertension

#### Mind & Brain

#### Articles

- ▶ Mediterranean diet
- ▶ Drunkenness
- ▶ Coronary heart disease



August 15  
08:00 PM E

In Texas it is illegal to take more than three sips of beer at a time while standing. / AP / FILE

41 Comments / f Shares / t Tweets / Stumble / @ Email

More +

Mortality  
io, MD, PhD; and Frank

Size: A A A

(2)

ocial and detrimental health effects,



**Science News***... from universities, journals, and other research organizations***One Drink Of Red Wine Is Good For You. Two Are Stressful.**

Feb. 13, 2008 — One drink of red wine relaxes your heart and improves circulation, but two or more can increase stress levels, according to new research from Peter Munk Cardiac Centre at Toronto General Hospital.

INSIDE THE EMERGENCY ROOM  
OF BOYS

Home | News

Current Issue /

**Relaxing To Circulation, But Two Drinks Are Stressful****Well**ASK WELL  
Ask Well: Nighttime Urination

apps



photos

in diet



part

Bal Harbour Shops



DAVID YURMAN



BVLGARI



HARRY WINSTON



uses use cloud services →

Subscribe

Obama's Race for the

What Boys Want

rohar: Yellen for America

Why China Needs More Children

1 effects,

**Why I**One of the  
the considera

By John C



FACEBOOK



TWITTER



GOOGLE+



SAVE



EMAIL



SHARE



PRINT


In Texas it is illegal to take

41 Comments / f Shares /

August 15<sup>th</sup>  
08:00 PM E

The patient didn't overdose on medication. She overdosed on grapefruit juice.

The 42-year-old was barely responding when her husband brought her to the emergency room. Her heart rate was slowing, and her blood pressure was falling. Doctors had to insert a breathing tube, and then a pacemaker, to revive her.



ACRA Video | US | World | Politics | Entertainment | Health

CBSNEWS

Science News

from universities, journals, and other research organizations

## One Drink Of Red Wine A Day Is Good For You. Two Drinks Are Stressful.

Feb. 13, 2008 — One drink of red wine a day may be good for you, but two or more are stressful, according to a new study.

INSIDE THE EMERGENCY ROOM  
OF BOYS

Home | News | Current Issue | /

THE CON  
Why I'm Grateful  
Re: By RON

Dr. Sanjay Gupta

One of the reasons the consumption of nuts has been associated with a reduced risk of major chronic diseases, including cardiovascular disease and type 2 diabetes mellitus. However, the association between nut consumption and mortality remains unclear.

Full Text of Background...

METHODS We examined the association between nut consumption and

In Texas it is illegal to take

41 Comments / f Shares /

The NEW ENGLAND JOURNAL of MEDICINE

from universities, journals, and other research organizations

## Two Nuts A Day May Be Good For You. Three To Circulation, But Two

### Association of Nut Consumption with Total and Cause-Specific Mortality

Ying Bao, M.D., Sc.D., Jiali Han, Ph.D., Frank B. Hu, M.D., Ph.D., Edward L. Giovannucci, M.D., Sc.D., Meir J. Stampfer, M.D., Dr.P.H., Walter C. Willett, M.D., Dr.P.H., and Charles S. Fuchs, M.D., M.P.H.  
N Engl J Med 2013; 369:2001-2011 / November 21, 2013 / DOI: 10.1056/NEJMoa1307352

Abstract Article References

Share: f t g+ in +

MEDIA IN THIS ARTICLE  
Video

Nuts and Death.

# Other Applications

---

- Evaluating the impact of a bilingual education program

# Other Applications

---

- Evaluating the impact of a bilingual education program
- Evaluating the impact of a new taxation program

# Other Applications

---

- Evaluating the impact of a bilingual education program
- Evaluating the impact of a new taxation program
- Making a robot to perform a complex task in a new, unforeseen environment

# Other Applications

---

- Evaluating the impact of a bilingual education program
- Evaluating the impact of a new taxation program
- Making a robot to perform a complex task in a new, unforeseen environment
- Diagnosing a patient and planning her/his optimal treatment

# Other Applications

---

- Evaluating the impact of a bilingual education program
- Evaluating the impact of a new taxation program
- Making a robot to perform a complex task in a new, unforeseen environment
- Diagnosing a patient and planning her/his optimal treatment
- Evaluating the effectiveness of a new drug

# Other Applications

---

- Evaluating the impact of a bilingual education program
- Evaluating the impact of a new taxation program
- Making a robot to perform a complex task in a new, unforeseen environment
- Diagnosing a patient and planning her/his optimal treatment
- Evaluating the effectiveness of a new drug
- Assessing the strength of the causal relationship between a set of genes and a certain type of cancer

# Other Applications

---

- Evaluating the impact of a bilingual education program
- Evaluating the impact of a new taxation program
- Making a robot to perform a complex task in a new, unforeseen environment
- Diagnosing a patient and planning her/his optimal treatment
- Evaluating the effectiveness of a new drug
- Assessing the strength of the causal relationship between a set of genes and a certain type of cancer



Tangled web of cause-effect relationships



# Why study causality?

---

# Why study causality?

---

- Cognitive observation

# Why study causality?

---

- Cognitive observation
  - human: sensors + agency + retrospection
  - ability to observe the world and infer the ‘next’ stage
  - ability to intervene in the world and change it
  - ability to reflect about what we did and infer new behavior

# Why study causality?

---

- Cognitive observation
  - human: sensors + agency + retrospection
  - ability to observe the world and infer the ‘next’ stage
  - ability to intervene in the world and change it
  - ability to reflect about what we did and infer new behavior

# Why study causality?

---

- Cognitive observation
  - human: sensors + agency + retrospection
  - ability to observe the world and infer the ‘next’ stage
  - ability to intervene in the world and change it
  - ability to reflect about what we did and infer new behavior
- Scientific perspective
  - to be able to understand nature and the human being
  - relies on empirical data, logic/math, and scientific method

# Why study causality?

---

- Cognitive observation
  - human: sensors + agency + retrospection
  - ability to observe the world and infer the ‘next’ stage
  - ability to intervene in the world and change it
  - ability to reflect about what we did and infer new behavior
- Scientific perspective
  - to be able to understand nature and the human being
  - relies on empirical data, logic/math, and scientific method
- Pragmatic perspective (e.g., AI, Econ, Medicine)
  - to be able to create “truly” intelligent systems
  - to be able to create and implement policies optimizing certain {social, economical, political} goals

# OUTLINE (PART I)

---

- Introduction
  - \* Data, data, data...
  - \* Basic definitions
- Technical Results
  - \* The truncated product formula
  - \* The back-door adjustment formula
  - \* The front-door adjustment formula
  - \* The do-calculus
- Tasks
  - \* Policy evaluation
  - \* Generalizability & Robustness
  - \* Decision-making & RL

# OUTLINE (PART I)

---

- Introduction
  - \* Data, data, data...
  - \* Basic definitions
- Technical Results
  - \* The truncated product formula
  - \* The back-door adjustment formula
  - \* The front-door adjustment formula
  - \* The do-calculus
- Tasks
  - \* Policy evaluation
  - \* Generalizability & Robustness
  - \* Decision-making & RL

Data, data, data...

# Ex 1. Data analysis in a 2x2 table...

(Simpson, 1951; Ch. 6, Pearl, 2000)

---

	surv. (Y)	not-surv. ( $\neg$ Y)		Recovery Rate
drug (X)	20	20	40	50%
no-drug ( $\neg$ X)	16	24	40	40%
	36	44		

# Stratifying by sex...

	surv. (Y)	not-surv. ( $\neg$ Y)		Recovery Rate
drug (X)	20	20	40	50%
no-drug ( $\neg$ X)	16	24	40	40%
	36	44		



male ( $\neg$ F)	surv. (Y)	not-surv.		Recovery Rate
drug (X)	18	12	30	60%
no-drug ( $\neg$ X)	7	3	10	70%
	25	15	40	

female (F)	surv. (Y)	not-surv. ( $\neg$ Y)		Recovery Rate
drug (X)	2	8	10	20%
no-drug ( $\neg$ X)	9	21	30	30%
	11	29	40	

# Which table should the physician consult?

	surv. (Y)	not-surv. ( $\neg Y$ )		Recovery Rate
drug (X)	20	20	40	50%
no-drug ( $\neg X$ )	16	24	40	40%
	36	44		

$$P(Y | F, X) < P(Y | F, \neg X)$$

$$P(Y | \neg F, X) < P(Y | \neg F, \neg X)$$

male ( $\neg F$ )	surv. (Y)	not-surv.		Recovery Rate
drug (X)	18	12	30	60%
no-drug ( $\neg X$ )	7	3	10	70%
	25	15	40	

↔

female (F)	surv. (Y)	not-surv. ( $\neg Y$ )		Recovery Rate
drug (X)	2	8	10	20%
no-drug ( $\neg X$ )	9	21	30	30%
	11	29	40	

# Which table should the physician consult?

	surv. (Y)	not-surv. ( $\neg Y$ )		Recovery Rate
drug (X)	20	20	40	50%
no-drug ( $\neg X$ )	16	24	40	40%
	36	44		



	surv. (Y)	not-surv.		Recovery Rate
male ( $\neg F$ )				
drug (X)	18	12	30	60%
no-drug ( $\neg X$ )	7	3	10	70%
	25	15	40	

	surv. (Y)	not-surv. ( $\neg Y$ )		Recovery Rate
female (F)				
drug (X)	2	8	10	20%
no-drug ( $\neg X$ )	9	21	30	30%
	11	29	40	

$P(Y | F, X) < P(Y | F, \neg X)$   
 $P(Y | \neg F, X) < P(Y | \neg F, \neg X)$   
but  
 $P(Y | X) > P(Y | \neg X) !!$

# WOULD THE RESULT BE DIFFERENT WITH A DIFFERENT STORY?

---

male (¬F)	surv. (Y)	not- surv.		Recovery Rate
drug (X)	18	12	30	60%
no- drug	7	3	10	70%
	25	15	40	


female (F)	surv. (Y)	not- surv.		Recovery Rate
drug (X)	2	8	10	20%
no- drug	9	21	30	30%
	11	29	40	

# WOULD THE RESULT BE DIFFERENT WITH A DIFFERENT STORY?

male ( $\neg F$ )	surv. (Y)	not-surv.		Recovery Rate
drug (X)	18	12	30	60%
no-drug	7	3	10	70%
	25	15	40	

female (F)	surv. (Y)	not-surv.		Recovery Rate
drug (X)	2	8	10	20%
no-drug	9	21	30	30%
	11	29	40	


$G_1$



# WOULD THE RESULT BE DIFFERENT WITH A DIFFERENT STORY?

male (¬F)	surv. (Y)	not- surv.		Recovery Rate
drug (X)	18	12	30	60%
no- drug	7	3	10	70%
	25	15	40	


female (F)	surv. (Y)	not- surv.		Recovery Rate
drug (X)	2	8	10	20%
no- drug	9	21	30	30%
	11	29	40	



# WOULD THE RESULT BE DIFFERENT WITH A DIFFERENT STORY?

male ( $\neg F$ )	surv. (Y)	not-surv.		Recovery Rate
drug (X)	18	12	30	60%
no-drug	7	3	10	70%
	25	15	40	

female (F)	surv. (Y)	not-surv.		Recovery Rate
drug (X)	2	8	10	20%
no-drug	9	21	30	30%
	11	29	40	



## Ex 2. Trivial data analysis (Thrun)?


---

#Firefighters (X)	#Size Fire (Y)
10	100
40	400
60	600
200	2000

Is  
 $Y = \beta X + \epsilon$  a good fit?

# Simple linear regression...


#Firefighters (X)	#Size Fire (Y)
10	100
40	400
60	600
200	2000



Is  
 $Y = \beta X + \epsilon$  a good fit?

# Simple linear regression...

#Firefighters (X)	#Size Fire (Y)
10	100
40	400
60	600
200	2000




Is  
 $Y = \beta X + \epsilon$  a good fit?

X and Y are perfectly correlated...

but does X cause Y?

# Simple linear regression...

#Firefighters (X)	#Size Fire (Y)
10	100
40	400
60	600
200	2000




Is

$Y = \beta X + \epsilon$  a good fit?

after all,  
what's the meaning of  $\beta$ ?

# Simple linear regression...

#Firefighters (X)	#Size Fire (Y)
10	100
40	400
60	600
200	2000



Is


$Y = \beta X + \epsilon$  a good fit?

Lasso?

after all,  
what's the meaning of  $\beta$ ?

# Simple linear regression...

#Firefighters (X)	#Size Fire (Y)
10	100
40	400
60	600
200	2000



Is

$Y = \beta X + \epsilon$  a good fit?




after all,


what's the meaning of  $\beta$ ?

Lasso?  
DNN?

## Ex 2b. Trivial data analysis (?)



# Ex 2b. Trivial data analysis (?)



Sources: Organic Trade Association, 2011 Organic Industry Survey; U.S. Department of Education, Office of Special Education Programs, Data Analysis System (DANS), OMB# 1820-0043: "Children with Disabilities Receiving Special Education Under Part B of the Individuals with Disabilities Education Act"


What's going on here...?

What's going on here...?




# TRADITIONAL STATISTICAL/ MACHINE LEARNING PARADIGM

---



# TRADITIONAL STATISTICAL/ MACHINE LEARNING PARADIGM


---



e.g., Infer whether customers who bought product  $A$  would also buy product  $B$  — or, compute  $Q = P(B | A)$ .

# TRADITIONAL STATISTICAL/ MACHINE LEARNING PARADIGM


---




e.g., Infer whether customers who bought product  $A$  would also buy product  $B$  — or, compute  $Q = P(B | A)$ .

- capturing regularities of the original  $P$  (LLN, CLT)

# FROM STATISTICAL TO CAUSAL ANALYSIS



# FROM STATISTICAL TO CAUSAL ANALYSIS




e.g., Estimate  $P'(\text{sales})$  if we double the price.

Estimate  $P'(\text{cancer})$  if we ban smoking.

How does  $P$  change to  $P'$ ? New oracle

# FROM STATISTICAL TO CAUSAL ANALYSIS



e.g., Estimate  $P'(sales)$  if we double the price.


Estimate  $P'(cancer)$  if we ban smoking.

What  
about  
the CLT?

How does  $P$  change to  $P'$ ? New oracle


# FROM STATISTICS TO COUNTERFACTUALS

Probability and statistics deal with static relations



# FROM STATISTICS TO COUNTERFACTUALS

Probability and statistics deal with static relations




What happens when  $P$  changes?

e.g., Estimate the probability that a patient who took a drug and died would be alive had he not taken the drug.

# FROM STATISTICS TO COUNTERFACTUALS

Probability and statistics deal with static relations



What happens when  $P$  changes?

e.g., Estimate the probability that a patient who took a drug and died would be alive had he not taken the drug.


Violating  
the laws  
of physics?

Instead of thinking about  
multiple individual distributions,  
let's dig deeper...




# THE STRUCTURAL MODEL PARADIGM

---



# THE STRUCTURAL MODEL PARADIGM


---



*SCM*  $M$  – Invariant strategy (mechanism, recipe, law, protocol) by which Nature assigns values to variables in the analysis.

# THE STRUCTURAL MODEL PARADIGM

---



*SCM M* – Invariant strategy (mechanism, recipe, law, protocol) by which Nature assigns values to variables in the analysis.

$P$  – model of data

$M$  – model of reality

# REPRESENTING THE DATA GENERATING MODEL

---

- Processes

# REPRESENTING THE DATA GENERATING MODEL

---

- Processes

$$\text{Edu} \leftarrow f_E(\text{SES}, U_e)$$

# REPRESENTING THE DATA GENERATING MODEL

---

- Processes

$$\text{Edu} \leftarrow f_E(\text{SES}, U_e)$$

$$\text{Income} \leftarrow f_I(\text{Edu}, \text{SES}, U_i)$$

# REPRESENTING THE DATA GENERATING MODEL


---

- Processes

$$\text{Edu} \leftarrow f_E(\text{SES}, U_e)$$

$$\text{Income} \leftarrow f_I(\text{Edu}, \text{SES}, U_i)$$

**G =**



# REPRESENTING THE DATA GENERATING MODEL


---

- Processes

$$\text{Edu} \leftarrow f_E(\text{SES}, U_e)$$

$$\text{Income} \leftarrow f_I(\text{Edu}, \text{SES}, U_i)$$

- Intervention



# REPRESENTING THE DATA GENERATING MODEL

---

- Processes


$$\text{Edu} \leftarrow f_E(\text{SES}, U_e)$$

$$\text{Income} \leftarrow f_I(\text{Edu}, \text{SES}, U_i)$$

- Intervention

$$\text{Edu} \leftarrow \text{Yes}$$

**G =**



# REPRESENTING THE DATA GENERATING MODEL

---

- Processes

$$\text{Edu} \leftarrow f_E(\text{SES}, U_e)$$


$$\text{Income} \leftarrow f_I(\text{Edu}, \text{SES}, U_i)$$

- Intervention

$$\text{Edu} \leftarrow \text{Yes}$$

$$\text{Income} \leftarrow f_I(\text{Edu}, \text{SES}, U_i)$$

**G =**



# REPRESENTING THE DATA GENERATING MODEL

---

- Processes

$$\text{Edu} \leftarrow f_E(\text{SES}, U_e)$$


$$\text{Income} \leftarrow f_I(\text{Edu}, \text{SES}, U_i)$$

- Intervention


$$\text{Edu} \leftarrow \text{Yes}$$

$$\text{Income} \leftarrow f_I(\text{Edu}, \text{SES}, U_i)$$

**G =**



**$G_{do(X)}$  =**



# REPRESENTING THE DATA GENERATING MODEL

---

- Processes

$$\text{Edu} \leftarrow f_E(\text{SES}, U_e)$$


$$\text{Income} \leftarrow f_I(\text{Edu}, \text{SES}, U_i)$$

- Intervention


$$\text{Edu} \leftarrow \text{Yes}$$

$$\text{Income} \leftarrow f_I(\text{Edu}, \text{SES}, U_i)$$

**G =**



**$G_{do(X)}$  =**



# REPRESENTING THE DATA GENERATING MODEL

---

# REPRESENTING THE DATA GENERATING MODEL

---

Definition: A **structural causal model M** (or, data generating model) is a tuple  $(V, U, F, P(u))$ , where

# REPRESENTING THE DATA GENERATING MODEL

---

Definition: A **structural causal model M** (or, data generating model) is a tuple  $(V, U, F, P(u))$ , where

- $V = \{V_1, \dots, V_n\}$  are endogenous variables,

# REPRESENTING THE DATA GENERATING MODEL

---

Definition: A **structural causal model M** (or, data generating model) is a tuple  $(V, U, F, P(u))$ , where

- $V = \{V_1, \dots, V_n\}$  are endogenous variables,
- $U = \{U_1, \dots, U_m\}$  are exogenous variables,

# REPRESENTING THE DATA GENERATING MODEL

---

Definition: A **structural causal model M** (or, data generating model) is a tuple  $(V, U, F, P(u))$ , where

- $V = \{V_1, \dots, V_n\}$  are endogenous variables,
- $U = \{U_1, \dots, U_m\}$  are exogenous variables,
- $F = \{f_1, \dots, f_n\}$  are functions determining  $V$ ,  
for each  $V_i$ ,  $V_i \leftarrow f_i(Pa_i, U_i)$ , where  $Pa_i \subset V$ ,  $U_i \subset U$ .

# REPRESENTING THE DATA GENERATING MODEL

---

Definition: A **structural causal model M** (or, data generating model) is a tuple  $(V, U, F, P(u))$ , where

- $V = \{V_1, \dots, V_n\}$  are **endogenous variables**,
- $U = \{U_1, \dots, U_m\}$  are **exogenous variables**,
- $F = \{f_1, \dots, f_n\}$  are functions determining  $V$ ,  
for each  $V_i$ ,  $V_i \leftarrow f_i(Pa_i, U_i)$ , where  $Pa_i \subset V, U_i \subset U$ .

# REPRESENTING THE DATA GENERATING MODEL

---

Definition: A **structural causal model M** (or, data generating model) is a tuple  $(V, U, F, P(u))$ , where

- $V = \{V_1, \dots, V_n\}$  are endogenous variables,
- $U = \{U_1, \dots, U_m\}$  are exogenous variables,
- $F = \{f_1, \dots, f_n\}$  are functions determining  $V$ ,  
for each  $V_i$ ,  $V_i \leftarrow f_i(Pa_i, U_i)$ , where  $Pa_i \subset V$ ,  $U_i \subset U$ .
- $P(u)$  is a distribution over  $U$ .

# REPRESENTING THE DATA GENERATING MODEL

---

Definition: A **structural causal model M** (or, data generating model) is a tuple  $(V, U, F, P(u))$ , where

- $V = \{V_1, \dots, V_n\}$  are endogenous variables,
- $U = \{U_1, \dots, U_m\}$  are exogenous variables,
- $F = \{f_1, \dots, f_n\}$  are functions determining  $V$ ,  
for each  $V_i$ ,  $V_i \leftarrow f_i(Pa_i, U_i)$ , where  $Pa_i \subset V$ ,  $U_i \subset U$ .
- $P(u)$  is a distribution over  $U$ .


(Axiomatic characterization [Halpern, Galles, Pearl, 1998].)

# COUNTERFACTUALS IN A NUTSHELL

---

## Definition:

Given a SCM model  $M$ , the potential outcome  $Y_x(u)$  for unit  $u$  is equal to the solution for  $Y$  in a mutilated model  $M_x$ , in which the equation for  $X$  is replaced by  $X = x$ .




# COUNTERFACTUALS IN A NUTSHELL

---

## Definition:

Given a SCM model  $M$ , the potential outcome  $Y_x(u)$  for unit  $u$  is equal to the solution for  $Y$  in a mutilated model  $M_x$ , in which the equation for  $X$  is replaced by  $X = x$ .



## The Law of Structural Counterfactuals:


$$Y_X(u) \triangleq Y_{M_X}(u)$$

# COUNTERFACTUALS IN A NLU

If  $Y = V_i$ ,  
and  $X = V \setminus Y$ ,  
 $Y_x(u) = f_i(pa_i, u)$

## Definition:

Given a SCM model  $M$ , the potential outcome  $Y_x(u)$  for unit  $u$  is equal to the solution for  $Y$  in a mutilated model  $M_x$ , in which the equation for  $X$  is replaced by  $X = x$ .



## The Law of Structural Counterfactuals:

$$Y_X(u) \triangleq Y_{M_x}(u)$$

SCM → PEARL (CAUSAL) HIERARCHY

# SCM → CAUSAL HIERARCHY

---

Level (Symbol)	Typical Activity	Typical Question	Examples
1 ○○ $P(y   x)$	Seeing	What is? How would seeing X change my belief in Y?	What does a symptom tell us about the disease?

# SCM → CAUSAL HIERARCHY

---

Level (Symbol)	Typical Activity	Typical Question	Examples
1 ○○ $P(y   x)$	Seeing ML - (Un)Supervised DNN, Bayes Net Regression	What is? How would seeing X change my belief in Y?	What does a symptom tell us about the disease?

# SCM → CAUSAL HIERARCHY

---

Level (Symbol)	Typical Activity	Typical Question	Examples
1 Associations 	Seeing ML - (Un)Supervised DNN, Bayes Net Regression	What is? How would seeing X change my belief in Y?	What does a symptom tell us about the disease?
2 Interventional 	Doing	What if? What if I do X?	What if I take aspirin, will my headache be cured?

# SCM → CAUSAL HIERARCHY

---

Level (Symbol)	Typical Activity	Typical Question	Examples
1 Associations 	Seeing ML - (Un)Supervised DNN, Bayes Net Regression	What is? How would seeing X change my belief in Y?	What does a symptom tell us about the disease?
2 Interventional 	Doing ML - Reinforcement Causal Bayes Net	What if? What if I do X?	What if I take aspirin, will my headache be cured?

# SCM → CAUSAL HIERARCHY

---

Level (Symbol)	Typical Activity	Typical Question	Examples
1 Associations 	Seeing ML - (Un)Supervised DNN, Bayes Net Regression	What is? How would seeing X change my belief in Y?	What does a symptom tell us about the disease?
2 Interventional 	Doing ML - Reinforcement Causal Bayes Net	What if? What if I do X?	What if I take aspirin, will my headache be cured?
3 Counterfactual 	Imagining, Retrospection	Why? What if I had acted differently?	Was it the aspirin that stopped my headache?

# SCM → CAUSAL HIERARCHY

---

Level (Symbol)	Typical Activity	Typical Question	Examples
1 Associations 	Seeing ML - (Un)Supervised DNN, Bayes Net Regression	What is? How would seeing X change my belief in Y?	What does a symptom tell us about the disease?
2 Interventional 	Doing ML - Reinforcement Causal Bayes Net	What if? What if I do X?	What if I take aspirin, will my headache be cured?
3 Counterfactual 	Imagining, Retrospection Structural Causal Model	Why? What if I had acted differently?	Was it the aspirin that stopped my headache?

# SCM → CAUSAL HIERARCHY

Level (Symbol)	Typical Activity	Typical Question	Examples
1 Associations 	Seeing		part 1
	ML - (Un)Supervised DNN, Bayes Net Regression		
2 Interventional 	Doing	What if? What if I do X?	What if I take aspirin, will my headache be cured?
	ML - Reinforcement Causal Bayes Net		
3 Counterfactual 	Imagining, Retrospection	Why? What if I had acted differently?	Was it the aspirin that stopped my headache?
	Structural Causal Model		

# SCM → CAUSAL HIERARCHY

Level (Symbol)	Typical Activity	Typical Question	Examples
1 👀 $P(y   x)$	Seeing ML - (Un)Supervised DNN, Bayes Net Regression		part 1
2 💪 $P(y   \text{do}(x), c)$	Doing ML - Reinforcement Causal Bayes Net		part 2
3 🧠 $P(y_x   x', y')$	Imagining, Retrospection Structural Causal Model	Why? What if I had acted differently?	Was it the aspirin that stopped my headache?

# SCM → CAUSAL HIERARCHY

---

Level (Symbol)	Typical Activity	Typical Question	Examples
1 Associations 	Seeing		part 1
	ML - (Un)Supervised DNN, Bayes Net Regression		
2 Interventional 	Doing		part 2
	ML - Reinforcement Causal Bayes Net		
3 Counterfactual 	Imagining, Retrospection Structural Causal Mod		part 3


# FIRST LAYER OF THE CAUSAL HIERARCHY

## ASSOCIATIONAL

(What if I see  $X=x$ ?)


# THE EMERGENCE OF THE FIRST LAYER

---



# THE EMERGENCE OF THE FIRST LAYER

---



Theorem (PV, 1991). Every **Markovian** structural causal model  $M$  (**recursive, with independent disturbances**) induces an observational distribution  $P(v_1, \dots, v_n)$  that can be factorized as

$$P(v_1, v_2, \dots, v_n) = \prod_i P(v_i \mid pa_i)$$

where  $pa_i$  are the (values of) the parents of  $V_i$  in the causal diagram associated with  $M$ .

# ENCODING PROBABILISTIC INVARIANCE (GRAPHHOIDS)

---

# ENCODING PROBABILISTIC INVARIANCE (GRAPHHOIDS)

---

Statements of the form:

# ENCODING PROBABILISTIC INVARIANCE (GRAPHHOIDS)

---

Statements of the form:

“X is **irrelevant** to Y given that **we know Z**”,

# ENCODING PROBABILISTIC INVARIANCE (GRAPHHOIDS)

---

Statements of the form:

“X is **irrelevant** to Y given that **we know** Z”,  
where X, Y, and Z are sets of variables.

# ENCODING PROBABILISTIC INVARIANCE (GRAPHHOIDS)

---

Statements of the form:

“X is **irrelevant** to Y given that **we know** Z”,  
where X, Y, and Z are sets of variables.

# ENCODING PROBABILISTIC INVARIANCE (GRAPHHOIDS)

---

Statements of the form:

“X is **irrelevant** to Y given that **we know Z**”,  
where X, Y, and Z are sets of variables.

The notion of irrelevance can be considered under different interpretations, including probabilistic, relational, and correlational.

# ENCODING PROBABILISTIC INVARIANCE (GRAPHHOIDS)

---

Statements of the form:

“X is **irrelevant** to Y given that **we know Z**”,  
where X, Y, and Z are sets of variables.

The notion of irrelevance can be considered under different interpretations, including probabilistic, relational, and correlational.

# ENCODING PROBABILISTIC INVARIANCE (GRAPHHOIDS)

---

Statements of the form:

“X is **irrelevant** to Y given that **we know Z**”,  
where X, Y, and Z are sets of variables.

The notion of irrelevance can be considered under different interpretations, including probabilistic, relational, and correlational.

(See also J. Pearl (1988) and P. David (1979). )

# THE LAW OF CONDITIONAL INDEPENDENCE

---

**SCM ( $M$ )**

$$C \leftarrow f_c(U_c)$$


$$S \leftarrow f_S(C, U_S)$$

$$R \leftarrow f_R(C, U_R)$$

$$W \leftarrow f_W(S, R, U_W)$$

# THE LAW OF CONDITIONAL INDEPENDENCE

**Graph ( $G$ )**




**SCM ( $M$ )**

$$\begin{aligned}C &\leftarrow f_c(U_c) \\S &\leftarrow f_s(C, U_s) \\R &\leftarrow f_R(C, U_R) \\W &\leftarrow f_W(S, R, U_W)\end{aligned}$$

# THE LAW OF CONDITIONAL INDEPENDENCE

**Graph ( $G$ )**



**SCM ( $M$ )**

$$\begin{aligned} C &\leftarrow f_c(U_c) \\ S &\leftarrow f_s(C, U_s) \\ R &\leftarrow f_R(C, U_R) \\ W &\leftarrow f_W(S, R, U_W) \end{aligned}$$


\* **Surprising result:**

If the  $U$ 's are independent, the observed distribution  $P(c, r, s, w)$  satisfies constraints that are:

- (1) independent of the  $f$ 's and of  $P(U)$ ,
- (2) readable from the graph.

# D-SEPARATION: READING CONSTRAINTS IMPLIED IN THE DATA

## Graph ( $G$ )




## SCM ( $M$ )

$$\begin{aligned}C &\leftarrow f_c(U_c) \\S &\leftarrow f_s(C, U_s) \\R &\leftarrow f_R(C, U_R) \\W &\leftarrow f_W(S, R, U_W)\end{aligned}$$

Every missing arrow advertises an independency, conditional on a separating set.

# D-SEPARATION: READING CONSTRAINTS IMPLIED IN THE DATA

## Graph ( $G$ )



## SCM ( $M$ )


$$\begin{aligned}C &\leftarrow f_c(U_c) \\S &\leftarrow f_s(C, U_s) \\R &\leftarrow f_R(C, U_R) \\W &\leftarrow f_W(S, R, U_W)\end{aligned}$$

Every missing arrow advertises an independency, conditional on a separating set.

$$C \perp\!\!\!\perp W \mid (S, R)$$

# D-SEPARATION: READING CONSTRAINTS IMPLIED IN THE DATA

## Graph ( $G$ )



## SCM ( $M$ )


$$\begin{aligned} C &\leftarrow f_c(U_c) \\ S &\leftarrow f_s(C, U_s) \\ R &\leftarrow f_R(C, U_R) \\ W &\leftarrow f_W(S, R, U_W) \end{aligned}$$

Every missing arrow advertises an independency, conditional on a separating set.

$$\begin{aligned} C \perp\!\!\!\perp W \mid (S, R) \\ S \perp\!\!\!\perp R \mid C \end{aligned}$$

# D-SEPARATION: READING CONSTRAINTS IMPLIED IN THE DATA

## Graph ( $G$ )



## SCM ( $M$ )

$$\begin{aligned} C &\leftarrow f_c(U_c) \\ S &\leftarrow f_s(C, U_s) \\ R &\leftarrow f_R(C, U_R) \\ W &\leftarrow f_W(S, R, U_W) \end{aligned}$$


Every missing arrow advertises an independency, conditional on a separating set.

$$\begin{aligned} C \perp\!\!\!\perp W \mid (S, R) \\ S \perp\!\!\!\perp R \mid C \end{aligned}$$

## Applications:

# D-SEPARATION: READING CONSTRAINTS IMPLIED IN THE DATA

## Graph ( $G$ )



## SCM ( $M$ )

$$\begin{aligned} C &\leftarrow f_c(U_c) \\ S &\leftarrow f_s(C, U_s) \\ R &\leftarrow f_R(C, U_R) \\ W &\leftarrow f_W(S, R, U_W) \end{aligned}$$


Every missing arrow advertises an independency, conditional on a separating set.

$$\begin{aligned} C \perp\!\!\!\perp W \mid (S, R) \\ S \perp\!\!\!\perp R \mid C \end{aligned}$$

## Applications:

# D-SEPARATION: READING CONSTRAINTS IMPLIED IN THE DATA

## Graph ( $G$ )



## SCM ( $M$ )

$$\begin{aligned} C &\leftarrow f_c(U_c) \\ S &\leftarrow f_s(C, U_s) \\ R &\leftarrow f_R(C, U_R) \\ W &\leftarrow f_W(S, R, U_W) \end{aligned}$$

Every missing arrow advertises an independency, conditional on a separating set.


$$\begin{aligned} C \perp\!\!\!\perp W \mid (S, R) \\ S \perp\!\!\!\perp R \mid C \end{aligned}$$

## Applications:

1. Structural learning

# D-SEPARATION: READING CONSTRAINTS IMPLIED IN THE DATA

## Graph ( $G$ )



## SCM ( $M$ )

$$\begin{aligned}C &\leftarrow f_c(U_c) \\S &\leftarrow f_s(C, U_s) \\R &\leftarrow f_R(C, U_R) \\W &\leftarrow f_W(S, R, U_W)\end{aligned}$$

Every missing arrow advertises an independency, conditional on a separating set.


$$\begin{aligned}C \perp\!\!\!\perp W \mid (S, R) \\S \perp\!\!\!\perp R \mid C\end{aligned}$$

## Applications:

1. Structural learning
2. Model testing

# D-SEPARATION: READING CONSTRAINTS IMPLIED IN THE DATA

## Graph ( $G$ )



## SCM ( $M$ )

$$\begin{aligned} C &\leftarrow f_c(U_c) \\ S &\leftarrow f_s(C, U_s) \\ R &\leftarrow f_R(C, U_R) \\ W &\leftarrow f_W(S, R, U_W) \end{aligned}$$

Every missing arrow advertises an independency, conditional on a separating set.

$$\begin{aligned} C \perp\!\!\!\perp W \mid (S, R) \\ S \perp\!\!\!\perp R \mid C \end{aligned}$$

## Applications:

1. Structural learning
2. Model testing
3. (Efficient) Probabilistic inference

# GRAPH SEPARATION (D-SEPARATION)

---

# GRAPH SEPARATION (D-SEPARATION)

---


	normal valve	abnormal valve	
$(X \perp\!\!\!\perp Y   Z)$	$x \leftarrow z \rightarrow y$	$x \rightarrow z \leftarrow y$	$(X \perp\!\!\!\perp Y)$
	$x \rightarrow z \rightarrow y$	$x \rightarrow z \leftarrow y$	$(X \not\perp\!\!\!\perp Y   Z)$
	$x \leftarrow z \leftarrow y$	$\downarrow$ $w$	

# GRAPH SEPARATION (D-SEPARATION)

---

	normal valve	abnormal valve	
	$x \leftarrow z \rightarrow y$	$x \rightarrow z \leftarrow y$	$(X \perp\!\!\!\perp Y   Z)$
$(X \perp\!\!\!\perp Y   Z)$	$x \rightarrow z \rightarrow y$	$x \rightarrow z \leftarrow y$	$(X \not\perp\!\!\!\perp Y   Z)$
	$x \leftarrow z \leftarrow y$	$\downarrow w$	


---



# GRAPH SEPARATION (D-SEPARATION)

	normal valve	abnormal valve	
$(X \perp\!\!\!\perp Y   Z)$	$x \leftarrow z \rightarrow y$	$x \rightarrow z \leftarrow y$	$(X \perp\!\!\!\perp Y)$
	$x \rightarrow z \rightarrow y$	$x \rightarrow z \leftarrow y$	$(X \not\perp\!\!\!\perp Y   Z)$
	$x \leftarrow z \leftarrow y$	$\downarrow$ $w$	


$C_{I_1} : (Wet \perp\!\!\!\perp Sprinkler)$



# GRAPH SEPARATION (D-SEPARATION)

	normal valve	abnormal valve	
	$x \leftarrow z \rightarrow y$	$x \rightarrow z \leftarrow y$	$(X \perp\!\!\!\perp Y   Z)$
$(X \perp\!\!\!\perp Y   Z)$	$x \rightarrow z \rightarrow y$	$x \rightarrow z \leftarrow y$	$(X \not\perp\!\!\!\perp Y   Z)$
	$x \leftarrow z \leftarrow y$	$\downarrow w$	


✗  $C_{I_1} : (Wet \perp\!\!\!\perp Sprinkler)$



# GRAPH SEPARATION (D-SEPARATION)

	normal valve	abnormal valve	
$(X \perp\!\!\!\perp Y   Z)$	$x \leftarrow z \rightarrow y$	$x \rightarrow z \leftarrow y$	$(X \perp\!\!\!\perp Y)$
	$x \rightarrow z \rightarrow y$	$x \rightarrow z \leftarrow y$	$(X \not\perp\!\!\!\perp Y   Z)$
	$x \leftarrow z \leftarrow y$	$\downarrow$ $w$	


- ✗ CI<sub>1</sub> : (Wet  $\perp\!\!\!\perp$  Sprinkler)
- CI<sub>2</sub> : (Wet  $\perp\!\!\!\perp$  Climate | Sprinkler)



# GRAPH SEPARATION (D-SEPARATION)

	normal valve	abnormal valve	
$(X \perp\!\!\!\perp Y   Z)$	$x \leftarrow z \rightarrow y$	$x \rightarrow z \leftarrow y$	$(X \perp\!\!\!\perp Y)$
	$x \rightarrow z \rightarrow y$	$x \rightarrow z \leftarrow y$	$(X \not\perp\!\!\!\perp Y   Z)$
	$x \leftarrow z \leftarrow y$	$\downarrow$ $w$	


- ✗ CI<sub>1</sub> : (Wet  $\perp\!\!\!\perp$  Sprinkler)
- ✗ CI<sub>2</sub> : (Wet  $\perp\!\!\!\perp$  Climate | Sprinkler)



# GRAPH SEPARATION (D-SEPARATION)

	normal valve	abnormal valve	
$(X \perp\!\!\!\perp Y   Z)$	$x \leftarrow z \rightarrow y$	$x \rightarrow z \leftarrow y$	$(X \perp\!\!\!\perp Y)$
	$x \rightarrow z \rightarrow y$	$x \rightarrow z \leftarrow y$	$(X \not\perp\!\!\!\perp Y   Z)$
	$x \leftarrow z \leftarrow y$	$\downarrow w$	


- ✗ CI<sub>1</sub> : (Wet  $\perp\!\!\!\perp$  Sprinkler)
- ✗ CI<sub>2</sub> : (Wet  $\perp\!\!\!\perp$  Climate | Sprinkler)
- CI<sub>3</sub> : (Rain  $\perp\!\!\!\perp$  Slippery | Wet)



# GRAPH SEPARATION (D-SEPARATION)

	normal valve	abnormal valve	
$(X \perp\!\!\!\perp Y   Z)$	$x \leftarrow z \rightarrow y$	$x \rightarrow z \leftarrow y$	$(X \perp\!\!\!\perp Y)$
	$x \rightarrow z \rightarrow y$	$x \rightarrow z \leftarrow y$	$(X \not\perp\!\!\!\perp Y   Z)$
	$x \leftarrow z \leftarrow y$	$\downarrow$ $w$	


- ✗ CI<sub>1</sub> : (Wet  $\perp\!\!\!\perp$  Sprinkler)
- ✗ CI<sub>2</sub> : (Wet  $\perp\!\!\!\perp$  Climate | Sprinkler)
- ✓ CI<sub>3</sub> : (Rain  $\perp\!\!\!\perp$  Slippery | Wet)



# GRAPH SEPARATION (D-SEPARATION)

	normal valve	abnormal valve	
$(X \perp\!\!\!\perp Y   Z)$	$x \leftarrow z \rightarrow y$	$x \rightarrow z \leftarrow y$	$(X \perp\!\!\!\perp Y)$
	$x \rightarrow z \rightarrow y$	$x \rightarrow z \leftarrow y$	$(X \not\perp\!\!\!\perp Y   Z)$
	$x \leftarrow z \leftarrow y$	$\downarrow$ $w$	


- ✗ CI<sub>1</sub> : (Wet  $\perp\!\!\!\perp$  Sprinkler)
- ✗ CI<sub>2</sub> : (Wet  $\perp\!\!\!\perp$  Climate | Sprinkler)
- ✓ CI<sub>3</sub> : (Rain  $\perp\!\!\!\perp$  Slippery | Wet)
- CI<sub>4</sub> : (Season  $\perp\!\!\!\perp$  Wet | Sprinkler, Rain)



# GRAPH SEPARATION (D-SEPARATION)

	normal valve	abnormal valve	
$(X \perp\!\!\!\perp Y   Z)$	$x \leftarrow z \rightarrow y$	$x \rightarrow z \leftarrow y$	$(X \perp\!\!\!\perp Y)$
	$x \rightarrow z \rightarrow y$	$x \rightarrow z \leftarrow y$	$(X \not\perp\!\!\!\perp Y   Z)$
	$x \leftarrow z \leftarrow y$	$\downarrow$ $w$	


- ✗ CI<sub>1</sub> : (Wet  $\perp\!\!\!\perp$  Sprinkler)
- ✗ CI<sub>2</sub> : (Wet  $\perp\!\!\!\perp$  Climate | Sprinkler)
- ✓ CI<sub>3</sub> : (Rain  $\perp\!\!\!\perp$  Slippery | Wet)
- ✓ CI<sub>4</sub> : (Season  $\perp\!\!\!\perp$  Wet | Sprinkler, Rain)



# GRAPH SEPARATION (D-SEPARATION)

	normal valve	abnormal valve	
$(X \perp\!\!\!\perp Y   Z)$	$x \leftarrow z \rightarrow y$	$x \rightarrow z \leftarrow y$	$(X \perp\!\!\!\perp Y)$
	$x \rightarrow z \rightarrow y$	$x \rightarrow z \leftarrow y$	$(X \not\perp\!\!\!\perp Y   Z)$
	$x \leftarrow z \leftarrow y$	$\downarrow$ $w$	


- ✗ CI<sub>1</sub> : (Wet  $\perp\!\!\!\perp$  Sprinkler)
- ✗ CI<sub>2</sub> : (Wet  $\perp\!\!\!\perp$  Climate | Sprinkler)
- ✓ CI<sub>3</sub> : (Rain  $\perp\!\!\!\perp$  Slippery | Wet)
- ✓ CI<sub>4</sub> : (Season  $\perp\!\!\!\perp$  Wet | Sprinkler, Rain)
- CI<sub>5</sub> : (Sprinkler  $\perp\!\!\!\perp$  Rain | Climate, Wet)



# GRAPH SEPARATION (D-SEPARATION)

	normal valve	abnormal valve	
$(X \perp\!\!\!\perp Y   Z)$	$x \leftarrow z \rightarrow y$	$x \rightarrow z \leftarrow y$	$(X \perp\!\!\!\perp Y)$
	$x \rightarrow z \rightarrow y$	$x \rightarrow z \leftarrow y$	$(X \not\perp\!\!\!\perp Y   Z)$
	$x \leftarrow z \leftarrow y$	$\downarrow$ $w$	

- ✗ CI<sub>1</sub> : (Wet  $\perp\!\!\!\perp$  Sprinkler)
- ✗ CI<sub>2</sub> : (Wet  $\perp\!\!\!\perp$  Climate | Sprinkler)
- ✓ CI<sub>3</sub> : (Rain  $\perp\!\!\!\perp$  Slippery | Wet)
- ✓ CI<sub>4</sub> : (Season  $\perp\!\!\!\perp$  Wet | Sprinkler, Rain)
- ✗ CI<sub>5</sub> : (Sprinkler  $\perp\!\!\!\perp$  Rain | Climate, Wet)




# THE SECOND LAYER ON CAUSAL HIERARCHY: INTERVENTIONAL (What if I do $X=x$ ? )

# OUTLINE

---

- Introduction
  - \* Data, data, data...
  - \* Basic definitions
- Technical Results
  - \* The truncated product formula
  - \* The back-door adjustment formula
  - \* The front-door adjustment formula
  - \* The do-calculus
- Tasks
  - \* Policy evaluation
  - \* Generalizability & Robustness
  - \* Decision-making & RL

# CAUSAL INFERENCE: MOVING ACROSS REGIMES



- What happens when  $P$  changes?  
e.g., Infer whether less people would get cancer if we ban smoking.

$$Q = P(Cancer = \text{true} \mid do(Smoking = \text{no})) \text{ Not an aspect of } P^{45}$$

## Observation 1:

The distribution alone tells us nothing about change; it just describes static conditions of a population (under a specific regime).

## Observation 1:


The distribution alone tells us nothing about change; it just describes static conditions of a population (under a specific regime).

## Observation 2:

We need to be able to represent “change”, or how the population reacts when it undergoes change in regimes.


# THE BIG PICTURE: THE CHALLENGE OF CAUSAL INFERENCE

Real world



$$P(z, x, w, y)$$

Alternative world




$$P(y \mid \text{do}(x))$$

change

Z: age, sex  
X: action  
W: mediator  
Y: outcome


# THE BIG PICTURE: THE CHALLENGE OF CAUSAL INFERENCE

Real world



$$P(z, x, w, y)$$

Alternative world



$$P(y \mid \text{do}(x))$$


change

Z: age, sex  
X: action  
W: mediator  
Y: outcome

- Goal: how much Y **changes** with X if we **vary** X between two different **constants** free from the influence of Z.
- This is the definition of **causal effect**.


# METHOD FOR COMPUTING CAUSAL EFFECTS: RANDOMIZED EXPERIMENTS

## Real world



## change


# Alternative world




Z: age, sex  
X: action  
W: mediator  
Y: outcome

# METHOD FOR COMPUTING CAUSAL EFFECTS: RANDOMIZED EXPERIMENTS

Real world




Alternative world




change

$\text{do}(X_0)$




$\text{do}(X_1)$




Z: age, sex  
X: action  
W: mediator  
Y: outcome

# METHOD FOR COMPUTING CAUSAL EFFECTS: RANDOMIZED EXPERIMENTS

Real world




Alternative world




change

$\text{do}(X_0)$



$\text{do}(X_1)$



Z: age, sex  
X: action  
W: mediator  
Y: outcome


Randomization:

$$P(y \mid \text{do}(X_0))$$

$$P(y \mid \text{do}(X_1))$$

# COMPUTING CAUSAL EFFECTS FROM OBSERVATIONAL DATA


## Real world



$$P(z, x, w, y)$$

## change


# Alternative world



$$P(y \mid do(x))$$

Z: age, sex  
X: action  
W: mediator  
Y: outcome

# COMPUTING CAUSAL EFFECTS FROM OBSERVATIONAL DATA



# Questions:

- \* What is the relationship between  $P(z, x, w, y)$  and  $P(y | \text{do}(x))$ ?
  - \* Is  $P(y | \text{do}(x)) = P(y | x)$ ?

# EFFECT OF INTERVENTIONS

---

# EFFECT OF INTERVENTIONS

---

- Def. 3.2.1 (Causal Effect)

Given two disjoint sets of variables,  $X$  and  $Y$ , the **causal effect** of  $X$  on  $Y$ , denoted as  $P(y | \text{do}(x))$ , is a function from  $X$  to the space of probability distributions of  $Y$ .

# EFFECT OF INTERVENTIONS

---

- Def. 3.2.1 (Causal Effect)

Given two disjoint sets of variables,  $X$  and  $Y$ , the **causal effect** of  $X$  on  $Y$ , denoted as  $P(y | \text{do}(x))$ , is a function from  $X$  to the space of probability distributions of  $Y$ .

For each realization  $x$  of  $X$ ,  $P(y | \text{do}(x))$  gives the probability  $Y = y$  induced by **deleting** from the model all equations corresponding to variables in  $X$  and **substituting**  $X = x$  in the remaining equations.


# COMPUTING CAUSAL EFFECTS FROM OBSERVATIONAL DATA

---

**Queries:**

$$Q_1 = \Pr(\text{wet} \mid \text{Sprinkler} = \text{on})$$

$$Q_2 = \Pr(\text{wet} \mid \text{do}(\text{Sprinkler} = \text{on}))$$




# COMPUTING CAUSAL EFFECTS FROM OBSERVATIONAL DATA

---

**Queries:**

$$\begin{aligned} Q_1 &= \Pr(\text{wet} \mid \text{Sprinkler} = \text{on}) \\ &= P(p_1) + P(p_2) \end{aligned}$$

$$Q_2 = \Pr(\text{wet} \mid \text{do}(\text{Sprinkler} = \text{on}))$$




# COMPUTING CAUSAL EFFECTS FROM OBSERVATIONAL DATA

---

**Queries:**

$$\begin{aligned} \mathbf{Q}_1 &= \Pr(\text{wet} \mid \text{Sprinkler} = \text{on}) \\ &= P(p_1) + P(p_2) \end{aligned}$$

$$\mathbf{Q}_2 = \Pr(\text{wet} \mid \text{do}(\text{Sprinkler} = \text{on}))$$




# COMPUTING CAUSAL EFFECTS FROM OBSERVATIONAL DATA

---

**Queries:**

$$\begin{aligned} Q_1 &= \Pr(\text{wet} \mid \text{Sprinkler} = \text{on}) \\ &= P(p_1) + P(p_2) \end{aligned}$$

$$Q_2 = \Pr(\text{wet} \mid \text{do}(\text{Sprinkler} = \text{on}))$$




# COMPUTING CAUSAL EFFECTS FROM OBSERVATIONAL DATA

---

**Queries:**

$$\begin{aligned} Q_1 &= \Pr(\text{wet} \mid \text{Sprinkler} = \text{on}) \\ &= P(p_1) + P(p_2) \end{aligned}$$

$$Q_2 = \Pr(\text{wet} \mid \text{do}(\text{Sprinkler} = \text{on}))$$




# COMPUTING CAUSAL EFFECTS FROM OBSERVATIONAL DATA

---

**Queries:**

$$\begin{aligned} Q_1 &= \Pr(\text{wet} \mid \text{Sprinkler} = \text{on}) \\ &= P(p_1) + P(p_2) \end{aligned}$$

$$\begin{aligned} Q_2 &= \Pr(\text{wet} \mid \text{do}(\text{Sprinkler} = \text{on})) \\ &= P(p_1) \end{aligned}$$




# COMPUTING CAUSAL EFFECTS FROM OBSERVATIONAL DATA

---

**Queries:**

$$\begin{aligned} Q_1 &= \Pr(\text{wet} \mid \text{Sprinkler} = \text{on}) \\ &= P(p_1) + P(p_2) \end{aligned}$$

$$\begin{aligned} Q_2 &= \Pr(\text{wet} \mid \text{do}(\text{Sprinkler} = \text{on})) \\ &= P(p_1) \end{aligned}$$




# COMPUTING CAUSAL EFFECTS FROM OBSERVATIONAL DATA

Queries:

$$\begin{aligned} Q_1 &= \Pr(\text{wet} \mid \text{Sprinkler} = \text{on}) \\ &= P(p_1) + P(p_2) \end{aligned}$$

$$\begin{aligned} Q_2 &= \Pr(\text{wet} \mid \text{do}(\text{Sprinkler} = \text{on})) \\ &= P(p_1) \end{aligned}$$

$$\sum_{\text{Se}, \text{Ra}, \text{Sl}} P(\text{Se}) P(\text{Sp} \mid \text{Se}) P(\text{Ra} \mid \text{Se}) P(\text{We} \mid \text{Sp}, \text{Ra}) P(\text{Sl} \mid \text{We})$$




# COMPUTING CAUSAL EFFECTS FROM OBSERVATIONAL DATA

Queries:

$$\begin{aligned} Q_1 &= \Pr(\text{wet} \mid \text{Sprinkler} = \text{on}) \\ &= P(p_1) + P(p_2) \end{aligned}$$

$$\begin{aligned} Q_2 &= \Pr(\text{wet} \mid \text{do}(\text{Sprinkler} = \text{on})) \\ &= P(p_1) \end{aligned}$$

$$\sum_{\text{Se}, \text{Ra}, \text{Sl}} P(\text{Se}) P(\text{Sp} \mid \text{Se}) P(\text{Ra} \mid \text{Se}) P(\text{We} \mid \text{Sp, Ra}) P(\text{Sl} \mid \text{We})$$



# TRUNCATED FACTORIZATION PRODUCT (OPERATIONALIZING INTERVENTIONS)

---

Corollary (Truncated Factorization, Manipulation Thm., G-comp.):  
The distribution generated by an intervention  $do(X=x)$   
(in a **Markovian** model  $M$ ) is given by the truncated factorization:

$$P(v_1, v_2, \dots, v_n \mid do(x)) = \prod_{i \mid V_i \notin X} P(v_i \mid pa_i) \quad \Bigg| \quad X = x$$

# THE IDENTIFIABILITY PROBLEM

---

# THE IDENTIFIABILITY PROBLEM

---

- Def. 3.2.2 (Causal Effect Identifiability)

The **causal effect** of X on Y is said to be **identifiable** from a graph G if the quantity  $P(y | \text{do}(x))$  can be computed uniquely from a positive probability of the observed variables.

# THE IDENTIFIABILITY PROBLEM

---

- Def. 3.2.2 (Causal Effect Identifiability)

The **causal effect** of  $X$  on  $Y$  is said to be **identifiable** from a graph  $G$  if the quantity  $P(y | \text{do}(x))$  can be computed uniquely from a positive probability of the observed variables.


That is, if  $P_{M_1}(y | \text{do}(x)) = P_{M_2}(y | \text{do}(x))$  for every pair of models  $M_1$  and  $M_2$  with  $P_{M_1}(v) = P_{M_2}(v) > 0$  and  $G(M_1) = G(M_2) = G$ .

# THE IDENTIFIABILITY PROBLEM

- Def. 3.2.2 (Causal Effect Identifiability)

The **causal effect** of  $X$  on  $Y$  is said to be **identifiable** from a graph  $G$  if the quantity  $P(y | \text{do}(x))$  can be computed uniquely from a positive probability of the observed variables.


That is, if  $P_{M_1}(y | \text{do}(x)) = P_{M_2}(y | \text{do}(x))$  for every pair of models  $M_1$  and  $M_2$  with  $P_{M_1}(v) = P_{M_2}(v) > 0$  and  $G(M_1) = G(M_2) = G$ .



# MEANING OF IDENTIFIABILITY

---


$P(y|do(x))$  is  
identifiable in  $G$



# MEANING OF IDENTIFIABILITY

---

$P(y|do(x))$  is  
identifiable in  $G$




*All models*

# MEANING OF IDENTIFIABILITY

---


$P(y|do(x))$  is  
identifiable in  $G$



# MEANING OF IDENTIFIABILITY

---


$P(y|do(x))$  is  
identifiable in  $G$



# MEANING OF IDENTIFIABILITY

---


$P(y|do(x))$  is  
identifiable in  $G$



# MEANING OF IDENTIFIABILITY

---


$P(y|do(x))$  is  
identifiable in  $G$



# MEANING OF IDENTIFIABILITY

---

$P(y|do(x))$  is  
identifiable in  $G$




Formally,  
for any two causal  
models (encoding the  
unobserved nature),  
 $N_1, N_2$ , s.t.

# MEANING OF IDENTIFIABILITY

---

$P(y|do(x))$  is  
identifiable in  $G$




Formally,  
for any two causal  
models (encoding the  
unobserved nature),  
 $N_1, N_2$ , s.t.  
 $G(N_1) = G(N_2) = G$ ,

# MEANING OF IDENTIFIABILITY

---


$P(y|do(x))$  is  
identifiable in  $G$



Formally,  
for any two causal  
models (encoding the  
unobserved nature),  
 $N_1, N_2$ , s.t.  
 $G(N_1) = G(N_2) = G$ ,  
 $P_1(v) = P_2(v)$

# MEANING OF IDENTIFIABILITY

$P(y|do(x))$  is  
identifiable in  $G$




Formally,  
for any two causal  
models (encoding the  
unobserved nature),  
 $N_1, N_2$ , s.t.  
 $G(N_1) = G(N_2) = G$ ,

$P_1(v) = P_2(v)$ , then  
 $P_1(y | do(x)) =$   
 $P_2(y | do(x))$ .

# MEANING OF IDENTIFIABILITY

$P(y|do(x))$  is  
identifiable in  $G$




Formally,  
for any two causal  
models (encoding the  
unobserved nature),  
 $N_1, N_2$ , s.t.  
 $G(N_1) = G(N_2) = G$ ,

$P_1(v) = P_2(v)$ , then  
 $P_1(y | do(x)) =$   
 $P_2(y | do(x))$ .

# MEANING OF IDENTIFIABILITY (II)


$P(y|do(x))$  is  
identifiability in  $G$



# MEANING OF IDENTIFIABILITY (II)

$P(y|do(x))$  is  
identifiability in  $G$


$P(y|do(x))$  is  
**not identifiable** in  $G$



# MEANING OF IDENTIFIABILITY (II)


$P(y|do(x))$  is  
identifiability in  $G$

$P(y|do(x))$  is  
**not identifiable** in  $G$



# MEANING OF IDENTIFIABILITY (II)

$P(y|do(x))$  is  
identifiability in  $G$




Models  
inducing  
 $P(v)$

Same  
 $P(y | do(x))$

Models  
with  $G$

*All models*

$P(y|do(x))$  is  
not identifiable in  $G$



Different  
 $P(y | do(x))$  !

# ADJUSTMENT BY DIRECT PARENTS

---

- Theorem 3.2.5

Given a causal diagram  $G$  of any Markovian model in which a subset  $V$  of variables are measured, the causal effect  $Q = P(y | \text{do}(x))$  is identifiable whenever  $\{X, Y, \text{Pa}_x\} \subseteq V$ , that is, whenever  $X, Y$ , and all parents of variables  $X$  are measured. The expression of  $Q$  is then obtained by adjustment for  $\text{PA}_x$ , or

# ADJUSTMENT BY DIRECT PARENTS

---

- Theorem 3.2.5

Given a causal diagram  $G$  of any Markovian model in which a subset  $V$  of variables are measured, the causal effect  $Q = P(y | \text{do}(x))$  is identifiable whenever  $\{X, Y, \text{Pa}_x\} \subseteq V$ , that is, whenever  $X, Y$ , and all parents of variables  $X$  are measured. The expression of  $Q$  is then obtained by adjustment for  $\text{PA}_x$ , or

$$Q = \sum_{\text{PA}_x} P(y | x, \text{PA}_x = \text{pa}_x) P(\text{PA}_x = \text{pa}_x)$$

# IDENTIFICATION IN MARKOVIAN MODELS

---

- Corollary 3.2.6

Given the causal diagram  $G$  of any Markovian model in which all variables are measured, the causal effect  $Q = P(y | do(x))$  is identifiable for every subsets of variables  $X$  and  $Y$  and is obtained from the truncated factorization, i.e.,

# IDENTIFICATION IN MARKOVIAN MODELS

---

- Corollary 3.2.6

Given the causal diagram  $G$  of any Markovian model in which all variables are measured, the causal effect  $Q = P(y | \text{do}(x))$  is identifiable for every subsets of variables  $X$  and  $Y$  and is obtained from the truncated factorization, i.e.,



$$P(y | \text{do}(x)) = \sum_{\forall Y \setminus X} \prod_{\{i \mid x_i \notin X\}} P(x_i | \text{pa}_i)$$

# IF SEASON IS LATENT, IS THE EFFECT STILL COMPUTABLE?

Queries:

$$\begin{aligned} Q_1 &= \Pr(\text{wet} \mid \text{Sprinkler} = \text{on}) \\ &= P(p_1) + P(p_2) \end{aligned}$$

$$\begin{aligned} Q_2 &= \Pr(\text{wet} \mid \text{do}(\text{Sprinkler} = \text{on})) \\ &= P(p_1) \end{aligned}$$




# IF SEASON IS LATENT, IS THE EFFECT STILL COMPUTABLE?

Queries:

$$\begin{aligned} Q_1 &= \Pr(\text{wet} \mid \text{Sprinkler} = \text{on}) \\ &= P(p_1) + P(p_2) \end{aligned}$$

$$\begin{aligned} Q_2 &= \Pr(\text{wet} \mid \text{do}(\text{Sprinkler} = \text{on})) \\ &= P(p_1) \end{aligned}$$

$$\sum_{\text{Se}, \text{Ra}, \text{Sl}} P(\text{Se}) P(\text{Sp} \mid \text{Se}) P(\text{Ra} \mid \text{Se}) P(\text{We} \mid \text{Sp, Ra}) P(\text{Sl} \mid \text{We})$$




# IF SEASON IS LATENT, IS THE EFFECT STILL COMPUTABLE?

Queries:

$$\begin{aligned} Q_1 &= \Pr(\text{wet} \mid \text{Sprinkler} = \text{on}) \\ &= P(p_1) + P(p_2) \end{aligned}$$

$$\begin{aligned} Q_2 &= \Pr(\text{wet} \mid \text{do}(\text{Sprinkler} = \text{on})) \\ &= P(p_1) \end{aligned}$$

$$\sum_{\text{Se}, \text{Ra}, \text{Sl}} P(\text{Se}) P(\text{Sp} \mid \text{Se}) P(\text{Ra} \mid \text{Se}) P(\text{We} \mid \text{Sp, Ra}) P(\text{Sl} \mid \text{We})$$




# IF SEASON IS LATENT, IS THE EFFECT STILL COMPUTABLE?

**Queries:**

$$\begin{aligned} Q_1 &= \Pr(\text{wet} \mid \text{Sprinkler} = \text{on}) \\ &= P(p_1) + P(p_2) \end{aligned}$$

$$\begin{aligned} Q_2 &= \Pr(\text{wet} \mid \text{do}(\text{Sprinkler} = \text{on})) \\ &= P(p_1) \end{aligned}$$

$$\sum_{\text{Se}, \text{Ra}, \text{Sl}} P(\text{Se}) P(\text{Sp} \mid \text{Se}) P(\text{Ra} \mid \text{Se}) P(\text{We} \mid \text{Sp, Ra}) P(\text{Sl} \mid \text{We})$$




# IF SEASON IS LATENT, IS THE EFFECT STILL COMPUTABLE?

Queries:

$$\begin{aligned} Q_1 &= \Pr(\text{wet} \mid \text{Sprinkler} = \text{on}) \\ &= P(p_1) + P(p_2) \end{aligned}$$

$$\begin{aligned} Q_2 &= \Pr(\text{wet} \mid \text{do}(\text{Sprinkler} = \text{on})) \\ &= P(p_1) \end{aligned}$$


$$\sum_{\text{Se}, \text{Ra}, \text{Sl}} P(\text{Se}) P(\text{Sp} \mid \text{Se}) P(\text{Ra} \mid \text{Se}) P(\text{We} \mid \text{Sp, Ra}) P(\text{Sl} \mid \text{We})$$



# IF SEASON IS LATENT, IS THE EFFECT STILL COMPUTABLE?

Queries:

$$\begin{aligned} Q_1 &= \Pr(\text{wet} \mid \text{Sprinkler} = \text{on}) \\ &= P(p_1) + P(p_2) \end{aligned}$$



$$\begin{aligned} Q_2 &= \Pr(\text{wet} \mid \text{do}(\text{Sprinkler} = \text{on})) \\ &= P(p_1) \end{aligned}$$


$$\sum_{\text{Se}, \text{Ra}, \text{Sl}} P(\text{Se}) P(\text{Sp} \mid \text{Se}) P(\text{Ra} \mid \text{Se}) P(\text{We} \mid \text{Sp}, \text{Ra}) P(\text{Sl} \mid \text{We})$$

$$= \sum_{\text{Se}} P(\text{We} \mid \text{Sp}, \text{Se}) P(\text{Se})$$

# IF SEASON IS LATENT, IS THE EFFECT STILL COMPUTABLE?

Queries:

$$\begin{aligned} Q_1 &= \Pr(\text{wet} \mid \text{Sprinkler} = \text{on}) \\ &= P(p_1) + P(p_2) \end{aligned}$$



$$\begin{aligned} Q_2 &= \Pr(\text{wet} \mid \text{do}(\text{Sprinkler} = \text{on})) \\ &= P(p_1) \end{aligned}$$


$$\sum_{\text{Se}, \text{Ra}, \text{Sl}} P(\text{Se}) P(\text{Sp} \mid \text{Se}) P(\text{Ra} \mid \text{Se}) P(\text{We} \mid \text{Sp}, \text{Ra}) P(\text{Sl} \mid \text{We})$$

$$= \sum_{\text{Se}} P(\text{We} \mid \text{Sp}, \text{Se}) P(\text{Se}) = \sum_{\text{Ra}} P(\text{We} \mid \text{Sp}, \text{Ra}) P(\text{Ra})$$

# IF SEASON IS LATENT, IS THE EFFECT STILL COMPUTABLE?

Queries:

$$\begin{aligned} Q_1 &= \Pr(\text{wet} \mid \text{Sprinkler} = \text{on}) \\ &= P(p_1) + P(p_2) \end{aligned}$$



$$\begin{aligned} Q_2 &= \Pr(\text{wet} \mid \text{do}(\text{Sprinkler} = \text{on})) \\ &= P(p_1) \end{aligned}$$

$$\sum_{\text{Se}, \text{Ra}, \text{Sl}} P(\text{Se}) P(\text{Sp} \mid \text{Se}) P(\text{Ra} \mid \text{Se}) P(\text{We} \mid \text{Sp}, \text{Ra}) P(\text{Sl} \mid \text{We})$$


$$= \sum_{\text{Se}} P(\text{We} \mid \text{Sp}, \text{Se}) P(\text{Se}) = \sum_{\text{Ra}} P(\text{We} \mid \text{Sp}, \text{Ra}) P(\text{Ra})$$

Adjustment formula

# THE PROBLEM OF CONFOUNDING

---

**Goal:** Find the effect of  $X$  on  $Y$ ,  $Q = P(y|do(x))$ , given measurements on auxiliary variables  $Z_1, \dots, Z_k$



Can  $Q$  be estimated if only a subset  $Z$  can be measured?

# THE BACK-DOOR CRITERION

---

# THE BACK-DOOR CRITERION

---

- Definition 3.3.1 (Back-door)

A set  $Z$  satisfies the **back-door criterion** relative to an ordered pair of variables  $(X, Y)$  in a DAG  $G$  if:

- (i) no node in  $Z$  is a descendent of  $X$ ; and
- (ii)  $Z$  blocks every path between  $X$  and  $Y$  that contains an arrow into  $X$ .

# THE BACK-DOOR CRITERION

---

- **Definition 3.3.1 (Back-door)**

A set  $Z$  satisfies the **back-door criterion** relative to an ordered pair of variables  $(X, Y)$  in a DAG  $G$  if:

- (i) no node in  $Z$  is a descendent of  $X$ ; and
- (ii)  $Z$  blocks every path between  $X$  and  $Y$  that contains an arrow into  $X$ .

- **Theorem 3.3.2 (Back-door Adjustment)**

If a set  $Z$  satisfies the bd-c relative to  $(X, Y)$  then the causal effect of  $X$  on  $Y$  is identifiable and given by:

$$P(y | \text{do}(x)) = \sum_z P(y | x, z) P(z)$$

# THE BACK-DOOR CRITERION

## (graphical condition)


---

$P(y | do(x))$  is estimable if there is  
a set  $Z$  of variables that  **$d$ -separates  $X$  from  $Y$  in  $G_x$**

# THE BACK-DOOR CRITERION

(graphical condition)



$P(y | do(x))$  is estimable if there is  
a set  $Z$  of variables that  **$d$ -separates  $X$  from  $Y$  in  $G_x$**



# THE BACK-DOOR CRITERION


(graphical condition)

$P(y | do(x))$  is estimable if there is  
a set  $Z$  of variables that  **$d$ -separates  $X$  from  $Y$  in  $G_x$**




# GOING BEYOND ADJUSTMENT: THE FRONT-DOOR CRITERION

---



# GOING BEYOND ADJUSTMENT: THE FRONT-DOOR CRITERION

---



Suppose  $Z_1, Z_2, Z_3, Z_4$ , and  $Z_5$  are not measured.

Can we compute  $Q = P(y \mid do(x))$ ?

# THE FRONT-DOOR CRITERION

---

# THE FRONT-DOOR CRITERION

---

- Definition 3.3.3 (Front-door)

A set  $Z$  satisfies is said to satisfy the **front-door criterion** relative to an ordered pair of variables  $(X, Y)$  if:

- (i)  $Z$  intercepts all directed paths from  $X$  to  $Y$ ;
- (ii) There is no back-door path from  $X$  to  $Z$ ; and
- (iii) All back-door paths from  $Z$  to  $Y$  are blocked by  $X$ .

# THE FRONT-DOOR CRITERION

---

- Definition 3.3.3 (Front-door)

A set  $Z$  satisfies is said to satisfy the **front-door criterion** relative to an ordered pair of variables  $(X, Y)$  if:

- (i)  $Z$  intercepts all directed paths from  $X$  to  $Y$ ;
- (ii) There is no back-door path from  $X$  to  $Z$ ; and
- (iii) All back-door paths from  $Z$  to  $Y$  are blocked by  $X$ .


- Theorem 3.3.2 (Front-door Adjustment)

If a set  $Z$  satisfies the fd-c relative to  $(X, Y)$ , then the causal effect of  $X$  on  $Y$  is identifiable and given by:

$$P(y | \text{do}(x)) = \sum_z P(z | x) \sum_{x'} P(y | x', z) P(x')$$


# THE FRONT-DOOR CRITERION

---



# THE FRONT-DOOR CRITERION


---



$$P(x, y, z, u) = P(u) P(x | u) P(z | x) P(y | z, u)$$

# THE FRONT-DOOR CRITERION

---




$$P(x, y, z, u) = P(u) P(x | u) P(z | x) P(y | z, u)$$

$$P(y, z, u | \text{do}(x)) = P(z | x) P(y | z, u) P(u)$$

# THE FRONT-DOOR CRITERION

---



$$P(x, y, z, u) = P(u) P(x | u) P(z | x) P(y | z, u)$$

$$P(y, z, u | \text{do}(x)) = P(z | x) P(y | z, u) P(u)$$

$$P(y | \text{do}(x)) = \sum_z P(z | x) \sum_u P(y | z, u) P(u)$$

# CAUSAL CALCULUS (IDENTIFIABILITY REDUCED TO CALCULUS)

---

(or, do-calculus)

Theorem 3.4.1. The following transformations are valid for every interventional distribution generated by a **structural causal model  $M$** :

# CAUSAL CALCULUS (IDENTIFIABILITY REDUCED TO CALCULUS)

---

(or, do-calculus)

Theorem 3.4.1. The following transformations are valid for every interventional distribution generated by a **structural causal model**  $M$ :

**Rule 1:** Ignoring observations

$$P(y \mid do(x), z, w) = P(y \mid do(x), w), \quad \text{if } (Y \perp\!\!\!\perp Z \mid X, W)_{G_{\overline{X}}}$$

# CAUSAL CALCULUS (IDENTIFIABILITY REDUCED TO CALCULUS)

---

(or, do-calculus)

Theorem 3.4.1. The following transformations are valid for every interventional distribution generated by a **structural causal model**  $M$ :

**Rule 1:** Ignoring observations

$$P(y \mid \text{do}(x), z, w) = P(y \mid \text{do}(x), w), \quad \text{if } (Y \perp\!\!\!\perp Z \mid X, W)_{G_{\overline{X}}}$$

**Rule 2:** Action/observation exchange

$$P(y \mid \text{do}(x), \text{do}(z), w) = P(y \mid \text{do}(x), z, w), \quad \text{if } (Y \perp\!\!\!\perp Z \mid X, W)_{G_{\overline{X}Z}}$$

# CAUSAL CALCULUS (IDENTIFIABILITY REDUCED TO CALCULUS)

---

(or, do-calculus)

Theorem 3.4.1. The following transformations are valid for every interventional distribution generated by a **structural causal model**  $M$ :

**Rule 1:** Ignoring observations

$$P(y \mid \text{do}(x), z, w) = P(y \mid \text{do}(x), w), \quad \text{if } (Y \perp\!\!\!\perp Z \mid X, W)_{G_{\overline{X}}}$$

**Rule 2:** Action/observation exchange

$$P(y \mid \text{do}(x), \text{do}(z), w) = P(y \mid \text{do}(x), z, w), \quad \text{if } (Y \perp\!\!\!\perp Z \mid X, W)_{G_{\overline{X}Z}}$$


**Rule 3:** Ignoring actions

$$P(y \mid \text{do}(x), \text{do}(z), w) = P(y \mid \text{do}(x), w), \quad \text{if } (Y \perp\!\!\!\perp Z \mid X, W)_{G_{\overline{X}Z(W)}}$$

where  $Z(W)$  is the set of  $Z$ -nodes that are not ancestors of any  $W$ -node in  $G_x$ .


# DERIVATION IN CAUSAL CALCULUS

---



# DERIVATION IN CAUSAL CALCULUS


---



$$P(c \mid do(s))$$

# DERIVATION IN CAUSAL CALCULUS

---




$$P(c | \text{do}(s)) = \sum_t P(c | \text{do}(s), t) P(t | \text{do}(s))$$

Probability Axioms


$$= \sum_t P(c | \text{do}(s), \text{do}(t)) P(t | \text{do}(s))$$

Rule 2



$$= \sum_t P(c | \text{do}(s), \text{do}(t)) P(t | s)$$

Rule 2



$$= \sum_t P(c | \text{do}(t)) P(t | s)$$

Rule 3



$$= \sum_{s'} \sum_t P(c | \text{do}(t), s') P(s' | \text{do}(t)) P(t | s)$$

Probability Axioms


$$= \sum_{s'} \sum_t P(c | t, s') P(s' | \text{do}(t)) P(t | s)$$

Rule 2




$$= \sum_{s'} \sum_t P(c | t, s') P(s') P(t | s)$$

Rule 3



# DERIVATION IN CAUSAL CALCULUS




$$P(c | do(s)) = \sum_t P(c | do(s), t) P(t | do(s))$$

Probability Axioms


$$= \sum_t P(c | do(s), do(t)) P(t | do(s))$$

Rule 2



$$= \sum_t P(c | do(s), do(t)) P(t | s)$$

Rule 2



$$= \sum_t P(c | do(t)) P(t | s)$$

Rule 3



$$= \sum_{s'} \sum_t P(c | do(t), s') P(s' | do(t)) P(t | s)$$

Probability Axioms

$$= \sum_{s'} \sum_t P(c | t, s') P(s' | do(t)) P(t | s)$$

Rule 2




$$= \sum_{s'} \sum_t P(c | t, s') P(s') P(t | s)$$

Rule 3



# DERIVATION IN CAUSAL CALCULUS




$$P(c | do(s)) = \sum_t P(c | do(s), t) P(t | do(s))$$

Probability Axioms


$$= \sum_t P(c | do(s), do(t)) P(t | do(s))$$

Rule 2




$$= \sum_t P(c | do(s), do(t)) P(t | s)$$

Rule 2



$$= \sum_t P(c | do(t)) P(t | s)$$

Rule 3




$$= \sum_{s'} \sum_t P(c | do(t), s') P(s' | do(t)) P(t | s)$$

Probability Axioms


$$= \sum_{s'} \sum_t P(c | t, s') P(s' | do(t)) P(t | s)$$

Rule 2



$$= \sum_{s'} \sum_t P(c | t, s') P(s') P(t | s)$$

Rule 3



*estimand* ( $f(P(z, x, y))$ )

# CBNs WITH LATENT VARIABLES

---

**Definition (Semi-Markovian Model):**

The distribution generated by an intervention  $do(X=x)$  in a **Semi-Markovian** model  $M$  is given by the (extended) truncated factorization product, namely,

$$P(v_1, v_2, \dots, v_n \mid do(x)) = \sum_u \prod_i P(v_i \mid pa_i, u_i) \prod_i P(u_i)$$

**Example:**

\* Assume that in our original graph  $Z$  was unobserved.



# CBNs WITH LATENT VARIABLES


**Definition (Semi-Markovian Model):**

The distribution generated by an intervention  $do(X=x)$  in a **Semi-Markovian** model  $M$  is given by the (extended) truncated factorization product, namely,

$$P(v_1, v_2, \dots, v_n \mid do(x)) = \sum_u \prod_i P(v_i \mid pa_i, u_i) \prod_i P(u_i)$$


**Example:**

\* Assume that in our original graph  $Z$  was unobserved.



# NON-IDENTIFIABLE MODELS

$P(y|do(x))$  is  
identifiability in  $G$




Models  
inducing  
 $P(v)$

Same  
 $P(y | do(x))$

Models  
with  $G$

*All models*


$P(y|do(x))$  is  
not identifiable in  $G$




Different  
 $P(y | do(x)) !$

# NON-IDENTIFIABLE MODELS (EXAMPLES)

---




# NON-IDENTIFIABLE MODELS (EXAMPLES)




- Bow-arc observation

# NON-IDENTIFIABLE MODELS (EXAMPLES)




- Bow-arc observation
- $P(y \mid \text{do}(x, z_2))$  versus  $P(y \mid \text{do}(x), z_2)$


# NON-IDENTIFIABLE MODELS (EXAMPLES)




(a)




(b)




(c)




(d)




(e)



(f)




(g)




(h)


# IDENTIFIABLE MODELS (EXAMPLES)




(a)




(b)




(c)




(d)



(e)



(f)



(g)

# IDENTIFYING THE EFFECT OF CONDITIONAL ACTIONS

---

# IDENTIFYING THE EFFECT OF CONDITIONAL ACTIONS

---

$P(y | \text{do}(X = g(z)))$  = the distribution of Y under the policy  
 $\text{do}(X = g(z))$ .

# IDENTIFYING THE EFFECT OF CONDITIONAL ACTIONS

---

$P(y | \text{do}(X = g(z)))$  = the distribution of  $Y$  under the policy  $\text{do}(X = g(z))$ .

To compute  $P(y | \text{do}(X = g(z)))$ , we condition on  $Z$  and write,

# IDENTIFYING THE EFFECT OF CONDITIONAL ACTIONS

---

$P(y | \text{do}(X = g(z)))$  = the distribution of  $Y$  under the policy  $\text{do}(X = g(z))$ .

To compute  $P(y | \text{do}(X = g(z)))$ , we condition on  $Z$  and write,

$$\begin{aligned} P(y | \text{do}(X = g(z))) &= \sum_z P(y | \text{do}(X = g(z)), z) P(z | \text{do}(X = g(z))) \\ &= \sum_z P(y | \text{do}(X = g(z)), z) P(z) \\ &= E_z[P(y | \text{do}(x), z) | x = g(z)] \end{aligned}$$

# IDENTIFYING THE EFFECT OF CONDITIONAL ACTIONS

---

$P(y | \text{do}(X = g(z)))$  = the distribution of  $Y$  under the policy  $\text{do}(X = g(z))$ .

To compute  $P(y | \text{do}(X = g(z)))$ , we condition on  $Z$  and write,

$$\begin{aligned} P(y | \text{do}(X = g(z))) \\ &= \sum_z P(y | \text{do}(X = g(z)), z) P(z | \text{do}(X = g(z))) \\ &= \sum_z P(y | \text{do}(X = g(z)), z) P(z) \\ &= E_z[P(y | \text{do}(x), z) | x = g(z)] \end{aligned}$$

(Conditioning on  $Z$  might create dependencies that will prevent the successful reduction of  $P(y | \text{do}(x), z)$  to a do-free expression. )

# IDENTIFYING THE EFFECT OF STOCHASTIC POLICIES

---

# IDENTIFYING THE EFFECT OF STOCHASTIC POLICIES

---

**Stochastic policy:** enforce the intervention  $\text{do}(X = x)$  with probability  $P^*(x | z)$

# IDENTIFYING THE EFFECT OF STOCHASTIC POLICIES

---

**Stochastic policy:** enforce the intervention  $\text{do}(X = x)$  with probability  $P^*(x | z)$

Given  $Z = z$ , the intervention  $\text{do}(X = x)$  will occur with probability  $P^*(x | z)$  and will produce a causal effect given by  $P(y | \text{do}(x), z)$ . Averaging over  $x$  and  $z$  gives

# IDENTIFYING THE EFFECT OF STOCHASTIC POLICIES

---

**Stochastic policy:** enforce the intervention  $\text{do}(X = x)$  with probability  $P^*(x | z)$

Given  $Z = z$ , the intervention  $\text{do}(X = x)$  will occur with probability  $P^*(x | z)$  and will produce a causal effect given by  $P(y | \text{do}(x), z)$ . Averaging over  $x$  and  $z$  gives

$$P(y)_{|P^*(x | z)} = \sum_x \sum_z P(y | \text{do}(x), z) P^*(x | z) P(z)$$

# IDENTIFYING THE EFFECT OF STOCHASTIC POLICIES

---

**Stochastic policy:** enforce the intervention  $\text{do}(X = x)$  with probability  $P^*(x | z)$




Given  $Z = z$ , the intervention  $\text{do}(X = x)$  will occur with probability  $P^*(x | z)$  and will produce a causal effect given by  $P(y | \text{do}(x), z)$ . Averaging over  $x$  and  $z$  gives

$$P(y)_{| P^*(x | z)} = \sum_x \sum_z P(y | \text{do}(x), z) P^*(x | z) P(z)$$


$P^*(x | z)$  is specified externally. Therefore, the identifiability of  $P(y | \text{do}(x), z)$  is a necessary and sufficient condition for the identifiability of any stochastic policy that shapes the distribution of  $X$  by the outcome  $Z$ .

# IDENTIFYING THE EFFECT OF PLANS

---




# IDENTIFYING THE EFFECT OF PLANS




$$P(y \mid \text{do}(x_1, x_2)) = P(y \mid x_1, \text{do}(x_2))$$

# IDENTIFYING THE EFFECT OF PLANS




$$\begin{aligned} P(y | \text{do}(x_1, x_2)) &= P(y | x_1, \text{do}(x_2)) \\ &= \sum_z P(y | x_1, \text{do}(x_2), z) P(z | x_1, \text{do}(x_2)) \end{aligned}$$

# IDENTIFYING THE EFFECT OF PLANS



$$\begin{aligned} P(y | \text{do}(x_1, x_2)) &= P(y | x_1, \text{do}(x_2)) \\ &= \sum_z P(y | x_1, \text{do}(x_2), z) P(z | x_1, \text{do}(x_2)) \\ &= \sum_z P(y | x_1, \text{do}(x_2), z) P(z | x_1) \end{aligned}$$

# IDENTIFYING THE EFFECT OF PLANS



$$\begin{aligned} P(y \mid \text{do}(x_1, x_2)) &= P(y \mid x_1, \text{do}(x_2)) \\ &= \sum_z P(y \mid x_1, \text{do}(x_2), z) P(z \mid x_1, \text{do}(x_2)) \\ &= \sum_z P(y \mid x_1, \text{do}(x_2), z) P(z \mid x_1) \\ &= \sum_z P(y \mid x_1, x_2, z) P(z \mid x_1) \end{aligned}$$

# Simpson's Paradox

(Pearson, 1899, Yule, 1903, Simpson 1951)

---

# Simpson's Paradox

## (Pearson, 1899, Yule, 1903, Simpson 1951)

---

- Pearson's shock: “spurious correlation”

We are thus forced to the conclusion that a mixture of heterogenous groups, each of which exhibits in itself no organic correlation, will exhibit a greater or less amount of correlation. This correlation may properly be called spurious, yet as it is almost impossible to guarantee the absolute homogeneity of any community, our results for correlation are always liable to an error, the amount of which cannot be foretold. To those who persist on looking upon all correlation as cause and effect, the fact that correlation can be produced between two quite uncorrelated characters A and B by taking an artificial mixture of two closely allied races, must come as rather a shock.

[Pearson, Lee, Brandy-Moore (1899)]

- Causation = perfect correlation
- “Not all correlations are causal” (Aldrich, 1994)

# Simpson's Paradox

(Pearson, 1899, Yule, 1903, Simpson 1951)

---

- e.g., UC Berkeley's alleged sex bias in graduate admission (Science, 1975)

Overall, data showed a higher rate of admission among male applicants, but, broken by departments, data showed a slight bias in favor of admitting female applicants.

- e.g., “reverse regression” (1970-1980)

Should one, in salary discrimination cases, compare salaries of equally qualified men and women, or, instead, compare qualifications of equally paid men and woman.  
(Opposite conclusions.)

# Simpson's Paradox

## (Robot: seeing and making sense)

	surv. (Y)	not-surv. ( $\neg Y$ )		Recovery Rate
drug (X)	20	20	40	50%
no-drug ( $\neg X$ )	16	24	40	40%
	36	44		



male ( $\neg F$ )	surv. (Y)	not-surv.		Recovery Rate
drug (X)	18	12	30	60%
no-drug ( $\neg X$ )	7	3	10	70%
	25	15	40	

female (F)	surv. (Y)	not-surv. ( $\neg Y$ )		Recovery Rate
drug (X)	2	8	10	20%
no-drug ( $\neg X$ )	9	21	30	30%
	11	29	40	

$P(Y | F, X) < P(Y | F, \neg X)$   
 $P(Y | \neg F, X) < P(Y | \neg F, \neg X)$   
but  
 $P(Y | X) > P(Y | \neg X) !!$

# Simpson's paradox

---

- Our tables implied what was deemed a paradox:

# Simpson's paradox

---

- Our tables implied what was deemed a paradox:

$$P(Y | F, X) < P(Y | F, \neg X)$$

# Simpson's paradox

---

- Our tables implied what was deemed a paradox:

$$P(Y | F, X) < P(Y | F, \neg X)$$

$$P(Y | \neg F, X) < P(Y | \neg F, \neg X)$$

# Simpson's paradox

---

- Our tables implied what was deemed a paradox:

$$P(Y | F, X) < P(Y | F, \neg X)$$

$$P(Y | \neg F, X) < P(Y | \neg F, \neg X)$$

and

# Simpson's paradox

---

- Our tables implied what was deemed a paradox:

$$P(Y | F, X) < P(Y | F, \neg X)$$

$$P(Y | \neg F, X) < P(Y | \neg F, \neg X)$$

and

$$P(Y | X) > P(Y | \neg X)$$

# Simpson's paradox

---

- Our tables implied what was deemed a paradox:

$$P(Y | F, X) < P(Y | F, \neg X)$$

$$P(Y | \neg F, X) < P(Y | \neg F, \neg X)$$

and

$$P(Y | X) > P(Y | \neg X)$$

# Dissolving the “paradox” with do-operator

---

- Our tables implied a deemed paradox:

$$P(Y | F, X) < P(Y | F, \neg X)$$

$$P(Y | \neg F, X) < P(Y | \neg F, \neg X)$$

and

$$P(Y | X) > P(Y | \bar{X})$$

- But if we re-write this using the do-notation:

$$P(Y | F, \text{do}(X)) < P(Y | F, \text{do}(\neg X))$$

$$P(Y | \neg F, \text{do}(X)) < P(Y | \neg F, \text{do}(\neg X))$$

# Dissolving the “paradox” with do-operator

---

- Our tables implied a deemed paradox:

$$P(Y | F, X) < P(Y | F, \neg X)$$

$$P(Y | \neg F, X) < P(Y | \neg F, \neg X)$$

and

$$P(Y | X) > P(Y | \neg X)$$

- But if we re-write this using the do-notation:

$$P(Y | F, \text{do}(X)) < P(Y | F, \text{do}(\neg X))$$

$$P(Y | \neg F, \text{do}(X)) < P(Y | \neg F, \text{do}(\neg X))$$

- It will not be possible to obtain:

$$P(Y | \text{do}(X)) > P(Y | \text{do}(\neg X))$$

# Proof

---

# Proof

---

- $P(Y | \text{do}(X)) = P(Y | \text{do}(X), F) P(F | \text{do}(X))$   
+  $P(Y | \text{do}(X), \neg F) P(\neg F | \text{do}(X))$

# Proof

---

- $$\begin{aligned} P(Y | \text{do}(X)) &= P(Y | \text{do}(X), F) P(F | \text{do}(X)) \\ &\quad + P(Y | \text{do}(X), \neg F) P(\neg F | \text{do}(X)) \\ &= P(Y | \text{do}(X), F) P(F) + P(Y | \text{do}(X), \neg F) P(\neg F) \end{aligned}$$

# Proof

---

- $$\begin{aligned} P(Y \mid \text{do}(X)) &= P(Y \mid \text{do}(X), F) P(F \mid \text{do}(X)) \\ &\quad + P(Y \mid \text{do}(X), \neg F) P(\neg F \mid \text{do}(X)) \\ &= P(Y \mid \text{do}(X), F) P(F) + P(Y \mid \text{do}(X), \neg F) P(\neg F) \end{aligned}$$
- $P(Y \mid \text{do}(\neg X)) = P(Y \mid \text{do}(\neg X), F) P(F) + P(Y \mid \text{do}(\neg X), \neg F) P(\neg F)$

# Proof

---

- $$\begin{aligned} P(Y | \text{do}(X)) &= P(Y | \text{do}(X), F) P(F | \text{do}(X)) \\ &\quad + P(Y | \text{do}(X), \neg F) P(\neg F | \text{do}(X)) \\ &= P(Y | \text{do}(X), F) P(F) + P(Y | \text{do}(X), \neg F) P(\neg F) \end{aligned}$$
- $P(Y | \text{do}(\neg X)) = P(Y | \text{do}(\neg X), F) P(F) + P(Y | \text{do}(\neg X), \neg F) P(\neg F)$
- And if assumptions hold:

# Proof

---

- $$\begin{aligned} P(Y | \text{do}(X)) &= P(Y | \text{do}(X), F) P(F | \text{do}(X)) \\ &\quad + P(Y | \text{do}(X), \neg F) P(\neg F | \text{do}(X)) \\ &= P(Y | \text{do}(X), F) P(F) + P(Y | \text{do}(X), \neg F) P(\neg F) \end{aligned}$$
- $P(Y | \text{do}(\neg X)) = P(Y | \text{do}(\neg X), F) P(F) + P(Y | \text{do}(\neg X), \neg F) P(\neg F)$
- And if assumptions hold:
$$\begin{aligned} P(Y | \text{do}(X), F) &< P(Y | \text{do}(\neg X), F) \\ P(Y | \text{do}(X), \neg F) &< P(Y | \text{do}(\neg X), \neg F) \end{aligned}$$

# Proof

---

- $$\begin{aligned} P(Y | \text{do}(X)) &= P(Y | \text{do}(X), F) P(F | \text{do}(X)) \\ &\quad + P(Y | \text{do}(X), \neg F) P(\neg F | \text{do}(X)) \\ &= P(Y | \text{do}(X), F) P(F) + P(Y | \text{do}(X), \neg F) P(\neg F) \end{aligned}$$
- $P(Y | \text{do}(\neg X)) = P(Y | \text{do}(\neg X), F) P(F) + P(Y | \text{do}(\neg X), \neg F) P(\neg F)$
- And if assumptions hold:
$$\begin{aligned} P(Y | \text{do}(X), F) &< P(Y | \text{do}(\neg X), F) \\ P(Y | \text{do}(X), \neg F) &< P(Y | \text{do}(\neg X), \neg F) \end{aligned}$$
- Then,
$$P(Y | \text{do}(X)) < P(Y | \text{do}(\neg X))$$


# WOULD DECISION-MAKING BE DIFFERENT WITH A DIFFERENT STORY?

---

# WOULD DECISION-MAKING BE DIFFERENT WITH A DIFFERENT STORY?


---

G<sub>1</sub>:




# WOULD DECISION-MAKING BE DIFFERENT WITH A DIFFERENT STORY?

---




# WOULD DECISION-MAKING BE DIFFERENT WITH A DIFFERENT STORY?

---



# WOULD DECISION-MAKING BE DIFFERENT WITH A DIFFERENT STORY?

---



# THE SURE THING PRINCIPLE

(formalizing Savage, 1972)

---

# THE SURE THING PRINCIPLE

(formalizing Savage, 1972)

---

Theorem 6.1.1. An **action X** that increases the probability of an event **Y** in **each subpopulation** must also increase the probability of **Y** in **the population as a whole**, provided that the action does not change the distribution of the sub-populations.

# THE SURE THING PRINCIPLE

(formalizing Savage, 1972)

---

Theorem 6.1.1. An **action X** that increases the probability of an event **Y** in **each subpopulation** must also increase the probability of **Y** in **the population as a whole**, provided that the action does not change the distribution of the sub-populations.

Summary.

- **Any** statistical relationship can be reversed by including additional factors in the analysis
- “Reversal” is possible in the calculus of proportions, but impossible in the calculus of causes
- People thinks causes, not proportions

# SUMMARY

---

# SUMMARY

---

- The do-calculus provides a syntactical characterization to the problem of policy evaluation for static and dynamic interventions.
- The problem of confounding and identification is essentially solved, non-parametrically.
- Simpson's Paradox is mathematized and dissolved.
- Applications are pervasive in the social and health sciences as well as in statistics, machine learning, and artificial intelligence.

# OUTLINE

---

- Introduction
  - \* Data, data, data...
  - \* Basic definitions
- Technical Results
  - \* The truncated product formula
  - \* The back-door adjustment formula
  - \* The front-door adjustment formula
  - \* The do-calculus
- Tasks
  - \* Policy evaluation
  - \* Generalizability and Robustness
  - \* Decision-making & RL

# OUTLINE

---

# SUMMARY - BIG PICTURE

---

1 query

2 model

3 data

# SUMMARY - BIG PICTURE

---

## 1 query

$$Q = P^*(y \mid \text{do}(x))$$

## 2 model

## 3 data


# SUMMARY - BIG PICTURE

---

1 query

$$Q = P^*(y \mid \text{do}(x))$$

2 model



3 data


# SUMMARY - BIG PICTURE

---

1 query

$$Q = P^*(y \mid \text{do}(x))$$

2 model



3 data

$$P(x, y, z)$$


# SUMMARY - BIG PICTURE

---

1 query

$$Q = P^*(y \mid \text{do}(x))$$

2 model




3 data

$$P(x, y, z)$$

inference  
engine

# SUMMARY - BIG PICTURE

---




# SUMMARY - BIG PICTURE

1 query

$$Q = P^*(y \mid \text{do}(x))$$

2 model



3 data

$$P(x, y, z)$$

With the current scientific knowledge about the problem (2) and the available data (3), is it possible to answer the research question (1)?

inference  
engine


→ solution  
(yes / no)

# SUMMARY - BIG PICTURE

1 query

$$Q = P^*(y \mid \text{do}(x))$$

2 model



3 data

$$P(x, y, z)$$

With the current scientific knowledge about the problem (2) and the available data (3), is it possible to answer the research question (1)?

inference  
engine


solution  
(yes / no)

# SUMMARY - BIG PICTURE

1 query

$$Q = P^*(y | \text{do}(x))$$

2 model



3 data

$$P(x, y, z)$$

With the current scientific knowledge about the problem (2) and the available data (3), is it possible to answer the research question (1)?

inference  
engine

solution  
(yes / no)

$$\begin{aligned} P(y | \text{do}(x)) &= \\ &\sum_z P(y | x, z) P(z) \end{aligned}$$

# MOTIVATION FOR DATA-FUSION

---

# MOTIVATION FOR DATA-FUSION

---

Target population  $\Pi^*$

(a) **US**

Census data  
available

Query of interest  $Q = P^*(y \mid \text{do}(x))$

# MOTIVATION FOR DATA-FUSION

---

Target population  $\Pi^*$

(a) **US**

Census data  
available

Query of interest  $Q = P^*(y \mid \text{do}(x))$

(b) **New York**

Survey data  
resembling target

# MOTIVATION FOR DATA-FUSION

---

Target population  $\Pi^*$

(a) **US**

Census data  
available

Query of interest  $Q = P^*(y \mid do(x))$

(b) **New York**

Survey data  
resembling target

(c) **Los Angeles**

Survey data  
younger population

# MOTIVATION FOR DATA-FUSION

---

Target population  $\Pi^*$

Query of interest  $Q = P^*(y \mid \text{do}(x))$

(a) **US**

Census data  
available

(b) **New York**

Survey data  
resembling target

(c) **Los Angeles**

Survey data  
younger population

(d) **Boston**

Age not recorded  
Mostly successful  
lawyers

(e) **San Francisco**

High post-treatment  
blood pressure

(f) **Texas**

Mostly Spanish  
subjects  
High attrition

(g) **Arkansas**

Randomized trial  
College students

(h) **Utah**

RCT, paid  
volunteers, mainly  
unemployed

(i) **Wyoming**

Natural experiment  
young athletes

# CAUSAL DATA SCIENCE

(“All data is not created equal”)

---

# CAUSAL DATA SCIENCE

## (“All data is not created equal”)

---

- Heterogenous datasets are pervasive in the empirical sciences since the data is collected:

# CAUSAL DATA SCIENCE

## (“All data is not created equal”)

---

- Heterogenous datasets are pervasive in the empirical sciences since the data is collected:
  - under different experimental conditions,

# CAUSAL DATA SCIENCE

## (“All data is not created equal”)

---

- Heterogenous datasets are pervasive in the empirical sciences since the data is collected:
  - under different experimental conditions,
  - the underlying populations are different,

# CAUSAL DATA SCIENCE

## (“All data is not created equal”)

---

- Heterogenous datasets are pervasive in the empirical sciences since the data is collected:
  - under different experimental conditions,
  - the underlying populations are different,
  - the sampling procedure is not random,

# CAUSAL DATA SCIENCE

## (“All data is not created equal”)

---

- Heterogenous datasets are pervasive in the empirical sciences since the data is collected:
  - under different experimental conditions,
  - the underlying populations are different,
  - the sampling procedure is not random,
  - the treatment assignment is not random,

# CAUSAL DATA SCIENCE

## (“All data is not created equal”)

---

- Heterogenous datasets are pervasive in the empirical sciences since the data is collected:
  - under different experimental conditions,
  - the underlying populations are different,
  - the sampling procedure is not random,
  - the treatment assignment is not random,
  - many variables are not measured.

# CAUSAL DATA SCIENCE

## (“All data is not created equal”)

---

- Heterogenous datasets are pervasive in the empirical sciences since the data is collected:
  - under different experimental conditions,
  - the underlying populations are different,
  - the sampling procedure is not random,
  - the treatment assignment is not random,
  - many variables are not measured.
- All these dimensions are now formalized.

# CAUSAL DATA SCIENCE

## (“All data is not created equal”)

---

- Heterogenous datasets are pervasive in the empirical sciences since the data is collected:
  - under different experimental conditions,
  - the underlying populations are different,
  - the sampling procedure is not random,
  - the treatment assignment is not random,
  - many variables are not measured.
- All these dimensions are now formalized.
- And there are conditions and algorithms to decide what is “entailed” from a certain data collection.

# CAUSAL DATA FUSION

## (“All data is noisy”)

- Heterogenous datasets from different empirical sciences sit side-by-side:
  - under different experiments,
  - the underlying populations are different,
  - the sampling procedure is not random,
  - the treatment assignment is not random,
  - many variables are not measured.
- All these dimensions are now formalized.
- And there are conditions and algorithms to decide what is “entailed” from a certain data collection.

PNAS  
Proceedings of the National Academy of Sciences of the United States of America

Home Articles Front Matter News Podcasts Authors

NEW RESEARCH IN Physical Sciences Social Sciences

Causal inference and the data-fusion problem

Elias Bareinboim and Judea Pearl

PNAS July 5, 2016 113 (27) 7345-7352; published ahead of print July 5, 2016 <https://doi.org/10.1073/pnas.1510507113>

Edited by Richard M. Shiffrin, Indiana University, Bloomington, IN, and approved March 15, 2016 (received for review June 29, 2015)

Check for updates

# CAUSAL DATA SCIENCE

## (“All data is not created equal”)

---


- Heterogenous datasets are pervasive in the empirical sciences since the data is collected:
  - under different experimental conditions,
  - the underlying populations are different,
  - the sampling procedure is not random,
  - the treatment assignment is not random,
  - many variables are not measured.
- All these dimensions are now formalized.
- And there are conditions and algorithms to decide what is “entailed” from a certain data collection.

# HETEROGENEOUS DATASETS

	Dataset 1	Dataset 2	...	Dataset n
<b>Target</b> $Q = P^*(y \mid \text{do}(x))$				


Population	Los Angeles	New York	Texas
Obs. / Exp.	Experimental	Observational	Experimental
Treat. Assign.	Randomized $Z_1$	-	Randomized $Z_2$
Sampling	Selection on Age	Selection on $\subseteq \subseteq$	-
Measured	$X_1, Z_1, W, M, Y_1$	$X_1, X_2, Z_1, N, Y_2$	$X_2, Z_1, W, L, M,$ $Y_1$

# HETEROGENEOUS DATASETS



d <sub>1</sub>	Population	Los Angeles	New York	Texas
Obs. / Exp.	Experimental	Observational	Experimental	
Treat. Assign.	Randomized Z <sub>1</sub>	-	Randomized Z <sub>2</sub>	
Sampling	Selection on Age	Selection on SES	-	
Measured	X <sub>1</sub> , Z <sub>1</sub> , W, M, Y <sub>1</sub>	X <sub>1</sub> , X <sub>2</sub> , Z <sub>1</sub> , N, Y <sub>2</sub>	X <sub>2</sub> , Z <sub>1</sub> , W, L, M, Y <sub>1</sub>	

# HETEROGENEOUS DATASETS



d <sub>1</sub>	Population	Los Angeles	New York	Texas
Obs. / Exp.	Experimental	Observational	Experimental	
Treat. Assign.	Randomized Z <sub>1</sub>	-	Randomized Z <sub>2</sub>	
Sampling	Selection on Age	Selection on SES	-	
Measured	X <sub>1</sub> , Z <sub>1</sub> , W, M, Y <sub>1</sub>	X <sub>1</sub> , X <sub>2</sub> , Z <sub>1</sub> , N, Y <sub>2</sub>	X <sub>2</sub> , Z <sub>1</sub> , W, L, M, Y <sub>1</sub>	

# HETEROGENEOUS DATASETS

	Dataset 1	Dataset 2	...	Dataset n
Target				
$Q = P^*(y   do(x))$				

$d_1$	Population	Los Angeles	New York	Texas
$d_2$	Obs. / Exp.	Experimental	Observational	Experimental
	Treat. Assign.	Randomized $Z_1$	-	Randomized $Z_2$
	Sampling	Selection on Age	Selection on $CES$	-
	Measured	$X_1, Z_1, W, M, Y_1$	$X_1, X_2, Z_1, N, Y_2$	$X_2, Z_1, W, L, M, Y_1$

# HETEROGENEOUS DATASETS

	Dataset 1	Dataset 2	...	Dataset n
d <sub>1</sub>	Population	Los Angeles	New York	Texas
d <sub>2</sub>	Obs. / Exp.	Experimental	Observational	Experimental
	Treat. Assign.	Randomized Z <sub>1</sub>	-	Randomized Z <sub>2</sub>
	Sampling	Selection on Age	Selection on SES	-
	Measured	X <sub>1</sub> , Z <sub>1</sub> , W, M, Y <sub>1</sub>	X <sub>1</sub> , X <sub>2</sub> , Z <sub>1</sub> , N, Y <sub>2</sub>	X <sub>2</sub> , Z <sub>1</sub> , W, L, M, Y <sub>1</sub>

# HETEROGENEOUS DATASETS

	Dataset 1	Dataset 2	...	Dataset n
	Target $Q = P^*(y   do(x))$			
d <sub>1</sub>	Population	Los Angeles	New York	Texas
d <sub>2</sub>	Obs. / Exp.	Experimental	Observational	Experimental
d <sub>3</sub>	Treat. Assign.	Randomized $Z_1$	-	Randomized $Z_2$
d <sub>3</sub>	Sampling	Selection on Age	Selection on $\subseteq \subseteq$	-
	Measured	$X_1, Z_1, W, M, Y_1$	$X_1, X_2, Z_1, N, Y_2$	$X_2, Z_1, W, L, M, Y_1$

# HETEROGENEOUS DATASETS

	Dataset 1	Dataset 2	...	Dataset n
<b>Target</b> $Q = P^*(y \mid \text{do}(x))$				
$d_1$	Population	Los Angeles	New York	Texas
$d_2$	Obs. / Exp.	Experimental	Observational	Experimental
$d_3$	Treat. Assign.	Randomized $Z_1$	-	Randomized $Z_2$
	Sampling	Selection on Age	Selection on SES	-
	Measured	$X_1, Z_1, W, M, Y_1$	$X_1, X_2, Z_1, N, Y_2$	$X_2, Z_1, W, L, M, Y_1$

# HETEROGENEOUS DATASETS

	Dataset 1	Dataset 2	...	Dataset n
<b>Target</b> $Q = P^*(y \mid \text{do}(x))$				

d <sub>1</sub>	Population	Los Angeles	New York	Texas
d <sub>2</sub>	Obs. / Exp.	Experimental	Observational	Experimental
d <sub>3</sub>	Treat. Assign.	Randomized Z <sub>1</sub>	-	Randomized Z <sub>2</sub>
d <sub>3</sub>	Sampling	Selection on Age	Selection on SES	-
d <sub>4</sub>	Measured	X <sub>1</sub> , Z <sub>1</sub> , W, M, Y <sub>1</sub>	X <sub>1</sub> , X <sub>2</sub> , Z <sub>1</sub> , N, Y <sub>2</sub>	X <sub>2</sub> , Z <sub>1</sub> , W, L, M, Y <sub>1</sub>

# HETEROGENEOUS DATASETS

	Dataset 1	Dataset 2	...	Dataset n
<b>Target</b> $Q = P^*(y \mid \text{do}(x))$				

$d_1$	Population	Los Angeles	New York	Texas
$d_2$	Obs. / Exp.	Experimental	Observational	Experimental
$d_3$	Treat. Assign.	Randomized $Z_1$	-	Randomized $Z_2$
$d_4$	Sampling	Selection on Age	Selection on $\subseteq$	-
	Measured	$X_1, Z_1, W, M, Y_1$	$X_1, X_2, Z_1, N, Y_2$	$X_2, Z_1, W, L, M,$ $Y_1$

# DATA-FUSION CHALLENGES

---

Dimensions

# DATA-FUSION CHALLENGES

---

Description of each dataset: tuple  $(d_1, d_2, d_3, d_4)$   
(population, obs./exp., sampling, measure.)

Dimensions

# DATA-FUSION CHALLENGES

---

Description of each dataset: tuple  $(d_1, d_2, d_3, d_4)$   
(population, obs./exp., sampling, measure.)

1. *Causal Inference* (observational studies)  
 $(d_1, \text{Observ.}, d_3, d_4) \rightarrow (d_1, \text{Experiment}(X), d_3, d_4)$

# DATA-FUSION CHALLENGES

---

Description of each dataset: tuple  $(d_1, d_2, d_3, d_4)$   
(population, obs./exp., sampling, measure.)

Dimensions

1. *Causal Inference* (observational studies)  
 $(d_1, \text{Observ.}, d_3, d_4) \rightarrow (d_1, \text{Experiment}(X), d_3, d_4)$
2. *Transportability* (External Validity)  
 $(\text{Bonobos}, d_2, d_3, d_4) \rightarrow (\text{Humans}, d_2, d_3, d_4)$

# DATA-FUSION CHALLENGES

Description of each dataset: tuple  $(d_1, d_2, d_3, d_4)$   
(population, obs./exp., sampling, measure.)

1. *Causal Inference* (observational studies)  
 $(d_1, \text{Observ.}, d_3, d_4) \rightarrow (d_1, \text{Experiment}(X), d_3, d_4)$
2. *Transportability* (External Validity)  
 $(\text{Bonobos}, d_2, d_3, d_4) \rightarrow (\text{Humans}, d_2, d_3, d_4)$
3. *Sampling Selection Bias*  
 $(d_1, d_2, \text{Select}(Salary), d_4) \rightarrow (d_1, d_2, \{\}, d_4)$

# DATA-FUSION CHALLENGES

Description of each dataset: tuple  $(d_1, d_2, d_3, d_4)$   
(population, obs./exp., sampling, measure.)

1. *Causal Inference* (observational studies)  
 $(d_1, \text{Observ.}, d_3, d_4) \rightarrow (d_1, \text{Experiment}(X), d_3, d_4)$
2. *Transportability* (External Validity)  
 $(\text{Bonobos}, d_2, d_3, d_4) \rightarrow (\text{Humans}, d_2, d_3, d_4)$
3. *Sampling Selection Bias*  
 $(d_1, d_2, \text{Select}(Salary), d_4) \rightarrow (d_1, d_2, \{\}, d_4)$

*Data-fusion:*

# DATA-FUSION CHALLENGES

Description of each dataset: tuple  $(d_1, d_2, d_3, d_4)$   
(population, obs./exp., sampling, measure.)

1. *Causal Inference* (observational studies)  
 $(d_1, \text{Observ.}, d_3, d_4) \rightarrow (d_1, \text{Experiment}(X), d_3, d_4)$
2. *Transportability* (External Validity)  
 $(\text{Bonobos}, d_2, d_3, d_4) \rightarrow (\text{Humans}, d_2, d_3, d_4)$
3. *Sampling Selection Bias*  
 $(d_1, d_2, \text{Select}(Salary), d_4) \rightarrow (d_1, d_2, \{\}, d_4)$

*Data-fusion:*

$$\{ (d_1, d_2, d_3, d_4) \} \rightarrow (d'_1, d'_2, d'_3, d'_4)$$

# DATA-FUSION IN 3-D

---

## 1. Experimental conditions

- Preliminary. Observational to Experimental regime

## 2. Environmental conditions

## 3. Sampling conditions

# DATA-FUSION IN 3-D

---


# THE DATA-FUSION PROBLEM

EXPERIMENTAL CONDITIONS ( $D = 1$ ) —  
GENERAL IDENTIFIABILITY

# THE CHALLENGE OF EXPERIMENTAL IDENTIFIABILITY

---

G:



(also called  $z$ -ID)

Z: Diet

X: Cholesterol level

Y: Heart Attack

Measured:


Observational study:  $P(x, y, z)$

Needed: Q =  $P(y | \text{do}(x)) = ?$

# WHAT CAN EXPERIMENTS ON DIET REVEAL ABOUT THE EFFECT OF CHOLESTEROL ON HEART ATTACK?

---

G:



Z: Diet


X: Cholesterol level

Y: Heart Attack

# WHAT CAN EXPERIMENTS ON DIET REVEAL ABOUT THE EFFECT OF CHOLESTEROL ON HEART ATTACK?

---

G:




- Z: Diet  
X: Cholesterol level  
Y: Heart Attack

Measured:

# WHAT CAN EXPERIMENTS ON DIET REVEAL ABOUT THE EFFECT OF CHOLESTEROL ON HEART ATTACK?

---

G:




- Z: Diet  
X: Cholesterol level  
Y: Heart Attack

Measured:

# WHAT CAN EXPERIMENTS ON DIET REVEAL ABOUT THE EFFECT OF CHOLESTEROL ON HEART ATTACK?

---

G:



Z: Diet  
X: Cholesterol level  
Y: Heart Attack


Measured:

Observational study:  $P(x, y, z)$

# WHAT CAN EXPERIMENTS ON DIET REVEAL ABOUT THE EFFECT OF CHOLESTEROL ON HEART ATTACK?

---

G:



Z: Diet  
X: Cholesterol level  
Y: Heart Attack

Measured:


Observational study:  $P(x, y, z)$

Experimental study:  $P(x, y | do(z))$

# WHAT CAN EXPERIMENTS ON DIET REVEAL ABOUT THE EFFECT OF CHOLESTEROL ON HEART ATTACK?

---

G:



Z: Diet  
X: Cholesterol level  
Y: Heart Attack

Measured:


Observational study:  $P(x, y, z)$

Experimental study:  $P(x, y | do(z))$

# WHAT CAN EXPERIMENTS ON DIET REVEAL ABOUT THE EFFECT OF CHOLESTEROL ON HEART ATTACK?

---

G:



Z: Diet  
X: Cholesterol level  
Y: Heart Attack

Measured:


Observational study:  $P(x, y, z)$

Experimental study:  $P(x, y | do(z))$

# WHAT CAN EXPERIMENTS ON DIET REVEAL ABOUT THE EFFECT OF CHOLESTEROL ON HEART ATTACK?

---

G:



Z: Diet  
X: Cholesterol level  
Y: Heart Attack

Measured:


Observational study:  $P(x, y, z)$

Experimental study:  $P(x, y | do(z))$

# WHAT CAN EXPERIMENTS ON DIET REVEAL ABOUT THE EFFECT OF CHOLESTEROL ON HEART ATTACK?

---

G:



Z: Diet  
X: Cholesterol level  
Y: Heart Attack

Measured:

Observational study:  $P(x, y, z)$


Experimental study:  $P(x, y | do(z))$

Needed:  $Q = P(y | do(x)) = ?$

# WHAT CAN EXPERIMENTS ON DIET REVEAL ABOUT THE EFFECT OF CHOLESTEROL ON HEART ATTACK?

---

G:



Z: Diet  
X: Cholesterol level  
Y: Heart Attack

Measured:

Observational study:  $P(x, y, z)$


Experimental study:  $P(x, y | do(z))$

Needed:  $Q = P(y | do(x)) = ? = \frac{P(x, y | do(z))}{P(x | do(z))}$

# WHAT CAN EXPERIMENTS ON DIET REVEAL ABOUT THE EFFECT OF CHOLESTEROL ON HEART ATTACK?

---

G:



Z: Diet

X: Cholesterol level

Y: Heart Attack

Measured:

Observational study:  $P(x, y, z)$


Experimental study:  $P(x, y | do(z))$

Needed:  $Q = P(y | do(x)) = ? = \frac{P(x, y | do(z))}{P(x | do(z))}$


(i.e.,  $\exists f, f: P(v), P(v | do(z)) \rightarrow P(y | do(x))$ )

# WHICH MODEL LICENSES THE $z$ -IDENTIFICATION OF THE CAUSAL EFFECT $X \rightarrow Y$ ?

(a)




(b)




(c)




(d)



(e)



(f)



# SUMMARY D=1: POLICY EVALUATION RESULTS

---

# SUMMARY D=1: POLICY EVALUATION RESULTS

---

- The estimability of any expression of the form

$$Q = P(y_1, y_2, \dots, y_n | do(x_1, x_2, \dots, x_m), z_1, z_2, \dots, z_k)$$

can be determined given any causal graph  $G$  containing measured and unmeasured variables.

# SUMMARY D=1: POLICY EVALUATION RESULTS

---

- The estimability of any expression of the form

$$Q = P(y_1, y_2, \dots, y_n | do(x_1, x_2, \dots, x_m), z_1, z_2, \dots, z_k)$$

can be determined given any causal graph  $G$  containing measured and unmeasured variables.

- If  $Q$  is estimable, then its estimand can be derived in polynomial time — by estimable we mean by a combination of observational and experimental studies.)
- The causal calculus is complete for this task.

# SUMMARY D=1: POLICY EVALUATION RESULTS

---

- The estimability of any expression of the form

$$Q = P(y_1, y_2, \dots, y_n | do(x_1, x_2, \dots, x_m), z_1, z_2, \dots, z_k)$$

can be determined given any causal graph  $G$  containing measured and unmeasured variables.

- If  $Q$  is estimable, then its estimand can be derived in polynomial time — by estimable we mean by a combination of observational and experimental studies.)
- The causal calculus is complete for this task.
- The algorithm is complete.

# SUMMARY D=1: POLICY EVALUATION RESULTS

---

- The estimability of any expression of the form

$$Q = P(y_1, y_2, \dots, y_n | do(x_1, x_2, \dots, x_m), z_1, z_2, \dots, z_k)$$

can be determined given any causal graph  $G$  containing measured and unmeasured variables.

- If  $Q$  is estimable, then its estimand can be derived in polynomial time — by estimable we mean by a combination of observational and experimental studies.)
- The causal calculus is complete for this task.
- The algorithm is complete.
- References

Pearl (1993, 1995) – Back-door criterion & *do*-calculus

Tian & Pearl (2002) – General ID algorithm

Shpitser & Pearl (2006) – Completeness (observational)

Bareinboim & Pearl (2012) – Completeness (obs. + experimental)

Lee, Correa & Bareinboim (2019) – Completeness (arbitrary exp.)

# THE DATA-FUSION PROBLEM

ENVIRONMENTAL CONDITIONS ( $D = 2$ ) —  
TRANSPORTABILITY, EXTERNAL VALIDITY, META-ANALYSIS

# THE TRANSPORTABILITY PROBLEM

---

Question:

Is it possible to predict the effect of  $X$  on  $Y$  in a target population  $\Pi^*$  (where no experiments are feasible), using experimental findings from a different population  $\Pi$ ?

Answer: Sometimes yes.

# THE TRANSPORTABILITY PROBLEM

---

Question:

Is it possible to predict the effect of  $X$  on  $Y$  in a target population  $\Pi^*$  (where no experiments are feasible), using experimental findings from a different population  $\Pi$ ?


Answer: Sometimes yes.

Our goal is to formally characterize when and how.

# MOVING FROM THE “LAB” TO THE “REAL WORLD”...

---


Lab



# MOVING FROM THE “LAB” TO THE “REAL WORLD”...

---

Lab




\* The lab stands for any environment, population, domain, setting.

# MOVING FROM THE “LAB” TO THE “REAL WORLD”...


---

Lab




# MOVING FROM THE “LAB” TO THE “REAL WORLD”...

---



# MOVING FROM THE “LAB” TO THE “REAL WORLD”...


---



Everything is assumed to be  
the same, trivially transportable!


# MOVING FROM THE “LAB” TO THE “REAL WORLD”...

---



Everything is assumed to be  
the same, trivially transportable!


# MOVING FROM THE “LAB” TO THE “REAL WORLD”...



Everything is assumed to be  
the same, trivially transportable!

# MOVING FROM THE “LAB” TO THE “REAL WORLD”...


---



Everything is assumed to be  
the same, trivially transportable!


# MOVING FROM THE “LAB” TO THE “REAL WORLD”...

---



# MOVING FROM THE “LAB” TO THE “REAL WORLD”...


---



# MOTIVATION

## WHAT CAN EXPERIMENTS IN LA TELL US ABOUT US?


---



$\Pi \rightarrow \Pi^*$   
LA      US

# MOTIVATION


## WHAT CAN EXPERIMENTS IN LA TELL US ABOUT US?



# MOTIVATION

## WHAT CAN EXPERIMENTS IN LA TELL US ABOUT US?


---



$\Pi \rightarrow \Pi^*$   
LA      US

# MOTIVATION


## WHAT CAN EXPERIMENTS IN LA TELL US ABOUT US?



$$\begin{matrix} \Pi & \rightarrow & \Pi^* \\ \text{LA} & & \text{US} \end{matrix}$$

# MOTIVATION

## WHAT CAN EXPERIMENTS IN LA TELL US ABOUT US?




$$\begin{matrix} \Pi & \rightarrow & \Pi^* \\ \text{LA} & & \text{US} \end{matrix}$$

Experimental study in LA

Measured:  $P(x, y, z)$   
Measured:  $P(y | do(x), z)$

# MOTIVATION

## WHAT CAN EXPERIMENTS IN LA TELL US ABOUT US?



$$\begin{matrix} \Pi & \rightarrow & \Pi^* \\ \text{LA} & & \text{US} \end{matrix}$$

Experimental study in LA


Measured:  $P(x, y, z)$   
Measured:  $P(y | do(x), z)$

Observational study in US

Measured:  $P^*(x, y, z)$   
[  $P^*(z) \neq P(z)$  ]

# MOTIVATION

## WHAT CAN EXPERIMENTS IN LA TELL US ABOUT US?



$$\begin{matrix} \Pi & \rightarrow & \Pi^* \\ \text{LA} & & \text{US} \end{matrix}$$

Experimental study in LA

Measured:  $P(x, y, z)$   
Measured:  $P(y | do(x), z)$


Observational study in US

Measured:  $P^*(x, y, z)$   
[  $P^*(z) \neq P(z)$  ]

Needed:  $Q = P^*(y | do(x)) = ?$

# MOTIVATION

## WHAT CAN EXPERIMENTS IN LA TELL US ABOUT US?



$$\begin{matrix} \Pi & \rightarrow & \Pi^* \\ \text{LA} & & \text{US} \end{matrix}$$

Experimental study in LA

Measured:  $P(x, y, z)$   
Measured:  $P(y | do(x), z)$


Observational study in US

Measured:  $P^*(x, y, z)$   
[  $P^*(z) \neq P(z)$  ]

Output  
Needed:  $Q = P^*(y | do(x)) = ?$

# MOTIVATION

## WHAT CAN EXPERIMENTS IN LA TELL US ABOUT US?



$$\begin{matrix} \Pi & \rightarrow & \Pi^* \\ \text{LA} & & \text{US} \end{matrix}$$

Experimental study in LA

Measured:  $P(x, y, z)$   
Measured:  $P(y | do(x), z)$


Observational study in US

Measured:  $P^*(x, y, z)$   
[  $P^*(z) \neq P(z)$  ]

Needed:  $Q = P^*(y | do(x)) = ?$

# MOTIVATION

## WHAT CAN EXPERIMENTS IN LA TELL US ABOUT US?



$$\begin{matrix} \Pi & \rightarrow & \Pi^* \\ \text{LA} & & \text{US} \end{matrix}$$

Experimental study in LA

Measured:  $P(x, y, z)$   
Measured:  $P(y | do(x), z)$


Observational study in US

Measured:  $P^*(x, y, z)$   
[  $P^*(z) \neq P(z)$  ]

Needed:  $Q = P^*(y | do(x)) = \sum_z P(y | do(x), z) P^*(z)$

# MOTIVATION

## WHAT CAN EXPERIMENTS IN LA TELL US ABOUT US?



$$\begin{matrix} \Pi & \rightarrow & \Pi^* \\ \text{LA} & & \text{US} \end{matrix}$$

Experimental study in LA

Measured:  $P(x, y, z)$   
Measured:  $P(y | do(x), z)$


Observational study in US

Measured:  $P^*(x, y, z)$   
[  $P^*(z) \neq P(z)$  ]

Needed:  $Q = P^*(y | do(x)) = \sum_z P(y | do(x), z) P^*(z)$

# MOTIVATION

## WHAT CAN EXPERIMENTS IN LA TELL US ABOUT US?



$$\begin{matrix} \Pi & \rightarrow & \Pi^* \\ \text{LA} & & \text{US} \end{matrix}$$

Experimental study in LA

$$\begin{aligned} & P(x, y, z) \\ \text{Measured: } & P(y | do(x), z) \end{aligned}$$

Observational study in US

$$\begin{aligned} & \text{Measured: } P^*(x, y, z) \\ & [P^*(z) \neq P(z)] \end{aligned}$$

$$\text{Needed: } Q = P^*(y | do(x)) = \sum_z P(y | do(x), z) P^*(z)$$


Transport Formula (recalibration):  $Q = F(P, P_{do}, P^*)$

# TRANSPORT FORMULAS DEPEND ON THE CAUSAL STORY

---


# TRANSPORT FORMULAS DEPEND ON THE CAUSAL STORY

---




# TRANSPORT FORMULAS DEPEND ON THE CAUSAL STORY

$z \leftarrow f_z(u_z, u_{xz})$   
 $x \leftarrow f_x(x, u_x, u_{xz}, u_{xy})$   
 $y \leftarrow f_y(x, z, u_y, u_{xy})$




# TRANSPORT FORMULAS DEPEND ON THE CAUSAL STORY

---



# TRANSPORT FORMULAS DEPEND ON THE CAUSAL STORY

---



LA:  $P(y \mid \text{do}(x), z)$


Z	X	Y

US:  $P^*(x, y, z)$

Z	X	Y

# TRANSPORT FORMULAS DEPEND ON THE CAUSAL STORY

---



LA:  $P(y \mid do(x), z)$

a)  $Z$  represents age


$$P^*(y \mid do(x)) = \sum_z P(y \mid do(x), z) P^*(z)$$

Z	X	Y

US:  $P^*(x, y, z)$

Z	X	Y

# TRANSPORT FORMULAS DEPEND ON THE CAUSAL STORY



LA:  $P(y \mid do(x), z)$

a)  $Z$  represents age

$$P^*(y \mid do(x)) = \sum_z P(y \mid do(x), z) P^*(z)$$


Z	X	Y

US:  $P^*(x, y, z)$

Z	X	Y

# TRANSPORT FORMULAS DEPEND ON THE CAUSAL STORY

---



LA:  $P(y | \text{do}(x), z)$

a)  $Z$  represents age

$$P^*(y | \text{do}(x)) = \sum_z P(y | \text{do}(x), z) P^*(z)$$

Z	X	Y

b)  $Z$  represents language skill


$$P^*(y | \text{do}(x)) = ?$$

US:  $P^*(x, y, z)$

Z	X	Y

# TRANSPORT FORMULAS DEPEND ON THE CAUSAL STORY

---



LA:  $P(y | \text{do}(x), z)$

a)  **$Z$  represents age**

$$P^*(y | \text{do}(x)) = \sum_z P(y | \text{do}(x), z) P^*(z)$$

Z	X	Y

b)  **$Z$  represents language skill**


$$P^*(y | \text{do}(x)) = P(y | \text{do}(x))$$

US:  $P^*(x, y, z)$

Z	X	Y

# TRANSPORT FORMULAS DEPEND ON THE CAUSAL STORY

---



LA:  $P(y | \text{do}(x), z)$

a) **Z represents age**

$$P^*(y | \text{do}(x)) = \sum_z P(y | \text{do}(x), z) P^*(z)$$

Z	X	Y

b) **Z represents language skill**

$$P^*(y | \text{do}(x)) = P(y | \text{do}(x))$$

US:  $P^*(x, y, z)$


c) **Z represents a bio-marker**

$$P^*(y | \text{do}(x)) = ?$$

Z	X	Y

# TRANSPORT FORMULAS DEPEND ON THE CAUSAL STORY

---



LA:  $P(y \mid do(x), z)$

a) **Z represents age**

$$P^*(y \mid do(x)) = \sum_z P(y \mid do(x), z) P^*(z)$$

Z	X	Y

b) **Z represents language skill**

$$P^*(y \mid do(x)) = P(y \mid do(x))$$

US:  $P^*(x, y, z)$


c) **Z represents a bio-marker**

$$P^*(y \mid do(x)) = \sum_z P(y \mid do(x), z) P^*(z \mid x)$$

Z	X	Y


# SELECTION DIAGRAMS

- How to encode disparities and commonalities about domains?



# SELECTION DIAGRAMS

- How to encode disparities and commonalities about domains?



# RESULT 1: TRANSPORTABILITY REDUCED TO CALCULUS

---

Theorem:

A causal relation  $Q$  is transportable from  $\Pi$  to  $\Pi^*$  iff  $Q(\Pi^*)$  is reducible, using the rules of causal calculus, to an expression in which  $S$  is separated from  $do()$ .

# RESULT 1: TRANSPORTABILITY REDUCED TO CALCULUS


---

Theorem:

A causal relation  $Q$  is transportable from  $\Pi$  to  $\Pi^*$  iff  $Q(\Pi^*)$  is reducible, using the rules of causal calculus, to an expression in which  $S$  is separated from  $do()$ .

Ex:

$$\begin{aligned} Q &= P^*(y \mid do(x)) = P(y \mid do(x), s) \\ &= \sum_w P(y \mid do(x), s, w) P(w \mid do(x), s) \\ &= \sum_w P(y \mid do(x), w) P(w \mid do(x), s) \\ &= \sum_w P(y \mid do(x), w) P(w \mid s) \\ &= \sum_w P(y \mid do(x), w) P^*(w) \end{aligned}$$




# RESULT 1: TRANSPORTABILITY REDUCED TO CALCULUS

Theorem:

A causal relation  $Q$  is transportable from  $\Pi$  to  $\Pi^*$  iff  $Q(\Pi^*)$  is reducible, using the rules of causal calculus, to an expression in which  $S$  is separated from  $do()$ .

Ex: query

$$\begin{aligned} Q &= P^*(y \mid do(x)) = P(y \mid do(x), s) \\ &= \sum_w P(y \mid do(x), s, w) P(w \mid do(x), s) \\ &= \sum_w P(y \mid do(x), w) P(w \mid do(x), s) \\ &= \sum_w P(y \mid do(x), w) P(w \mid s) \\ &= \sum_w P(y \mid do(x), w) P^*(w) \end{aligned}$$



# RESULT 1: TRANSPORTABILITY REDUCED TO CALCULUS


Theorem:

A causal relation  $Q$  is transportable from  $\Pi$  to  $\Pi^*$  iff  $Q(\Pi^*)$  is reducible, using the rules of causal calculus, to an expression in which  $S$  is separated from  $do()$ .

Ex: query

$$\begin{aligned} Q &= P^*(y \mid do(x)) = P(y \mid do(x), s) \\ &= \sum_w P(y \mid do(x), s, w) P(w \mid do(x), s) \\ &= \sum_w P(y \mid do(x), w) P(w \mid do(x), s) \\ &= \sum_w P(y \mid do(x), w) P(w \mid s) \\ &= \sum_w P(y \mid do(x), w) P^*(w) \end{aligned}$$

data



# RESULT 1: TRANSPORTABILITY REDUCED TO CALCULUS


Theorem:

A causal relation  $Q$  is transportable from  $\Pi$  to  $\Pi^*$  iff  $Q(\Pi^*)$  is reducible, using the rules of causal calculus, to an expression in which  $S$  is separated from  $\text{do}()$ .

Ex: query

$$\begin{aligned} Q &= P^*(y \mid \text{do}(x)) = P(y \mid \text{do}(x), s) \\ &= \sum_w P(y \mid \text{do}(x), s, w) P(w \mid \text{do}(x), s) \\ &= \sum_w P(y \mid \text{do}(x), w) P(w \mid \text{do}(x), s) \\ &= \sum_w P(y \mid \text{do}(x), w) P(w \mid s) \\ &= \sum_w P(y \mid \text{do}(x), w) P^*(w) \end{aligned}$$


data



\* Deterministic sep.: (Dawid, 1979, 2002; Constantinou & Dawid, 2016)

# RESULT 2: ALGORITHM TO DETERMINE IF AN EFFECT IS TRANSPORTABLE


---



# RESULT 2: ALGORITHM TO DETERMINE IF AN EFFECT IS TRANSPORTABLE


---

INPUT: Annotated Causal Graph



# RESULT 2: ALGORITHM TO DETERMINE IF AN EFFECT IS TRANSPORTABLE


---



INPUT: Annotated Causal Graph

$S \rightarrow$  Factors creating differences

# RESULT 2: ALGORITHM TO DETERMINE IF AN EFFECT IS TRANSPORTABLE




INPUT: Annotated Causal Graph

$S \rightarrow$  Factors creating differences

$$P^*(y | do(x)) = \sum_z P(y | do(x), z) \sum_w P^*(z | w) \sum_t P(w | do(x), t) P^*(t)$$

# RESULT 2: ALGORITHM TO DETERMINE IF AN EFFECT IS TRANSPORTABLE



INPUT: Annotated Causal Graph







$S \rightarrow$  Factors creating differences

OUTPUT:

1. Transportable or not?
2. Yes = Transport formula
  1. Measurements to be taken in the experimental domain
  2. Measurements to be taken in the target domain
3. No = Not possible

$$P^*(y | do(x)) = \sum_z P(y | do(x), z) \sum_w P^*(z | w) \sum_t P(w | do(x), t) P^*(t)$$

# RESULT 3: GRAPHICAL CONDITION FOR TRANSPORTABILITY




# THE CHALLENGE OF GENERAL TRANSPORTABILITY

Assume we want to transport  $Q = P^*(y | \text{do}(x))$  from  $\{\prod^a, \prod^b\}$  with experimental information  $\{X, Z\}$  in  $\prod^a$ ,  $\{X, Z\}$  in  $\prod^b$ .

LA:

$$\begin{aligned}P(y, z | \text{do}(x)) \\P(y, x | \text{do}(z))\end{aligned}$$

Z	X	Y




$\prod^a (= \text{LA})$

:NY

$$\begin{aligned}P(y, z | \text{do}(x)) \\P(y, x | \text{do}(z))\end{aligned}$$

Z	X	Y



$\prod^b (= \text{NYC})$

# THE CHALLENGE OF GENERAL TRANSPORTABILITY


Assume we want to transport  $Q = P^*(y \mid \text{do}(x))$  from  $\{\prod^a, \prod^b\}$  with experimental information  $\{X, Z\}$  in  $\prod^a$ ,  $\{X, Z\}$  in  $\prod^b$ .

LA:


$$P(y, z \mid \text{do}(x))$$

$$P(y, x \mid \text{do}(z))$$

Z	X	Y



$\prod^a (=LA)$



$\prod^b (=NYC)$


:NY  
 $P(y, z \mid \text{do}(x))$   
 $P(y, x \mid \text{do}(z))$

Z	X	Y


If we apply the result from limited transportability (1D),  
 $Q$  is (provably) **not** transportable from  $\prod^a$  and  $\prod^b$  separately.

# DERIVATION OF GENERAL TRANSPORTABILITY

---



$$\prod^{\mathbf{a}} (=LA)$$




$$\prod^{\mathbf{b}} (=NYC)$$

$$\begin{aligned}
 Q &= P^*(y \mid \text{do}(x)) = P(y \mid \text{do}(x), s^a, s^b) && \text{Meta-TR Def.} \\
 &= \sum_z P(y \mid \text{do}(x), z, s^a, s^b) P(z \mid \text{do}(x), s^a, s^b) && \text{Probability Axioms} \\
 &= \sum_z P(y \mid \text{do}(x), \text{do}(z), s^a, s^b) P(z \mid \text{do}(x), s^a, s^b) && \text{Rule 2} \\
 &= \sum_z P(y \mid \text{do}(z), s^a, s^b) P(z \mid \text{do}(x), s^a, s^b) && \text{Rule 3} \\
 &= \sum_z P(y \mid \text{do}(z), s^a) P(z \mid \text{do}(x), s^a, s^b) && \text{Rule 1* in } D_b \\
 &= \sum_z P(y \mid \text{do}(z), s^a) P(z \mid \text{do}(x), s^b) && \text{Rule 1* in } D_a \\
 &= \sum_z P^{(b)}(y \mid \text{do}(z)) P^{(a)}(z \mid \text{do}(x))
 \end{aligned}$$


# DERIVATION OF GENERAL TRANSPORTABILITY

---



query

$\prod^a (=LA)$



$\prod^b (=NYC)$

$$Q = P^*(y \mid \text{do}(x)) = P(y \mid \text{do}(x), s^a, s^b)$$

Meta-TR Def.

$$= \sum_z P(y \mid \text{do}(x), z, s^a, s^b) P(z \mid \text{do}(x), s^a, s^b)$$

Probability Axioms

$$= \sum_z P(y \mid \text{do}(x), \text{do}(z), s^a, s^b) P(z \mid \text{do}(x), s^a, s^b)$$

Rule 2

$$= \sum_z P(y \mid \text{do}(z), s^a, s^b) P(z \mid \text{do}(x), s^a, s^b)$$

Rule 3

$$= \sum_z P(y \mid \text{do}(z), s^a) P(z \mid \text{do}(x), s^a, s^b)$$

Rule 1\* in  $D_b$


$$= \sum_z P(y \mid \text{do}(z), s^a) P(z \mid \text{do}(x), s^b)$$

Rule 1\* in  $D_a$

$$= \sum_z P^{(b)}(y \mid \text{do}(z)) P^{(a)}(z \mid \text{do}(x))$$


Meta-TR Def.

# DERIVATION OF GENERAL TRANSPORTABILITY



query

$\prod^a (=LA)$



$\prod^b (=NYC)$

$$Q = P^*(y \mid do(x)) = P(y \mid do(x), s^a, s^b)$$

Meta-TR Def.

$$= \sum_z P(y \mid do(x), z, s^a, s^b) P(z \mid do(x), s^a, s^b)$$

Probability Axioms

$$= \sum_z P(y \mid do(x), do(z), s^a, s^b) P(z \mid do(x), s^a,$$

Rule 2

$$= \sum_z P(y \mid do(z), s^a, s^b) P(z \mid do(x), s^a, s^b)$$

Rule 3

$$= \sum_z P(y \mid do(z), s^a) P(z \mid do(x), s^a, s^b)$$

Rule 1\* in  $D_b$

$$= \sum_z P(y \mid do(z), s^a) P(z \mid do(x), s^b)$$

Rule 1\* in  $D_a$

$$= \sum_z P^{(b)}(y \mid do(z)) P^{(a)}(z \mid do(x))$$

Meta-TR Def.

data

NYC

LA


## A MORE CHALLENGING INSTANCE ...

---

Assume we want to transport  $Q = P^*(y | \text{do}(x))$  from  $\{\prod^a, \prod^b\}$  by intervening over  $\{Z_1\}$  in  $\prod^*$ ,  $\{Z_2\}$  in  $\prod^a$ ,  $\{Z_1\}$  in  $\prod^b$ .


## A MORE CHALLENGING INSTANCE ...

Assume we want to transport  $Q = P^*(y \mid do(x))$  from  $\{\Pi^a, \Pi^b\}$  by intervening over  $\{Z_1\}$  in  $\Pi^*$ ,  $\{Z_2\}$  in  $\Pi^a$ ,  $\{Z_1\}$  in  $\Pi^b$ .




# A MORE CHALLENGING INSTANCE ...

Assume we want to transport  $Q = P^*(y | \text{do}(x))$  from  $\{\prod^a, \prod^b\}$  by intervening over  $\{Z_1\}$  in  $\prod^*$ ,  $\{Z_2\}$  in  $\prod^a$ ,  $\{Z_1\}$  in  $\prod^b$ .



$\prod^a (=LA)$




$\prod^b (=NYC)$

$$\begin{aligned}
 P_x^*(y) &= \sum_{z_1, z_3, w, u} P^*(z_1) (w_1^{(1)} \sum_{z_2'} P_{z_1}^*(z_3|x, Z_2') P_{z_1}^*(Z_2') + w_2^{(1)} \sum_{z_2'} P_{z_1}^{(b)}(z_3|x, Z_2') P_{z_1}^{(b)}(Z_2')) \\
 &\quad (w_1^{(2)} (\sum_{z_2'} P_{z_1}^*(u|w, z_3, x, Z_2') P_{z_1}^*(z_3|x, Z_2') P_{z_1}^*(Z_2')) / (\sum_{z_2''} P_{z_1}^*(z_3|x, Z_2'') P_{z_1}^*(Z_2''))) \\
 &\quad + w_2^{(2)} P_{z_2}^{(a)}(u|w, z_3, x, Z_2') ) P^*(w|x, z_1, z_2, z_3) P^*(y|x, z_1, z_2, z_3, w, u)
 \end{aligned}$$

# IS THE GOLD STANDARD SO GOLD? (GENERALIZABILITY FROM EXPERIMENTS)

---


Before randomization




# IS THE GOLD STANDARD SO GOLD? (GENERALIZABILITY FROM EXPERIMENTS)

---

Before randomization




After randomization




# IS THE GOLD STANDARD SO GOLD? (GENERALIZABILITY FROM EXPERIMENTS)

Before randomization



After randomization



The gold standard is not absolute, and strong assumptions are required to validate claims from trials.

# SUMMARY D=2: POLICY GENERALIZATION

---

# SUMMARY D=2: POLICY GENERALIZATION

---

- Nonparametric transportability can be determined provided that the problem instance is encoded in selection diagrams.

# SUMMARY D=2: POLICY GENERALIZATION

---

- Nonparametric transportability can be determined provided that the problem instance is encoded in selection diagrams.
- When transportability is feasible, the transport formula can be derived in polynomial time.

# SUMMARY D=2: POLICY GENERALIZATION

---

- Nonparametric transportability can be determined provided that the problem instance is encoded in selection diagrams.
- When transportability is feasible, the transport formula can be derived in polynomial time.
- The causal calculus and the algorithm are complete.

# SUMMARY D=2: POLICY GENERALIZATION

---

- Nonparametric transportability can be determined provided that the problem instance is encoded in selection diagrams.
- When transportability is feasible, the transport formula can be derived in polynomial time.
- The causal calculus and the algorithm are complete.
- These results solve long-standing problems in the empirical sciences including external validity, meta-analysis, and heterogeneity.

# SUMMARY D=2: POLICY GENERALIZATION

---

- Nonparametric transportability can be determined provided that the problem instance is encoded in selection diagrams.
- When transportability is feasible, the transport formula can be derived in polynomial time.
- The causal calculus and the algorithm are complete.
- These results solve long-standing problems in the empirical sciences including external validity, meta-analysis, and heterogeneity.
- References:

# SUMMARY D=2: POLICY GENERALIZATION

---

- Nonparametric transportability can be determined provided that the problem instance is encoded in selection diagrams.
- When transportability is feasible, the transport formula can be derived in polynomial time.
- The causal calculus and the algorithm are complete.
- These results solve long-standing problems in the empirical sciences including external validity, meta-analysis, and heterogeneity.
- References:

Pearl & Bareinboim (2011) – semantics for transportability

Bareinboim & Pearl (2012) – completeness of limited TR

Bareinboim et al (2013) – multiple & limited + extensions

Bareinboim & Pearl (2014) – completeness of general TR

# THE DATA-FUSION PROBLEM

SAMPLING CONDITIONS ( $D = 3$ )

# THE SELECTION BIAS PROBLEM

---

- Selection bias, caused by preferential exclusion of samples from the data, is a major obstacle to valid causal and statistical inferences;


# THE SELECTION BIAS PROBLEM

---

- Selection bias, caused by **preferential exclusion** of samples from the data, is a major obstacle to valid causal and **statistical** inferences;


# THE SELECTION BIAS PROBLEM

- Selection bias, caused by **preferential exclusion** of samples from the data, is a major obstacle to valid **causal** and **statistical** inferences;




# THE SELECTION BIAS PROBLEM

- Selection bias, caused by **preferential exclusion** of samples from the data, is a major obstacle to valid **causal** and **statistical** inferences;



# THE SELECTION BIAS PROBLEM


- Selection bias, caused by **preferential exclusion** of samples from the data, is a major obstacle to valid **causal** and **statistical** inferences;



- Under what conditions can we estimate the query (e.g.,  $P(y | x)$ ) from  $P(v | S = 1)$ .

# THE ORIGIN OF SELECTION BIAS

G:



Under selection ( $S=1$ ):

$P(X=1, Y=1 | S=1) = 1/2$   
 $P(X=0, Y=0 | S=1) = 1/2$   
(correlation=1)

$$P(Y=y|X=1, S=1) - P(Y=y|X=0, S=1) = 1$$


M:

$$\begin{aligned} X &= U_x \\ Y &= U_y \\ S &= \neg(X \oplus Y) \end{aligned}$$

$$P(U_x, U_y) = 1/4$$

# THE ORIGIN OF SELECTION BIAS

G:



Under selection ( $S=1$ ):

$$\begin{aligned} P(X=1, Y=1 | S=1) &= 1/2 \\ P(X=0, Y=0 | S=1) &= 1/2 \\ (\text{correlation}=1) \end{aligned}$$

$$P(Y=y | X=1, S=1) - P(Y=y | X=0, S=1) = 1$$

but

$$P(Y = y, X = x) = 1/4$$

$$P(Y=y | X=1) - P(Y=y | X=0) = 0$$

M:


$$\begin{aligned} X &= U_x \\ Y &= U_y \\ S &= \neg(X \oplus Y) \end{aligned}$$

$$P(U_x, U_y) = 1/4$$


# CONFOUNDING BIAS vs SELECTION BIAS

---

- Unblockable “flow” of information between treatment and outcome — spurious correlation.




$G_1$



$G_2$


# SELECTION VERSUS TRANSPORTABILITY (MAN-MADE VERSUS NATURE-MADE)

Selection Bias



Z = sample-producing factors  
S = selection indicator

Transportability



Z = disparities across populations  
S = difference-producing factors

# LITERATURE ON SELECTION

---

# LITERATURE ON SELECTION

---

1. Qualitative assumptions about the data-generating model (treatment-dependent, outcome-dependent).

# LITERATURE ON SELECTION

---

1. Qualitative assumptions about the data-generating model (treatment-dependent, outcome-dependent).

# LITERATURE ON SELECTION

---

1. Qualitative assumptions about the data-generating model (treatment-dependent, outcome-dependent).

[Cornfield'51, Whittemore'78, Geng'92, Didelez'10]

# LITERATURE ON SELECTION

---

1. Qualitative assumptions about the data-generating model (treatment-dependent, outcome-dependent).

[Cornfield'51, Whittemore'78, Geng'92, Didelez'10]

# LITERATURE ON SELECTION

---

1. Qualitative assumptions about the data-generating model (treatment-dependent, outcome-dependent).

[Cornfield'51, Whittemore'78, Geng'92, Didelez'10]

2. Parametric assumptions about the data-generating model (e.g., linear, monotonic).

# LITERATURE ON SELECTION

---

1. Qualitative assumptions about the data-generating model (treatment-dependent, outcome-dependent).

[Cornfield'51, Whittemore'78, Geng'92, Didelez'10]

2. Parametric assumptions about the data-generating model (e.g., linear, monotonic).

# LITERATURE ON SELECTION

---

1. Qualitative assumptions about the data-generating model (treatment-dependent, outcome-dependent).

[Cornfield'51, Whittemore'78, Geng'92, Didelez'10]

2. Parametric assumptions about the data-generating model (e.g., linear, monotonic).

[Heckman'79, Angrist'91, Pearl'13]

# LITERATURE ON SELECTION

---

1. Qualitative assumptions about the data-generating model (treatment-dependent, outcome-dependent).

[Cornfield'51, Whittemore'78, Geng'92, Didelez'10]

2. Parametric assumptions about the data-generating model (e.g., linear, monotonic).

[Heckman'79, Angrist'91, Pearl'13]

# LITERATURE ON SELECTION

---

1. Qualitative assumptions about the data-generating model (treatment-dependent, outcome-dependent).  
[Cornfield'51, Whittemore'78, Geng'92, Didelez'10]
2. Parametric assumptions about the data-generating model (e.g., linear, monotonic).  
[Heckman'79, Angrist'91, Pearl'13]
3. Quantitive assumptions about the data-generating model (i.e.  $P(S = 1 | Z)$ ).

# LITERATURE ON SELECTION

---

1. Qualitative assumptions about the data-generating model (treatment-dependent, outcome-dependent).  
[Cornfield'51, Whittemore'78, Geng'92, Didelez'10]
2. Parametric assumptions about the data-generating model (e.g., linear, monotonic).  
[Heckman'79, Angrist'91, Pearl'13]
3. Quantitive assumptions about the data-generating model (i.e.  $P(S = 1 | Z)$ ).

# LITERATURE ON SELECTION

---

1. Qualitative assumptions about the data-generating model (treatment-dependent, outcome-dependent).

[Cornfield'51, Whittemore'78, Geng'92, Didelez'10]

2. Parametric assumptions about the data-generating model (e.g., linear, monotonic).

[Heckman'79, Angrist'91, Pearl'13]

3. Quantitive assumptions about the data-generating model (i.e.  $P(S = 1 | Z)$ ).

[Elkan'01, Zadrozny'04, Cortes'08, Storkey'09]

# SELECTION BIAS -- VARIANTS

---

- **Selection bias without external data (Odds Ratio):**  
Input:  $P(v | S = 1)$ ;  
Output:  $OR(y,x)$  recoverable? if so, how?

# SELECTION BIAS -- VARIANTS

---

- **Selection bias without external data (Odds Ratio):**  
Input:  $P(v | S = 1)$ ;  
Output:  $OR(y, x)$  recoverable? if so, how?
- **Selection bias without external data:**  
Input:  $P(v | S = 1)$ ;  
Output:  $P(y | x)$  recoverable? if so, how?

# SELECTION BIAS -- VARIANTS

---

- **Selection bias without external data (Odds Ratio):**  
Input:  $P(v | S = 1)$ ;  
Output:  $OR(y, x)$  recoverable? if so, how?
- **Selection bias without external data:**  
Input:  $P(v | S = 1)$ ;  
Output:  $P(y | x)$  recoverable? if so, how?
- **Selection bias with external data:**  
Input:  $P(v | S = 1) + P(t), T \subseteq V$ ;  
Output:  $P(y | x)$  recoverable? if so, how?

# SELECTION BIAS -- VARIANTS

---

- **Selection bias without external data (Odds Ratio):**  
Input:  $P(v | S = 1)$ ;  
Output:  $OR(y, x)$  recoverable? if so, how?
- **Selection bias without external data:**  
Input:  $P(v | S = 1)$ ;  
Output:  $P(y | x)$  recoverable? if so, how?
- **Selection bias with external data:**  
Input:  $P(v | S = 1) + P(t), T \subseteq V$ ;  
Output:  $P(y | x)$  recoverable? if so, how?
- **Selection bias in causal inference:**  
Input:  $P(v | S = 1) + P(t), T \subseteq V$ ;  
Output:  $P(y | do(x))$  recoverable? if so, how?

# SELECTION BIAS -- VARIANTS

---


- **Selection bias without external data (Odds Ratio):**  
Input:  $P(v | S = 1)$ ;  
Output:  $OR(y, x)$  recoverable? if so, how?
- **Selection bias without external data:**  
Input:  $P(v | S = 1)$ ;  
Output:  $P(y | x)$  recoverable? if so, how?
- **Selection bias with external data:**  
Input:  $P(v | S = 1) + P(t), T \subseteq V$ ;  
Output:  $P(y | x)$  recoverable? if so, how?
- **Selection bias in causal inference:**  
Input:  $P(v | S = 1) + P(t), T \subseteq V$ ;  
Output:  $P(y | do(x))$  recoverable? if so, how?
- **Selection bias with instrumental variables:**  
Input:  $P(v | S = 1) + P(t), T \subseteq V$ ;  
Output: Bounds for  $P(y | do(x))$  recoverable? if so, how?

# SELECTION BIAS WITHOUT EXTERNAL INFORMATION


---

Question. Under what conditions can we estimate the distribution  $P(y | x)$  from  $P(v | S=1)$ .

Theorem.  $Q = P(y | x)$  is recoverable from selection biased data if and only if  $(S \perp\!\!\!\perp Y | X)$ .



$P(y | x)$  is **not recoverable**.




$P(y | x)$  is **recoverable**.

# SELECTION WITH EXTERNAL INFORMATION

---


Assume we want to estimate  $Q = P(y | x)$  from selection...



# SELECTION WITH EXTERNAL INFORMATION

---

Assume we want to estimate  $Q = P(y | x)$  from selection...




$Q$  is **not recoverable** by the previous theorem...

# SELECTION WITH EXTERNAL INFORMATION

---

Assume we want to estimate  $Q = P(y | x)$  from selection...




$Q$  is **not recoverable** by the previous theorem...  
but what if  $P(W_1, W_2)$  is available?

# RECOVERABILITY WITH EXTERNAL INFORMATION

Theorem.  $P(y | x)$  is recoverable if there is a set  $C$  such that  $(Y \perp\!\!\!\perp S | C, X)$  holds in  $G$  and  $P(C, X)$  is estimable.

Moreover,  $P(y | x) = \sum_c P(y | x, c, S = 1) P(c | x)$




$C = \{W_1, W_2\}?$   
 $= \{W_2, Z_1, Z_2\}?$   
 $= \{W_2, Z_3\}?$

# RECOVERABILITY WITH EXTERNAL INFORMATION

Theorem.  $P(y | x)$  is recoverable if there is a set  $C$  such that  $(Y \perp\!\!\!\perp S | C, X)$  holds in  $G$  and  $P(C, X)$  is estimable.

Moreover,  $P(y | x) = \sum_c P(y | x, c, S = 1) P(c | x)$




$C = \{W_1, W_2\}$ ? yes  
 $= \{W_2, Z_1, Z_2\}$ ?  
 $= \{W_2, Z_3\}$ ?

# RECOVERABILITY WITH EXTERNAL INFORMATION

Theorem.  $P(y | x)$  is recoverable if there is a set  $C$  such that  $(Y \perp\!\!\!\perp S | C, X)$  holds in  $G$  and  $P(C, X)$  is estimable.

Moreover,  $P(y | x) = \sum_c P(y | x, c, S = 1) P(c | x)$




$C = \{W_1, W_2\}$ ? yes  
 $= \{W_2, Z_1, Z_2\}$ ? no  
 $= \{W_2, Z_3\}\text{?}$

# RECOVERABILITY WITH EXTERNAL INFORMATION

Theorem.  $P(y | x)$  is recoverable if there is a set  $C$  such that  $(Y \perp\!\!\!\perp S | C, X)$  holds in  $G$  and  $P(C, X)$  is estimable.




Moreover,  $P(y | x) = \sum_c P(y | x, c, S = 1) P(c | x)$



$C = \{W_1, W_2\}$ ? yes  
 $= \{W_2, Z_1, Z_2\}$ ? no  
 $= \{W_2, Z_3\}$ ? yes

# SELECTION BIAS IN CAUSAL INFERENCE

We want to recover  $Q = P(y | \text{do}(x))$ .



$C = \{W_2\}$  **recoverable**  
 $= \{W_1\}$  **not recoverable**

$C = \{\}$  **not recoverable**  
 $= \{W\}$  **not recoverable**

$= \{W, X\}$  **recoverable**

**recoverable**  
 $(S \perp\!\!\!\perp Y | X, W_2)$   
does not hold

# SELECTION BIAS IN CAUSAL INFERENCE

---

Def. Let a set  $Z$  be partitioned into  $Z^+$  and  $Z^-$  such that  $Z^+$  contains all non-descendants of  $X$  and  $Z^-$  the descendants of  $X$ .  $Z$  is said to satisfy the s-backdoor criterion relative to an ordered pair of variables  $(X, Y)$  and an ordered pair  $(M, T)$  in a graph  $G_s$  if  $Z^+$  and  $Z^-$  satisfy the following conditions:

- (i)  $Z^+$  blocks all backdoor paths from  $X$  to  $Y$ ;
- (ii)  $X$  and  $Z^+$  block all paths between  $Z^-$  and  $Y$ ;
- (iii)  $X$  and  $Z$  block all paths between  $S$  and  $Y$ ;
- (iv)  $Z \cup \{X, Y\} \subseteq M, Z \subseteq T$ .

Thm. If a set  $Z$  satisfies the s-backdoor relative to the pairs  $(X, Y)$  and  $(M, T)$ , then the effect of  $X$  on  $Y$  is identifiable and recoverable and is given by the formula

$$P(y | \text{do}(x)) = \sum_z P(y | x, z, S = 1) P(z)$$

# SUMMARY (D=3): SELECTION BIAS

---

# SUMMARY (D=3): SELECTION BIAS

---

- Nonparametric recoverability of selection from causal and statistical settings can be determined provided that an augmented causal graph is available.

# SUMMARY (D=3): SELECTION BIAS

---

- Nonparametric recoverability of selection from causal and statistical settings can be determined provided that an augmented causal graph is available.
- When recoverability is feasible, the estimand can be derived in polynomial time.

# SUMMARY (D=3): SELECTION BIAS

---

- Nonparametric recoverability of selection from causal and statistical settings can be determined provided that an augmented causal graph is available.
- When recoverability is feasible, the estimand can be derived in polynomial time.
- The result is complete for pure recoverability and sufficient for recoverability with external information.

# SUMMARY (D=3): SELECTION BIAS

---

- Nonparametric recoverability of selection from causal and statistical settings can be determined provided that an augmented causal graph is available.
- When recoverability is feasible, the estimand can be derived in polynomial time.
- The result is complete for pure recoverability and sufficient for recoverability with external information.
- The back-door criterion can be generalized to handle selection bias.

# SUMMARY (D=3): SELECTION BIAS

---

- Nonparametric recoverability of selection from causal and statistical settings can be determined provided that an augmented causal graph is available.
- When recoverability is feasible, the estimand can be derived in polynomial time.
- The result is complete for pure recoverability and sufficient for recoverability with external information.
- The back-door criterion can be generalized to handle selection bias.
- Stronger results can be obtained for the OR recoverability.

# SUMMARY (D=3): SELECTION BIAS

---

- Nonparametric recoverability of selection from causal and statistical settings can be determined provided that an augmented causal graph is available.
- When recoverability is feasible, the estimand can be derived in polynomial time.
- The result is complete for pure recoverability and sufficient for recoverability with external information.
- The back-door criterion can be generalized to handle selection bias.
- Stronger results can be obtained for the OR recoverability.
- References:

Bareinboim, Pearl (2012) – OR completeness & IVs

Bareinboim, Tian, Pearl (2014) – Markovian completeness

Bareinboim, Tian (2015) – Causal queries

Correa, Bareinboim (2017) – Complete adjustment

# CONCLUSIONS

---

# CONCLUSIONS

---

- Data-fusion requires encoding of structural features of the data-generating model.

# CONCLUSIONS

---

- Data-fusion requires encoding of structural features of the data-generating model.
- Both population-level and individual-level causal inference and fusion can be performed once the proper formal machinery is employed; algorithms and conditions are available.

# CONCLUSIONS

---

- Data-fusion requires encoding of structural features of the data-generating model.
- Both population-level and individual-level causal inference and fusion can be performed once the proper formal machinery is employed; algorithms and conditions are available.

# CONCLUSIONS

---

- Data-fusion requires encoding of structural features of the data-generating model.
- Both population-level and individual-level causal inference and fusion can be performed once the proper formal machinery is employed; algorithms and conditions are available.
- Principled framework for data-fusion — pooling and aggregating observational, experimental, and counterfactual information spread throughout heterogenous domains for population- and individual-level causal inference (understanding) and decision-making.

“Development of Western Science is based on two great achievements, the invention of the formal logical system (in Euclidean geometry) by the Greek philosophers, and the discovery of the possibility to find out causal relationships by systematic experiment (during the Renaissance)”.  
Albert Einstein.

“Imagine how much harder physics would be if electrons had feelings!”. Richard Feynman.

THANK YOU!