

Special Features:

This rock is probably closely related to the previously described sample. The alteration is similar in nature and origin.

History:

Igneous extrusive. Severely altered.

Sample P503/67 (Top): TS19990

Location:

430 feet 1 inch - 430 feet 2 inches.

Rock Name:

Altered, amygdaloidal, porphyritic ?andesite.

Hand Specimen:

A fine-grained, amygdaloidal, mottled green rock. The sample is in contact with a darker clastic rock. The contact is at an angle of approximately 30° to the core axis.

Thin Section:

A visual estimate of the constituents gives the following:

	<u>%</u>
Clay minerals	30
Carbonate	20
Chlorite	30
Epidote	3
Hydrated-iron oxides	15
Opakes	2

The rock is composed of an altered mass of subradiating plagioclase laths that have been variously replaced by carbonate and/or clay minerals. The interstitial areas are occupied by chlorite, carbonate, disseminated opakes and minor epidote. There is a tendency for the grain-size to coarsen as the contact with a clastic sandstone is approached, and hydrated iron-oxides and opakes increase in abundance towards the contact. Amygdules are quite common and are usually formed of a rim of chlorite enclosing a core of carbonate. Occasionally a fibrous, radiating clay mineral occupies the amygdules.

At a distance of approximately 10 mm from the contact a structure becomes evident. Here there is an alignment of the plagioclase laths which parallels the plane of the contact. This structure becomes more marked as the contact is approached. Immediately at the contact the structure disappears and the rock assumes the habit of an altered devitrified glass. The contact is quite sharp.

History:

Igneous extrusive and subsequently altered. The contact appears concordant although evidence for this is somewhat speculative.

Sample P503/67 (Bottom): TS19991

Location:

430 feet 2 inches - 430 feet 4 inches.

Rock Name:

Coarse-grained, volcaniclastic sandstone.

Hand Specimen:

A medium-grained, dark brownish-grey, massive clastic rock. Minor fractures randomly traverse the rock and are infilled with a secondary carbonate cement.

Thin Section:

A visual estimate of the constituents gives the following:

	<u>%</u>
Igneous rock fragments	60
Quartz	20
Carbonate	5
Clay minerals	10
Hydrated iron-oxides	5

In the main the rock is microscopically massive although there is a slight suggestion of a structure which is defined by the sub-parallel alignment of elongate fragments of quartz. Apart from some of the quartz the clastic components are rounded to sub-angular in shape and have a moderate sphericity. The rock fragments are derived from fine-grained, siliceous, rhyolitic rocks. Many show evidence of having a porphyritic texture. Phenocrysts of high-quartz are commonly included within the rock fragments. Phenocrysts of feldspar are much less common. The most common fragment type is

rhyolitic in composition and composed of a mottled aggregate of alkali feldspar and less commonly plagioclase. Other fragments tend towards the composition of rhyodacites and yet others, although minor, appear to have formed by the alteration and devitrification of an acid glass. Quartz fragments are common and it is possible that many, if not all, were derived from former phenocrysts.

The matrix is composed of hydrated iron-oxides, opaques, chlorite and clay minerals. Occasional flakes of detrital muscovite are found together with minor secondary calcite. Occasional, small fragments of detrital carbonate represent the only obviously sedimentary fragments.

There is a tendency for the grain-size to coarsen with depth down the hole and for the rock fragments to become more complete.

The rock is traversed by a few minor fractures that are infilled with carbonate.

There is no evidence to suggest that this rock may have been subjected to a metamorphic event.

This rock type is very significant in that it petrologically relates this formation to the Corunna Conglomerate. A few fragments of Moonachie Formation - Volcaniclastic phase are preserved in the detritus. These fragments are composed wholly of clastic rhyolitic rock fragments that are cemented together by a little fine-grained sericite. The fragments have been compared with rocks collected by Smale (1966) and are similar in composition and texture. They also appear identical to P526/67 at the bottom of Roopena DDH No.2 to the east of DDH No.1.

History:

Water laid sedimentary rock derived from a rhyolitic area of provenance.

Sample P504/67: TS19992

Location:

431 feet 6 inches - 431 feet 9 inches.

Rock Name:

Coarsely laminated arenaceous siltstone.

Hand Specimen:

A fine-grained, pale-grey, silty rock. The sample has a poor structure defined by variations in the grain-size of the detrital components.

Thin Section:

A visual estimate of the constituents gives the following:

	<u>5</u>
Quartz	35
Carbonate	8
Chlorite	5
Igneous rock fragments	15
Clay minerals	37

The rock is composed of scattered fragments of quartz, carbonate, chlorite and fine-grained igneous rocks the latter of which are identical to those described earlier. A few laths of fine-grained, detrital micas are scattered throughout. Variations in the ratio of fragments to matrix impart a structure which is further enhanced by the sub-parallel alignment of the long axes of the detrital grains. The matrix is essentially composed of clay minerals with lesser amounts of chlorite and very fine-grained quartz grains. In those zones in which there is a deficiency of the coarser-grained clastic components, the orientation of fibrous micas and clay minerals defines the structure. These laths are obviously detrital. All of the detrital components have been derived from a rhyolitic terrain: There is no evidence for any sedimentary fragments.

Sample P505/67: TS19993: TS19994

Location:

432 feet - f32 feet 9 inches.

Rock Name:

Lithic conglomerate.

Hand Specimen:

A conglomerate composed of angular to subrounded rock and mineral fragments set in a fine-grained matrix. Interbedded layers of fine-grained, laminated siltstone have a sub-parallel orientation that defines a prominent structure in the rock. This structure is approximately perpendicular to the core axis.

There is no positive evidence to suggest the rock was composed of igneous rocks or rock fragments.

History:

Sedimentary, deformed and recrystallized.

Sample P508/67: TS19997 - TS19998

Location:

490 feet - 490 feet 2 1/2 inches.

Rock Name:

Fine-grained, acid igneous rock.

Hand Specimen:

A dark reddish-brown mottled rock. The sample is completely massive although there is evidence for incipient deformation.

Thin Section:

The rock is composed of a fine-grained, felsitic mass of feldspar which has been partially reddened. The degree of reddening is variable and imparts the mottled colouration to the rock. The groundmass has a variable grain-size, but there is no consistent variation. Occasional phenocrysts of feldspar are found randomly distributed. Disseminated opaques are common accessory components together with lesser amounts of zircon and apatite. Fracture zones traverse the rock. These have been infilled with a mosaic of crystalline, magmatic quartz and small amounts of opaque minerals. The rock appears to have been subjected to mild deformation immediately after solidification and then been silicified by late-stage magmatic silica. Alteration is too intense to allow adequate identification of the feldspars but it is probable that the sample is rich in potash.

History:

Igneous extrusive.

Sample P509/67: TS19999

Location:

Beda Bore. Torrens 1 mile sheet.

Rock Name:

Pyroxene Basalt.

Hand Specimen:

Dark fine-grained fragments of a massive igneous rock.

Thin Section:

A visual estimate of the constituents from a fresh fragment gives the following:

	<u>%</u>
Plagioclase	60
Opagues	5
Pyroxene	25
Sphene	2
Mica	8

The freshest fragments are composed of fine- to medium-grained subradiating laths of plagioclase with the interstitial areas being occupied by a colourless clinopyroxene (augite). The feldspar is labradorite (An_{50} to An_{60}), crystals of which are partially replaced by muscovite and other colourless micas. Any interstitial areas are occupied by opaque minerals.

The submitted fragments vary in composition, as does the degree of alteration. Some fragments are almost devoid of mafic minerals, and composed wholly of plagioclase feldspar that is partially replaced by carbonate, chlorite and opagues (specular hematite). Although the fresher fragments predominate all gradations between the two extremes are found.

History:

Igneous.

crystals of opaques and angular quartz grains. Clay minerals are very fine-grained and show evidence for diagenetic recrystallization. Occasional mica laths are orientated in the plane of the bedding. Minor amounts of zircon were the only accessories identified. Two opaque mineral phases are present.

The first is pyrite which usually occurs as spherical or sub-spherical crystals which are usually associated with the darker layers in the rock. This implies a syngenetic origin for the pyrite (possibly biological). The second opaque mineral phase is chalcopyrite but this mineral is not closely related to the bedding. In one instance it is clearly related to cross fracturing. Occasionally it replaces pyrite spheroids. The chalcopyrite is definitely not syngenetic but is most likely of hypogene origin. Textural features suggest that the chalcopyrite is not the result of secondary enrichment.

History:

Sedimentary. There is no evidence to suggest a metamorphic event. The recrystallization can be ascribed only to diagenetic processes.

Sample P507/67: TS19996

Location:

477 feet 5 inches.

Rock Name:

A coarse-grained epiclastic lithic greywacke.

Hand Specimen:

A reddish brown, brecciated rock. The sample is completely massive.

Thin Section:

This rock is composed almost wholly of deformed and partially recrystallized quartz-rich sedimentary rock fragments. Each fragment has a pronounced structure defined by planes of deformation along which recrystallization has occurred. Protophase relics of the primary rock type are set in a fine-grained groundmass of recrystallized quartz and some sericite. The primary mineral components are severely strained. It is virtually impossible to distinguish the various fragments. The whole rock after consolidation has been subjected to slight deformation.

Thin Section:

A visual estimate of the constituents gives the following:

	<u>5</u>
Framework - ?Moonable Formation	15
- Acid igneous rocks	50
Matrix	35

This rock compositionally and texturally resembles P503 (volcaniclastic phase). The major differences include a marked increase in the number of fragments interpreted as having been derived from the Moonable Formation - volcaniclastic phase, the coarser grain-size, and the presence of graded bedding.

This rock is the first in which sedimentary rock fragments appear in significant proportions. Fragments of carbonate and deformed and recrystallized quartzites appear.

The siltstone interbeds have sharp contacts with the conglomerate. The fragments of which they are composed are consistent with their having been derived by the weathering of acid igneous rock fragments. Quartz fragments, some of which retain crystal faces, are scattered throughout. Zircon crystals are present in accessory amounts.

History:

Sedimentary.

Sample P506/67: TS19995: PS11251

Location:

465 feet 6 inches.

Rock Name:

Mineralized carbonaceous siltstone.

Hand Specimen:

A dark-grey, fine-grained, laminated siltstone. Sulphide minerals are distributed along parting fractures and bedding planes in the rock.

Thin Section:

The rock is composed of a laminated mass of clay minerals. The laminations are accentuated by sub-parallel precipitations of opaque minerals and/or carbonaceous material. Distributed through the clay minerals are finely disseminated and larger

LATITUDE LONGITUDE

BORE HOLE RECORD

INDEX NO.

39° 42' 51"

157° 24' 24"

NAME OF BORE/WELL: DOOPENA STRATIGRAPHIC D.D.H #1.

GEOLOGIC AREA : BARBER POINT FORM.

_____ :

_____ :

_____ :

NOTES ON LOCATION : S. H. Point 4.11.DATE COMPLETED :

0-430' Doopena Volcanics (basalt);

BACKY POINT FORMATION

ABBREVIATED LOG :REFERENCES : Trans. Doq. Soc. S. P. Vol. 94 (1973) p. 209-211 B. Thompson.REMARKS : 70 ft. Assays.

463-467 ft vert. depth.

width.

%

C.

467-469

2.

0.53

475-483-4

5.4

0.3

483.4-490

6.6

0.3

(S. H. Point 4.11 B. Thompson 8/17/71.

GEOCHEMICAL RESULTS ROOPENA DDH 1 409

Drilled by DM and analyzed by ASARCO.

INTERVAL (ft)	(PPM)				
	Cu	Pb	Zn	Ag	
0-10	55	190	270	< 1	
10-20	50	340	300		
20-30	40	270	350		
30-40	40	110	190		
40-50	40	135	100		
50-60	30	15	150		
60-70	40	10	90		
70-80	40	50	180		
80-90	45	20	170		
90-100	35	10	140		
100-110	25	15	130		
110-120	30	10	120		
120-130	170	480	190		
130-140	45	430	190		
140-150	45	45	280		
150-160	20	10	180		
160-170	35	15	120		
170-180	30	5	130		
180-190	35	5	110		
190-200	20	10	110		
200-210	35	55	150	↓	
210-220	30	30	160	3	
220-230	45	20	150	< 1	
230-240	40	120	150		
240-250	30	45	430		
250-260	45	15	130		
260-270	40	15	110		
270-280	50	30	120	↓	
280-290	30	35	150	3	

ROOPEENA DDH1
(ppm)

410

INTERVAL (ft)	C _n	P _g	Z _n	A _g	
290 - 300	30	160	330	21	
300 - 310	45	150	210		
310 - 320	45	160	190		
320 - 330	35	95	220		
330 - 340	55	340	380		
340 - 350	5	40	600		
350 - 360	35	1100	480		
360 - 370	60	5500	440		
370 - 380	60	1000	500		
380 - 390	5	25	460		
390 - 400	15	25	380		
400 - 410	60	250	280		
410 - 420	100	100	160		
420 - 430	120	15	210		
430 - 440	75	15	120		
440 - 450	30	15	75		
450 - 460	160	50	160		
460 - 470	4300	30	40		
470 - 480	900	160	25		
480 - 490	900	40	25		
490 - 500	85	5	15	5	
500 - 511	40	30	20	<1	



amdel

The Australian Mineral Development Laboratories

Flemington Street, Frewville, South Australia 5063
Phone 79 1662, telex AA82520

Please address all correspondence to the Director
In reply quote: **AN3/316/0 - 3961/72**

411

7 March 1972

The Manager
Asarco (Australia) Pty Limited
323 Wakefield Street
ADELAIDE SA 5000

REPORT AN3961/72

YOUR REFERENCE: Order A1743
IDENTIFICATION: A14701 to A14751
DATE RECEIVED: 23/2/72

Enquiries quoting AN3961/72 to Officer in Charge please.

Analysis by: A.E. Francis

Officer in Charge, Analytical Section: A.B. Timms

ATD
for F.R. Hartley
Director

Roopena No1 DDH.

plm

MAP		STATE	SA
GRID			
PROSPECT	Base Hole Survey		
SAMPLER	SBW	FILE	
TYPE	Core dust		

JOB 3961/12 AMDEL ANALYTICAL SERVICE
FORM 6

TT	Sample No.						Feet		
			Cu	Pb	Zn	Hg	From	To	
1	H 14701		55	190	270	<1	0	10	weather volcan
2	2		50	340	300	<1	10	20	"
3	3		40	270	350	<1	20	30	"
4	4		40	110	190	<1	30	40	"
5	5		40	35	100	<1	40	50	"
6	Str 51						5		"
7	6		30	15	150	<1	50	60	"
8	7		40	10	90	<1	60	70	"
9	8		40	50	180	<1	70	80	volcan
10	9		45	20	170	<1	80	90	"
11	10		35	10	140	<1	90	100	"
12	11		25	15	130	1	100	110	"
13	12x		30	10	120	<1	110	120	"
14	13		170	480	190	1	120	130	"
15	14		45	430	190	<1	130	140	"
16	15		45	45	200	<1	140	150	"
17	16		20	10	180	<1	150	160	? pyrit
18	17		35	15	120	<1	160	170	volcan
19	11 14718		30	5	130	<1	170	180	
20	12 ✓								

FORM 76

JOB

2961/12

AMDEL ANALYTICAL SERVICE

BATCH NO.

413

2

TT	Sample No.		Cu	Pb	Zn	Fe	From	Act To	
1	FA 12719		35	5	110	<1	130	190	volca
2	20		20	10	110	<1	190	200	"
3	1		35	55	150	<1	200	210	"
4	2		30	30	160	3	210	220	"
5	3		45	20	150	1	220	230	"
6	4		40	120	150	1	230	240	"
7	5x		30	45	430	<1	240	250	"
8	6		45	15	130	<1	250	260	"
9	7		40	15	110	<1	260	270	"
10	8		50	30	120	<1	270	280	"
11	9		30	35	150	3	280	290	"
12	10		30	160	330	1	290	300	"
13	1		45	150	210	<1	300	310	"
14	2		45	160	190	<1	310	320	"
15	3		35	95	220	<1	320	330	"
16	51								
17	4		55	340	380	<1	330	340	"
18	5		5	40	600	<1	340	350	"
19	12736		35	1100	480	<1	350	360	"
20	25x								

TT	Sample No.		Cu	Pb	Zn	Fe	From	To	
1	A 14731		60	5500	440	<1 VOLCS	360	370	sulphur volcanics
2	8		60	1000	500	<1 VOLCS	370	380	volcanics
3	9		5	25	460	<1 VOLCS	380	390	"
4	51051								
5	40		15	25	380	<1 ROOPENA	390	400	"
6	1		60	250	280	<1 ROOPENA	400	410	"
7	2		100	100	160	<1 ROOPENA	410	420	"
8	3		120	15	210	<1 ROOPENA	420	430	"
9	4		75	15	120	<1 ROOPENA	430	440	arkose
10	5		30	15	75	<1 ROOPENA	440	450	"
11	6		160	50	160	<1 ROOPENA	450	460	"
12	7	*	4300	30	40	1 BLACK POINT	460	470	siltstone
13	8		900	160	25	1 BLACK POINT	470	480	"
14	9		900	40	25	1 BLACK POINT	480	490	pyrite
15	10		85	5	15	5 Jc	490	500	siltstone
16	11 11151		40	30	20	1	500	511 ft.	"
17	45		NO OTHER SAMPLES TAKEN				Cu Pb Zn - Code C		
18	BLANK		—	—	—	—	Ag - Code CV		
19							All Results in PPM		
20									

High Pb - several - Highest 5500 ppm 360 to 370 feet.

High Cu 4300 ppm From 460 to 470 feet.

* DPM records give:

463 to 467	0.35% Cu
467 to 469	0.52% Cu
478 to 483.4	0.3% Cu
483.4 to 490	0.2% Cu

BORE AND WELL SPOIL SAMPLES

Obtained

By

D. KER, 1963..

WESTERN STUART SHELF AREA.

Sample No.	Grid Reference	Latitude: Longitude:	Bore Ref. No.	Name of Bore or Well	Palaeo Sample No.	Remarks
✓ 26502	J/3	28° 29' 138° 41'	23	Apatoongannie Well	F43/63	
26501				1/4 Mile SSW of Mungeranie H/S (Shale)	F44/63	Surface Spoil Sample (not plotted)

WESTERN STUART SHELF AREA.

Sample No.	Grid Reference	Latitude: Longitude:	Bore Ref. No.	Name of Bore or Well	Palaeo Sample No.	Remarks
✓ 26508	E/3	28° 50' 133° 53'	2	Paragon Bore	F12/63	18S, 14N Mt. Willoughby (near Homestead)
✓ 26509	E/3	28° 57' 133° 59'	6	Watchercallem Bore	F11/63	
✓ 26510	E/4	29° 01' 133° 56'	2	Corkscrew Bore	F13/63	
✓ 26513	F/2	27° 57' 134° 27'	9	Timothy Bore	F16/63	Copper Hill Stn.
✓ 26512	F/2	28° 00' 134° 23'	8	Patricia Bore	F15/63	Appreetina South Copper Hill Stn.
✓ 26511	F/2	27° 57' 134° 16'	5	No. 2 Bore	F14/63	Copper Hill Stn. Appreetina South
✓ 26514	F/3	28° 47' 134° 19'	10	No. 5 Bore	F17/63	
✓ 26515	F/3	28° 51' 134° 22'	14	Mt Clarence H/S Deep Bore	F18/63	
✓ 26516	F/3	28° 53' 134° 37'	18	Sputnick Bore	F19/63	
✓ 26517	F/3	28° 29' 134° 11'	31	Pooramungie Bore	F20/63	
✓ 26518	F/3	28° 50' 134° 41'	40	Giddy Giddina Bore	F21/63	

WESTERN STUART SHELF AREA.

Sample No.	Grid Reference	Latitude: Longitude:	Bore Ref. No.	Name of Bore or Well	Palaeo Sample No.	Remarks
/ 26519	F/3	28° 22' 134° 11'	32	Lesley's Bore	F22/63	Murloocoppie West (Willoughby)
/ 26520	F/3	28° 56' 134° 10'	44	Craigs Bore	F23/63	
/ 26521	F/3	28° 55' 134° 15'	45	West Point Bore	F24/63A	
/ 26522	F/3	28° 55' 134° 15'	45	West Point	F24/63B	
/ 26523	F/3	28° 42' 134° 24'	58	Broken Bit Bore	F26/63	
/ 26524	F/3	28° 06' 134° 06'	47	Dead Finish Bore	F27/63	Mount Willoughby
/ 26527	F/3	28° 12' 134° 23'	52	Bridget Bore	F28/63	Evelyn Downs
/ 26506	F/3	28° 50' 134° 27'	16	Yellow Bullock Bore	F29/63	
/ 26505	F/3	28° 57' 134° 39'	29	Willow No. 3 Bore	F30/63	
/ 26504	F/3	28° 56' 134° 20'	41	Abandoned Bore H/S	F31/63	
/ 26525	F/4	29° 05' 134° 36'	129	Box Hole Bore	F53/63	14N, 14S

· WESTERN STUART SHELF AREA.

Sample No.	Grid Reference	Latitude: Longitude:	Bore Ref. No.	Name of Bore or Well	Palaeo Sample No.	Remarks
✓ 26526	F/4	29° 07' 134° 29'	131	Abandoned	F51/63	
✓ 26527	F/4	29° 11' 134° 20'	12	Abandoned Bore	F52/63	Green Sand
✓ 26528	F/4	29° 11' 134° 20'	12	Abandoned Bore	F57/63	Blue Sand
✓ 26529	F/4	29° 22' 134° 24'	38	Abandoned Bore	F55/63	
✓ 26530	F/4	29° 05' 134° 20'	4	No. 2 Bore	F56/63	
✓ 26531	F/4	29° 39' 135° 00'	152	Abandoned Well	F50/63	
✓ 26532	F/4	29° 10' 134° 16'	11	Abandoned Well	F54/63	Mabel Creek
✓ 26503	F/4	29° 03' 134° 16'	3	Abandoned Bedoura Bore	F32/63	Woorong Creek
✓ 26533	F/4	29° 06' 134° 28'	130	Abandoned Bore	F49/63	Mount Clarence
✓ 26534	F/4	29° 24' 134° 46'	135	Abandoned Bore	F48/63	
✓ 26535	F/4	29° 32' 134° 55'	147	Abandoned Bore	F47/63	

WESTERN STUART SHELF AREA.

Sample No.	Grid Reference	Latitude: Longitude:	Bore Ref. No.	Name of Bore or Well	Palaeo Sample No.	Remarks
26536	F/4	29° 02' 134° 18'	180	Ping Pong Bore	F46/63	Woorong Creek
26537	F/5	30° 15' 134° 46'	93	Glynn Bore	F61/63	Bulgunnia
26538	F/5	30° 12' 134° 40'	75	Abandoned Bore	F60/63	Bulgunnia
26539	F/5	° °	94	Sextabyng Well	F59/63	Bulgunnia
26540	F/5	30° 26' 134° 58'	133	New Bore East 34 Pdk.	F58/63	Wilgena
26541	G/4	29° 52' 135° 13'	12	Dept. Works Mirikita	F69/63	13N, 13S The Twins
26542	G/4	° °	15	Barbies Bore	F71/63	The Twins
26543	G/4	29° 58' 135° 08'	18	Coronation Bore	F70/63	
26544	G/4	29° 57' 135° 55'	34	No. 6 Bore	F67/63	
26545	G/4	° °	42	Blue Duck Well	F72/63	Millers Creek
26546	G/4	29° 49' 135° 49'	26	Beetram Downs No. 1	F68/63	

WESTERN STUART SHELF AREA.

Sample No.	Grid Reference	Latitude: Longitude:	Bore REF. No.	Name of Bore or Well	Palaeo Sample No.	Remarks
26547	G/4	29° 05' 135° 42'	60	Pages Well No. 2	F73/63	
26548	G/4	29° 40' 135° 01'	7	New Year Well	F66/63	
26549	G/4	29° 58' 135° 30'	30	Beetram Downs No. 4 Bore	F65/63	
26550	G/4	29° 47' 135° 55'	32	No. 15 Bore	F64/63	Millers Creek
26551	G/4	29° 40' 135° 01'	45	Abandoned Well	F63/63	
26552	G/4	29° 01' 135° 49'	53	New Bore	F62/63	
X 26553	G/5	30° 34' 135° 53'	211	Clemens Bore	F96/63	13S
X 26554	G/5	30° 14' 135° 35'	274	Cavanaugh Bore	F90/63	
X 26555	G/5	30° 17' 135° 47'	278	Birthday Bore	F91/63	
X 26556	G/5	30° 07' 135° 33'	253	Backlands Bore (old)	F100/63	
X 26557	G/5	30° 04' 135° 34'	264	Backlands Bore (new)	F87/63	

WESTERN STUART SHELF AREA.

Sample No.	Grid Reference	Latitude: Longitude:	Bore Ref. No.	Name of Bore or Well	Palaeo Sample No.	Remarks
x/ 26558	G/5	30° 04' 135° 45'	265	Abandoned Gondalloo Bore (old)	F88/63	/
x/ 26559	G/5	30° 18' 135° 48'	279	Buckley Bore	F92/63	/
✓ 26560	G/5	30° 19' 135° 52'	61	Abandoned Bore	F94/63	/
✓ 26561	G/5	30° 07' 135° 49'	57	Paisley Well	F109/63	
✓ 26562	G/5	30° 11' 135° 44'	56	Russell Bore	F101/63	Mt. Eba
✓ 26563	G/5	30° 08' 135° 10'	40	Gondalloo Bore (new)	F93/63	
x/ 26564	G/5	30° 00' 135° 23'	257	Banjo Bore	F89/63	Nth of Mt. Eba
x/ 26565	G/5	30° 01' 135° 23'	258	Twins Stn. Bore	F108/63	
/ 26566	G/5	° °	191	Bore No. 2	F107/63	Not plotted. Near Bore 308 in G/5
x/ 26567	G/5	30° 26' 135° 50'	308	Bore No. 2	F106/63	
x/ 26568	G/5	30° 23' 135° 50'	181	Abandoned Bore Nth. Vivian Well Hut	F96/63	

Sample No.	Grid Reference	Latitude: Longitude:	Bore Ref. No.	Name of Bore or Well	Palaeo Sample No.	Remarks
✓ 26569	G/5	30° 25' 135° 50'	297	Abandoned Bore	F105/63	
✓ 26570	G/5	30° 26' 135° 50'	315	Abandoned "	F103/63	
✓ 26571	G/5	30° 25' 135° 52'	317	Deep Bore	F99/63	
✓ 26572	G/5	30° 25' 135° 50'	314	Abandoned Bore	F104/63	
✓ 26573	G/5	30° 48' 135° 48'	295	Adj. to Pyra- mid Bore	F98/63	
✓ 26574	G/5	30° 57' 135° 39'	237	Abandoned Well (McKinnons Dam)	F85/63	
✓ 26575	G/5	30° 34' 135° 44'	204	Southern Cross Well	F110/63	
✓ 26576	G/5	30° 22' 135° 10'	177	Bluff Paddock Bore	F97/63	
✓ 26577	G/5	30° 49' 135° 27'	154	Sludge	F86/63	
✓ 26578	G/5	30° 24' 135° 50'	193	Police Camp Paddock	F102/63	
✓ 26579	G/5	30° 04' 135° 19'	17	Crombies Bore	F111/63	
✓ 26580	G/5	30° 05' 135° 21'	19	Bald Hill Salt Bore	F112/63	