

014

E X O T I C

Nipaecoccus nipae

Coconut Mealybug

Field Recognition

Body maroon to red, about 3 mm long, with few side (lateral) wax filaments. Dorsal wax filaments occurring. Body round to oval. No ovisac produced, males common.

Known Southeastern Distribution

Florida and Louisiana.

Common Hosts

Polyphagous species, most common on palms.

Coconut Mealybug
Nipaecoccus nipae



014

E X O T I C

Nipaecoccus nipae

Coconut Mealybug

Field Recognition

Body maroon to red, about 3 mm long, with few side (lateral) wax filaments. Dorsal wax filaments occurring. Body round to oval. No ovisac produced, males common.

Known Southeastern Distribution

Florida and Louisiana.

Common Hosts

Polyphagous species, most common on palms.

Coconut Mealybug
Nipaecoccus nipae

Male Puparia



014

E X O T I C

Nipaecoccus nipae

Coconut Mealybug

Field Recognition

Body maroon to red, about 3 mm long, with few side (lateral) wax filaments. Dorsal wax filaments occurring. Body round to oval. No ovisac produced, males common.

Known Southeastern Distribution

Florida and Louisiana.

Common Hosts

Polyphagous species, most common on palms.

Coconut Mealybug
Nipaecoccus nipae

