

Project 2: Variational Quantum Eigensolver: Constructing Potential Energy Surfaces for Small Molecules

Introduction

Currently, the Variational Quantum Eigensolver (VQE) [1] is the most feasible technique for solving the electronic structure problem on a near-term noisy quantum computer. Yet, one needs to understand that VQE is a quite general framework that requires several choices to be made by the user to be efficient. In what follows, we will illustrate state-of-the-art techniques to make VQE efficient for obtaining potential energy surfaces (PESs) for small molecules.

A set of molecules suggested for investigation is: H_2 , LiH , H_4 , H_2O , N_2 . They are of the increasing level of difficulty. A simple qualitative rule from chemistry is: the system PES is harder to calculate accurately if more chemical bonds are destroyed at the same time. Therefore, H_2 and LiH are the easiest systems (single bond), H_4 and H_2O are harder (two bonds), and N_2 is the hardest (triple bond). (Actually, 1-2% of the entire worldwide energy is spent on breaking N_2 bonds to make ammonia (NH_3) that is turned into fertilizers. Thus being able to model such processes is a necessary step for computational development of new catalysts for the ammonia production.) As a smaller correction to this qualitative picture of computational hardness, one can add a number of electrons in the system, the larger the number of electrons the more computationally demanding is the study. Thus, LiH is harder than H_2 and H_2O is harder than H_4 .

Another qualitative indicator of hardness is whether classical computing approximate methods for electronic structure that have a polynomial scaling are able to treat the system with chemical accuracy (1 kcal/mol). Turns out, if one breaks more than a single bond state-of-the-art approaches like coupled cluster single and double (CCSD) cannot achieve chemical accuracy for all points of PES. Thus, quantum computing methods based on VQE were thought as a viable alternative to treat such so-called strongly correlated systems.

Your task

Using H_2 and H_2O examples, we will illustrate main steps of the VQE process from the setup of the Hamiltonian till obtaining the circuit for the IBM quantum computer. Your task will be to repeat those steps for one or two other molecules so that you can obtain the ground state PES of your molecules both on a classical computer using classical methods and VQE simulator, and on a quantum computer.

Recommended bonds elongation for H_4 and H_2O are depicted in Fig. 1. Tequila is a free software package where all calculations will be illustrated.




Figure 1: Recommended distortions: H_2O , symmetric OH bond length changes around $R(\text{O}-\text{H})=0.75 \text{ \AA}$ and fixed $\angle \text{HOH} = 107.6^\circ$; H_4 , change of β in the range $[85^\circ, 95^\circ]$ keeping planar configuration with fixed $R = 1.738 \text{ \AA}$.

Step #1: Generating PES using classical methods

Hartree-Fock (HF), Configuration Interaction Singles and Doubles (CISD), Coupled Cluster Singles and Doubles (CCSD), and the exact answer within the chosen basis Full Configuration Interaction (FCI) [2]. All these methods can easily be used for all molecules in minimal atomic basis, STO-3G. Comparing accuracy of PES generated for different species using different methods you can see the difference between weakly correlated (H_2 , LiH) and strongly correlated (H_4 , H_2O , N_2) systems. Even though the FCI method provides the exact answer it scales exponentially with the number of basis functions and thus cannot be used for larger systems (system size can be either correlated with the number of electrons or number of basis functions). All approximated methods (HF, CISD, CCSD) scale polynomially with the system size but fail to deliver chemical accuracy along PESs.

Illustrative code: `S1_Classical_Methods.ipynb`.

Step #2: Generating the qubit Hamiltonian

To proceed to VQE one needs to generate the qubit Hamiltonian, the easiest path is via first generating the electronic Hamiltonian in the second quantized form and then transform it into the qubit form using one of the fermion-to-qubit transformations: Jordan-Wigner or Bravyi-Kitaev [3]. Next, some qubit operators can be substituted by numbers (± 1) because their states are stationary for the specific electronic state (e.g. ground state). This reduction is very useful for fitting larger problem in a fewer qubit description and is based on Hamiltonian symmetries, which are discussed in Refs. [4, 5, 6]

Illustrative code: `S2_Hamiltonian_gen.ipynb`.

Step #3: Unitary transformations

There are three main paths : 1) historically first, the unitary coupled cluster (UCC) approach has originated from the ideas of the classical electronic structure theory [7, 8], 2) hardware-efficient, direct optimization of gates and their sequence [9], and 3) the qubit coupled cluster (QCC) based approach [10]. We will illustrate the first and third approaches, since the second one frequently runs into difficulties with energy optimization [11]. Among the two, even though UCC follows directly from a classical molecular orbital picture, it doesn't provide an efficient framework for generating gates later. In contrast, QCC can efficiently optimize energy and generate efficient circuits.

In this step, we will look at how to generate elementary unitary operations in the UCC and QCC schemes for model systems and to optimize their continuous parameters (amplitudes).

Illustrative code: `S3_U_ansatz.ipynb`.

Step #4: Hamiltonian measurements

To obtain the expectation value of the qubit Hamiltonian it needs to be measured at the end of the VQE circuit. The problem is that the entire Hamiltonian cannot be measured using current hardware. In this step, we show how to partition the Hamiltonian to a minimal number of groups, whose elements can be all measured simultaneously [12, 13].

Illustrative code: `S4_Measurement.ipynb`.

Step #5: Use of quantum hardware

To carry out the VQE algorithm on the actual quantum hardware one needs to present unitary transformations found in steps 3 and 4 as a sequence of gates (i.e. circuit). Here, we will illustrate circuit generation using compilers. Obtained circuits will allow one to either carry out the entire VQE optimization procedure by optimizing amplitudes of step 3 unitaries or its single step using amplitudes for step 3 obtained on a classical simulator (easier route).

Illustrative code: `S5_Circuits.ipynb`.

Further Challenges

Now you're on your own. Discuss ideas for expanding on the above steps with your group. Some possible directions for more advanced treatment:

1. How to obtain excited electronic states of the same or different symmetry? This task is highly relevant to any spectroscopic application of quantum computing because in real experiments and applications not absolute energy but their differences are measured and are important.
2. Hamiltonian measurements (Step #4): There are more advanced techniques to consider that are based on partitioning in the fermionic operator space and then bringing the fragments into the qubit space [14]
3. Unitary transformations (Step #3): Some of the unitary transformations can be applied on the Hamiltonian instead of the wavefunction, this increases accuracy without increasing the circuit depth, but it increases the size of the Hamiltonian and thus introduces the measurement overhead. See Refs. [15, 16] for details.
4. Qubit Hamiltonian preparation (Step #2): Typical fermion-qubit transformations substitute a single molecular spin-orbit by a qubit, which restricts calculations to relatively small basis sets because of the current limitations in the number of qubits. There were several techniques proposed recently to compress larger basis sets into smaller number of qubits [17, 18].
5. What are alternatives to VQE for the electronic structure problem using quantum computers with shallow circuits without error-correction?

Business-related questions in the context of quantum entrepreneurship:

1. What other molecular properties one can obtain using VQE for the purpose of rational material and drug design?
2. What are the systems (molecules / materials) which are challenging for classical computing and whose modelling is valuable ?
3. What businesses can benefit from more accurate electronic structure calculations ?

References

- [1] Alberto Peruzzo, Jarrod McClean, Peter Shadbolt, Man-Hong Yung, Xiao-Qi Zhou, Peter J. Love, Alán Aspuru-Guzik, and Jeremy L. O’Brien. A variational eigenvalue solver on a photonic quantum processor. *Nat. Commun.*, 5(1):4213, dec 2014.
- [2] Trygve Helgaker, Poul Jørgensen, and Jeppe Olsen. *Molecular Electronic-Structure Theory*. John Wiley & Sons, Ltd, Chichester, UK, aug 2000.
- [3] Jacob T. Seeley, Martin J. Richard, and Peter J. Love. The bravyi-kitaev transformation for quantum computation of electronic structure. *J. Chem. Phys.*, 137:224109–224109, Dec 2012.
- [4] Sergey Bravyi, Jay M Gambetta, Antonio Mezzacapo, and Kristan Temme. Tapering off qubits to simulate fermionic Hamiltonians. *arXiv.org*, January 2017.
- [5] Tzu-Ching Yen, Robert A Lang, and Artur F Izmaylov. Exact and approximate symmetry projectors for the electronic structure problem on a quantum computer. *J. Chem. Phys.*, 151(16):164111, October 2019.
- [6] Kanav Setia, Richard Chen, Julia E Rice, Antonio Mezzacapo, Marco Pistoia, and James Whitfield. Reducing qubit requirements for quantum simulation using molecular point group symmetries. *arXiv.org*, October 2019.
- [7] Jonathan Romero, Ryan Babbush, Jarrod R McClean, Cornelius Hempel, Peter J Love, and Alán Aspuru-Guzik. Strategies for quantum computing molecular energies using the unitary coupled cluster ansatz. *Quantum Sci. and Technol.*, 4(1):014008, oct 2018.
- [8] Cornelius Hempel, Christine Maier, Jonathan Romero, Jarrod McClean, Thomas Monz, Heng Shen, Petar Jurcevic, Ben P. Lanyon, Peter Love, Ryan Babbush, Alán Aspuru-Guzik, Rainer Blatt, and Christian F. Roos. Quantum Chemistry Calculations on a Trapped-Ion Quantum Simulator. *Phys. Rev. X*, 8(3):31022, 2018.
- [9] Abhinav Kandala, Antonio Mezzacapo, Kristan Temme, Maika Takita, Markus Brink, Jerry M. Chow, and Jay M. Gambetta. Hardware-efficient variational quantum eigensolver for small molecules and quantum magnets. *Nature*, 549:242–246, Sep 2017.
- [10] Ilya G. Ryabinkin, Tzu-Ching Yen, Scott N. Genin, and Artur F. Izmaylov. Qubit coupled cluster method: A systematic approach to quantum chemistry on a quantum computer. *J. Chem. Theory Comput.*, 14(12):6317–6326, 2018.

- [11] Jarrod R. McClean, Sergio Boixo, Vadim N. Smelyanskiy, Ryan Babbush, and Hartmut Neven. Barren plateaus in quantum neural network training landscapes. *Nat. Commun.*, 9:1–6, 2018.
- [12] Vladyslav Verteletskyi, Tzu-Ching Yen, and Artur F Izmaylov. Measurement optimization in the variational quantum eigensolver using a minimum clique cover. *J. Chem. Phys.*, 152(12):124114, March 2020.
- [13] Tzu-Ching Yen, Vladyslav Verteletskyi, and Artur F. Izmaylov. Measuring all compatible operators in one series of a single-qubit measurements using unitary transformations. *J. Chem. Theory Comput.*, 16:2400–2409, Jul 2020.
- [14] William J. Huggins, Jarrod McClean, Nicholas Rubin, Zhang Jiang, Nathan Wiebe, K. Birgitta Whaley, and Ryan Babbush. Efficient and Noise Resilient Measurements for Quantum Chemistry on Near-Term Quantum Computers. *arXiv e-prints*, Jul 2019.
- [15] Ilya G. Ryabinkin, Robert A. Lang, Scott N. Genin, and Artur F. Izmaylov. Iterative qubit coupled cluster approach with efficient screening of generators. *J. Chem. Theory Comput.*, 16(2):1055–1063, 2020.
- [16] Robert A Lang, Ilya G Ryabinkin, and Artur F Izmaylov. Iterative Qubit Coupled Cluster method with involutory linear combinations of Pauli products. *arXiv.org*, February 2020.
- [17] Tyler Takeshita, Nicholas C Rubin, Zhang Jiang, Eunseok Lee, Ryan Babbush, and Jarrod R McClean. Increasing the representation accuracy of quantum simulations of chemistry without extra quantum resources. *arXiv.org*, February 2019.
- [18] Nicholas P. Bauman, Guang Hao Low, and Karol Kowalski. Quantum simulations of excited states with active-space downfolded Hamiltonians. *arXiv e-prints*, Sep 2019.