
Actual graphs and paired predictions

BigEarthNet-19 | trained seed42 models | identical inputs | all four archived cases


RGB input	Fixed 3 x 3 (K=9)	Fixed 4 x 4 (K=16)

(a) Validation row 97389



Pastures (y=1): 29.89% -> 46.36% | still wrong


(b) Validation row 14680



Natural grassland / sparse vegetation (y=1): 42.85% -> 51.59% | corrected

Pastures (y=0): 64.38% -> 69.97% | still wrong


(c) Validation row 28450



Transitional woodland / shrub (y=1): 44.74% -> 91.82% | corrected

Marine waters (y=1): 64.96% -> 5.84% | new miss

(d) Validation row 4439



Urban fabric (y=1): 41.38% -> 76.57% | corrected

Beaches / dunes / sands (y=1): 17.67% -> 49.32% | still wrong


Orange arrows: captured feature edges only (18 for K9, 32 for K16); neighbor -> query.
Region colors and IDs indicate spatial bins. Spatial edges and all 19 scores are retained in HTML.
At threshold 0.5: 3 corrected label decisions, 1 new miss. Four cases are not a population sample.

Within-region pooling weights

BigEarthNet-19 | trained seed42 models | identical inputs | all four archived cases

RGB input	Fixed 3 x 3 (K=9)	Fixed 4 x 4 (K=16)

(a) Validation row 97389




Regions with max pooling mass >=0.90: 3x3 6/9; 4x4 12/15 (non-singleton only).

(b) Validation row 14680




Regions with max pooling mass >=0.90: 3x3 7/9; 4x4 15/15 (non-singleton only).

(c) Validation row 28450



Regions with max pooling mass >=0.90: 3x3 8/9; 4x4 14/15 (non-singleton only).

(d) Validation row 4439



Regions with max pooling mass >=0.90: 3x3 4/9; 4x4 7/15 (non-singleton only).

Red: above uniform mass in the same region. Blue: below uniform. Transparent: uniform.
Mass is normalized within each region; heat is not class attribution or a semantic segmentation.
K16 has one single-token region: mass=1 there is compulsory. No new model training or inference.