



LONDON
MATHEMATICAL
SOCIETY
EST. 1865



DEPARTMENT OF
STATISTICS

The oriented contact process and the East model

Patrik Gerber supervised by Prof. Paul Chleboun

Supported by LMS and the Mathematical Institute, University of Oxford.

Department of Statistics
University of Oxford

4 October 2018

East process

Front propagation

Mixing time

One-sided contact process

East vs contact

Results for one-sided contact process

Percolation construction

References

Let me first give you an overview of what my project was about. It started out primarily concerned with the East-process. The idea was to apply a recent comparison based method that yielded new results for a different process. The corresponding results for the East process are already known, but an alternative, shorter proof could give indication that this comparison method could be useful for other KCMs of oriented nature, the class of processes that the East process belongs to. Upon reviewing the literature it became apparent that the necessary results for the oriented contact process were never precisely stated or proven even though the machinery had been developed. In the past 6 weeks we created a self contained proof of the necessary results and apply them to prove a weaker version of the known mixing time results for the East process.

1. So let me introduce the East process to you.
2. read slide
3. Its state space is the set of configurations of 0s and 1s on the Integer lattice. Talk about powerset.
4. Talk about graphical representation. Point out cross that doesn't kill spin.

The East process is a continuous time Markov chain

1. So let me introduce the East process to you.
2. read slide
3. Its state space is the set of configurations of 0s and 1s on the Integer lattice. Talk about powerset.
4. Talk about graphical representation. Point out cross that doesn't kill spin.

East process



The East process is a continuous time Markov chain .

- ▶ State space: $\Omega := \{0, 1\}^{\mathbb{Z}}$, or equivalently, $\mathcal{P}(\mathbb{Z})$.

1. So let me introduce the East process to you.
2. read slide
3. Its state space is the set of configurations of 0s and 1s on the Integer lattice. Talk about powerset.
4. Talk about graphical representation. Point out cross that doesn't kill spin.

East process



The East process is a continuous time Markov chain .

- ▶ State space: $\Omega := \{0, 1\}^{\mathbb{Z}}$, or equivalently, $\mathcal{P}(\mathbb{Z})$.
- ▶ Each site $x \in \mathbb{Z}$ updates with rate 1 to $B \sim \text{Ber}(p)$, $p \in (0, 1)$ **iff** $x - 1$ is occupied.

1. So let me introduce the East process to you.
2. read slide
3. Its state space is the set of configurations of 0s and 1s on the Integer lattice. Talk about powerset.
4. Talk about graphical representation. Point out cross that doesn't kill spin.

East process

The East process is a continuous time Markov chain .

- ▶ State space: $\Omega := \{0, 1\}^{\mathbb{Z}}$, or equivalently, $\wp(\mathbb{Z})$.
- ▶ Each site $x \in \mathbb{Z}$ updates with rate 1 to $B \sim \text{Ber}(p)$, $p \in (0, 1)$ **iff** $x - 1$ is occupied.




Figure: East process started from $\{0\}$

1. So let me introduce the East process to you.
2. read slide
3. Its state space is the set of configurations of 0s and 1s on the Integer lattice. Talk about powerset.
4. Talk about graphical representation. Point out cross that doesn't kill spin.

1. The first step towards the proof of the linearity of the mixing time is a result about the front of the East process
2. Read slide
3. Read theorem. What this says is
4. We give an alternative proof but for now, assuming this result, we can go on to show the desired linearity of the mixing time.

Definition (Front)

Let $(\sigma_t)_{t \geq 0}$ be an East process started from $\{0\}$. The front of $(\sigma_t)_{t \geq 0}$ is the rightmost occupied site, written $X(\sigma_t) := \sup \sigma_t$.

1. The first step towards the proof of the linearity of the mixing time is a result about the front of the East process
2. Read slide
3. Read theorem. What this says is
4. We give an alternative proof but for now, assuming this result, we can go on to show the desired linearity of the mixing time.

Definition (Front)

Let $(\sigma_t)_{t \geq 0}$ be an East process started from $\{0\}$. The front of $(\sigma_t)_{t \geq 0}$ is the rightmost occupied site, written $X(\sigma_t) := \sup \sigma_t$.

Theorem (Blondel, 2013, Lemma 3.2)

There exist constants $0 < \alpha < v$ and $\gamma, C > 0$ such that

$$\mathbb{P}(X(\sigma_t) \in (\alpha t, vt)) \geq 1 - Ce^{-\gamma t} \quad \forall t \geq 0.$$

1. The first step towards the proof of the linearity of the mixing time is a result about the front of the East process
2. Read slide
3. Read theorem. What this says is
4. We give an alternative proof but for now, assuming this result, we can go on to show the desired linearity of the mixing time.

Definition (Front)

Let $(\sigma_t)_{t \geq 0}$ be an East process started from $\{0\}$. The front of $(\sigma_t)_{t \geq 0}$ is the rightmost occupied site, written $X(\sigma_t) := \sup \sigma_t$.

Theorem (Blondel, 2013, Lemma 3.2)

There exist constants $0 < \alpha < v$ and $\gamma, C > 0$ such that

$$\mathbb{P}(X(\sigma_t) \in (\alpha t, vt)) \geq 1 - Ce^{-\gamma t} \quad \forall t \geq 0.$$

Proof.

We propose an alternative proof (under restrictions on p) detailed later.



1. The first step towards the proof of the linearity of the mixing time is a result about the front of the East process
2. Read slide
3. Read theorem. What this says is ...
4. We give an alternative proof but for now, assuming this result, we can go on to show the desired linearity of the mixing time.

1. I hope you are all familiar with the mixing time of a Markov chain.
2. Talk about restriction to $[0, L]$
3. Read theorem.
4. Talk about how to use front to prove this. Talk about restriction on p .

Restricted to $\{0, 1, \dots, L\}$ with a fixed 1 at origin East process converges to equilibrium.

1. I hope you are all familiar with the mixing time of a Markov chain.
2. Talk about restriction to $[0, L]$
3. Read theorem.
4. Talk about how to use front to prove this. Talk about restriction on p .

Restricted to $\{0, 1, \dots, L\}$ with a fixed 1 at origin East process converges to equilibrium.

Theorem (Cancrini et al., 2008, Theorem 6.1)

The mixing time of the East process on $\{0, 1, \dots, L\}$ with a fixed 1 at the origin is $\Theta(L)$.

1. I hope you are all familiar with the mixing time of a Markov chain.
2. Talk about restriction to $[0, L]$
3. Read theorem.
4. Talk about how to use front to prove this. Talk about restriction on p .

Restricted to $\{0, 1, \dots, L\}$ with a fixed 1 at origin East process converges to equilibrium.

Theorem (Cancrini et al., 2008, Theorem 6.1)

The mixing time of the East process on $\{0, 1, \dots, L\}$ with a fixed 1 at the origin is $\Theta(L)$.

Alternative proof using only front propagation.

Upper bound: Coupling time vs. propagation of the front.

Lower bound: Hitting times of large sets (Peres and Sousi, 2015).



1. I hope you are all familiar with the mixing time of a Markov chain.
2. Talk about restriction to $[0, L]$
3. Read theorem.
4. Talk about how to use front to prove this. Talk about restriction on p .

1. So I explained how the front propagation result can be used to prove the mixing time. But how do we prove the front propagation result? This is where the comparison technique comes in. Instead of studying the spectral gap of the East process, we compare it to a somewhat simpler process - the one-sided/oriented contact process.
2. Explain how basic and oriented contact process works.
3. Point out that cross always implies death.

One-sided contact process



The one-sided contact process is similar to the East process, but 'recovery' is unconstrained.

1. So I explained how the front propagation result can be used to prove the mixing time. But how do we prove the front propagation result? This is where the comparison technique comes in. Instead of studying the spectral gap of the East process, we compare it to a somewhat simpler process - the one-sided/oriented contact process.
2. Explain how basic and oriented contact process works.
3. Point out that cross always implies death.

One-sided contact process

The one-sided contact process is similar to the East process, but 'recovery' is unconstrained.




Figure: One-sided contact process started from $\{0\}$

1. So I explained how the front propagation result can be used to prove the mixing time. But how do we prove the front propagation result? This is where the comparison technique comes in. Instead of studying the spectral gap of the East process, we compare it to a somewhat simpler process - the one-sided/oriented contact process.
2. Explain how basic and oriented contact process works.
3. Point out that cross always implies death.

1. How do we actually compare the two?
2. Basic coupling
3. Partial order and how its preserved
4. Front
5. Critical value and why its important

Same graphical structure for East and contact processes \Rightarrow
'basic coupling'.

1. How do we actually compare the two?
2. Basic coupling
3. Partial order and how its preserved
4. Front
5. Critical value and why its important

Same graphical structure for East and contact processes \Rightarrow
'basic coupling'.

Let $(\eta_t)_{t \geq 0}$ be the one-sided contact process started from $\{0\}$.
Then under the basic coupling

$$\eta_t \leq \sigma_t \quad \forall t \geq 0.$$

1. How do we actually compare the two?
2. Basic coupling
3. Partial order and how its preserved
4. Front
5. Critical value and why its important

Same graphical structure for East and contact processes \Rightarrow
'basic coupling'.

Let $(\eta_t)_{t \geq 0}$ be the one-sided contact process started from $\{0\}$.
Then under the basic coupling

$$\eta_t \leq \sigma_t \quad \forall t \geq 0.$$

In particular,

$$X(\eta_t) \leq X(\sigma_t) \quad \forall t \geq 0.$$

1. How do we actually compare the two?
2. Basic coupling
3. Partial order and how its preserved
4. Front
5. Critical value and why its important

Same graphical structure for East and contact processes \Rightarrow
'basic coupling'.

Let $(\eta_t)_{t \geq 0}$ be the one-sided contact process started from $\{0\}$.
Then under the basic coupling

$$\eta_t \leq \sigma_t \quad \forall t \geq 0.$$

In particular,

$$X(\eta_t) \leq X(\sigma_t) \quad \forall t \geq 0.$$

For this to be useful, we need $p > p_c$ where
 $p_c := \sup\{p : |\eta| \rightarrow 0 \text{ a.s.}\}$. From here on consider only such p .

1. How do we actually compare the two?
2. Basic coupling
3. Partial order and how its preserved
4. Front
5. Critical value and why its important

1. There are some results that we need for the oriented contact process.
2. Extinction time
3. Read theorem and explain
4. These results were never written down or proven in detail before.

Results for one-sided contact process



Let $\tau(\eta_*)$ be the extinction time. Then

$$\mathbb{P}(\tau(\eta_*) = \infty) > 0 \quad \forall p > p_c.$$

1. There are some results that we need for the oriented contact process.
2. Extinction time
3. Read theorem and explain
4. These results were never written down or proven in detail before.

Let $\tau(\eta_.)$ be the extinction time. Then

$$\mathbb{P}(\tau(\eta_.) = \infty) > 0 \quad \forall p > p_c.$$

Theorem

If $p > p_c$, there exists $\alpha > 0$ such that for all $a < \alpha$ there exist constants $\gamma, C > 0$ such that

$$\mathbb{P}(X(\eta_t) < at \mid \tau(\eta_.) = \infty) \leq Ce^{-\gamma t} \quad \forall t \geq 0.$$

Furthermore there exist constants $\gamma, C > 0$ such that

$$\mathbb{P}(t < \tau(\eta_.) < \infty) \leq Ce^{-\gamma t} \quad \forall t \geq 0.$$

1. There are some results that we need for the oriented contact process.
2. Extinction time
3. Read theorem and explain
4. These results were never written down or proven in detail before.

Let $\tau(\eta_*)$ be the extinction time. Then

$$\mathbb{P}(\tau(\eta_*) = \infty) > 0 \quad \forall p > p_c.$$

Theorem

If $p > p_c$, there exists $\alpha > 0$ such that for all $a < \alpha$ there exist constants $\gamma, C > 0$ such that

$$\mathbb{P}(X(\eta_t) < at \mid \tau(\eta_*) = \infty) \leq Ce^{-\gamma t} \quad \forall t \geq 0.$$

Furthermore there exist constants $\gamma, C > 0$ such that

$$\mathbb{P}(t < \tau(\eta_*) < \infty) \leq Ce^{-\gamma t} \quad \forall t \geq 0.$$

Remark

Hinted at in Durrett and Griffeath, 1983 but never proved in detail.

1. There are some results that we need for the oriented contact process.
2. Extinction time
3. Read theorem and explain
4. These results were never written down or proven in detail before.




Figure: Red path showing how oriented percolation to infinity occurs.

- Blondel, Oriane (2013). “Front progression in the East model”. In: *Stochastic Processes and their Applications* 123.9, pp. 3430–3465.
- Cancrini, Nicoletta et al. (2008). “Kinetically constrained spin models”. In: *Probability Theory and Related Fields* 140.3-4, pp. 459–504.
- Durrett, Richard and David Griffeath (1983). “Supercritical contact processes on \mathbb{Z} ”. In: *The Annals of Probability*, pp. 1–15.
- Peres, Yuval and Perla Sousi (2015). “Mixing times are hitting times of large sets”. In: *Journal of Theoretical Probability* 28.2, pp. 488–519.