



LONDON  
MATHEMATICAL  
SOCIETY  
EST. 1865



DEPARTMENT OF  
**STATISTICS**

# The oriented contact process and the East model

Patrik Gerber supervised by Prof. Paul Chleboun

Supported by LMS and the Mathematical Institute, University of Oxford.

Department of Statistics  
University of Oxford

4 October 2018

East process

Front propagation

Mixing time

One-sided contact process

East vs contact

Results for one-sided contact process

Percolation construction

References



---

The East process is a continuous time Markov chain

The East process is a continuous time Markov chain .

- ▶ State space:  $\Omega := \{0, 1\}^{\mathbb{Z}}$ , or equivalently,  $\mathcal{P}(\mathbb{Z})$ .

The East process is a continuous time Markov chain .

- ▶ State space:  $\Omega := \{0, 1\}^{\mathbb{Z}}$ , or equivalently,  $\mathcal{P}(\mathbb{Z})$ .
- ▶ Each site  $x \in \mathbb{Z}$  updates with rate 1 to  $B \sim \text{Ber}(p)$ ,  $p \in (0, 1)$   
iff  $x - 1$  is occupied.

The East process is a continuous time Markov chain .

- ▶ State space:  $\Omega := \{0, 1\}^{\mathbb{Z}}$ , or equivalently,  $\wp(\mathbb{Z})$ .
- ▶ Each site  $x \in \mathbb{Z}$  updates with rate 1 to  $B \sim \text{Ber}(p)$ ,  $p \in (0, 1)$   
iff  $x - 1$  is occupied.




Figure: East process started from  $\{0\}$





## Definition (Front)

Let  $(\sigma_t)_{t \geq 0}$  be an East process started from  $\{0\}$ . The front of  $(\sigma_t)_{t \geq 0}$  is the rightmost occupied site, written  $X(\sigma_t) := \sup \sigma_t$ .

## Definition (Front)

Let  $(\sigma_t)_{t \geq 0}$  be an East process started from  $\{0\}$ . The front of  $(\sigma_t)_{t \geq 0}$  is the rightmost occupied site, written  $X(\sigma_t) := \sup \sigma_t$ .

## Theorem (Blondel, 2013, Lemma 3.2)

*There exist constants  $0 < \alpha < v$  and  $\gamma, C > 0$  such that*

$$\mathbb{P}(X(\sigma_t) \in (\alpha t, vt)) \geq 1 - Ce^{-\gamma t} \quad \forall t \geq 0.$$

## Definition (Front)

Let  $(\sigma_t)_{t \geq 0}$  be an East process started from  $\{0\}$ . The front of  $(\sigma_t)_{t \geq 0}$  is the rightmost occupied site, written  $X(\sigma_t) := \sup \sigma_t$ .

## Theorem (Blondel, 2013, Lemma 3.2)

*There exist constants  $0 < \alpha < v$  and  $\gamma, C > 0$  such that*

$$\mathbb{P}(X(\sigma_t) \in (\alpha t, vt)) \geq 1 - Ce^{-\gamma t} \quad \forall t \geq 0.$$

## Proof.

We propose an alternative proof (under restrictions on  $p$ ) detailed later. □



Restricted to  $\{0, 1, \dots, L\}$  with a fixed 1 at origin East process converges to equilibrium.

Restricted to  $\{0, 1, \dots, L\}$  with a fixed 1 at origin East process converges to equilibrium.

Theorem (Cancrini et al., 2008, Theorem 6.1)

*The mixing time of the East process on  $\{0, 1, \dots, L\}$  with a fixed 1 at the origin is  $\Theta(L)$ .*

Restricted to  $\{0, 1, \dots, L\}$  with a fixed 1 at origin East process converges to equilibrium.

Theorem (Cancrini et al., 2008, Theorem 6.1)

*The mixing time of the East process on  $\{0, 1, \dots, L\}$  with a fixed 1 at the origin is  $\Theta(L)$ .*

Alternative proof using only front propagation.

Upper bound: Coupling time vs. propagation of the front.

Lower bound: Hitting times of large sets (Peres and Sousi, 2015).



# One-sided contact process

---



# One-sided contact process

The one-sided contact process is similar to the East process, but 'recovery' is unconstrained.

# One-sided contact process

The one-sided contact process is similar to the East process, but 'recovery' is unconstrained.




Figure: One-sided contact process started from  $\{0\}$



Same graphical structure for East and contact processes  $\Rightarrow$  'basic coupling'.

Same graphical structure for East and contact processes  $\Rightarrow$  'basic coupling'.

Under the basic coupling

$$(\text{Oriented contact process}) \quad \eta_t \leq \sigma_t \quad (\text{East process}) \quad \forall t \geq 0.$$

Same graphical structure for East and contact processes  $\Rightarrow$  'basic coupling'.

Under the basic coupling

(Oriented contact process)  $\eta_t \leq \sigma_t$  (East process)  $\forall t \geq 0$ .

In particular,

$$X(\eta_t) \leq X(\sigma_t) \quad \forall t \geq 0.$$

Same graphical structure for East and contact processes  $\Rightarrow$  'basic coupling'.

Under the basic coupling

$$(\text{Oriented contact process}) \quad \eta_t \leq \sigma_t \quad (\text{East process}) \quad \forall t \geq 0.$$

In particular,

$$X(\eta_t) \leq X(\sigma_t) \quad \forall t \geq 0.$$

For this to be useful, we need  $p > p_c$  where  $p_c := \sup\{p : |\eta| \rightarrow 0 \text{ a.s.}\}$ . From here on consider only such  $p$ .

# Results for one-sided contact process

---



# Results for one-sided contact process

Let  $\tau(\eta_.)$  be the extinction time. Then

$$\mathbb{P}(\tau(\eta_.) = \infty) > 0 \qquad \forall p > p_c.$$

# Results for one-sided contact process

Let  $\tau(\eta_.)$  be the extinction time. Then

$$\mathbb{P}(\tau(\eta_.) = \infty) > 0 \quad \forall p > p_c.$$

## Theorem

*If  $p > p_c$ , there exists  $\alpha > 0$  such that for all  $a < \alpha$  there exist constants  $\gamma, C > 0$  such that*

$$\mathbb{P}(X(\eta_t) < at \mid \tau(\eta_.) = \infty) \leq Ce^{-\gamma t} \quad \forall t \geq 0.$$

*Furthermore there exist constants  $\gamma, C > 0$  such that*

$$\mathbb{P}(t < \tau(\eta_.) < \infty) \leq Ce^{-\gamma t} \quad \forall t \geq 0.$$

# Results for one-sided contact process

Let  $\tau(\eta_.)$  be the extinction time. Then

$$\mathbb{P}(\tau(\eta_.) = \infty) > 0 \quad \forall p > p_c.$$

## Theorem

*If  $p > p_c$ , there exists  $\alpha > 0$  such that for all  $a < \alpha$  there exist constants  $\gamma, C > 0$  such that*

$$\mathbb{P}(X(\eta_t) < at \mid \tau(\eta_.) = \infty) \leq Ce^{-\gamma t} \quad \forall t \geq 0.$$

*Furthermore there exist constants  $\gamma, C > 0$  such that*

$$\mathbb{P}(t < \tau(\eta_.) < \infty) \leq Ce^{-\gamma t} \quad \forall t \geq 0.$$

## Remark

*Hinted at in Durrett and Griffeath, 1983 but never proved in detail.*

# Percolation construction 1

---

## Definition

$\{(z(r), r)\}_{r \in [t, s]} \subset \mathbb{Z} \times \mathbb{R}_{\geq 0}$  is a *path* if it follows arrows and goes through no cross.

## Definition

$\{(z(r), r)\}_{r \in [t, s]} \subset \mathbb{Z} \times \mathbb{R}_{\geq 0}$  is a *path* if it follows arrows and goes through no cross.





Figure: Red and blue lines are examples of a path.

# Percolation construction 2

---



**Figure:** Red path showing how oriented percolation to infinity occurs.



- Blondel, Oriane (2013). “Front progression in the East model”. In: *Stochastic Processes and their Applications* 123.9, pp. 3430–3465.
- Cancrini, Nicoletta et al. (2008). “Kinetically constrained spin models”. In: *Probability Theory and Related Fields* 140.3-4, pp. 459–504.
- Durrett, Richard and David Griffeath (1983). “Supercritical contact processes on  $\mathbb{Z}$ ”. In: *The Annals of Probability*, pp. 1–15.
- Peres, Yuval and Perla Sousi (2015). “Mixing times are hitting times of large sets”. In: *Journal of Theoretical Probability* 28.2, pp. 488–519.