

## Graphical Abstract

### **Collective sedimentation from buoyant, particle-laden gravity currents**

Paul A. Jarvis, Allan Fries, Jonathan Lemus, Costanza Bonadonna, Amanda Clarke, Irene Manzella, Jeremy Phillips

## Highlights

### **Collective sedimentation from buoyant, particle-laden gravity currents**

Paul A. Jarvis, Allan Fries, Jonathan Lemus, Costanza Bonadonna, Amanda Clarke, Irene Manzella, Jeremy Phillips

- Research highlight 1
- Research highlight 2

# Collective sedimentation from buoyant, particle-laden gravity currents

Paul A. Jarvis<sup>a,\*</sup>, Allan Fries<sup>a</sup>, Jonathan Lemus<sup>a</sup>, Costanza Bonadonna<sup>a</sup>,  
Amanda Clarke<sup>b</sup>, Irene Manzella<sup>c</sup>, Jeremy Phillips<sup>d</sup>

*<sup>a</sup>Department of Earth Sciences, University of Geneva, Rue des Marichers, Geneva, 1205, Switzerland*

*<sup>b</sup>School of Earth and Space Exploration, Arizona State University, ISTB4-BLDG75, 781 E Terrance Mall, Tempe, AZ, 85287-6004, USA*

*<sup>c</sup>School of Geography, Earth and Environmental Sciences, University of Plymouth, Drake Circus, Plymouth, PL4 8AA, UK*

*<sup>d</sup>School of Earth Sciences, University of Bristol, Wills Memorial Building, Queens Road, Bristol, BS8 1RJ, UK*

---

## Abstract

*Keywords:*

---

## 1. Introduction

## 2. Methods

Experiments were performed in a perspex flume of internal length 353 cm, width  $(12.2 \pm 0.5)$  cm and depth 50 cm. The uncertainty on the width is due to bowing of the tank walls. During the experimental setup, two removable gates can be placed at 24 and 53 cm, respectively, from the left-hand end, creating three sections. The left-most section takes no part in the experiment, whilst the short middle section is where the particle suspension is prepared, and is referred to as the gated section (length 27 cm). The remaining length of the flume is called the environment (length 3m).

---

\*Corresponding author: paul.jarvis@unige.ch

The flume sits within a metal framework. Three vertical supports, each 3cm thick, at distances of 87 cm, 173.5 cm and 260 cm from the left-hand end prevent bowing. This effectively separates the flume into four, nearly equal, sections. Behind each section a backing board is placed. For experiments with no particles, red or blue food colouring is added to the current and the backing board is white. Otherwise the boards are black. The top 5 cm of each board is a row of  $(5 \times 5)$  cm<sup>2</sup> squares, alternating in colour from black-to-white. Meanwhile, at the base of the tank, tape is used to create a similar scale.

The day before an experiment, the flume is filled up to a depth of 47.5 cm. Separately, the desired mass of sugar is completely dissolved in approximately 15 l of water. The flume and sugar solution are then both allowed to equilibrate to room temperature overnight. The next day, the flume is imaged (single frame) in its current configuration. In some experiments, the whole flume was imaged using a Canon Legria HFG40. In others, this camera was positioned closer to the tank, but only imaged the left-hand half of the flume, whilst the right-hand half was imaged using a Nikon D700 (DSLR). The captured frame(s) are used as reference images using the top(back) and bottom(front) scales. By capturing the image with the back scale partially submerged, it is possible to correct position measurements for distortion due to the refractive index (RI) of the water.

The flume water level is then lowered until it is of a depth of approximately 36.5 cm. Gates 1 and 2 are then put in place. The sugar solution is then added to the environment section. The gated section is then topped up with water until the water depth there is the same as the environment.

This procedure results in a water depth of approximately 40 cm. The environment section is rigorously stirred. The RI of fluid from four positions in the environment section (top and bottom, near and far from the gate) is measured using a refractometer to ensure a uniform density. The RI of the gated section is also measured to check that there has been no significant leakage of sugar solution through the gate. A calibrated digital thermometer is used to measure the temperature of both sections. In all experiments, the maximum difference in temperature between the sections was 0.1 °C.

Recording of the experiment then begins. The required mass of particles is added to the gated section which is thoroughly stirred to ensure a uniform particle concentration. Finally, gate 2 is removed and the particle suspension spreads along the free surface as a buoyant gravity current. Once the current head reaches the end of the tank, gate 2 is returned to its position and recording stops.

### **3. Results**

### **4. Discussion**

### **5. Conclusions**

### **References**