

# Extensions of the Axiom of Determinacy

Paul B. Larson

May 18, 2022



# Contents

0.1	Introduction . . . . .	5
0.2	Notation . . . . .	8
0.3	Prerequisites . . . . .	9
0.4	Forms of Choice . . . . .	10
0.5	Partial orders . . . . .	10
<b>I</b>	<b>Preliminaries</b>	<b>13</b>
<b>1</b>	<b>Determinacy</b>	<b>15</b>
1.1	The Axiom of Determinacy . . . . .	15
1.2	Turing Determinacy . . . . .	17
<b>2</b>	<b>The Wadge hierarchy</b>	<b>21</b>
2.1	Lipschitz determinacy . . . . .	21
2.2	Lipschitz degrees and Wadge degrees . . . . .	24
2.3	Pointclasses . . . . .	28
2.4	Universal sets . . . . .	31
2.5	The cardinal $\Theta$ . . . . .	36
<b>3</b>	<b>Coding lemmas</b>	<b>39</b>
<b>4</b>	<b>Properties of pointclasses</b>	<b>45</b>
4.1	Separation and reduction . . . . .	45
4.2	The prewellordering property . . . . .	50
4.3	Prewellorderings and wellfounded relations . . . . .	55
4.4	Closure under wellordered unions . . . . .	57
<b>5</b>	<b>Strong partition cardinals</b>	<b>61</b>
<b>6</b>	<b>Suslin sets and Uniformization</b>	<b>69</b>
6.1	Suslin sets . . . . .	69
6.2	Uniformization . . . . .	75
6.3	The Solovay sequence . . . . .	77

<b>II</b>	<b><math>\text{AD}^+</math></b>	<b>81</b>
<b>7</b>	<b><math>&lt;\Theta</math>-Determinacy</b>	<b>83</b>
<b>8</b>	<b><math>\infty</math>-Borel sets</b>	<b>89</b>
8.1	$\infty$ -Borel codes . . . . .	89
8.2	Local $\infty$ -Borel codes . . . . .	96
<b>9</b>	<b>Cone measure ultrapowers</b>	<b>101</b>
9.1	$S$ -cones . . . . .	102
9.2	Measurable cardinals from cone measures . . . . .	105
9.3	The degree order on sets of ordinals . . . . .	107
9.4	Pointed trees . . . . .	111
9.5	Coding ultrapowers . . . . .	112
9.6	Forcing with positive sets . . . . .	114
<b>10</b>	<b>Vopěnka algebras</b>	<b>119</b>
10.1	The Vopěnka algebra . . . . .	119
10.2	Codes for projections, and Uniformization . . . . .	125
10.3	The Vopěnka algebra for $\infty$ -Borel sets . . . . .	129
<b>11</b>	<b>Suslin sets and strong codes</b>	<b>135</b>
11.1	Generic codes . . . . .	135
11.2	Producing strong generic codes . . . . .	139
11.3	$\infty$ -Borel representations from Uniformization . . . . .	142
11.4	Closure of the Suslin cardinals . . . . .	145
<b>12</b>	<b>Scales from Uniformization</b>	<b>149</b>
12.1	Ordinal determinacy in the codes . . . . .	149
12.2	Boundedness for $\infty$ -Borel relations . . . . .	156
12.3	Becker's argument . . . . .	157
<b>13</b>	<b><math>\text{AD}_{\mathbb{R}}</math> from scales</b>	<b>165</b>
13.1	Weakly homogeneous trees . . . . .	166
13.2	Normal measures on $\mathcal{P}_{\aleph_1}(\lambda)$ . . . . .	171
13.3	$\text{AD}$ implies $\text{AD}_{\mathbb{R}}$ if all sets of reals are Suslin . . . . .	174
<b>14</b>	<b>Questions</b>	<b>179</b>

This draft includes notes to myself, in footnotes. Aside from these, the book should be close to its final form. Comments and corrections are more than welcome.

## 0.1 Introduction

The Axiom of Determinacy (AD) is the statement that all length- $\omega$  integer games of perfect information are determined. The beginning of Chapter 1 contains a more precise definition, but we expect the reader to be familiar with the classical theory of determinacy, as found in [8, 10], for instance. The axiom  $\text{AD}^+$  is a generalization, due to W. Hugh Woodin, of the Axiom of Determinacy. We define it as the conjunction of three statements.

**0.1.1 Definition.** The axiom  $\text{AD}^+$  is the conjunction of the following three statements.

1.  $\text{DC}_{\mathbb{R}}$
2. Every subset of  $\omega^\omega$  is  $\infty$ -Borel.
3. ( $<\Theta$ -Determinacy) For all  $\lambda < \Theta$ , every  $A \subseteq \omega^\omega$  and every continuous function  $\pi: {}^\omega\lambda \rightarrow \omega^\omega$ ,  $\pi^{-1}(A)$  is determined (as a subset of  ${}^\omega\lambda$ ).

Our definition of  $\text{AD}^+$  diverges from Woodin's own terminology, as he defines  $\text{AD}^+$  to be the conjunction of items (2) and (3) above. As  $\text{AD}^+$  is usually considered only in the context of  $\text{DC}_{\mathbb{R}}$  the choice of definition does not make a significant difference. One difference between  $\text{DC}_{\mathbb{R}}$  and the other two parts of  $\text{AD}^+$  as we have defined it is that  $\text{DC}_{\mathbb{R}}$  is witnessed by real numbers, so, assuming the Axiom of Choice,  $\text{DC}_{\mathbb{R}}$  holds in any inner model containing the reals. The other two parts of  $\text{AD}^+$  however are witnessed by bounded subsets of  $\Theta$ .

The  $\infty$ -Borel sets are defined in Definition 8.1.1 for subsets of  $2^\omega$ , and in Remark 8.1.9 for subsets of  $\omega^\omega$ . We define  $<\Theta$ -Determinacy (which Woodin calls Ordinal Determinacy) in Chapter 7. The restriction of  $<\Theta$ -Determinacy to the case where  $\pi$  is the identity function on  $\omega^\omega$  is exactly AD.

It is an open question whether AD implies any or all of the parts of  $\text{AD}^+$ , and in fact whether AD plus any two parts of  $\text{AD}^+$  imply the third. It also open whether  $\text{AD}_{\mathbb{R}}$  implies  $\text{AD}^+$ . The issue in this case is whether  $\text{AD}_{\mathbb{R}}$  implies  $<\Theta$ -Determinacy, as  $\text{DC}_{\mathbb{R}}$  is easily seen to follow from  $\text{AD}_{\mathbb{R}}$ , and the second part of  $\text{AD}^+$  follows from  $\text{AD}_{\mathbb{R}}$  (moreover, from  $\text{AD} + \text{Uniformization}$ ) by Theorem 11.3.4.

If  $M \subseteq N$  are models of AD with the same reals, and every set of reals in  $M$  is Suslin in  $N$ , then  $M \models \text{AD}^+$  (this is Theorem 8.2.8). In fact, this is the context which the axiom  $\text{AD}^+$  was designed to describe; its original name was “within scales” (a scale being a certain means of coding a Suslin representation). Since (assuming AD) the set of  $A \subseteq \omega^\omega$  for which  $L(A, \mathbb{R}) \models \text{AD}^+$  is a nonempty

initial segment of the Wadge hierarchy (see the list below),  $\text{AD}^+$  is also useful for a bottom-up analysis of models of determinacy.

Here is a list of the major results of the book (all due to Woodin, except where noted):

- In  $L(\mathbb{R})$ ,  $\text{AD}$  implies  $\text{AD}^+$ . This is Corollary 8.2.6.
- If  $\text{AD}$  holds and every true  $\Sigma_1^2$  sentence is witnessed by a Suslin, co-Suslin set, then  $\text{AD}^+$  holds. This is Theorem 8.2.7.
- If  $\text{AD}^+$  holds then every inner model containing the reals satisfies  $\text{AD}^+$ . This is Theorem 8.2.8.
- If  $\text{AD}$  holds then for all  $A \subseteq \omega^\omega$ , either  $\mathcal{P}(\omega^\omega) \subseteq L(A, \mathbb{R})$  or  $A^\#$  exists. This is Theorem 10.1.8.
- Assuming  $\text{AD} + \text{DC}_{\mathbb{R}}$ , a subset of  $\omega^\omega$  is  $\infty$ -Borel if and only if there is a set of ordinals  $S$  such that  $A \in L(S, \mathbb{R})$ . This follows from part (1) of Corollary 10.3.7.
- Assuming  $\text{AD}$ , if  $A \subseteq \omega^\omega$  is  $\infty$ -Borel, then so is every set of reals in  $L(A, \mathbb{R})$ . See Remark 10.3.9.
- Assuming  $\text{AD}^+$ ,  $\text{AD}_{\mathbb{R}}$  is equivalent to the assertion that the Solovay sequence has limit length. This is Theorem 11.2.4.
- If  $<\Theta$ -Determinacy and Uniformization hold, then every subset of  $\omega^\omega$  is Suslin. This follows from Theorem 11.3.4, Corollary 10.2.7 and Theorem 11.2.1.
- If  $\text{AD}^+$  holds then the set of Suslin cardinals is closed below  $\Theta$ . This follows from Theorem 11.4.1.
- If  $\text{AD}$  holds, then the set of Suslin cardinals is closed below its supremum. This is Theorem ?? (joint with John Steel).
- If  $\text{AD}^+$  holds, then  $\text{AD}_{\mathbb{R}}$  fails if and only if there is a set of ordinals  $T$  such that  $L(\mathcal{P}(\mathbb{R})) = L(T, \mathbb{R})$ . This is Theorem 13.3.3.

In addition, we prove the follow theorem illustrating the relationships among  $\text{AD}$ ,  $\text{AD}_{\mathbb{R}}$ , the Suslin property and Uniformization. The first sentence of the theorem is due to Woodin, aside from the implication from (1) to (3), which is due independently to Donald Martin. The implication from (4) to (1) under  $\text{DC}$  combines results of Woodin and Becker (see Chapter 12).

**Theorem 0.1.2.** *Each of the following statements implies the ones below it, and the first two statements are equivalent. If  $\text{DC}$  holds, then all four statements are equivalent.*

1.  $\text{AD} + \text{“every subset of } \omega^\omega \text{ is Suslin”}$

2.  $\text{AD}^+$  + “every subset of  $\omega^\omega$  is Suslin”
3.  $\text{AD}_{\mathbb{R}}$
4.  $\text{AD}$  + Uniformization

*Proof.* Since  $\text{AD}$  is the restriction of  $<\Theta$ -Determinacy to the case  $\lambda = \omega$ , (2) implies (1). Theorem 6.2.1 says that Suslin sets can be uniformized, and Remark 6.2.2 shows that Uniformization implies  $\text{DC}_{\mathbb{R}}$ . The implication from (1) to (2) then follows from Theorems 5.0.4 and 7.0.1, and Remark 6.1.7. Theorem 13.0.1 says that (1) implies (3). The implication from (3) to (4) is covered in Remark 6.2.2. That (4) implies (1) under  $\text{DC}$  is Theorem 12.3.1.  $\square$

Part I of the book collects various facts (primarily from the Cabal seminar) which will be needed in Part II. It is not intended as a comprehensive introduction to the topics discussed there (for which, see [32, 15, 16, 17]). Part I also significantly overlaps [5]. A natural way to read the book might be to start at the beginning of Part II and refer back to Part I as needed.

The material in this book covers the development of the theory of  $\text{AD}^+$  up to some point in the early 1990’s. We plan to continue the story in future volumes, starting with a book on derived models and reversals. In particular, the following results of Woodin are not included in this book, although we hope to prove them in the next volume.

- $\text{AD}^+$  implies  $\Sigma_1$  reflection into the Suslin, co-Suslin sets.
- $\text{AD}^+$  implies that the ultrapower of  $V$  by the Turing measure is well-founded.
- Suppose that  $\text{AD}$  holds, and  $\text{AD}^+$  fails, and let  $\Gamma$  be the pointclass of Suslin, co-Suslin sets. Then  $L(\Gamma, \mathbb{R})$  satisfies  $\text{AD}_{\mathbb{R}}$  and contains every  $A \subseteq \omega^\omega$  for which  $L(A, \mathbb{R}) \models \text{AD}^+$ , and  $\mathcal{P}(\omega^\omega) \cap L(\Gamma, \mathbb{R}) \subseteq \Gamma$ .
- If  $\text{AD}$  holds and there is a largest Suslin cardinal, then  $\text{AD}^+$  holds.
- If  $\text{AD}$  holds, and  $A \subseteq \omega^\omega$  is Suslin, then  $L(A, \mathbb{R}) \models \text{AD}^+$ .
- $\text{AD} + \text{DC}_{\mathbb{R}}$  implies that for all ordinals  $\chi$ , every subset of  $\omega^\omega$  in  $L(\mathcal{P}(\chi))$  is  $\infty$ -Borel.

The statement that the ultrapower of the  $V$  by the Turing measure is well-founded was originally one of the axioms of  $\text{AD}^+$ , before Woodin proved that it follows from the other axioms. This statement is easily seen to follow from the assumption that Turing Determinacy and  $\text{DC}$  hold, which in turn follows from the assumption that Turing Determinacy and  $\text{DC}_{\mathbb{R}}$  hold, along with the assumption that there is some set  $X$  with the property that every set is definable from  $X$ , a real and an ordinal. This is the approach taken in this book, although in some instances we include the wellfoundedness of the Turing ultrapower (or some other related ultrapower) as an explicit assumption.

This book has a great deal of overlap with many sources, notably [6, 21, 22].

The writing of the book was supported by NSF grants DMS-1201494 and DMS-1764320. The author thanks the Mathematics Department at Harvard University for its hospitality during the Fall semester of 2021, when a significant portion of this book was written.

## 0.2 Notation

We reserve the symbol  $\mathbb{R}$  for the real line, which is never used directly. However, we use traditional notation such as  $\text{AD}_{\mathbb{R}}$ ,  $\text{DC}_{\mathbb{R}}$ ,  $L(\mathbb{R})$  and so on, as these terms are equivalent to more relevant forms such as  $\text{AD}_{\omega^\omega}$ ,  $\text{DC}_{\omega^\omega}$  and  $L(\omega^\omega)$ . We informally use the word “real” to mean an element of the Baire space  $\omega^\omega$ .

We use the following notational conventions.

- We write  $\text{Ord}$  for the class of ordinals.
- We write  $X^Y$  to mean the set of functions from  $Y$  to  $X$ , and, when  $\gamma$  is an ordinal  $X^{<\gamma}$  to mean  $\bigcup_{\alpha \in \gamma} X^\alpha$ .
- For an ordinal  $\alpha$  and a set of ordinals  $X$ ,  $[X]^\alpha$  denotes the collection of subsets of  $X$  of ordertype  $\alpha$ , and  $[X]^{<\alpha}$  denotes the collection of subsets of  $X$  of ordertype less than  $\alpha$ .
- Given a set  $X$ , we write  $\exists^X$  and  $\forall^X$  for existential and universal quantification over  $X$ , respectively.
- A *preorder* on a set is a binary relation which is reflexive and transitive. A preorder is *wellfounded* if every nonempty subset of its domain has a minimal element.
- If  $\leq$  is a wellfounded preorder on a set  $X$ , the *canonical rank function*  $\text{rank}_\leq$  associated to  $\leq$  assigns to each  $x \in X$  the least ordinal  $\alpha$  such that  $\alpha > \text{rank}_\leq(y)$  for all  $y \in X$  such that  $y \leq x$  and  $x \not\leq y$ .
- In any preorder, especially the natural order on the ordinals, the *strict supremum* of a set (if it exists) is the least element in the domain of the order which is strictly greater than every element of the set.
- Formally, a partial order  $\mathbb{P}$  is a pair  $(\text{dom}(\mathbb{P}), \leq_{\mathbb{P}})$ , where  $\text{dom}(\mathbb{P})$  is the domain or underlying set of  $\mathbb{P}$ , and  $\leq_{\mathbb{P}}$  is a partial order on  $\text{dom}(\mathbb{P})$ .
- Given a partial order  $\mathbb{P}$ , a *nice*  $\mathbb{P}$ -name for an element of  $\omega^\omega$  is a set  $\tau$  such that
  - each element of  $\tau$  is a pair  $(p, (n, m))$ , where  $p \in \mathbb{P}$  and  $n, m \in \omega$  (formally we should write  $(p, \tilde{x})$ , where  $\tilde{x}$  is the pair  $(n, m)$ );
  - for each  $p \in \mathbb{P}$  and each  $n \in \omega$ , there exist  $q \leq p$  and  $m \in \omega$  such that  $(q, (n, m)) \in \tau$ ;



- for all  $p, p' \in \mathbb{P}$  and all  $n, m, m' \in \omega$ , if  $(p, (n, m))$  and  $(p', (n, m'))$  are both in  $\tau$ , and  $p$  and  $p'$  are compatible, then  $m = m'$ .
- Given a nice  $\mathbb{P}$ -name  $\tau$  and a set  $g \subseteq \mathbb{P}$ , we let  $\tau_g$  be the set of pairs  $(n, m)$  for which there exists a  $p \in g$  with  $(p, (n, m)) \in \tau$ .
- Given a partial order  $\mathbb{P}$  and a set (or class)  $B$ , we say that a set  $g \subseteq \mathbb{P}$  is *B-generic* if  $g$  intersects every dense open subset of  $\mathbb{P}$  in  $B$ .
- We let  $\leq_{\text{Gö}}$  be the Gödel order, i.e., the order on pairs of ordinals defined as follows :  $(\alpha, \beta) \leq_{\text{Gö}} (\delta, \gamma)$  if any of the following holds:
  - $\max\{\alpha, \beta\} < \max\{\delta, \gamma\}$ ;
  - $\max\{\alpha, \beta\} = \max\{\delta, \gamma\}$  and  $\alpha < \delta$ ;
  - $\max\{\alpha, \beta\} = \max\{\delta, \gamma\}$ ,  $\alpha = \delta$  and  $\beta \leq \gamma$ .
- Given ordinals  $\alpha$  and  $\beta$ , we write  $\prec \alpha, \beta \succ$  for the ordinal ordertype of the set of predecessors of  $(\alpha, \beta)$  in the order  $\leq_{\text{Gö}}$ .
- For a set  $X$ , we let  $\text{TC}(X)$  denote the transitive closure of  $X$ .
- Given a cardinal  $\kappa$ ,  $H(\kappa)$  denotes the collection of sets of hereditary cardinality less than  $\kappa$ , i.e., the set of  $x$  for which  $|\text{TC}(x)| < \kappa$ .

We sometimes write HF for  $H(\aleph_0)$ . The members of HF are said to be *hereditarily finite*. A set  $x \subseteq \text{HF}$  is *semi-recursive* if  $x$  is  $\Sigma_1$ -definable over HF, and *recursive* if  $x$  and  $\text{HF} \setminus x$  are both semi-recursive. Given  $A, B \subseteq \text{HF}$ , we say that  $A$  is *Turing reducible* to  $B$ , and write  $A \leq_{\text{TU}} B$  if there is a semi-recursive function  $f : \text{HF} \rightarrow \text{HF}$  such that  $A = f^{-1}[B]$ . We say that  $A$  is *Turing equivalent* to  $B$  if  $A$  and  $B$  are Turing-reducible to each other. We write  $\leq_{\text{TU}}$  and  $\equiv_{\text{TU}}$  respectively for the restrictions of Turing reducibility and Turing equivalence to  $\omega^\omega$ .

We also sometimes write HC for  $H(\aleph_1)$ . The members of HC are said to be *hereditarily countable*. We say that a set  $x \in \omega^\omega$  *HC-codes* a set  $y \in \text{HC}$  if  $(\omega, \{(n, m) \in \omega \times \omega : x(2^n 3^m) = 0\})$  is isomorphic to  $(\text{TC}(\{y\}), \in)$ .

### 0.3 Prerequisites

We expect the reader to be familiar with Zermelo-Fraenkel set theory, Gödel numbering of formulas, relative constructibility, ordinal definability, inner models of the form  $\text{HOD}_X$ , ultrapowers, sharps and forcing, along with some elementary notions from descriptive set theory (e.g., Borel sets and projective sets). We do not expect the reader to be familiar with everything in the books [8, 24, 10, 19, 32, 35], but they make good references.

## 0.4 Forms of Choice

Our base theory in this book is Zermelo-Fraenkel set theory (ZF). Additional axioms will be stated as used. Although we will sometimes consider models of the Axiom of Choice (AC), our main interest is in models of the Axiom of Determinacy (AD), which contradicts AC. Weak forms of AC can (and do) hold in models of AD, however.

Given a set  $X$ , the principle of **Dependent Choice** for  $X$  ( $\text{DC}_X$ ) is the statement that whenever  $T \subseteq {}^{<\omega}X$  is a nonempty tree (i.e., a subset of  ${}^{<\omega}X$  closed under initial segments) with the property that every element of  $T$  has a proper extension in  $T$ , there exists an infinite path through  $T$ . The principle of **Countable Choice** for  $X$  ( $\text{CC}_X$ ) is the statement that for all countable sets  $Y$ , if  $\langle A_y : y \in Y \rangle$  is a sequence of nonempty subsets of  $X$ , then there exists a function  $f : Y \rightarrow X$  such that  $f(y)$  is in  $A_y$ , for each  $y \in Y$ . The principle  $\text{CC}_X$  follows immediately from  $\text{DC}_X$ . A classical argument (due to Mycielski; see Remark 1.1.2) shows that  $\text{CC}_{\mathbb{R}}$  follows from AD. Whether or not  $\text{DC}_{\mathbb{R}}$  follows from AD is an open question. Note however that if  $\text{DC}_{\mathbb{R}}$  holds, then any inner model containing  $\mathcal{P}(\omega)$  satisfies  $\text{DC}_{\mathbb{R}}$ .

The following theorem is part of the result that, in  $L(\mathbb{R})$ , AD implies  $\text{AD}^+$ .

**Theorem 0.4.1** (Kechris [12]). *Assuming  $V=L(\mathbb{R})$ , AD implies  $\text{DC}_{\mathbb{R}}$ .*

The axiom of **Dependent Choice** (DC) asserts that  $\text{DC}_X$  holds for every set  $X$ . Similarly, the axiom of **Countable Choice** (CC) asserts that  $\text{CC}_X$  holds for every set  $X$ .

**0.4.2 Remark.** We make frequent use of the standard fact that if every set is definable from an ordinal and a member of  $X$ , then  $\text{DC}_X$  implies DC.

## 0.5 Partial orders

We list here some classical partial orders which are used as forcing notions throughout the book.

- $\text{Col}(\kappa, X)$ , where  $\kappa$  is an infinite cardinal and  $X$  is a set. Conditions are functions  $f : \alpha \rightarrow X$ , where  $\alpha < \kappa$ . The order is extension.
- $\text{Col}^*(\kappa, X)$ , where  $\kappa$  is an infinite cardinal and  $X$  is a set. Conditions are injective functions  $f : \alpha \rightarrow X$ , where  $\alpha < \kappa$ . The order is extension.
- Given a set  $X$  consisting of infinite subsets of  $\omega$ , the classical *almost-disjoint coding* forcing for  $X$  (due to Jensen and Solovay [9]) consists of pairs  $(a, B)$ , where  $a$  is a finite subset of  $\omega$  and  $B$  is a finite subset of  $X$ , with the order  $(a, B) \leq (c, D)$  if
  - $c$  is either the emptyset or  $a \cap (\max(c) + 1)$ ,
  - $D \subseteq B$  and

- $(a \setminus c) \cap r = \emptyset$  for all  $r \in D$ .

This partial order is c.c.c. and adds a subset of  $\omega$  having finite intersection with each member of  $X$  and infinite intersection with each element of  $\mathcal{P}(\omega)$  not contained mod-finite in the union of a finite subset of  $X$ .



**Part I**

**Preliminaries**



# Chapter 1

## Determinacy

In Chapter 1 we review the Axiom of Determinacy (AD) and the axiom of Turing Determinacy (TD). We expect that most readers will be familiar with the material in this chapter, and include it mostly for easy reference. Remark 1.1.2 below lists four classical consequences of AD which we will be using throughout the book. We make some original definitions in Section 1.2 to formalize the process of lifting the ultrafilter property from one cone measure to another.

### 1.1 The Axiom of Determinacy

Given a set  $X$ , a set  $A \subseteq X^\omega$  is *determined* (as a subset of  $X^\omega$ ) if there is a function  $\pi: X^{<\omega} \rightarrow X$  such that one of the two following statements holds.

1. For every  $x \in X^\omega$ , if  $x(2n) = \pi(x \upharpoonright 2n)$  holds for all  $n \in \omega$ , then  $x \in A$ .
2. For every  $x \in X^\omega$ , if  $x(2n+1) = \pi(x \upharpoonright (2n+1))$  hold for all  $n \in \omega$ , then  $x \notin A$ .

We let  $\text{AD}_X$  denote the statement that every subset of  $X^\omega$  is determined (as a subset of  $X^\omega$ ). The Axiom of Determinacy (AD) is the statement  $\text{AD}_\omega$ .

**1.1.1 Remark.** The following easily verifiable assertions (for arbitrary sets  $X$  and  $Y$ ) show that AD implies that  $\text{AD}_X$  holds for each countable set  $X$ .

- If there is an injection from  $X$  to  $Y$ , then  $\text{AD}_Y$  implies  $\text{AD}_X$ .
- If  $X$  is wellorderable and there is a surjection from  $X$  to  $Y$ , then  $\text{AD}_X$  implies  $\text{AD}_Y$ .
- If AD holds, then so does  $\text{AD}_X$  for each countable set  $X$ .

It is convenient and conventional to rephrase determinacy in terms of games. A set  $A \subseteq X^\omega$  corresponds to a game  $\mathcal{G}_A$  between players  $I$  and  $II$ , who alternate picking members of  $X$ , with  $I$  winning if and only if the induced member of  $X^\omega$  is in  $A$ . The set  $A$  is called the *payoff set* for  $\mathcal{G}_A$ .

I	$x(0)$	$x(2)$	$x(4)$	$\dots$
II	$x(1)$	$x(3)$	$\dots$	

A run of the game  $\mathcal{G}_A$ ;  $I$  wins if and only if  $x$  is in  $A$ .

A function  $\pi: X^{<\omega} \rightarrow X$  is then said to be a *strategy* in the game  $\mathcal{G}_A$ . A function  $\pi$  as in case (1) above is a *winning strategy* for player  $I$ ; in case (2) it is a winning strategy for player  $II$ . If  $\sigma$  is a strategy and  $x$  is in  $X^\omega$ , then we write  $\sigma * x$  for combined output of the two players when  $I$  plays according to  $\sigma$  and  $II$  plays  $x$ , that is, the unique  $y \in X^\omega$  such that

- $y(2n+1) = x(n)$  for all  $n \in \omega$ ;
- $y(n) = \sigma(y \upharpoonright n)$  for all even  $n \in \omega$ .

Similarly, we write  $x * \sigma$  for the unique  $y \in X^\omega$  such that

- $y(2n) = x(n)$  for all  $n \in \omega$ ;
- $y(n) = \sigma(y \upharpoonright n)$  for all odd  $n \in \omega$ .

The statement that a set  $A \subseteq X^\omega$  is determined can then be rephrased as asserting the existence of a strategy  $\sigma$  such that one of the two following statements holds.

- For every  $x \in X^\omega$ ,  $\sigma * x$  is in  $A$ .
- For every  $x \in X^\omega$ ,  $x * \sigma$  is not in  $A$ .

**1.1.2 Remark.** We list four classical consequences of AD, the details of which are presented in Chapter 33 of [8] and Sections 27 and 28 of [10]. An abbreviated history of determinacy axioms can be found in [26].

1. Jan Mycielski observed that AD implies  $\text{CC}_\mathbb{R}$  (as defined in Section 0.4). To see this, let  $A_i$  ( $i \in \omega$ ) be nonempty subsets of  $\omega^\omega$ , and consider the game where  $I$  plays  $i \in \omega$  and then  $II$  must list the values of a member of  $A_i$ .
2. Morton Davis proved that AD implies that every uncountable subset of  $\omega^\omega$  contains a perfect set. To see this, consider a set  $A \subseteq \omega^\omega$ , and the game where players  $I$  and  $II$  collaborate to build an  $x \in \omega^\omega$ , with player  $II$  choosing individual digits as usual, but player  $I$  allowed to play finite sequences from  $\omega$ , with  $I$  winning if the concatenation of these moves is in  $A$ . Player  $II$  has a winning strategy if and only if  $A$  is countable, and a winning strategy for  $I$  induces a perfect subset of  $A$ . It follows from ZF that if every uncountable subset of  $\omega^\omega$  contains a perfect set then there is no injection from  $\omega_1$  into  $\omega^\omega$  (such an injection would give a wellordering of  $\omega^\omega$  in ordertype  $\omega_1$ ). The nonexistence of an injection from  $\omega_1$  into  $\omega^\omega$



(which we will denote by writing  $\aleph_1 \not\leq 2^{\aleph_0}$ ) is equivalent to the assertion that for any inner model  $M$  satisfying Choice (equivalently, for all models of the form  $L[S]$ , for  $S$  a subset of  $\omega_1$ ), and any countable ordinal  $\alpha$ ,  $\mathcal{P}(\alpha) \cap M$  is countable. Since ZF implies the existence of partition of  $\mathcal{P}(\omega)$  into  $\aleph_1$  many sets, this shows that the statement  $\text{AD}_{\omega_1}$  is inconsistent with ZF.

3. Stefan Banach proved that AD implies that every subset of  $\omega^\omega$  (similarly, every subset of  $2^\omega$ ) has the property of Baire, i.e., that for each  $A \subseteq \omega^\omega$  there exists an open  $U \subseteq \omega^\omega$  such that the symmetric difference  $A \Delta U$  is meager. To see this, fix a bijection  $\pi: \omega \rightarrow \omega^{<\omega}$  and, given  $A \subseteq \omega^\omega$ , let  $B$  be the set of  $x \in \omega^\omega$  for which the concatenation of the values  $\pi(x(i))$  ( $i \in \omega$ ) is in  $A$ . If player  $I$  was a winning strategy in  $\mathcal{G}_B$ , then  $A$  is relatively comeager in some open set; if player  $II$  does, then  $A$  is meager. Waclaw Sierpiński that nonprincipal ultrafilters on  $\omega$  (considered as subsets of  $2^\omega$ ) do not have the property of Baire. It follows that nonprincipal ultrafilters on  $\omega$  don't exist under AD, and from this that, under AD, every nonprincipal ultrafilter on any set is countably complete.
4. Robert Solovay proved AD implies that the club filter on  $\omega_1$  is an ultrafilter (which is countably complete by either part (1) or (3) above). This gives another proof that (under AD) there is no injection from  $\omega_1$  into  $\mathcal{P}(\omega)$ . We present a proof that AD implies the measurability of  $\omega_1$  (not Solovay's original proof) in Remark 1.2.7.

Given  $\Gamma \subseteq \mathcal{P}(\omega^\omega)$ , we will write  $\text{Baire}(\Gamma)$  for the assertion that every element of  $\Gamma$  has the property of Baire.

## 1.2 Turing Determinacy

In this section we prove Martin's theorem that under AD the cone measure on the Turing degrees is an ultrafilter. We specify a class of relations for which the notion of cone measure applies, and call these *ordered equivalence relations*. We define a relative coarseness relation (being *as thick as*) on ordered equivalence relations for which the property of the associated cone measure being an ultrafilter is preserved upwards. We prove Martin's theorem for the smallest relation for which his original proof applies, which we call  $\leq_{\text{Ma}}$ . In practice we will often use his theorem with Turing equivalence and coarser relations, in the sense introduced in Definition 1.2.3.

Fixing an enumeration  $\langle \sigma_n : n < \omega \rangle$  of  $\omega^{<\omega}$ , we can associate to each  $x \in \omega^\omega$  a function (strategy)  $\pi_x: \omega^{<\omega} \rightarrow \omega$  defined by the formula  $\pi_x(\sigma_n) = x(n)$ .

Let  $\text{even}: \omega^\omega \rightarrow \omega^\omega$  be the function defined by letting  $\text{even}(y)(n) = y(2n)$  for each  $n \in \omega$  and let  $\text{odd}: \omega^\omega \rightarrow \omega^\omega$  be the function defined by letting  $\text{odd}(y)(n) = y(2n+1)$  for each  $n \in \omega$ . Given  $x \in \omega^\omega$ , let

- $I_x^*: \omega^\omega \rightarrow \omega^\omega$  be the function defined by letting  $I_x^*(y)$  be  $\pi_y * x$ , i.e., the result of playing  $x$  for player  $II$  against the strategy  $\pi_y$  for player  $I$ ;

- $II_x^*: \omega^\omega \rightarrow \omega^\omega$  be the function defined by letting  $II_x^*(y)$  be  $x * \pi_y$ , i.e., the result of playing  $x$  for player  $I$  against the strategy  $\pi_y$  for player  $II$ ;
- $\mathcal{F}_x$  be the smallest set of functions on  $\omega^\omega$  which is closed under composition and contains the identity function, even, odd and the functions  $I_{f(x)}^*$  and  $II_{f(x)}^*$  for each  $f \in \mathcal{F}_x$ .

Let  $\leq_{\text{Ma}}$  be the reflexive binary relation on  $\omega^\omega$  defined by setting  $y \leq_{\text{Ma}} x$  if  $y = f(x)$  for some  $f \in \mathcal{F}_x$ . Observe that if  $y \leq_{\text{Ma}} x$  then  $\mathcal{F}_y \subseteq \mathcal{F}_x$ ; it follows from this that  $\leq_{\text{Ma}}$  is transitive. The functions even and odd can be used to show that  $\leq_{\text{Ma}}$  is countably directed.

We let  $\equiv_{\text{Ma}}$  denote the equivalence relation  $\leq_{\text{Ma}} \cap \geq_{\text{Ma}}$ . The ordered pairs  $(\equiv_{\text{Tu}}, \leq_{\text{Tu}})$  and  $(\equiv_{\text{Ma}}, \leq_{\text{Ma}})$  are instances of the following definition, with  $X$  as  $\omega^\omega$ .

**1.2.1 Definition.** An *ordered equivalence relation* is a pair  $(E, \leq_E)$  where  $E$  is an equivalence relation on a set  $X$  and  $\leq_E$  is a preorder order on  $X$  such that  $E = \leq_E \cap \geq_E$ . We say that  $(E, \leq_E)$  is an ordered equivalence relation on  $X$ . We write  $[x]_E$  for the  $E$ -equivalence class of  $x$ , and  $\mathcal{C}_E$  for the set of  $E$ -equivalence classes (or *E-classes*).

Given an ordered equivalence relation  $(E, \leq_E)$  on  $\omega^\omega$ , and  $x \in \omega^\omega$ , we define the *upward  $\leq_E$ -cone* of  $x$  to be  $\mathcal{U}_E(x) = \{[y]_E : y \geq_E x\}$  and the *downward  $\leq_E$ -cone* of  $x$  to be  $\mathcal{D}_E(x) = \{[y]_E : y \leq_E x\}$ . In each case we say that  $x$  is a *base* for the corresponding cone. Since we are typically interested in upward cones, we often write “cone” for “upward cone”. The  *$\leq_E$ -cone measure* is

$$\{A \subseteq \{[x]_E : x \in X\} : \exists x \in X \mathcal{U}_E(x) \subseteq A\}.$$

We will write  $[x]_{\text{Ma}}$ ,  $\mathcal{C}_{\text{Ma}}$ ,  $[x]_{\text{Tu}}$  and  $\mathcal{C}_{\text{Tu}}$  instead of the more cumbersome forms using  $\equiv_{\text{Ma}}$  and  $\equiv_{\text{Tu}}$ . We also write  $\mu_{\text{Ma}}$  and  $\mu_{\text{Tu}}$  for the corresponding cone measures, which are ultrafilters under AD. We call  $\mu_{\text{Tu}}$  the *Turing measure*. A *Turing cone* is an upward cone for  $\leq_{\text{Tu}}$ . We also use the expression *Turing cone* for the union of the members of a Turing cone as just defined (i.e., for the set of reals corresponding to the set of equivalence classes).

**Theorem 1.2.2 (Martin).** *Suppose that AD holds. Then  $\mu_{\text{Ma}}$  is a countably complete ultrafilter.*

*Proof.* Since AD implies  $\text{CC}_{\mathbb{R}}$ ,  $\mu_{\text{Ma}}$  is countably complete. It suffices then to show that each subset of  $\mathcal{C}_{\text{Ma}}$  either contains or is disjoint from a cone. Fix  $A \subseteq \mathcal{C}_{\text{Ma}}$  and consider the game  $\mathcal{G}_{A^*}$ , where  $A^*$  is the set of  $x \in \omega^\omega$  such that  $[x]_{\text{Ma}}$  is in  $A$ . Let  $\pi$  be a winning strategy (for either player), and let  $x \in \omega^\omega$  be such that  $\pi = \pi_x$ . Let  $y \in \omega^\omega$  be such that  $y \geq_{\text{Ma}} x$ . If  $\pi$  is a winning strategy for player  $I$ , then  $[\pi_x * y]_{\text{Ma}}$  is in  $A$ , and  $[y]_{\text{Ma}} = [\pi_x * y]_{\text{Ma}}$  (one direction of the equivalence uses the function  $I_y^* \circ g$ , where  $g \in \mathcal{F}_y$  is such that  $x = g(y)$ ; the other uses the function odd). If  $\pi$  is a winning strategy for player  $II$ , then  $[y * \pi_x]_{\text{Ma}}$  is not in  $A$ , and  $[y]_{\text{Ma}} = [y * \pi_x]_{\text{Ma}}$ .  $\square$

We will typically be considering equivalence relations satisfying the following definition, with  $\leq_{\text{Ma}}$  or  $\leq_{\text{TU}}$  in the role of  $F$ , especially those of the form of the second type listed in Example 1.2.4, with  $S$  a set of ordinals.

**1.2.3 Definition.** Given two ordered equivalence relations  $(E, \leq_E)$  and  $(F, \leq_F)$  on the same underlying set  $X$ , say that  $(E, \leq_E)$  is *as thick as*  $(F, \leq_F)$  if, for all  $x, y$  in  $X$ ,

1. if  $x \leq_F y$  then  $x \leq_E y$ ;
2. if  $x \leq_E y$ , then for some  $z \in [y]_E$ ,  $x \leq_F z$ .

Every ordered equivalence relation is trivially as thick as itself. We leave it as an exercise to check that the “as thick as” relation is transitive. Given the first condition in Definition 1.2.3, the second condition is equivalent to saying that for all  $x \in X$ ,

$$\mathcal{U}_E(x) = \{[y]_E : x \leq_F y\}.$$

If  $\text{CC}_{\mathbb{R}}$  holds and  $(E, \leq_E)$  and  $(F, \leq_F)$  are ordered equivalence relations, with  $F$  a subset of  $E$  and  $F$  countably directed, then  $E$  is also countably directed.

**1.2.4 Example.** The following are examples of ordered equivalence relations on  $\omega^\omega$  which are as thick as  $(\equiv_{\text{Ma}}, \leq_{\text{Ma}})$ .

1.  $(\equiv_{\text{TU}}, \leq_{\text{TU}})$
2. For any set  $S$ , the equivalence relation  $L(S, x) = L(S, y)$ , under the order  $x \in L(S, y)$ . When  $S = \emptyset$ , the corresponding equivalence classes are called the *constructibility degrees*.
3. For any set  $S$ , the equivalence relation  $\text{HOD}_{S \cup \{x\}} = \text{HOD}_{S \cup \{y\}}$ , under the order  $x \in \text{HOD}_{S \cup \{y\}}$ .

**1.2.5 Remark.** Suppose that  $(E, \leq_E)$  and  $(F, \leq_F)$  are ordered equivalence relations on a set  $X$ ,  $(E, \leq_E)$  is as thick as  $(F, \leq_F)$ , and  $\mu_F$  is an ultrafilter on the set of  $F$ -classes. Then  $\mu_E$  is an ultrafilter on the set of  $E$ -classes. To see this, let  $A$  be a set of  $E$ -classes, and let  $B$  be the set of  $F$ -classes contained in a member of  $A$ . Since  $\mu_F$  is an ultrafilter, we can fix an  $x \in \omega^\omega$  such that  $\mathcal{U}_F(x)$  is either contained in or disjoint from  $B$ . Since  $\mathcal{U}_E(x) = \{[y]_E : y \geq_F x\}$ ,  $\mathcal{U}_E(x)$  is either contained in or disjoint from  $A$ .

We let Turing Determinacy (TD) denote the statement that the  $\leq_{\text{TU}}$ -cone measure is an ultrafilter on the  $\equiv_{\text{TU}}$ -classes. A recent result of Peng and Yu [33] shows that TD implies  $\text{CC}_{\mathbb{R}}$ . Their proof gives the analogous result for the relation  $x \in L[S, y]$ , for  $S$  a set of ordinals and  $x, y \in \omega^\omega$ , if in addition the sharp of each such pair  $(S, y)$  exists. It is open whether either of AD and TD implies  $\text{DC}_{\mathbb{R}}$ , or whether either of TD and TD +  $\text{DC}_{\mathbb{R}}$  implies AD. Woodin has shown that TD +  $\text{DC}_{\mathbb{R}}$  +  $V=L(\mathbb{R})$  implies AD [2]. Remark 1.2.5 and Theorem 1.2.2 (or, its proof, in conjunction with the Peng-Yu theorem) give the following.

**Corollary 1.2.6** (ZF + TD). *If  $(E, \leq_E)$  is an ordered equivalence relation on  $\omega^\omega$  which is as thick as  $\equiv_M$ , then  $\mu_E$  is a countably complete ultrafilter on  $\mathcal{C}_E$ .*

The following remark gives (from TD) Solovay's theorem that, assuming AD,  $\omega_1$  is measurable. It also shows that TD implies that  $\aleph_1 \not\leq 2^{\aleph_0}$ .

**1.2.7 Remark.** Let  $(E, \leq_E)$  be an ordered equivalence relation on  $\omega^\omega$  such that each downward cone is countable, and suppose that  $\mu_E$  is a countably complete ultrafilter on  $\mathcal{C}_E$ . For each  $\alpha < \omega_1$ , let  $X_\alpha$  be the set of  $x \in \omega^\omega$  which HC-code  $\alpha$ , in the sense of Section 0.2. For each  $\sigma \in \mathcal{C}_E$ , let  $\alpha_\sigma$  be  $\sup\{\alpha < \omega_1 : \exists x \in X_\alpha \exists y \in \sigma \ x \leq_E y\}$ . Let  $U$  be the set of  $A \subseteq \omega_1$  for which  $\{\sigma \in \mathcal{C}_E : \alpha_\sigma \in A\} \in \mu_E$ . Then  $U$  is a countably complete ultrafilter on  $\omega_1$ .

The following example gives consequences of the Turing measure being an ultrafilter. Similar arguments will appear in Chapters 9, 10 and 11.

**1.2.8 Example.** Let  $S$  be a set of ordinals, let  $\leq_S$  be the binary relation on  $\omega^\omega$  given by the binary relation  $x \in L[S, y]$ , let  $\equiv_S$  be the corresponding equivalence relation and let  $\mu_S$  be the  $\equiv_S$ -cone measure. Assume that  $\mu_S$  is an ultrafilter. It follows then by Remark 1.2.7 that  $\aleph_1 \not\leq 2^{\aleph_0}$ . For each  $x \in \omega^\omega$ , there exist  $y$  and  $z$  in  $\omega^\omega$  such that  $y$  is generic for Sacks forcing over  $L[S, x]$  and  $z$  codes a wellordering of  $\omega$  in ordertype  $\omega_1^{L[S, x]}$ . By standard facts about Sacks forcing [34],  $\bigcup_{n \in \omega} \{(2n, x(n)), (2n+1, y(n))\}$  is an immediate  $\leq_S$ -successor of  $x$ . It follows then that the set of  $\leq_S$ -successors contains a set in  $\mu_S$ . Let  $f: \omega^\omega \rightarrow \omega_1$  be the function defined by setting  $f(x)$  to be  $\omega_1^{L[S, x]}$ . Since Sacks forcing preserves  $\omega_1$ , we have that for every  $x \in \omega^\omega$ , there exist  $y, z$  in  $\omega^\omega$  both  $\leq_S$ -above  $x$ , such that  $f(x) = f(y)$  and  $f(x) < f(z)$ .

## Chapter 2

# The Wadge hierarchy

In this chapter we review some basic concepts from classical descriptive set theory. In Section 2.1 we introduce Lipschitz determinacy and prove Martin's theorem that the corresponding hierarchy is wellfounded under  $\text{AD} + \text{DC}_{\mathbb{R}}$ . In Section 2.2 we review the relationship between the Lipschitz degrees and the Wadge degrees, and show that a Wadge class whose rank is a limit ordinal is selfdual if and only if its rank has countable cofinality. In Section 2.3 we introduce boldface and lightface pointclasses. Section 2.4 is about universal sets for pointclasses and the s-m-n property. Finally, Section 2.5 introduces  $\Theta$ , the least (nonzero) ordinal which is not a surjective image of  $\omega^\omega$ .

### 2.1 Lipschitz determinacy

Given sets  $A, B \subseteq \omega^\omega$ , we say that  $A$  is *Wadge reducible* to  $B$  (and write  $A \leq_{\text{Wa}} B$ ) if there is a continuous function  $f: \omega^\omega \rightarrow \omega^\omega$  such that  $A = f^{-1}[B]$ . A function  $f: \omega^\omega \rightarrow \omega^\omega$  is *Lipschitz* if, for all  $x, y \in \omega^\omega$  and  $n \in \omega$ , if  $x \upharpoonright n = y \upharpoonright n$ , then  $f(x) \upharpoonright n = f(y) \upharpoonright n$ . We say that  $A$  is *Lipschitz reducible* to  $B$  (and write  $A \leq_{\text{L}} B$ ) if there is a Lipschitz function  $f: \omega^\omega \rightarrow \omega^\omega$  such that  $A = f^{-1}[B]$ . Since Lipschitz functions are continuous,  $A \leq_{\text{L}} B$  implies  $A \leq_{\text{W}} B$ .

**2.1.1 Remark.** Since  $A = f^{-1}[B]$  implies that  $(\omega^\omega \setminus A) = f^{-1}[\omega^\omega \setminus B]$ ,  $A \leq_{\text{W}} B$  implies  $(\omega^\omega \setminus A) \leq_{\text{W}} (\omega^\omega \setminus B)$  and  $A \leq_{\text{L}} B$  implies  $(\omega^\omega \setminus A) \leq_{\text{L}} (\omega^\omega \setminus B)$ .

The relations  $\leq_{\text{W}}$  and  $\leq_{\text{L}}$  are easily seen to be preorders, that is, reflexive and transitive. We write  $=_{\text{Wa}}$  and  $=_{\text{Li}}$  respectively for the equivalence relations  $\leq_{\text{W}} \cap \geq_{\text{Wa}}$  and  $\leq_{\text{L}} \cap \geq_{\text{Li}}$ , whose equivalence classes are respectively called *Wadge classes* and *Lipschitz classes*. Given  $A \subseteq \omega^\omega$ , we write  $[A]_{\text{Wa}}$  for the Wadge class of  $A$ , and  $[A]_{\text{Li}}$  for the Lipschitz class of  $A$ . We say that a Wadge class or Lipschitz class is *selfdual* if it contains a pair of complements (in which case it is closed under complements, by Remark 2.1.1), and *nonselfdual* otherwise. We write  $x <_{\text{Wa}} y$  for  $(x \leq_{\text{W}} y) \wedge \neg(y \leq_{\text{W}} x)$  and  $x <_{\text{Li}} y$  for  $(x \leq_{\text{L}} y) \wedge \neg(y \leq_{\text{L}} x)$ .

The following definition converts a pair  $A, B \subseteq \omega^\omega$  into a set  $A \oplus B \subseteq \omega^\omega$  such that a winning strategy for player  $II$  in  $\mathcal{G}_{A \oplus B}$  gives a Lipschitz reduction of  $A$  to  $B$ .

**2.1.2 Definition.** Let  $A, B$  be subsets of  $\omega^\omega$ . We write  $A \oplus B$  for the set of  $\langle x_i : i \in \omega \rangle$  for which  $\langle x_{2i} : i \in \omega \rangle \in A$  if and only if  $\langle x_{2i+1} : i \in \omega \rangle \notin B$ .

The game  $\mathcal{G}_{A \oplus B}$  can then be seen as a game where  $I$  plays  $x \in \omega^\omega$ ,  $II$  plays  $y \in \omega^\omega$ , and  $II$  wins if and only if  $(x \in A \Leftrightarrow y \in B)$ . We call this the *Lipschitz game* for the pair  $(A, B)$ .

**2.1.3 Remark.** The relation  $A \leq_L B$  is equivalent to the assertion that player  $II$  has a winning strategy in the game  $\mathcal{G}_{A \oplus B}$ ; it also follows from player  $I$  having a winning strategy in  $\mathcal{G}_{B \oplus (\omega^\omega \setminus A)}$ .

This observation leads to the following fundamental fact.

**Theorem 2.1.4 (Wadge).** *Let  $A$  and  $B$  be subsets of  $\omega^\omega$ . If player  $I$  has a winning strategy in  $\mathcal{G}_{A \oplus B}$ , then  $(\omega^\omega \setminus B) \leq_L A$ . If player  $II$  has a winning strategy in  $\mathcal{G}_{A \oplus B}$ , then  $A \leq_L B$ .*

Wadge's Theorem (along with Remarks 2.1.1 and 2.1.3) has the following consequences. Proposition 2.1.5 (which can be proved directly from Wadge's Theorem) shows that the ordering on the Lipschitz degrees induced by  $\leq_{Li}$  is almost linear, the exceptions being pairs of the form  $[A]_{Li}$ ,  $[\omega^\omega \setminus A]_{Li}$ , for nonselfdual classes  $[A]_{Li}$ . The conclusion of the proposition is sometimes called the Semi-Linear Ordering Principle for Lipschitz maps.

**Proposition 2.1.5.** *Let  $A$  and  $B$  be subsets of  $\omega^\omega$  such that  $A \oplus B$  and  $B \oplus A$  are both determined. If  $A \not\leq_L B$  and  $B \not\leq_L A$ , then  $A =_{Li} (\omega^\omega \setminus B)$ .*

**Proposition 2.1.6.** *Let  $A$  and  $B$  be subsets of  $\omega^\omega$  such that the sets  $B \oplus A$  and  $B \oplus (\omega^\omega \setminus A)$  are both determined. If  $A <_{Li} B$ , then player  $I$  wins both  $\mathcal{G}_{B \oplus A}$  and  $\mathcal{G}_{B \oplus (\omega^\omega \setminus A)}$ .*

*Proof.* In the case of  $B \oplus (\omega^\omega \setminus A)$ , one gets otherwise that  $A <_{Li} B \leq_L (\omega^\omega \setminus A)$ , and therefore by Remark 2.1.1 that  $A =_{Li} (\omega^\omega \setminus A)$  and  $B \leq_L A$ , giving a contradiction.  $\square$

**Proposition 2.1.7.** *Let  $A$  and  $B$  be subsets of  $\omega^\omega$  such that  $A \oplus B$  and  $B \oplus A$  are both determined. If  $A <_{Wa} B$  then  $A <_{Li} B$ .*

*Proof.* Since  $B \not\leq_W A$ ,  $B \not\leq_L A$ , so by Proposition 2.1.5, either  $A =_{Li} (\omega^\omega \setminus B)$  or  $A \leq_L B$ . The former is impossible, as then  $A \leq_W B$  would imply that  $(\omega^\omega \setminus B) \leq_W (\omega^\omega \setminus A)$  and thus  $B \leq_W A$ , giving a contradiction.  $\square$

We let *Lipschitz Determinacy* be the statement that  $A \oplus B$  is determined for all subsets  $A, B$  of  $\omega^\omega$ .

**2.1.8 Remark.** It is not hard to see that Lipschitz Determinacy implies  $\text{CC}_{\mathbb{R}}$  (see [25], for instance; the proof there combines the proof of Theorem 2.5.4 with a standard strategy selection argument, which appears also at the end of the proof of Theorem 13.2.3). Similarly, Lipschitz Determinacy implies that every uncountable subset of  $\omega^\omega$  contains a perfect set. To see this, consider the game  $\mathcal{G}_{A \oplus B}$ , where  $A$  is a Borel set which is not a countable union of closed sets, and  $B$  is non-Borel. A winning strategy for player II produces an uncountable analytic subset of  $B$ . It follows then that Lipschitz Determinacy implies  $\aleph_1 \not\leq 2^{\aleph_0}$ .

The following gives Martin's theorem showing that  $<_{\text{Li}}$  is wellfounded (assuming  $\text{AD} + \text{DC}_{\mathbb{R}}$ , for instance). By Proposition 2.1.7, theorem implies the corresponding version for  $<_{\text{Wa}}$ .

**Theorem 2.1.9** (Martin). *If Lipschitz Determinacy + Baire( $\mathcal{P}(\omega^\omega)$ ) holds, then there does not exist a sequence  $\langle A_i : i \in \omega \rangle$  consisting of subsets of  $\omega^\omega$ , such that  $A_{i+1} <_{\text{Li}} A_i$  for each  $i \in \omega$ .*

*Proof.* Suppose towards a contradiction that such a sequence  $\langle A_i : i < \omega \rangle$  does exist. Applying Proposition 2.1.6 and  $\text{CC}_{\mathbb{R}}$  we can fix winning strategies  $f_i^0, f_i^1$  ( $i \in \omega$ ) for player I in the games  $\mathcal{G}_{A_i \oplus A_{i+1}}$  and  $\mathcal{G}_{A_i \oplus (\omega^\omega \setminus A_{i+1})}$ , respectively.

For each  $x \in {}^\omega 2$ , define  $y_i(x) \in \omega^\omega$  ( $i \in \omega$ ) by letting  $y_i(x)(k)$  be

$$f_i^{x(i)}(\langle y_i(x)(0), y_{i+1}(x)(0), \dots, y_i(x)(k-1), y_{i+1}(x)(k-1) \rangle).$$

Note that the values  $y_i(x)(k)$  are defined recursively in  $k$ , for all  $x$  and  $i$  simultaneously. For each fixed  $x$ , each  $y_i(x)$  is the set of values given by the strategy  $f_i^{x(i)}$  when playing against the moves for player II listed by  $y_{i+1}(x)$ . It follows that for each  $i \in \omega$ , if  $x(i) = 0$  then  $y_i(x) \in A_i \Leftrightarrow y_{i+1}(x) \notin A_{i+1}$ , and if  $x(i) = 1$  then  $y_i(x) \in A_i \Leftrightarrow y_{i+1}(x) \in A_{i+1}$ .

Let  $Z$  be the set of  $x \in {}^\omega 2$  for which  $y_0(x) \in A_0$ . It suffices to show that whenever  $x, x' \in {}^\omega 2$  differ at exactly one point,  $x \in Z$  if and only if  $x' \notin Z$ , since no subset of  ${}^\omega 2$  with the property of Baire can have this property.

First observe that if  $x, x' \in {}^\omega 2$  and  $i \in \omega$  are such that  $x(j) = x'(j)$  for all  $j \geq i$ , then  $y_i(x) = y_i(x')$ . If  $i_0$  is the unique  $i \in \omega$  such that  $x(i) \neq x'(i)$ , then  $y_{i_0+1}(x) = y_{i_0+1}(x')$ . Since  $x(i_0) \neq x'(i_0)$ , it follows that  $y_{i_0}(x) \in A_{i_0}$  if and only if  $y_{i_0}(x') \notin A_{i_0}$ . Since  $x(i) = x'(i)$  for all  $i < i_0$ , it follows that  $y_i(x) \in A_i$  if and only if  $y_i(x') \notin A_i$  for all such  $i$ .  $\square$

Recall that a preorder is *wellfounded* if every nonempty subset of its domain has a minimal element. The additional assumption of  $\text{DC}_{\mathbb{R}}$  derives the wellfoundedness of  $<_{\text{Li}}$  from Theorem 2.1.9. Since every Wadge degree is a union of Lipschitz degrees, Corollary 2.1.10 holds for  $<_{\text{Wa}}$  as well. Theorem 2.2.8 shows (assuming Lipschitz Determinacy and Baire( $\mathcal{P}(\omega^\omega)$ )) but not  $\text{DC}_{\mathbb{R}}$  that  $<_{\text{Li}}$  is wellfounded if and only if  $<_{\text{Wa}}$  is.

**Corollary 2.1.10** (Martin). *If  $\text{DC}_{\mathbb{R}} + \text{Lipschitz Determinacy} + \text{Baire}(\mathcal{P}(\omega^\omega))$  holds, then  $<_{\text{Li}}$  and  $<_{\text{Wa}}$  are wellfounded.*

*Proof.* If  $<_{\text{Li}}$  is illfounded, then  $\text{DC}_{\mathbb{R}}$  gives a sequence  $\langle A_i : i \in \omega \rangle$  of subsets of  $\omega^\omega$  such that  $A_{i+1} <_{\text{Li}} A_i$  for each  $i \in \omega$ .  $\square$

As noted in Section 0.4, it is open whether AD implies  $\text{DC}_{\mathbb{R}}$ . It is also open whether  $\text{DC}_{\mathbb{R}}$  is needed for Corollary 2.1.10.

The *Lipschitz rank*  $\text{LR}(A)$  (respectively, *Wadge rank*  $\text{WR}(A)$ ) of a set  $A \subseteq \omega^\omega$  is recursively defined to be the least ordinal greater than the Lipschitz (Wadge) rank of every  $B \subseteq \omega^\omega$  with  $B <_{\text{Li}} A$  ( $B <_{\text{Wa}} A$ ), if this is defined (that is, the functions LR and WR are the canonical rank functions on the wellfounded initial segments of  $\leq_{\text{Li}}$  and  $\leq_{\text{Wa}}$ ). By Proposition 2.1.5, for each ordinal  $\alpha$ , the subsets of  $\omega^\omega$  of Lipschitz rank (Wadge rank)  $\alpha$  (if there are any) consist either of a single Lipschitz (Wadge) class or a pair of Lipschitz (Wadge) classes corresponding to complements. Theorem 2.2.8 says more about the relationship between the two sets of classes.

## 2.2 Lipschitz degrees and Wadge degrees

Following [40], we give some more details on the structure of the Lipschitz and Wadge hierarchies, using Lipschitz Determinacy and the assumption that all sets of reals have the Baire property, but not  $\text{DC}_{\mathbb{R}}$ . We begin by noting that  $\emptyset$  and  $\omega^\omega$  are the only subsets of  $\omega^\omega$  of Lipschitz or Wadge rank 0, and that they are Wadge (and thus Lipschitz) inequivalent.

We write  $s \frown x$  for the concatenation of  $s$  and  $x$ , where  $s$  is a finite sequence and  $x$  is either finite or infinite. Given  $A \subseteq \omega^\omega$  and  $i \in \omega$ , we write  $A^{(i)}$  for  $\{\langle i \rangle \frown x : x \in A\}$  and  $A_{(i)}$  for  $\{x : \langle i \rangle \frown x \in A\}$ . Note that  $(A^{(i)})_{(i)} = A$ , while  $(A_{(i)})^{(i)} = \{x \in A : x(0) = i\}$ .

Proposition 2.2.1 shows (among other things) that  $[A^{(i)}]_{\text{Li}}$  is the least Lipschitz class above  $[A]_{\text{Li}}$  when  $A$  is selfdual.

**Proposition 2.2.1** (ZF + Lipschitz Determinacy). *Let  $A \subseteq \omega^\omega$  be such that  $[A]_{\text{Li}}$  is selfdual, and let  $i$  be an element of  $\omega$ . Then*

1.  $[A^{(i)}]_{\text{Li}}$  is selfdual, and  $[A^{(i)}]_{\text{Li}}$  is the  $\leq_{\text{L}}$ -least Lipschitz class above  $[A]_{\text{Li}}$ ;
2.  $A_{(i)} <_{\text{Li}} A$ .

*Proof.* The function  $f(x) = \langle i \rangle \frown x$  shows that  $A \leq_{\text{L}} A^{(i)}$  and  $A_{(i)} \leq_{\text{L}} A$ . To see that  $A <_{\text{Li}} A^{(i)}$  and  $A_{(i)} <_{\text{Li}} A$ , note that whenever  $f : \omega^\omega \rightarrow \omega^\omega$  has the property that for each  $n \in \omega$ ,  $f(x) \restriction n$  depends only on  $x \restriction n$ , there is an  $x \in \omega^\omega$  such that  $x(0) = i$  and

$$f(x) = \langle x(1), x(2), x(3), \dots \rangle.$$

It follows that no such  $f$  witnesses that  $A^{(i)} \leq_{\text{L}} (\omega^\omega \setminus A)$  or  $(\omega^\omega \setminus A) \leq_{\text{L}} A_{(i)}$ , which is enough since  $[A]_{\text{Li}}$  is selfdual.

Using the assumption that  $A$  is selfdual, it is easy to see that  $[A^{(i)}]_{\text{Li}}$  is selfdual. Finally, if  $C \subseteq \omega^\omega$  is such that  $A <_{\text{Li}} C$ , then player  $I$  wins  $\mathcal{G}_{C \ominus (\omega^\omega \setminus A)}$ , by



Proposition 2.1.6. Any strategy witnessing this can be used to show that  $A^{(i)} \leq_L C$  (using the fact that  $C \neq \omega^\omega$  to deal with inputs whose first coordinates are not  $i$ ).  $\square$

**2.2.2 Remark.** For any  $A \subseteq \omega^\omega$  and  $i \in \omega$ ,  $[A]_{\text{Wa}} = [A^{(i)}]_{\text{Wa}}$ . For the direction left open by Proposition 2.2.1, consider the function on  $\omega^\omega$  which removes the first coordinate of its input.

**2.2.3 Remark.** An argument similar to the proof of Proposition 2.2.1 shows that  $[\bigcup_{i \in \omega} A^{(i)}]_{\text{Li}}$  is also the  $\leq_L$ -least Lipschitz class above  $[A]_{\text{Li}}$  (again assuming Lipschitz Determinacy). It is also not hard to show directly that  $A^{(j)} =_{\text{Li}} \bigcup_{i \in \omega} A^{(i)}$  for each  $j \in \omega$ .

**2.2.4 Remark.** Let  $\pi: \omega \times \omega \rightarrow \omega$  be a bijection, and suppose that  $A_i$  ( $i \in \omega$ ) are subsets of  $\omega^\omega$ . Let  $C \subseteq \omega^\omega$  be the set of functions of the form

$$\langle \pi(i, x(0)), x(1), x(2), \dots \rangle$$

for  $i \in \omega$  and  $x \in A_i$ . Then  $A_i \leq_L C$  for all  $i \in \omega$ . If  $\text{CC}_{\mathbb{R}}$  holds, then for all  $D \subseteq \omega^\omega$ , if  $A_i \leq_L D$  for all  $i \in \omega$  then  $C \leq_L D$ . If  $\{A_i : i \in \omega\}$  does not have a  $\leq_L$ -maximal element, then  $A_i <_{\text{Li}} C$  for all  $i \in \omega$ .

If Lipschitz Determinacy holds, then  $[C]_{\text{Li}}$  is nonselfdual exactly in the case where  $\{A_i : i \in \omega\}$  has a  $\leq_L$ -maximal element whose Lipschitz class is nonselfdual. To see this, note first of all that  $[C]_{\text{Li}}$  is clearly selfdual in the case where  $\{A_i : i \in \omega\}$  has a  $\leq_L$ -maximal element whose Lipschitz class is selfdual, as  $[C]_{\text{Li}}$  is equal to this class (similarly, if  $\{A_i : i \in \omega\}$  has a  $\leq_W$ -maximal element, then  $[C]_{\text{Wa}}$  is equal to this class). In the remaining case,  $\{[A_i]_{\text{Li}} : i \in \omega\}$  has the same supremum as

$$\{[A_i]_{\text{Li}} : i \in \omega\} \cup \{[\omega^\omega \setminus A_i]_{\text{Li}} : i \in \omega\}.$$

Running the construction of  $C$  above with the set  $\{A_i : i \in \omega\} \cup \{\omega^\omega \setminus A_i : i \in \omega\}$  clearly gives a selfdual Lipschitz class (equal to  $[C]_{\text{Li}}$ ). This argument gives the following facts, for any  $A \subseteq \omega^\omega$ .

- If  $[A]_{\text{Li}}$  is nonselfdual, then the pair  $[A]_{\text{Li}}, [\omega^\omega \setminus A]_{\text{Li}}$  has a  $\leq_L$ -least upper bound, and this upper bound is selfdual.
- If  $[A]_{\text{Li}}$  is the  $\leq_L$ -supremum of a countable set of Lipschitz classes strictly below it, and Lipschitz Determinacy holds, then  $[A]_{\text{Li}}$  is selfdual (Proposition 2.2.5 below gives the converse, for non-successor classes).

From Proposition 2.2.1 and Remarks 2.2.2 and 2.2.4 it follows that for all  $A \subseteq \omega^\omega$  such that  $[A]_{\text{Li}}$  is selfdual, the first  $\omega_1$  many Lipschitz classes above  $[A]_{\text{Li}}$  are all selfdual and contained in  $[A]_{\text{Wa}}$ . Proposition 2.2.5 shows (assuming Lipschitz Determinacy and that the Lipschitz rank of each subset of  $\omega^\omega$  exists) that the nonselfdual Lipschitz classes are exactly those whose Lipschitz rank is either 0 or an ordinal of uncountable cofinality.

**Proposition 2.2.5** (ZF + Lipschitz Determinacy). *Let  $A \subseteq \omega^\omega$  be such that  $[A]_{\text{Li}}$  is selfdual, and  $[A]_{\text{Li}}$  is not the  $\leq_L$ -least Lipschitz class above any other class. Then  $[A]_{\text{Li}}$  is the  $\leq_L$ -supremum of a countable set of Lipschitz classes strictly below it.*

*Proof.* Let  $\pi: \omega \times \omega \rightarrow \omega$  be a bijection. For each  $i \in \omega$ , let  $B_i$  be the set of sequences of the form

$$\langle \pi(n, x(0)), x(1), x(2), \dots \rangle$$

such that either  $n$  is even and  $\langle i, x(0), x(1), x(2), \dots \rangle$  is in  $A$ , or  $n$  is odd and  $\langle i, x(0), x(1), x(2), \dots \rangle$  is not in  $A$ . Then by Remark 2.2.4  $[B_i]_{\text{Li}}$  is the least upper bound of the pair  $\{[A_{(i)}]_{\text{Li}}, [(\omega^\omega \setminus A)_{(i)}]_{\text{Li}}\}$  and  $B_i$  is selfdual. By Proposition 2.2.1, and the assumption that  $[A]_{\text{Li}}$  is not the  $\leq_L$ -least Lipschitz class above any other class, we have that  $B_i <_{\text{Li}} A$ .

For each  $i \in \omega$ , let  $C_i$  be the set of  $x \in \omega^\omega$  such that  $x(0) = i$  and  $\langle x(1), x(2), \dots \rangle \in B_i$ . By Proposition 2.2.1, each  $[C_i]_{\text{Li}}$  is the  $<_{\text{Li}}$ -successor of the corresponding  $[B_i]_{\text{Li}}$ . Since  $[A]_{\text{Li}}$  is not a successor class, it follows that  $B_i <_{\text{Li}} C_i <_{\text{Li}} A$  holds, for each  $i \in \omega$ . Let  $D$  be the set constructed from  $\{C_i : i \in \omega\}$  as in Remark 2.2.4. Then  $[D]_{\text{Li}}$  is the supremum of  $\{[C_i]_{\text{Li}} : i < \omega\}$ , and it suffices to see that  $A \leq_L D$ .

Let  $g: {}^{<\omega}\omega \rightarrow \omega$  be such that for all  $n \in \omega \setminus 2$ ,

$$g(\langle i_0, \dots, i_n \rangle) = \langle \pi(i_0, i_0), \pi(0, i_1), i_2, \dots, i_n \rangle.$$

Let  $g^*: \omega^\omega \rightarrow \omega^\omega$  be the Lipschitz function induced by  $g$ . Let us see that  $g^*$  witnesses that  $A \leq_L D$ . Fix  $x \in \omega^\omega$ , and let  $g^*(x)$  have the form

$$\langle \pi(i_0, y(0)), y(1), y(2), \dots \rangle$$

for some  $i_0 \in \omega$  and some  $y \in \omega^\omega$ . Then  $i_0 = x(0) = y(0)$ ,  $y(1) = \pi(0, x(1))$  and  $y(i) = x(i)$  for all  $i \in \omega \setminus 2$ . We want to see that  $x \in A$  if and only if  $g^*(x) \in D$ . Now,  $g^*(x)$  is in  $D$  if and only if  $y \in C_{i_0}$ , which in turn happens if and only if  $\langle \pi(0, x(1)), x(2), x(3), \dots \rangle$  is in  $B_{i_0}$ , which happens if and only if  $x$  is in  $A$ .  $\square$

The following proposition completes the analysis of which Wadge classes are selfdual classes, as well as the relationship between the Lipschitz classes and the Wadge classes.

**Theorem 2.2.6** (Steel, Van Wesep). *Suppose that*

$$\text{Lipschitz Determinacy} + \text{Baire}(\mathcal{P}(\omega^\omega))$$

*holds. For all  $A \subseteq \omega^\omega$ , if  $[A]_{\text{Wa}}$  is selfdual, then so is  $[A]_{\text{Li}}$ .*

*Proof.* Suppose towards a contradiction that  $[A]_{\text{Wa}}$  is selfdual but  $[A]_{\text{Li}}$  is not. By Lipschitz Determinacy, if  $[A]_{\text{Li}}$  is not selfdual, then player  $I$  wins  $\mathcal{G}_{A \ominus (\omega^\omega \setminus A)}$ . Let  $\sigma$  be a strategy witnessing this.

Consider the following game between players  $I$  and  $II$  (which is called the *Wadge game*). For each  $i \in \omega$ , player  $I$  plays a value  $x(i)$ , and player  $II$  either

passes or plays a value  $y(j)$ , for  $j$  the least  $k \in i + 1$  for which a value for  $y(k)$  has not yet been chosen. If at the end of the game there is a  $j \in \omega$  for which  $y(j)$  has not been chosen, then  $II$  loses. Otherwise  $II$  wins if and only if  $x \in A \Leftrightarrow y \notin A$ . The statement that  $A \leq_W (\omega^\omega \setminus A)$  is equivalent to the existence of a winning strategy for  $II$ . Let  $\tau$  be such a strategy.

For each positive  $n \in \omega$  and each sequence  $\bar{c} = \langle c_m : m < n \rangle \in 3^n$ , there is a unique (possibly partial) function  $s_{\bar{c}}$  on  $n \times \omega$  satisfying the following conditions.

- For each  $m < n$ , the set of  $i$  for which  $(m, i) \in \text{dom}(s_{\bar{c}})$  is an ordinal  $\alpha_m \in \omega + 1$ . We let  $t_m$  be the function with domain  $\alpha_m$  such that  $t_m(i) = s_{\bar{c}}(m, i)$  for all  $i < \alpha_m$ .
- The largest  $m < n$  for which  $\alpha_m > 0$  is the largest  $m < n$  for which  $c_m = 2$ , and for this  $m$ ,  $\alpha_m = 1$  and  $s_{\bar{c}}(m, 0) = \sigma(\langle \rangle)$ .
- For each  $m < n - 1$  such that  $c_m = 2$ ,  $\alpha_m = \alpha_{m+1} + 1$ , and for each  $i < \alpha_m$ ,  $s_{\bar{c}}(m, i)$  is the response given by  $\sigma$  when player  $II$  plays  $t_m \upharpoonright i$ .
- For each  $m < n - 1$  such that  $c_m = 0$ ,  $\alpha_m = \alpha_{m+1}$ , and  $t_m = t_{m+1}$ .
- For each  $m < n - 1$  such that  $c_m = 1$ ,  $t_m$  is the longest sequence of nonpassing moves made by  $\tau$  in response to  $t_{m+1}$ .

Choose integers  $i_k$  ( $k \in \omega$ ) so that  $i_0 = 0$ ,  $i_{k+1} > i_k + 1$  for all  $k \in \omega$  and, for each sequence  $\bar{c} = \langle c_m : m < i_{k+1} \rangle$  in  $3^n$ , if  $c_m = 2$  for all

$$m \in i_{k+1} \setminus \{i_p : p \leq k\},$$

then the corresponding value  $\alpha_j$  is at least  $k$  for each  $j \leq i_k$ .

For each  $x \in {}^\omega 2$  and each  $m \in \omega$ , let  $c_m^x$  be  $x(k)$  if  $m = i_k$  for some  $k \in \omega$ , and let  $c_m^x$  be 2 otherwise. For each such  $x$  there is a unique sequence  $\langle y_m^x : m \in \omega \rangle \in (\omega^\omega)^\omega$  such that for each  $m \in \omega$ ,

- if  $c_m^x = 2$ , then  $y_m^x$  is the sequence produced by  $\sigma$  when player  $II$  plays  $y_{m+1}^x$ ;
- if  $c_m^x = 0$ , then  $y_m^x = y_{m+1}^x$ ;
- if  $c_m^x = 1$ , then  $y_m^x$  is the sequence produced by  $\tau$  when player  $I$  plays  $y_{m+1}^x$ .

Then as in the proof of Theorem 2.1.9, if  $x, x' \in 2^\omega$  and  $k \in \omega$  are such that  $x(p) = x'(p)$  for all  $p > k$ , then  $y_m^x = y_m^{x'}$  for all  $m > i_k$ . So again, if we let  $Z$  be the set of  $x$  for which  $y_0^x \in A$ ,  $Z$  cannot have the property of Baire, since whenever  $x$  and  $x'$  disagree at exactly one point, exactly one of them will be in  $Z$ .  $\square$

Theorem 2.2.6 has the following corollary, which shows that the nonselfdual Lipschitz classes and Wadge classes coincide.

**Corollary 2.2.7.** *Suppose that Lipschitz Determinacy + Baire( $\mathcal{P}(\omega^\omega)$ ) holds. Let  $A$  be a subset of  $\omega^\omega$ . If  $[A]_{\text{Li}}$  is nonselfdual, then  $[A]_{\text{Li}} = [A]_{\text{Wa}}$ .*

*Proof.* Supposing otherwise, fix  $B \in [A]_{\text{Wa}} \setminus [A]_{\text{Li}}$ . By Wadge's Theorem (Theorem 2.1.4), either  $(\omega^\omega \setminus A) \leq_L B$  or  $(\omega^\omega \setminus B) \leq_L A$ . Each of these implies that  $[A]_{\text{Wa}}$  is selfdual, giving a contradiction by Theorem 2.2.6.  $\square$

Summarizing, we have the following.

**Theorem 2.2.8.** *Suppose that Lipschitz Determinacy + Baire( $\mathcal{P}(\omega^\omega)$ ) holds.*

1. *The minimal Wadge classes consist of the singletons  $\{\emptyset\}$  and  $\{\omega^\omega\}$ .*
2. *Each selfdual Wadge class contains  $\aleph_1$  many selfdual Lipschitz classes, and these are ordered in ordertype  $\omega_1$  by  $\leq_L$ .*
3. *Each nonselfdual Wadge class is equivalent to the corresponding Lipschitz class, and has a selfdual class as an immediate successor.*
4. *A Wadge class which is neither minimal nor a successor is selfdual if and only if it is the supremum of a countable set of classes strictly below it.*

As we show in Proposition 2.5.4, a straightforward diagonal argument (from [36]) shows that there is no largest Wadge degree, if Lipschitz Determinacy holds.

**2.2.9 Remark.** The material in this section does not give a definition for the least Wadge class above a given selfdual class, or show (without assuming  $\text{DC}_{\mathbb{R}}$ ) that such a class exists. Adding  $\text{DC}_{\mathbb{R}}$  to the hypotheses of Theorem 2.2.8  $[A]_{\text{Wa}}$  has a pair of nonselfdual classes as immediate successors, the Wadge classes corresponding to the  $<_{\text{Li}}$ -least Lipschitz classes above all the Lipschitz classes contained in  $[A]_{\text{Wa}}$ .

## 2.3 Pointclasses

The notions of Wadge reducibility and Wadge rank naturally generalize to other topological spaces. In general, we say that for any pair of topological spaces  $X$  and  $Y$ , and any sets  $A \subseteq X$  and  $B \subseteq Y$ , that  $A \leq_{\text{Wa}} B$  if there is a continuous function  $f: X \rightarrow Y$  such that  $A = f^{-1}[B]$ . We will be concerned only with topological spaces of the form  $X_1 \times \cdots \times X_n$  for some positive  $n \in \omega$ , where at least one  $X_i$  is  $\omega^\omega$ , and each  $X_i$  is either  $\omega^\omega$  or  $\omega$ . Throughout this book, we let  $\mathcal{X}$  be the collection of such spaces; these spaces are all homeomorphic with  $\omega^\omega$ .

A *pointset* is a subset of a member of  $\mathcal{X}$ . A *pointclass* is a set of pointsets. A *boldface pointclass* is a pointclass closed under continuous preimages, i.e., the union of an initial segment of the Wadge hierarchy on pointsets.

The *cofinality* of a pointclass  $\Gamma$  is the least ordinal cardinality of a cofinal subset of  $\{[A]_{\text{Wa}} : A \in \Gamma \cap \mathcal{P}(\omega^\omega)\}$  under the order induced by  $<_{\text{Wa}}$ , if any such cardinality exists. We say that  $\Gamma$  has uncountably cofinality if its cofinality is not countable.

Given a finite sequence  $s \in {}^{<\omega}\omega$ , we let  $[s] = \{x \in \omega^\omega \mid x \restriction |s| = s\}$ . A *basic open interval* of a space  $X_1 \times \cdots \times X_n$  in  $\mathcal{X}$  is a product of the form  $a_1 \times \cdots \times a_n$ , where, for each  $i \in \{1, \dots, n\}$ ,

- $a_i$  is of the form  $[s]$ , for some  $s \in {}^{<\omega}\omega$ , if  $X_i = \omega^\omega$ ;
- $a_i$  is either  $\emptyset$ ,  $\omega$  or  $\{m\}$  for some  $m \in \omega$  if  $X_i = \omega$ .

We say that continuous functions  $f: X_1 \times \cdots \times X_n \rightarrow Y_1 \times \cdots \times Y_m$  between spaces in  $\mathcal{X}$  is *recursive* if the set of pairs of basic open intervals  $U \subseteq X$ ,  $V \subseteq Y$  for which  $f[U] \subseteq V$  is recursive (i.e.,  $\Delta_1$ -definable over HF). A *lightface pointclass* is a pointclass closed under preimages of continuous functions which are recursive. Under these definitions boldface pointclasses are also lightface. Our definitions may be nonstandard; in any case, lightface pointclasses are rarely mentioned in this book.

Given  $X \in \mathcal{X}$  and  $A \subseteq X$ , we write  $\check{A}$  for  $X \setminus A$ . Given a pointclass  $\Gamma$ , we write  $\check{\Gamma}$  for  $\{\check{A} : A \in \Gamma\}$ . We say that a pointclass  $\Gamma$  is *selfdual* if  $\Gamma = \check{\Gamma}$ ; otherwise it is *nonselfdual*. Observe that  $\Gamma$  is a boldface pointclass if and only if  $\check{\Gamma}$  is.

**2.3.1 Example.** The collection of analytic subsets of spaces in  $\mathcal{X}$  (i.e.,  $\Sigma_1^1$ ) is a nonselfdual boldface pointclass, as are the projective pointclasses  $\Sigma_n^1$  and  $\Pi_n^1$  for all  $n \in \omega$ . The pointclasses and  $\Delta_n^1 = \Sigma_n^1 \cap \Pi_n^1$  are boldface and selfdual. Assuming  $\text{CC}_{\mathbb{R}}$ , all of the classes mentioned in this example are closed under countable unions and intersections.

The following is an immediate, but useful, corollary of Theorem 2.1.4 (and Remark 2.1.1).

**Corollary 2.3.2** (ZF + Lipschitz Determinacy). *Let  $\Gamma$  be a nonselfdual boldface pointclass, and suppose that  $\Lambda$  is a boldface pointclass properly containing  $\Gamma$ . Then  $\check{\Gamma} \subseteq \Lambda$ .*

A member  $A$  of a pointclass  $\Gamma$  is *complete* for  $\Gamma$  (or  $\Gamma$ -*complete*) if every member of  $\Gamma$  is a continuous preimage of (e.g., Wadge below)  $A$ .

**Proposition 2.3.3.** *Suppose that Lipschitz Determinacy + Baire( $\mathcal{P}(\omega^\omega)$ ) holds. If  $\Gamma$  is a boldface pointclass, then every member of  $\Gamma \setminus \check{\Gamma}$  is  $\Gamma$ -complete.*

*Proof.* Fix a set

$$A \in \mathcal{P}(\omega^\omega) \cap (\Gamma \setminus \check{\Gamma}).$$

Then  $[A]_{\text{Li}}$  is nonselfdual, so  $[A]_{\text{Li}} = [A]_{\text{Wa}}$ , by Corollary 2.2.7. Suppose towards a contradiction that there is a  $B \in \Gamma$  such that  $B \not\leq_W A$ . Then, by Theorem 2.1.4,  $A \leq_W (\omega^\omega \setminus B)$ , contradicting our assumption that  $A \notin \check{\Gamma}$ .  $\square$

We write  $\omega^\omega \times \mathcal{X}$  for the set of  $X \in \mathcal{X}$  of the form

$$\omega^\omega \times X_1 \times \cdots \times X_n,$$

for some  $X_1, \dots, X_n$  (all equal to either  $\omega$  or  $\omega^\omega$ ). Given  $X \in \omega^\omega \times \mathcal{X}$  and  $A \subseteq X$ , we write  $\exists^{\omega^\omega} A$  for the set

$$\{(x_1, \dots, x_n) : \exists x_0 \in \omega^\omega (x_0, \dots, x_n) \in A\},$$

and  $\forall^{\omega^\omega} A$  for the set

$$\{(x_1, \dots, x_n) : \forall x_0 \in \omega^\omega (x_0, \dots, x_n) \in A\}.$$

Given a pointclass  $\Gamma$ , we write  $\exists^{\omega^\omega} \Gamma$  for

$$\{\exists^{\omega^\omega} A : A \in \Gamma \wedge \exists X \in \omega^\omega \times \mathcal{X} A \subseteq X\}$$

and  $\forall^{\omega^\omega} \Gamma$  for

$$\{\forall^{\omega^\omega} A : A \in \Gamma \wedge \exists X \in \omega^\omega \times \mathcal{X} A \subseteq X\}.$$

Similarly, for any ordinal  $\delta$ , we write  $\bigcup_\delta \Gamma$  for the collection of sets of the form  $\bigcup_{\alpha < \delta} A_\alpha$ , where each  $A_\alpha$  is in  $\Gamma$ , and the  $A_\alpha$ 's are all subsets of the same element of  $\mathcal{X}$ .

A pointclass  $\Gamma$  is  $\exists^{\omega^\omega}$ -closed if  $\exists^{\omega^\omega} \Gamma \subseteq \Gamma$ , and  $\forall^{\omega^\omega}$ -closed if  $\forall^{\omega^\omega} \Gamma \subseteq \Gamma$ .

**Proposition 2.3.4** (ZF +  $\text{CC}_\mathbb{R}$ ). *If  $\Gamma$  is an  $\exists^{\omega^\omega}$ -closed boldface pointclass with a complete set, then  $\Gamma$  is closed under countable unions. If in addition Lipschitz Determinacy holds, and  $\Delta$  is a selfdual  $\exists^{\omega^\omega}$ -closed boldface pointclass with uncountable cofinality, then  $\Delta$  is closed under countable unions and countable intersections.*

*Proof.* We prove the first part first. Let  $A \subseteq \omega^\omega$  be  $\Gamma$ -complete, and let  $B_i \subseteq \omega^\omega$  ( $i \in \omega$ ) be elements of  $\Gamma$ . Applying  $\text{CC}_\mathbb{R}$ , for each  $i$ , let  $f_i$  be a continuous function such that  $B_i = f_i^{-1}[A]$ . Let  $C$  be  $\{(x, y) \in \omega^\omega \times \omega^\omega : y \in B_{x(0)}\}$  and define  $g : \omega^\omega \times \omega^\omega \rightarrow \omega^\omega$  by setting  $g(x, y)$  to be  $f_{x(0)}(y)$ . Then  $C = g^{-1}[A]$ , so  $C \in \Gamma$ . Since  $\bigcup_{i \in \omega} B_i = \{y : \exists x \in \omega^\omega (x, y) \in C\}$ , we are done.

For the second part, the assumptions imply that every countable subset of  $\Delta$  is contained in a boldface pointclass  $\Gamma_0$  having a complete set. Then  $\exists^{\omega^\omega} \Gamma_0$  is contained in  $\Delta$  and satisfies the assumptions of the first part.  $\square$

**2.3.5 Remark.** The proof of Proposition 2.3.4 shows the following, without the assumption of  $\text{CC}_\mathbb{R}$

- If  $\Gamma$  is an  $\exists^{\omega^\omega}$ -closed boldface pointclass with a complete set, then  $\Gamma$  is closed under unions.
- If Lipschitz Determinacy holds and  $\Delta$  is a selfdual  $\exists^{\omega^\omega}$ -closed boldface pointclass, then either  $\Delta$  is closed under unions and intersections or there exists an  $A \subseteq \omega^\omega$  such that  $\Delta$  is the collection of pointsets which are Wadge reducible to either  $A$  or  $\omega^\omega \setminus A$ .

**2.3.6 Remark.** Let  $\Gamma$  be a boldface pointclass. If  $\Gamma$  is closed under unions, then so are  $\exists^{\omega^\omega} \Gamma$  and  $\forall^{\omega^\omega} \Gamma$  (the latter case can be verified by coding a pair of reals with a single real). Similarly, if  $\Gamma$  is closed under countable unions (and  $\text{CC}_\mathbb{R}$  holds, for the  $\forall^{\omega^\omega}$  case), then so are  $\exists^{\omega^\omega} \Gamma$  and  $\forall^{\omega^\omega} \Gamma$ .

## 2.4 Universal sets

Given an integer  $n \in \omega \setminus 2$ , a set  $A \subseteq (\omega^\omega)^n$  is *universal* for a pointclass  $\Gamma$  (or  $\Gamma$ -*universal*) if  $A \in \Gamma$ , and for every  $B \subseteq (\omega^\omega)^{n-1}$  in  $\Gamma$  there is an  $x \in \omega^\omega$  such that

$$B = \{(y_1, \dots, y_{n-1}) \mid (x, y_1, \dots, y_{n-1}) \in A\}$$

(we call this set  $A_x$ ). If  $\Gamma$  is closed under Wadge-equivalence, and  $n$  is in  $\omega \setminus 2$ , then there exists a universal subset of  $(\omega^\omega)^n$  in  $\Gamma$  if and only if there is a universal subset of  $(\omega^\omega)^2$  in  $\Gamma$ .

**2.4.1 Example.** Fixing an enumeration  $\langle \sigma_n : n \in \omega \rangle$  of  $\omega^{<\omega}$ , the set of  $(x, y)$  in  $(\omega^\omega)^2$  such that  $y \in [\sigma_n]$  for some  $n \in x^{-1}[\{0\}]$  is a universal open set. It follows that there exist a universal closed set  $C \subseteq (\omega^\omega)^3$ , and universal analytic and coanalytic subsets of  $(\omega^\omega)^2$ .

The usual diagonal argument shows that selfdual lightface pointclasses do not have universal sets.

**Proposition 2.4.2.** *If  $\Delta$  is a selfdual lightface pointclass, then  $\Delta$  does not have a universal set.*

*Proof.* Given a set  $A \subseteq (\omega^\omega)^2$  in  $\Delta$ , let  $B$  be the set of  $x \in \omega^\omega$  for which  $(x, x) \notin A$ . Then for any  $x \in \omega^\omega$ ,  $x \in B$  if and only if  $x \notin A_x$ .  $\square$

Nonselfdual boldface pointclasses have universal sets, under suitable determinacy assumptions.

**Theorem 2.4.3.** *Suppose that Lipschitz Determinacy + Baire( $\mathcal{P}(\omega^\omega)$ ) holds. If  $\Gamma$  is a boldface pointclass, then  $\Gamma$  has a universal set if and only if it is nonselfdual.*

*Proof.* The selfdual case follows from Proposition 2.4.2.

For the other case, suppose that  $\Gamma$  is nonselfdual, and fix a set

$$A \in \mathcal{P}(\omega^\omega) \cap (\Gamma \setminus \check{\Gamma}).$$

Then  $[A]_{\text{Li}}$  is nonselfdual, so  $[A]_{\text{Li}} = [A]_{\text{Wa}}$ , by Corollary 2.2.7, and  $A$  is  $\Gamma$ -complete by Proposition 2.3.3. Fix a recursive bijection  $\rho: \omega^{<\omega} \rightarrow \omega$ . For each  $x \in \omega^\omega$ , define  $f_x: \omega^\omega \rightarrow \omega^\omega$  by setting  $f_x(y)(n)$  to be  $x(\rho(y \upharpoonright (n+1)))$ .

Then each  $f_x$  is Lipschitz, and each Lipschitz function from  $\omega^\omega$  to  $\omega^\omega$  is equal to  $f_x$  for some  $x \in \omega^\omega$ . Now let  $U$  be the set of  $(x, y) \in (\omega^\omega)^2$  such that  $f_x(y) \in A$ . Then  $U =_{\text{Wa}} A$ , and  $U$  is  $\Gamma$ -universal.  $\square$

**2.4.4 Remark.** Suppose that  $\Gamma$  is a boldface pointclass and  $U \subseteq (\omega^\omega)^n$  is  $\Gamma$ -universal, for some  $n \in \omega \setminus 3$ . Then

$$\{(z_1, \dots, z_{n-2}) \in (\omega^\omega)^{n-2} : \exists y \in \omega^\omega (x, y, z_1, \dots, z_{n-2}) \in U\}$$

is universal for  $\exists^\omega \Gamma$  and

$$\{(z_1, \dots, z_{n-2}) \in (\omega^\omega)^{n-2} : \forall y \in \omega^\omega (x, y, z_1, \dots, z_{n-2}) \in U\}$$

is universal for  $\forall^{\omega^\omega} \Gamma$ . It follows from Theorem 2.4.3 that if  $\Gamma$  is nonselfdual, then so are  $\exists^{\omega^\omega} \Gamma$  and  $\forall^{\omega^\omega} \Gamma$  (although either or both of these pointclasses could be equal to  $\Gamma$ ).

In order to prove the Moschovakis Coding Lemma (in Chapter 3) we will need sequences of universal sets in different dimensions which are suitably coherent.

**2.4.5 Definition.** Let  $\bar{U} = \langle U_n : n \in \omega \setminus \{0\} \rangle$  be such that each  $U_n$  is a subset of  $(\omega^\omega)^{n+1}$ .

- The sequence  $\bar{U}$  has the *s-m-n property* if for each pair of positive integers  $n < m$ , there exists a continuous  $s_{m,n} : (\omega^\omega)^{n+1} \rightarrow \omega^\omega$  such that, for all  $x, y_1, \dots, y_m \in \omega^\omega$ ,

$$(x, y_1, \dots, y_m) \in U_m$$

if and only if  $(s_{m,n}(x, y_1, \dots, y_n), y_{n+1}, \dots, y_m) \in U_{m-n}$ .

- The sequence  $\bar{U}$  has the *recursion property* (with respect to a pointclass  $\Gamma$ ) if for each  $n \in \omega \setminus \{0\}$  and each

$$A \in \mathcal{P}((\omega^\omega)^{n+1}) \cap \Gamma,$$

there exists an  $x \in \omega^\omega$  such that for all  $y_1, \dots, y_n \in \omega^\omega$ ,

$$(x, y_1, \dots, y_n) \in U_n$$

if and only if  $(x, y_1, \dots, y_n) \in A$  (i.e., such that  $U_{n,x} = A_x$ ).

We will omit the phrase “with respect to  $\Gamma$ ” when talking about sequences  $\bar{U}$  with the recursion property, since  $\Gamma$  is recoverable from  $\bar{U}$ . Similarly, when we say that  $\langle U_n : n \in \omega \setminus \{0\} \rangle$  is a sequence of sets with the s-m-n property, we will mean that each  $U_n$  is a subset of  $(\omega^\omega)^{n+1}$ .

**2.4.6 Remark.** When  $U_n$  is a universal set, the recursion property as defined above is a fixed point property. By the universality of  $U_n$  there is, for each set  $A$  as in the definition, and each  $x \in \omega^\omega$ , a  $z \in \omega^\omega$  such that  $A_x = U_{n,z}$ . The point of the definition is that there is some  $x$  such that  $A_x = U_{n,x}$ .

Theorems 2.4.7 and 2.4.8 below show that if  $\Gamma$  is a boldface pointclass with a universal set, then there exists a sequence of  $\Gamma$ -universal sets with the s-m-n and recursion properties. The statement of Theorem 2.4.7, and its proof, are taken from [6].

**Theorem 2.4.7.** *If  $\Gamma$  is a boldface pointclass with a universal set then there exists a sequence of  $\Gamma$ -universal sets with the s-m-n property.*

*Proof.* Fix homeomorphisms  $\pi_n : \omega^\omega \rightarrow (\omega^\omega)^n$  for each  $n \in \omega \setminus 2$ , and let  $\pi_{m,n} : \omega^\omega \rightarrow \omega^\omega$  ( $n < m \in \omega \setminus 2$ ) be such that

$$\pi_m(x) = (\pi_{m,0}(x), \dots, \pi_{m,m-1}(x))$$



for all  $x \in \omega^\omega$ , so that  $\pi_{m,n}(\pi_m^{-1}(x_0, \dots, x_{m-1})) = x_n$  for all  $x_0, \dots, x_{m-1} \in \omega^\omega$ . Let  $U \subseteq (\omega^\omega)^2$  be a universal set for  $\Gamma$ . For each  $n \in \omega \setminus \{0\}$ , let  $U_n$  be the set of  $(x, y_1, \dots, y_n) \in (\omega^\omega)^{n+1}$  such that

$$(\pi_{2,0}(x), \pi_{n+1}^{-1}(\pi_{2,1}(x), y_1, \dots, y_n)) \in U.$$

We check first that each  $U_n$  is  $\Gamma$ -universal. Fix  $n \in \omega \setminus \{0\}$  and  $A \subseteq (\omega^\omega)^n$  in  $\Gamma$ . We want to find an  $x \in \omega^\omega$  such that  $U_{n,x} = A$ . Fixing any  $z \in \omega^\omega$ , we have that  $\pi_{n+1}^{-1}[\{(z, y_1, \dots, y_n) : (y_1, \dots, y_n) \in A\}]$  is in  $\Gamma$ . Since  $U$  is universal, there is a  $w \in \omega^\omega$  such that  $U_w = \pi_{n+1}^{-1}[\{(z, y_1, \dots, y_n) : (y_1, \dots, y_n) \in A\}]$ . Let  $x = \pi_2^{-1}(w, z)$ . Then for all  $y_1, \dots, y_n \in \omega^\omega$ ,  $(x, y_1, \dots, y_n)$  is in  $U_n$  if and only if  $(w, \pi_{n+1}^{-1}(z, y_1, \dots, y_n))$  is in  $U$ , which holds if and only if  $(y_1, \dots, y_n)$  is in  $A$ .

To check that the s-m-n property holds, fix  $m > n$  in  $\omega$ . Let  $W$  be the set of  $w \in \omega^\omega$  for which

$$(u(w), \pi_{m+1}^{-1}(v(w), r_1(w), \dots, r_n(w), t_1(w), \dots, t_{m-n}(w))) \in U,$$

where

- $u(w) = \pi_{2,0}(\pi_{n+1,0}(\pi_{m-n+1,0}(w)))$ ;
- $v(w) = \pi_{2,1}(\pi_{n+1,0}(\pi_{m-n+1,0}(w)))$ ;
- $r_i(w) = \pi_{n+1,i}(\pi_{m-n+1,0}(w))$  for  $i \in \{1, \dots, n\}$ ;
- $t_j(w) = \pi_{m-n+1,j}(w)$  for  $j \in \{1, \dots, m-n\}$ .

Then  $W \leq_{\text{Wa}} U$ , and, as  $U$  is universal for  $\Gamma$ , there exists a  $z \in \omega^\omega$  such that  $U_z = W$ .

Define  $s_{m,n} : (\omega^\omega)^{n+1} \rightarrow \omega^\omega$  by setting

$$s_{m,n}(x, y_1, \dots, y_n) = \pi_2^{-1}(z, \pi_{n+1}^{-1}(x, y_1, \dots, y_n))$$

for all  $x, y_1, \dots, y_n \in \omega^\omega$ . Now fix  $x, y_1, \dots, y_m \in \omega^\omega$ . Then  $(x, y_1, \dots, y_m) \in U_m$  if and only if

$$(\pi_{2,0}(x), \pi_{m+1}^{-1}(\pi_{2,1}(x), y_1, \dots, y_m)) \in U,$$

and  $(s_{m,n}(x, y_1, \dots, y_n), y_{n+1}, \dots, y_m) \in U_{m-n}$  if and only if

$$(\pi_{2,0}(s_{m,n}(x, y_1, \dots, y_n)), \pi_{m-n+1}^{-1}(\pi_{2,1}(s_{m,n}(x, y_1, \dots, y_n)), y_{n+1}, \dots, y_m)) \in U.$$

Now,  $\pi_{2,0}(s_{m,n}(x, y_1, \dots, y_n)) = z$  and

$$\pi_{2,1}(s_{m,n}(x, y_1, \dots, y_n)) = \pi_{n+1}^{-1}(x, y_1, \dots, y_n).$$

Since  $U_z = W$ , we have that

$$(z, \pi_{m-n+1}^{-1}(\pi_{n+1}^{-1}(x, y_1, \dots, y_n), y_{n+1}, \dots, y_m)) \in U$$

if and only if  $\pi_{m-n+1}^{-1}(\pi_{n+1}^{-1}(x, y_1, \dots, y_n), y_{n+1}, \dots, y_m) \in W$ , which, letting  $w$  be  $\pi_{m-n+1}^{-1}(\pi_{n+1}^{-1}(x, y_1, \dots, y_n), y_{n+1}, \dots, y_m)$ , holds if and only if

$$(u(w), \pi_{m+1}^{-1}(v(w), r_1(w), \dots, r_n(w), t_1(w), \dots, t_{m-n}(w))) \in U,$$

which holds if and only if

$$(\pi_{2,0}(x), \pi_{m+1}^{-1}(\pi_{2,1}(x), y_1, \dots, y_m)) \in U,$$

since

- $u(w) = \pi_{2,0}(x)$ ,
- $v(w) = \pi_{2,1}(x)$ ,
- $r_i(w) = y_i$  for  $i \in \{1, \dots, n\}$  and
- $t_j(w) = y_{n+i}$  for  $j \in \{1, \dots, m-n\}$ .

□

The following is a version of Kleene's Recursion Theorem, saying that a sequence of  $\Gamma$ -universal sets with the s-m-n property has the recursion property.

**Theorem 2.4.8** (Kleene). *If  $\Gamma$  is a pointclass and  $\bar{U} = \langle U_n : n \in \omega \setminus \{0\} \rangle$  is a sequence of  $\Gamma$ -universal sets with the s-m-n property, then  $\bar{U}$  has the recursion property.*

*Proof.* Fix  $n \in \omega \setminus \{0\}$  and  $A \in \mathcal{P}((\omega^\omega)^{n+1}) \cap \Gamma$ . Applying the assumption that  $U_{n+1}$  is  $\Gamma$ -universal, let  $y \in \omega^\omega$  be such that for all  $w, z_1, \dots, z_n$  in  $\omega^\omega$ ,  $(y, w, z_1, \dots, z_n) \in U_{n+1}$  if and only if  $(s(w, w), z_1, \dots, z_n) \in A$ , where  $s : (\omega^\omega)^2 \rightarrow \omega^\omega$  witnesses the s-m-n property for  $\bar{U}$  in the role of  $s_{n+1,1}$ . Then for all  $w, z_1, \dots, z_n \in \omega^\omega$ ,

$$(s(y, w), z_1, \dots, z_n) \in U_n$$

if and only if

$$(y, w, z_1, \dots, z_n) \in U_{n+1}$$

if and only if

$$(s(w, w), z_1, \dots, z_n) \in A.$$

Then  $x = s(y, y)$  is as desired. □

Putting together Theorems 2.4.3, 2.4.7 and 2.4.8 we get the following.

**Theorem 2.4.9.** *Suppose that Lipschitz Determinacy + Baire( $\mathcal{P}(\omega^\omega)$ ) holds. If  $\Gamma$  is a nonselfdual boldface pointclass, then there exists a sequence of universal  $\Gamma$ -sets with the s-m-n and recursion properties.*

Theorem 2.4.10 below is another version of the Recursion Theorem. A function  $f : \omega^\omega \rightarrow \omega^\omega$  is in  $\Sigma_1^1$  if it is  $\Sigma_1^1$  as a subset of  $\omega^\omega \times \omega^\omega$  (equivalently, if the  $f$ -preimage of each open set is in  $\Sigma_1^1$ ).

**Theorem 2.4.10** (Kleene). *Suppose that  $\Gamma$  is an  $\exists^{\omega^\omega}$ -closed pointclass and that  $\bar{U} = \langle U_n : n \in \omega \setminus \{0\} \rangle$  is a sequence of  $\Gamma$ -universal sets with the recursion property. For any  $m \in \omega \setminus \{0\}$  and any  $\Sigma_1^1$  function  $f : \omega^\omega \rightarrow \omega^\omega$ , there is an  $x \in \omega^\omega$  such that  $U_{m,x} = U_{m,f(x)}$ .*

*Proof.* Since  $\Gamma$  is  $\exists^{\omega^\omega}$ -closed, the set

$$A = \{(x, y_1, \dots, y_m) \in (\omega^\omega)^3 : (f(x), y_1, \dots, y_m) \in U_m\}$$

is in  $\Gamma$ . Then for each  $x \in \omega^\omega$ ,  $A_x = U_{m,f(x)}$ . Applying the recursion property for  $U_m$ , we get an  $x \in \omega^\omega$  such that  $U_{m,x} = A_x$ .  $\square$

Using a recursive bijection  $\pi : \omega \rightarrow \omega \times \omega$ , we can associate to each  $y \in \omega^\omega$  an  $\omega$ -sequence  $\langle (y)_n : n \in \omega \rangle$  of elements of  $\omega^\omega$  by setting  $(y)_n(m)$  to be  $y(\pi^{-1}(n, m))$ . Loosely following 7D.7 (page 430) of [32], we say that a set  $B \subseteq X$  (for some  $X$  in  $\mathcal{X}$ ) is in  $\text{pos-}\Sigma_1^1(A)$  (when  $A$  is a nonempty subset of  $\omega^\omega$ ) if

$$B = \{x \in X : \exists y \in \omega^\omega ((\forall n \in \omega (y)_n \in A) \wedge (x, y) \in S)\},$$

for some  $\Sigma_1^1$  set  $S \subseteq X \times \omega^\omega$ . We define  $\text{pos-}\Sigma_1^1(\emptyset)$  to be  $\Sigma_1^1$ . Then  $\text{pos-}\Sigma_1^1(A)$  is the pointclass of sets which are  $\Sigma_1^1$  using a predicate for  $A$  positively. If  $A$  is a subset of  $(\omega^\omega)^n$  for some positive  $n \in \omega$ , we say that a set  $B \subseteq X$  is in  $\text{pos-}\Sigma_1^1(A)$  if it is in  $\text{pos-}\Sigma_1^1(b[A])$  for some recursive bijection  $b : (\omega^\omega)^n \rightarrow \omega^\omega$ . If  $A_1, \dots, A_n$  are subsets of  $(\omega^\omega)^n$ , we write  $\text{pos-}\Sigma_1^1(A_1, \dots, A_n)$  for  $\text{pos-}\Sigma_1^1(A)$ , where  $A$  is the disjoint union of  $A_1, \dots, A_n$ , i.e., the set

$$\{\langle \pi(i, x(0)), x(1), x(2), \dots \rangle : i \in \{1, \dots, n\}, x \in A_i\},$$

which we write as  $A_1 \oplus \dots \oplus A_n$ .

**2.4.11 Remark.** For any  $X \in \mathcal{X}$  and  $A \subseteq \omega^\omega$ ,  $A$  is in  $\text{pos-}\Sigma_1^1(A)$ , and  $\text{pos-}\Sigma_1^1(A)$  is a boldface pointclass closed under  $\exists^{\omega^\omega}$ , countable unions and countable intersections (assuming  $\text{CC}_{\mathbb{R}}$ ). The existence of universal sets for  $\Sigma_1^1$  (as shown in Example 2.4.1) implies that each pointclass of the form  $\text{pos-}\Sigma_1^1(A)$  has a universal set.

Given  $A \subseteq \omega^\omega$ ,  $\Sigma_1^1(A)$  the collection of subsets of spaces in  $\mathcal{X}$  which can be defined by a  $\Sigma_1^1$  formula in a predicate for  $A$ . This is the same as the pointclass  $\text{pos-}\Sigma_1^1(A, \omega^\omega \setminus A)$ . Similarly, we write  $\Sigma_1^1(A_1, \dots, A_n)$  for

$$\text{pos-}\Sigma_1^1(A_1, \dots, A_n, \omega^\omega \setminus A_1, \dots, \omega^\omega \setminus A_n).$$

Given a positive  $n \in \omega$ ,  $\Pi_n^1(A)$  is the set of complements of sets in  $\Sigma_1^1(A)$ ,  $\Sigma_{n+1}^1(A)$  is the set of continuous images of sets in  $\Pi_n^1(A)$  and  $\Delta_n^1(A)$  is the intersection of  $\Sigma_n^1(A)$  and  $\Pi_n^1(A)$ . We say that a set  $B$  is *projective in  $A$*  if it is in  $\bigcup_{n \in \omega} \Sigma_{n+1}^1(A)$  and write  $\Delta_\omega(A)$  for the collection of sets projective in  $A$ . We say that a pointclass  $\Delta$  is *projectively closed* if for each  $A \in \Delta \cap \mathcal{P}(\omega^\omega)$ , every set projective in  $A$  is in  $\Delta$ .

**2.4.12 Remark.** Let  $\bar{U} = \langle U_n : n < \omega \rangle$  be a sequence of universal sets for  $\Sigma_1^1$ . For any  $A \subseteq \omega^\omega$ , let  $U_n(A)$  be the set

$$\{x \in (\omega^\omega)^{n+1} : \exists y \in \omega^\omega ((\forall n \in \omega (y)_n \in A) \wedge (x, y) \in U_{n+1})\}.$$

Then each  $U_n(A)$  is universal for  $\text{pos-}\Sigma_1^1(A)$ . Furthermore, if  $s_{m,n}$  ( $n < m < \omega$ ) are functions witnessing the s-m-n property for  $\bar{U}$ , then for all  $n < m < \omega$ , function  $s_{m+1,n}$  witnesses the s-m-n property for  $\bar{U}(A) = \langle U_n(A) : n < \omega \rangle$  for  $m$  and  $n$ . Similarly, given a function  $f: \omega^\omega \rightarrow \omega^\omega$  and  $n \in \omega \setminus \{0\}$ , if  $U_{n,x} = U_{n,f(x)}$ , then  $U_{n,x}(A) = U_n(A)_x = U_n(A)_{f(x)} = U_{n,f(x)}(A)$  for all  $A \subseteq \omega^\omega$ .

For  $\bar{U}$  as Remark 2.4.12,  $n \in \omega$  and  $A_1, \dots, A_m \subseteq \omega^\omega$ , we write  $U_n(A_1, \dots, A_m)$  for  $U_n(A_1 \oplus \dots \oplus A_m)$ .

## 2.5 The cardinal $\Theta$

This section presents results of Solovay on the height of the Wadge hierarchy, taken from [36].

**2.5.1 Definition.** The ordinal  $\Theta$  is defined to be the least nonzero ordinal which is not a surjective image of  $\mathcal{P}(\omega)$ .

It follows immediately from this definition that  $\Theta$  is a cardinal.

**2.5.2 Remark.** We will often making use of the fact that continuous functions on  $\omega^\omega$  can be coded by members of  $\omega^\omega$ . While there are many ways of doing this, we fix one for concreteness and convenience to use throughout the book. Recall from Example 2.4.1 that there is a universal set  $C \subseteq (\omega^\omega)^3$  for the pointclass of closed subsets of spaces in  $\mathcal{X}$ . For each continuous function  $f: \omega^\omega \rightarrow \omega^\omega$ , there is an  $x \in \omega^\omega$  such that  $C_x = f$ . Let  $\mathcal{F}^c$  be the set of  $x$  for which  $C_x$  is a continuous function from  $\omega^\omega$  to  $\omega^\omega$ , and for each  $x \in \mathcal{F}^c$  let  $f_x^c$  denote the set  $C_x$ . Then  $\mathcal{F}^c$  is in  $\Pi_1^1$  (the hardest part of this may be checking that the domain of  $f_x^c$  is total; see Exercise 33.1(i) of [19]). Using universal closed sets  $C_k \subseteq (\omega^\omega)^{k+3}$  ( $k \in \omega$ ), we can in a similar fashion fix for each pair  $(n, m) \in \omega \times \omega$  a set  $\mathcal{F}^{c,n,m}$  of codes  $x$  for all continuous functions  $f_x^{c,n,m}: (\omega^\omega)^n \rightarrow (\omega^\omega)^m$ .

**2.5.3 Remark.** The usual diagonal argument shows that it is not possible to have an association of each  $x \in \omega^\omega$  to a continuous function  $f_x^*: \omega^\omega \rightarrow \omega^\omega$  in such a way that the map  $(x, y) \mapsto f_x^*(y)$  is continuous on  $(\omega^\omega)^2$ . To see this, consider the function  $g: \omega^\omega \rightarrow \omega^\omega$  defined by setting  $g(x)(n) = f_x^*(x)(n) + 1$  for each  $x \in \omega^\omega$ , for any such map  $x \mapsto f_x^*$ .

Another diagonalization argument gives a Wadge class above a given selfdual class (recall from Section 2.2 that there is a natural way to define the least Wadge class above a pair of complementary nonselfdual classes).

**Theorem 2.5.4 (Solovay).** *There a function  $j: \mathcal{P}(\omega^\omega) \rightarrow \mathcal{P}(\omega^\omega)$  such that, for all  $A \subseteq \omega^\omega$ ,  $A <_{\text{Wa}} j(A)$ .*

*Proof.* Let  $C$  be the fixed universal set from Remark 2.5.2. Given  $A \subseteq \omega^\omega$ , let  $B$  be the set of  $x \in \mathcal{F}^c$  such that, for all  $y \in A$ ,  $(x, x, y) \notin C$  (so if  $x \in \mathcal{F}^c$ , then  $x \notin (f_x^c)^{-1}[A]$ ). Let  $j(A)$  be the disjoint union  $A \oplus B$ .  $\square$

**2.5.5 Remark.** The proof of Theorem 2.5.4 shows that if  $A \subseteq \omega^\omega$  and  $\Delta$  is a selfdual boldface pointclass containing  $A$  and  $C$ , then  $j(A)$  is in  $\forall^{\omega^\omega} \Delta$ .

The proof of Theorem 2.5.4 does not show how high the Wadge rank of  $j(A)$  is relative to  $A$ . In particular, we have the following question (one could ask a similar question about the limit stages of the construction in the proof of Theorem 2.5.9).

**2.5.6 Question.** Does AD imply that the Wadge rank of  $j(A)$  is defined whenever  $A$  is a subset of  $\omega^\omega$  whose Wadge rank is defined?

The main result of this section is the following fact. One direction is proved in Proposition 2.5.8, the other in Theorem 2.5.9.

**Theorem 2.5.7** (Solovay). *The ordinal  $\Theta$  is the supremum of the set of ordinals  $\gamma$  for which there exists a  $<_{\text{Wa}}$ -increasing sequence*

$$\langle A_\alpha : \alpha < \gamma \rangle.$$

If one assumes in addition that  $<_{\text{Wa}}$  is wellfounded, then one gets that  $\Theta$  is the supremum of  $\{\text{WR}(A) : A \subseteq \omega^\omega\}$ . This is Corollary 2.5.11.

The following proposition shows that if  $\langle A_\alpha : \alpha \leq \gamma \rangle$  is a  $\leq_{\text{Wa}}$ -increasing sequence, then there is a surjection from  $\omega^\omega$  to  $\gamma + 1$  defined by mapping  $x \in \omega^\omega$  to  $\alpha$  if

$$[A_\alpha]_{\text{Wa}} = [(f_x^c)^{-1}[A_\gamma]]_{\text{Wa}}$$

(if there exists such an  $\alpha$ , and 0 otherwise). It follows that  $\gamma < \Theta$ . It shows moreover that for each  $A \subseteq \omega^\omega$  whose Wadge rank is defined,  $\text{WR}(A) < \Theta$ .

**Proposition 2.5.8.** *Let  $A$  be a subset of  $\omega^\omega$ , and let  $\Delta$  be the smallest selfdual boldface pointclass containing  $A$  and  $\mathcal{F}^c$ . Let  $\leq$  be the order on  $\mathcal{F}^c$  defined by setting  $x \leq y$  if and only if*

$$(f_x^c)^{-1}[A] \leq_{\text{W}} (f_y^c)^{-1}[A].$$

*Then  $(\mathcal{F}^c, \leq)$  is isomorphic to  $(\{B \subseteq \omega^\omega : B \leq_{\text{W}} A\}, \leq_{\text{W}})$ , and  $\leq$  is in  $\exists^{\omega^\omega} \forall^{\omega^\omega} \Delta$ .*

*Proof.* That  $(\mathcal{F}^c, \leq)$  is isomorphic to  $(\{B \subseteq \omega^\omega : B \leq_{\text{W}} A\}, \leq_{\text{W}})$  is immediate from the definitions. That  $\leq$  is in  $\exists^{\omega^\omega} \forall^{\omega^\omega} \Delta$  follows from noting that  $x \leq y$  if and only if  $\{x, y\} \subseteq \mathcal{F}^c$  and there exists a  $u \in \mathcal{F}^c$  such that for all  $z, w, v, t$  in  $\omega^\omega$ , if  $w = f_x^c(z)$ ,  $v = f_u^c(z)$  and  $t = f_y^c(v)$ , then  $w \in A$  if and only if  $t \in A$  (that is,  $z \in (f_x^c)^{-1}[A]$  if and only if  $f_u^c(z) \in (f_y^c)^{-1}[A]$ ).  $\square$

Finally, we use Theorem 2.5.4 to build  $<_{\text{Wa}}$ -increasing sequences of any length less than  $\Theta$ .

**Theorem 2.5.9** (Solovay). *If  $\gamma$  is an ordinal and  $g: \omega^\omega \rightarrow \gamma$  is a surjection, then there is a  $<_{\text{Wa}}$ -increasing sequence  $\langle A_\alpha : \alpha < \gamma \rangle$  definable from  $g$ .*

*Proof.* Let  $g: \omega^\omega \rightarrow \gamma$  be a surjection, for some ordinal  $\gamma$ , and let  $\pi: \omega \times \omega \rightarrow \omega$  be a recursive bijection. Recursively define the sequence  $\langle A_\alpha : \alpha < \gamma \rangle$  by setting each  $A_\alpha$  to be

$$j(\pi[\{(x, y) \in \omega^\omega \times \omega^\omega \mid g(y) < \alpha \text{ and } x \in A_{g(y)}\}]),$$

where  $j$  is as in Theorem 2.5.4. □

**2.5.10 Remark.** Let  $\Delta$  be an  $\exists^{\omega^\omega}$ -closed selfdual pointclass containing  $\mathcal{F}^c$ , let  $\gamma$  be an ordinal such that  $\bigcup_\gamma \Delta \subseteq \Delta$  and let  $f: \omega^\omega \rightarrow \gamma$  be a surjection such that  $f^{-1}[\{\alpha\}] \in \Delta$  for all  $\alpha < \gamma$ . The proof of Theorem 2.5.9, along with Remark 2.5.5, shows that  $\Delta$  contains a set whose Wadge rank is at least  $\gamma$  (or undefined).

It follows that if the Wadge rank of each subset of  $\omega^\omega$  is defined, then  $\Theta$  is the supremum of the Wadge ranks of the elements of  $\mathcal{P}(\omega^\omega)$ .

**Corollary 2.5.11** (Solovay). *Suppose that Lipschitz Determinacy holds, and that  $\text{WR}(A)$  is defined for every  $A \subseteq \omega^\omega$ . Then  $\Theta = \{\text{WR}(A) \mid A \subseteq \omega^\omega\}$ .*

## Chapter 3

# Coding lemmas

This chapter presents the Moschovakis Coding Lemma, which is one of the fundamental theorems of the theory of determinacy and which will be used throughout this book. Among other things the Coding Lemma can be used to map  $\omega^\omega$  onto the powerset of any ordinal below  $\Theta$  (see Corollary 3.0.2). Theorem 3.0.1 is the basic form of the lemma. Theorem 3.0.3 is a uniform version, which will be needed in Chapter 5. We follow [22]. Section 7D of [32] also contains proofs of the theorems in this section.

Given a strategy  $\sigma: \omega^{<\omega} \rightarrow \omega$ , and  $x \in \omega^\omega$ , we let  $\sigma \circ x$  (similarly,  $x \circ \sigma$ ) be the sequence of moves played by player  $I$  ( $II$ ) when he plays according to  $\sigma$  and player  $II$  ( $I$ ) plays  $x$ .

**Theorem 3.0.1** (The Coding Lemma; Moschovakis). *Assume that AD holds. Let*

- $X$  be a subset of  $\omega^\omega$ ;
- $Z$  be a subset of  $X \times \omega^\omega$ ;
- $f$  be a function from  $X$  to the ordinals;
- $<_f$  be  $\{(y, z) \in X^2 : f(y) < f(z)\}$ .

For each  $y \in X$ , let  $[y]_f$  denote  $\{z \in X : f(y) = f(z)\}$ . Then there exists a  $\text{pos-}\Sigma_1^1(<_f)$  set  $A \subseteq Z$  such that for all  $y \in X$ ,

$$A \cap ([y]_f \times \omega^\omega) = \emptyset \text{ if and only if } Z \cap ([y]_f \times \omega^\omega) = \emptyset.$$

*Proof.* It suffices to consider the case where the range of  $f$  is an ordinal  $\gamma$ , and, proving the theorem by induction, we may assume that it holds for all smaller ordinals. As  $\text{pos-}\Sigma_1^1(<_f)$  is closed under unions and contains all countable subsets of  $\omega^\omega$ , we may assume that  $\gamma$  is an uncountable limit ordinal. It follows then that  $X$  is in  $\text{pos-}\Sigma_1^1(<_f)$ . Applying Theorem 2.4.7 and Remark 2.4.12, let  $\bar{U} = \langle U_n : n \in \omega \setminus \{0\} \rangle$  be a sequence of universal sets for  $\text{pos-}\Sigma_1^1(<_f)$  with the s-m-n property. We seek an  $x \in \omega^\omega$  such that

1.  $U_{2,x} \subseteq Z$ ;
2. for all  $y \in X$ , if  $Z \cap ([y]_f \times \omega^\omega) \neq \emptyset$  then  $U_{2,x} \cap ([y]_f \times \omega^\omega) \neq \emptyset$ .

Let  $Y = \{x \in \omega^\omega : U_{2,x} \subseteq Z\}$ . For each  $x \in \omega^\omega$ , let  $\alpha_x$  be the least value  $f(y)$  for any  $y \in X$  witnessing the failure of item (2) for  $x$ , if there exists such a  $y$ ; otherwise, let  $\alpha_x = \gamma$ . Consider the game between players  $I$  and  $II$  where player  $I$  builds  $x_1 \in \omega^\omega$ , player  $II$  builds  $x_2 \in \omega^\omega$ , and  $I$  wins if and only if  $x_1 \in Y$  and either  $x_2 \notin Y$  or  $\alpha_{x_1} \geq \alpha_{x_2}$ . We will show that a winning strategy for either player gives the desired conclusion.

First, suppose that  $\sigma$  is a winning strategy for player  $I$ . Then for all  $x \in \omega^\omega$ ,  $U_{2,\sigma \circ x} \subseteq Z$ . The set  $\bigcup_{x \in \omega^\omega} U_{2,\sigma \circ x}$  is in  $\text{pos-}\Sigma_1^1(<_f)$ , which is  $\exists^\omega$ -closed. By the assumption that the theorem holds for all ordinals smaller than  $\gamma$ , there exists for each  $\alpha < \gamma$  an  $x \in Y$  such that  $\alpha_x \geq \alpha$ . It follows then that  $\bigcup_{x \in \omega^\omega} U_{2,\sigma \circ x}$  is as desired.

Now suppose that  $\tau$  is a winning strategy for player  $II$ . For each  $y \in X$ , let  $[<y]_f$  denote the set of  $z \in X$  with  $f(z) < f(y)$ . The set of  $(x, y, z, w) \in (\omega^\omega)^4$  for which  $(x, z, w) \in U_2$ ,  $(y, z) \in X^2$  and  $z \in [<y]_f$  is in  $\text{pos-}\Sigma_1^1(<_f)$ . Fix  $a_0 \in \omega^\omega$  such that this set is  $U_{4,a_0}$ . Let  $s_{4,2}: (\omega^\omega)^3 \rightarrow \omega^\omega$  be a continuous function witnessing the s-m-n property of  $\bar{U}$  for  $n = 2$  and  $m = 4$ , and let  $h_0: (\omega^\omega)^2 \rightarrow \omega^\omega$  be defined by setting  $h_0(x, y) = s_{4,2}(a_0, x, y)$ . Then for all  $(x, y) \in \omega^\omega \times X$ ,

$$U_{2,h_0(x,y)} = U_{2,x} \cap ([<y]_f \times \omega^\omega). \quad (3.1)$$

Similarly, the set of  $(x, y, z) \in \omega^\omega \times X \times \omega^\omega$  for which  $(h_0(x, y) \circ \tau, y, z) \in U_2$  is in  $\text{pos-}\Sigma_1^1(<_f)$ . Fix  $a_1 \in \omega^\omega$  such that this set is  $U_{3,a_1}$ . Let  $s_{3,1}: (\omega^\omega)^2 \rightarrow \omega^\omega$  be a continuous function witnessing the s-m-n property of  $\bar{U}$  for  $n = 1$  and  $m = 3$ , and let  $h_1: \omega^\omega \rightarrow \omega^\omega$  be defined by setting  $h_1(x) = s_{3,1}(a_1, x)$ . Then for all  $x \in \omega^\omega$ ,

$$U_{2,h_1(x)} = \bigcup_{y \in X} (U_{2,h_0(x,y) \circ \tau} \cap (\{y\} \times \omega^\omega)). \quad (3.2)$$

By Theorem 2.4.10, there exists an  $x_1 \in \omega^\omega$  such that  $U_{2,x_1} = U_{2,h_1(x_1)}$ . We show now that  $x_1$  satisfies conditions (1) and (2). For condition (1), suppose toward a contradiction that the condition fails, and consider  $(y, z) \in U_{2,x_1} \setminus Z$  with  $f(y)$  minimal. Since  $U_{2,x_1} = U_{2,h_1(x_1)}$ ,  $(y, z) \in U_{2,h_0(x_1,y) \circ \tau}$ . Then by equation (3.1), for all  $(a, b) \in U_{2,h_0(x_1,y)}$ ,  $(a, b)$  is in  $U_{2,x_1}$  and  $f(a) < f(y)$ . By the minimality of  $f(y)$ , it follows that  $(a, b) \in Z$  and therefore that  $h_0(x_1, y) \in Y$ . Since  $\tau$  is a winning strategy for  $II$ ,  $h_0(x_1, y) \circ \tau \in Y$ , which means that  $(y, z) \in Z$ , giving a contradiction.

For condition (2), suppose toward a contradiction that  $\alpha_{x_1} < \gamma$ . Let  $y \in X$  be such that  $f(y) = \alpha_{x_1}$ . By equation (3.1), the fact that  $x_1$  is in  $Y$  and the definition of  $\alpha_{x_1}$ ,  $h_0(x_1, y) \in Y$ , and  $\alpha_{h_0(x_1,y)} = \alpha_{x_1}$ . Since  $\tau$  is a winning strategy for  $II$ ,  $\alpha_{h_0(x_1,y) \circ \tau} > \alpha_{h_0(x_1,y)} = \alpha_{x_1}$ , which is impossible, as

$$(U_{2,h_0(x_1,y) \circ \tau} \cap ([y]_f \times \omega^\omega)) \subseteq U_{2,x_1},$$

by equation (3.2) and the choice of  $x_1$ . This completes the proof.  $\square$



Corollary 3.0.2 lists two immediate consequences of the Coding Lemma. The first part of the corollary to the Coding Lemma follows from the fact that there is a surjection from  $\omega^\omega$  to  $\Sigma_1^1$ .

**Corollary 3.0.2.** *If AD holds, then each of the following hold.*

1. *For each  $\lambda < \Theta$  there is a surjection from  $\mathcal{P}(\omega)$  to  $\mathcal{P}(\lambda)$ .*
2.  *$\Theta$  is a limit cardinal.*

We now prove a uniform version of Theorem 3.0.1 which is used in the proof of Theorem 5.0.11, which in turn is used to prove that the strong partition cardinals are cofinal below  $\Theta$ . Given a sequence  $\langle U_n : n < \omega \rangle$  of universal sets for  $\Sigma_1^1$ , and an  $A \subseteq \omega^\omega$ , we let  $\langle U_n(A) : n < \omega \rangle$  be as in Remark 2.4.12.

**Theorem 3.0.3** (The Uniform Coding Lemma). *Assume that AD holds. Let*

- *$\bar{U} = \langle U_n : n < \omega \rangle$  be a sequence of universal sets for  $\Sigma_1^1$  with the  $s$ - $m$ - $n$  property;*
- *$X$  be a subset of  $\omega^\omega$ ;*
- *$Z$  be a subset of  $X \times \omega^\omega$ ;*
- *$f$  be a function from  $X$  to the ordinals.*

*For each  $y \in X$ , let  $[y]_f$  denote  $\{z \in X : f(y) = f(z)\}$ , let  $[<y]_f$  denote  $\{z \in X : f(z) < f(y)\}$  and let*

$$C_y = \omega^\omega \setminus ([y]_f \cup [<y]_f).$$

*Then there exists an  $x \in \omega^\omega$  such that for all  $y \in X$ ,*

1.  $U_{2,x}([y]_f, C_y) \subseteq Z \cap ([y]_f \times \omega^\omega),$
2.  $U_{2,x}([y]_f, C_y) \neq \emptyset$  if and only if  $Z \cap ([y]_f \times \omega^\omega) \neq \emptyset.$

*Proof.* It suffices to consider the case where the range of  $f$  is an ordinal  $\gamma$ , and, proving the theorem by induction, we may assume that it holds for all smaller ordinals. As  $\Sigma_1^1$  is closed under unions and contains all countable subsets of  $\omega^\omega$ , we may assume that  $\gamma$  is an uncountable limit ordinal. Let  $Y$  be the set

$$\{x \in \omega^\omega : \forall y \in X \ U_{2,x}([y]_f, C_y) \subseteq Z \cap ([y]_f \times \omega^\omega)\}.$$

For each  $x \in \omega^\omega$ , let  $\alpha_x$  be the least value  $f(y)$  for a  $y \in X$  witnessing the failure of conclusion (2) of the theorem, with respect to  $x$ , if there exists such a  $y$ ; otherwise, let  $\alpha_x = \gamma$ . Consider the game between players  $I$  and  $II$  where player  $I$  builds  $x_1 \in \omega^\omega$ , player  $II$  builds  $x_2 \in \omega^\omega$ , and  $I$  wins if and only if  $x_1 \in Y$  and either  $x_2 \notin Y$  or  $\alpha_{x_1} \geq \alpha_{x_2}$ . We will show that a winning strategy for either player gives the desired conclusion.

First, suppose that  $\sigma$  is a winning strategy for player  $I$ . Let  $z_\sigma \in \omega^\omega$  be such that  $U_{2,z_\sigma} = \bigcup_{x \in \omega^\omega} U_{2,\sigma \circ x}$ . Then for all  $P_1, P_2 \subseteq \omega^\omega$  (in particular, in the case  $P_1 = [y]_f$ ,  $P_2 = C_y$ , for some  $y \in X$ ),

$$U_{2,z_\sigma}(P_1, P_2) = \bigcup_{x \in \omega^\omega} U_{2,\sigma \circ x}(P_1, P_2).$$

Since  $\sigma$  is a winning strategy for  $I$ ,  $z_\sigma$  is in  $Y$ . By our assumption that the theorem holds for all ordinals smaller than  $\gamma$ , there exist for each  $\alpha < \gamma$  an  $x \in Y$  such that  $\alpha_x \geq \alpha$ . It follows then that  $z_\sigma$  is as desired.

Now suppose that  $\tau$  is a winning strategy for player  $II$ . Let  $\pi: \omega \times \omega \rightarrow \omega$  be the recursive bijection used to define the classes  $\text{pos-}\widetilde{\Sigma}_1^1(A)$  and the map  $y \mapsto \langle (y)_n : n \in \omega \rangle$  for  $y \in \omega^\omega$ . For each  $i \in \{1, 2\}$ , let  $Q_i$  be the set of pairs  $(y, r) \in \omega^\omega \times \omega^\omega$  such that

$$(r)_0 = \langle \pi(i, y(0)), y(1), y(2), \dots \rangle$$

(this corresponds to how we defined the class  $\text{pos-}\widetilde{\Sigma}_1^1(A_1, A_2)$  in terms of our definition of  $\text{pos-}\widetilde{\Sigma}_1^1(A)$  in Section 2.4). Then for any  $P, P' \subseteq \omega^\omega$  and  $z \in \omega^\omega$ ,

- $z \in P$  if and only if  $(z, r) \in Q_1$  holds for some  $r \in \omega^\omega$  with  $(r)_0 \in P \oplus P'$ .
- $z \in P'$  if and only if  $(z, r) \in Q_2$  holds for some  $r \in \omega^\omega$  with  $(r)_0 \in P \oplus P'$ .

For each  $r \in \omega^\omega$ , let  $r^- \in \omega^\omega$  be such that  $(r^-)_n = (r)_{n+1}$  for all  $n \in \omega$ .

Let  $a_0 \in \omega^\omega$  be such that  $U_{5,a_0}$  is the set of  $(x, y, z, w, r) \in (\omega^\omega)^5$  for which  $(x, z, w, r^-) \in U_3$  and  $(y, r) \in Q_2$ . Let  $s_{5,2}: (\omega^\omega)^3 \rightarrow \omega^\omega$  be a continuous function witnessing the s-m-n property of  $\bar{U}$  for  $n = 2$  and  $m = 5$ , and let  $h_0: (\omega^\omega)^2 \rightarrow \omega^\omega$  be defined by setting  $h_0(x, y) = s_{5,2}(a_0, x, y)$ . Then for all  $(x, y) \in \omega^\omega \times \omega^\omega$ , all  $P, P' \subseteq \omega^\omega$  and all  $(z, w) \in \omega^\omega \times \omega^\omega$ ,

$$(z, w) \in U_{2,h_0(x,y)}(P, P')$$

if and only if

$$\exists r \in \omega^\omega ((\forall n \in \omega (r)_n \in P \oplus P') \wedge (h_0(x, y), z, w, r) \in U_3)$$

if and only if

$$\exists r \in \omega^\omega ((\forall n \in \omega (r)_n \in P \oplus P') \wedge (a_0, x, y, z, w, r) \in U_5)$$

if and only if

$$\exists r \in \omega^\omega ((\forall n \in \omega (r)_n \in P \oplus P') \wedge (x, z, w, r^-) \in U_3 \wedge (y, r) \in Q_2)$$

if and only if

$$(z, w) \in U_{2,x}(P, P') \wedge y \in P'.$$

That is, we have the following fact (\*\*): for all  $(x, y) \in \omega^\omega \times \omega^\omega$ , and all  $P, P' \subseteq \omega^\omega$  (in particular, sets of the form  $[y]_f$  and  $C_y$ ),  $U_{2,h_0(x,y)}(P, P')$  is  $U_{2,x}(P, P')$  if  $y$  is in  $P'$  and  $\emptyset$  otherwise.

Now let  $a_1 \in \omega^\omega$  be such that  $U_{4,a_1}$  is the set of  $(x, z, w, r) \in (\omega^\omega)^4$  such that, for some  $y \in \omega^\omega$ ,  $(z, w, r^-) \in U_{3,h_0(x,y) \circ \tau}$  and  $(y, r) \in Q_1$  holds. Let  $s_{4,1}: (\omega^\omega)^2 \rightarrow \omega^\omega$  be a continuous function witnessing the s-m-n property of  $\bar{U}$  for  $n = 1$  and  $m = 3$ , and let  $h_1: \omega^\omega \rightarrow \omega^\omega$  be defined by setting  $h_1(x) = s_{4,1}(a_1, x)$ .

Then for all  $x \in \omega^\omega$ , all  $P, P' \subseteq \omega^\omega$  and all  $(z, w) \in \omega^\omega \times \omega^\omega$ ,

$$(z, w) \in U_{2,h_1(x)}(P, P')$$

if and only if

$$\exists r \in \omega^\omega ((\forall n \in \omega (r)_n \in P \oplus P') \wedge (h_1(x), z, w, r) \in U_3)$$

if and only if

$$\exists r \in \omega^\omega ((\forall n \in \omega (r)_n \in P \oplus P') \wedge (a_1, x, z, w, r) \in U_4)$$

if and only if

$$\exists r \in \omega^\omega ((\forall n \in \omega (r)_n \in P \oplus P') \wedge \exists y \in \omega^\omega ((z, w, r^-) \in U_{3,h_0(x,y) \circ \tau} \wedge (y, r) \in Q_1))$$

if and only if

$$\exists y \in P (z, w) \in U_{2,h_0(x,y) \circ \tau}(P, P').$$

Then for all  $x \in \omega^\omega$  and all  $P, P' \subseteq \omega^\omega$ ,

$$U_{2,h_1(x)}(P, P') = \bigcup_{y \in P} U_{2,h_0(x,y) \circ \tau}(P, P'). \quad (3.3)$$

By Theorem 2.4.10 and Remark 2.4.12, there exists an  $x_1 \in \omega^\omega$  such that  $U_{2,x_1}(P, P') = U_{2,h_1(x_1)}(P, P')$  for all  $P, P' \subseteq \omega^\omega$ . We show now that  $x_1$  satisfies both conditions in the conclusion of the theorem. For condition (1) (i.e., the assertion that  $x_1 \in Y$ ), fix  $y^* \in X$  with

$$U_{2,x_1}([y^*]_f, C_{y^*}) \not\subseteq Z \cap ([y^*]_f \times \omega^\omega)$$

and  $f(y^*)$  minimal. Since

$$U_{2,x_1}([y^*]_f, C_{y^*}) = U_{2,h_1(x_1)}([y^*]_f, C_{y^*}),$$

there exists by equation (3.3) a  $y' \in [y^*]_f$  such that

$$U_{2,h_0(x_1,y') \circ \tau}([y^*]_f, C_{y^*}) \not\subseteq Z \cap ([y^*]_f \times \omega^\omega).$$

We want to see that  $h_0(x_1, y') \in Y$ , since then, as  $\tau$  is a winning strategy for  $II$ , we will have that  $h_0(x_1, y') \circ \tau \in Y$ , giving a contradiction. For each  $y \in X$ , we have by (\*\*) that

$$U_{2,h_0(x_1,y') \circ \tau}([y]_f, C_y) = U_{2,x_1}([y]_f, C_y) \subseteq Z \cap ([y]_f \times \omega^\omega)$$

if  $f(y) < f(y')$ , by the minimality of  $f(y^*)$ , and  $U_{2,h_0(x_1,y')}([y^*]_f, C_y) = \emptyset$  otherwise. In either case,  $U_{2,h_0(x_1,y')}([y]_f, C_y) \subseteq Z \cap ([y]_f \times \omega^\omega)$ , which shows that  $h_0(x_1, y') \in Y$  as desired.

For condition (2), suppose toward a contradiction that  $\alpha_{x_1} < \gamma$ . Let  $y^* \in X$  be such that  $f(y^*) = \alpha_{x_1}$ . By (\*\*), the fact that  $x_1$  is in  $Y$  and the definition of  $\alpha_{x_1}$ ,  $h_0(x_1, y^*) \in Y$ , and  $\alpha_{h_0(x_1,y^*)} = \alpha_{x_1}$ . Since  $\tau$  is a winning strategy for *II*,  $\alpha_{h_0(x_1,y^*) \circ \tau} > \alpha_{h_0(x_1,y^*)} = \alpha_{x_1}$ , which is impossible, as

$$U_{2,h_0(x_1,y^*) \circ \tau}([y^*]_f, C_{y^*}) \subseteq U_{2,x_1}([y^*]_f, C_{y^*}),$$

by equation (3.3) and the choice of  $x_1$ . This completes the proof. □

## Chapter 4

# Properties of pointclasses

In this chapter we develop various properties of boldface pointclasses under AD, including closure and separation properties, and the prewellordering property. Our immediate goal is preparing for the proof of the existence of strong partition cardinals in Chapter 5. Some of the results in this chapter will also be used in Chapter 6. The material in this chapter is part of a very deep and general theory. However, we are mostly interested in taking the shortest path to proving the existence of strong partition cardinals. Readers interested in this general theory are directed to [15, 16, 17] for an introduction. Much of this chapter is taken from [5, 6] and conversations with their author.

### 4.1 Separation and reduction

Given a pointclass  $\Gamma$ , and disjoint subsets  $A$  and  $B$  of the same space in  $\mathcal{X}$ , we say that  $A$  and  $B$  are  $\Gamma$ -separable if there exists a set  $C \in \Gamma \cap \check{\Gamma}$  such that  $A \subseteq C$  and  $B \cap C = \emptyset$  (and  $\Gamma$ -inseparable otherwise). We say that a pointclass  $\Gamma$  satisfies the *separation property* if all pairs of disjoint subsets of  $\omega^\omega$  in  $\Gamma$  are  $\Gamma$ -separable. We write  $\text{Sep}(\Gamma)$  to indicate that  $\Gamma$  has the separation property.

We follow [38]. We say that a function  $f: \omega^\omega \rightarrow \omega^\omega$  is *strongly Lipschitz* if for all  $x, y \in \omega^\omega$  and  $n \in \omega$ , if  $x \upharpoonright n = y \upharpoonright n$ , then  $f(x)(n) = f(y)(n)$ . Strategies for player I in Lipschitz games induce strongly Lipschitz functions.

The following lemma is used in the proof of Theorem 4.1.2.

**Lemma 4.1.1** (ZF + AD; Steel). *Let  $\Gamma$  be a boldface pointclass and let  $(A_0, A_1)$  be a  $\Gamma$ -inseparable pair of subsets of  $\omega^\omega$ . Let  $B_0$  and  $B_1$  be disjoint subsets of  $\omega^\omega$ , both in  $\Gamma$  or both in  $\check{\Gamma}$ . Then there is a strongly Lipschitz map  $f$  so that  $f[B_0] \subseteq A_0$  and  $f[B_1] \subseteq A_1$ .*

*Proof.* Let  $\Delta = \Gamma \cap \check{\Gamma}$ . Consider the game where I plays  $x$ , II plays  $y$ , and I wins if  $y \in B_0$  implies  $x \in A_0$ , and  $y \in B_1$  implies  $x \in A_1$ . It suffices to see that II cannot have a winning strategy. Supposing that  $\sigma$  were a winning strategy for II, let  $g$  be the function  $x \mapsto x \circ \sigma$  ( $x \circ \sigma$  is the sequence of moves made

by player  $II$  when he plays according to  $\sigma$  and player  $I$  plays  $x$ ; this notation was defined at the beginning of Chapter 3). Then  $g$  is continuous, and  $g^{-1}[B_0]$  is in  $\Delta$ , since  $\omega^\omega \setminus g^{-1}[B_0] = g^{-1}[B_1]$ . Furthermore,  $g^{-1}[B_0]$  separates  $A_0$  and  $A_1$ .  $\square$

**Theorem 4.1.2** (ZF+AD; Steel). *If  $\Gamma$  is a nonselfdual boldface pointclass, then at least one of  $\Gamma$  and  $\check{\Gamma}$  satisfies the separation property.*

*Proof.* Suppose toward a contradiction that the pairs  $(A_0, A_1)$  and  $(C_0, C_1)$  form counterexamples to  $\text{Sep}(\Gamma)$  and  $\text{Sep}(\check{\Gamma})$ , respectively. By Lemma 4.1.1, there is a strongly Lipschitz function  $f$  mapping  $A_0$  into  $C_0$  and  $A_1$  into  $C_1$ . Let  $B_0 = f^{-1}[C_0]$  and let  $B_1 = f^{-1}[C_1]$ . Then  $B_0$  and  $B_1$  are disjoint and both in  $\check{\Gamma}$ , and  $A_0 \subseteq B_0$  and  $A_1 \subseteq B_1$ . Applying Lemma 4.1.1 again, there exist strongly Lipschitz functions  $f_0, f_1$  and  $f_2$  such that

- $f_0[A_0] \subseteq A_1$ ;
- $f_0[A_1] \subseteq A_0$ ;
- $f_1[A_0] \subseteq A_0$ ;
- $f_1[\omega^\omega \setminus B_0] \subseteq A_1$ ;
- $f_2[A_1] \subseteq A_1$ ;
- $f_2[\omega^\omega \setminus B_1] \subseteq A_0$ .

Since  $f_0, f_1$  and  $f_2$  are all strongly Lipschitz, for each  $r \in 3^\omega$  there is a unique sequence  $\langle x_n^r : n < \omega \rangle$  of elements of  $\omega^\omega$  such that  $x_n^r = f_{r(n)}(x_{n+1}^r)$  for all  $n \in \omega$ . Let  $3^\omega$  have its induced topology as a subspace of  $\omega^\omega$ . By Baire( $\mathcal{P}(\omega^\omega)$ ), every subset of  $3^\omega$  has the property of Baire, in particular the set  $C = \{r \in 3^\omega : x_0^r \notin A_0 \cup A_1\}$ .

Suppose first that  $C$  is nonmeager. By the Baire property, we may fix an  $s \in 3^{<\omega}$  such that  $\{r \in 3^\omega : x_0^{s \hat{\ } r} \notin A_0 \cup A_1\}$  is comeager. However, for each  $s \in 3^{<\omega}$  and  $r \in 3^\omega$ , if  $x_0^r \in A_0 \cup A_1$  then  $x_0^{s \hat{\ } r} \in A_0 \cup A_1$ . It follows then that  $x_0^r \notin A_0 \cup A_1$ , for comeagerly many  $r \in 3^\omega$ . Since  $B_0 \cap B_1 = \emptyset$ , we may fix an  $i^* \in \{0, 1\}$  such that  $x_0^r \notin B_{i^*}$ , for nonmeagerly many  $r \in 3^\omega$ . Inspecting  $f_0, f_1$  and  $f_2$ , we see that for each  $i \in \{0, 1\}$  and each  $r \in 3^\omega$ , if  $x_0^r \notin B_i$  then  $x_0^{(i+1) \hat{\ } r} \in A_{1-i}$ . It follows then that  $x_0^r \in A_0 \cup A_1$  for nonmeagerly many  $r \in 3^\omega$ , which gives a contradiction.

Suppose on the other hand that  $C$  is comeager. Choose  $i^* \in \{0, 1\}$  so that  $\{r \in 3^\omega : x_0^r \in A_{i^*}\}$  is nonmeager. Choose  $s \in 3^{<\omega}$  such that the set  $\{r \in 3^\omega : x_0^{s \hat{\ } r} \in A_{i^*}\}$  is comeager. For any  $r \in 3^\omega$  such that  $x_0^r \in A_{i^*}$ ,  $x_0^{s \hat{\ } (0) \hat{\ } r} \in A_{1-i^*}$ . It follows that  $x_0^{s \hat{\ } r} \in A_{1-i^*}$  for nonmeagerly many  $r \in 3^\omega$ , giving another contradiction.  $\square$

**4.1.3 Remark.** Theorems 5.2 and 5.3 of [40] show that if AD holds then for any nonselfdual pointclass  $\Gamma$ , at most one of  $\Gamma$  and  $\check{\Gamma}$  has the separation property.

A pointclass  $\Gamma$  is said to have the *reduction property* if for all  $A, B \subseteq \omega^\omega$  in  $\Gamma$ , there exist disjoint  $A', B' \in \Gamma$  such that  $A' \subseteq A$ ,  $B' \subseteq B$  and  $A' \cup B' = A \cup B$ . In this case we say that the sets  $A'$  and  $B'$  *reduce* the sets  $A$  and  $B$ . We write  $\text{Red}(\Gamma)$  to indicate that the pointclass  $\Gamma$  has the reduction property. It follows easily from the definitions that  $\text{Red}(\Gamma)$  implies  $\text{Sep}(\check{\Gamma})$ . Along with Remark 4.1.3, this shows that, under AD,  $\text{Red}(\Gamma)$  and  $\text{Sep}(\Gamma)$  cannot both hold for a nonselfdual pointclass. The following (taken from 4B.12 of [32]) gives a direct proof.

**Theorem 4.1.4.** *Assume that Lipschitz Determinacy + Baire( $\mathcal{P}(\omega^\omega)$ ) holds. If  $\Gamma$  is a nonselfdual boldface pointclass and  $\text{Red}(\Gamma)$  holds, then  $\text{Sep}(\Gamma)$  does not hold.*

*Proof.* By Theorem 2.4.3 there exist a  $\Gamma$ -universal set  $U \subseteq (\omega^\omega)^2$ . Let  $\pi: \omega^\omega \rightarrow (\omega^\omega)^2$  be a continuous bijection, and let  $\pi_0, \pi_1$  be the functions on  $\omega^\omega$  such that  $\pi(x) = (\pi_0(x), \pi_1(x))$  for all  $x \in \omega^\omega$ . Let  $A$  be the set of  $x \in \omega^\omega$  such that  $(\pi_0(x), x) \in U$  and let  $B$  be the set of  $x \in \omega^\omega$  such that  $(\pi_1(x), x) \in U$ . Let  $A'$  and  $B'$  in  $\Gamma$  witness the reduction property for  $A$  and  $B$ .

Suppose now that  $C \in \Gamma \cap \check{\Gamma}$  witnesses the separation property for  $A'$  and  $B'$  (i.e.,  $C$  contains  $A'$  and is disjoint from  $B'$ ). Let  $x \in \omega^\omega$  be such that  $U_{\pi_0(x)} = \omega^\omega \setminus C$  and  $U_{\pi_1(x)} = C$ . If  $x \in C$ , then  $x \in B \setminus A$ , so  $x \in B'$ , giving a contradiction. Similarly, if  $x \notin C$ , then  $x \in A \setminus B$ , so  $x \in A'$ , giving another contradiction.  $\square$

**4.1.5 Remark.** Theorems 4.1.2 and 4.1.4 together imply, assuming AD, that if  $\Gamma$  is a nonselfdual boldface pointclass, then at most one of  $\Gamma$  and  $\check{\Gamma}$  has the reduction property. Theorem 4.1.6, which is Theorem 5.5 of [40], shows that if in addition  $\Gamma$  is closed under finite unions and intersections, then the separation property holds for exactly one of  $\Gamma$  and  $\check{\Gamma}$ , and the reduction property holds for the other.

**Theorem 4.1.6** (ZF + AD; Van Wesep). *If  $\Gamma$  is a boldface pointclass closed under finite intersections, and  $\text{Sep}(\check{\Gamma})$  holds, then  $\text{Red}(\Gamma)$  holds.*

The proof of Theorem 4.1.6 (which we give below) uses the following lemma.

**Lemma 4.1.7.** *Suppose that AD holds, and that  $\Gamma$  is a boldface pointclass closed under finite intersections. If  $\text{Red}(\Gamma)$  does not hold, then for all  $A, B$  in  $\mathcal{P}(\omega^\omega) \cap \check{\Gamma}$  there exist  $C, D \in \Gamma$  such that*

- $A \setminus B \subseteq C \setminus D$ ;
- $B \setminus A \subseteq D \setminus C$ ;
- $C \cup D = \omega^\omega$ .

*Proof.* Let  $(E, F)$  be subsets of  $\omega^\omega$  witnessing  $\neg \text{Red}(\Gamma)$ , and fix sets  $A, B$  in  $\mathcal{P}(\omega^\omega) \cap \check{\Gamma}$ . Consider the game  $\mathcal{G}(E, F, A, B)$  in which player  $I$  produces  $x \in \omega^\omega$ , player  $II$  produces  $y \in \omega^\omega$  and player  $II$  wins if and only if the following conditions are met:

- $y \in E \cup F$ ;
- if  $x \in A \setminus B$  then  $y \in E \setminus F$ ;
- if  $x \in B \setminus A$  then  $y \in F \setminus E$ .

Suppose toward a contradiction that  $\sigma$  is a winning strategy for player  $I$ . Let  $E'' = \{y \in \omega^\omega : \sigma \circ y \notin A\}$  and let  $F'' = \{y \in \omega^\omega : \sigma \circ y \notin B\}$ . Then we have the following:

- $E \cup F \subseteq E'' \cup F''$ ;
- $(E \cup F) \cap (E'' \cap F'') = \emptyset$ ;
- $E \setminus F \subseteq E''$ ;
- $F \setminus E \subseteq F''$ ;
- $E'', F'' \in \Gamma$ .

Let  $E' = E \cap E''$  and let  $F' = F \cap F''$ . Then  $E'$  and  $F'$  are disjoint sets in  $\Gamma$ , and  $E \cup F = E' \cup F'$ . This contradicts the assumption that  $E$  and  $F$  witness the failure of  $\text{Red}(\Gamma)$ .

It follows then that there is a winning strategy  $\tau$  for player  $II$ . Let  $C$  be  $\{x \in \omega^\omega : x \circ \tau \in E\}$  and let  $D$  be  $\{x \in \omega^\omega : x \circ \tau \in F\}$ . Then  $C$  and  $D$  are as desired.  $\square$

*Proof of Theorem 4.1.6.* Supposing that  $\text{Red}(\Gamma)$  fails, we show that  $\text{Red}(\check{\Gamma})$  holds, contradicting Theorem 4.1.4. Let  $A$  and  $B$  be subsets of  $\omega^\omega$  in  $\check{\Gamma}$ . Let  $C$  and  $D$  be as given by Lemma 4.1.7. Applying  $\text{Sep}(\check{\Gamma})$ , let  $E \in \Gamma \cap \check{\Gamma}$  be such that  $\omega^\omega \setminus C \subseteq E$  and  $E \cap (\omega^\omega \setminus D) = \emptyset$ . Let  $C' = C \setminus E$  and let  $D' = D \cap E$ . Then

$$D' \cap (A \setminus B) = C' \cap (B \setminus A) = C' \cap D' = \emptyset$$

and  $A \cup B \subseteq C' \cup D'$ , so  $A \setminus D'$  and  $B \setminus C'$  satisfy the conclusion of the reduction property with respect to  $A$  and  $B$ .  $\square$

The following result (which we need for Theorem 6.1.10) is sometimes called the 0th Periodicity Theorem. Recall that  $x \leq_{\text{Tu}} y$  and  $x \equiv_{\text{Tu}} y$  refer to Turing reducibility and were defined in Section 0.2.

**Theorem 4.1.8** (ZF + AD; Kechris). *Suppose that  $\Gamma$  is a boldface pointclass closed under countable intersections and countable unions, and that  $\text{Red}(\Gamma)$  holds.*

- If  $\exists^{\omega^\omega} \Gamma \subseteq \Gamma$ , then  $\text{Red}(\forall^{\omega^\omega} \Gamma)$  holds.
- If  $\forall^{\omega^\omega} \Gamma \subseteq \Gamma$ , then  $\text{Red}(\exists^{\omega^\omega} \Gamma)$  holds.



*Proof.* Let  $A$  and  $B$  be subsets of  $(\omega^\omega)^2$  in  $\Gamma$ . For the first part we will find a reduction for  $A^* = \{x : \forall y \in \omega^\omega (x, y) \in A\}$  and  $B^* = \{x : \forall y \in \omega^\omega (x, y) \in B\}$ . Let

$$A' = \{(x, z) \in (\omega^\omega)^2 : \forall y \leq_{\text{Tu}} z (x, y) \in A\}$$

and let

$$B' = \{(x, z) \in (\omega^\omega)^2 : \forall y \leq_{\text{Tu}} z (x, y) \in B\}.$$

Since  $\Gamma$  is closed under countable intersections,  $A'$  and  $B'$  are in  $\Gamma$ ; fix sets  $C \subseteq A'$  and  $D \subseteq B'$  reducing them. Let

$$C' = \{(x, z) \in (\omega^\omega)^2 : \exists z' \equiv_{\text{Tu}} z (x, z') \in C\}$$

and let

$$D' = \{(x, z) \in (\omega^\omega)^2 : \forall z' \equiv_{\text{Tu}} z (x, z') \in D\}.$$

Then  $C'$  and  $D'$  are in  $\Gamma$ , since  $\Gamma$  is closed under countable intersections and countable unions, and they reduce  $A'$  and  $B'$ .

Let  $\mathcal{G}_A(x)$  (for  $x \in \omega^\omega$ ) be the game where players  $I$  and  $II$  collaborate to build  $z \in \omega^\omega$ , where  $I$  wins if and only if  $(x, z) \in C'$ , and let  $E$  be the set of  $x$  for which  $I$  has a winning strategy in  $\mathcal{G}_A(x)$ . Let  $\mathcal{G}_B(x)$  (for  $x \in \omega^\omega$ ) be the game where players  $I$  and  $II$  collaborate to build  $z \in \omega^\omega$ , where  $I$  wins if and only if  $(x, z) \in D'$ , and let  $F$  be the set of  $x$  for which  $I$  has a winning strategy in  $\mathcal{G}_B(x)$ . Then  $E$  and  $F$  are in  $\forall^{\omega^\omega} \Gamma$  (via the assertion that player  $II$  does not have a winning strategy, and the assumption that  $\exists^{\omega^\omega} \Gamma \subseteq \Gamma$ ), and they are disjoint, since  $C'$  and  $D'$  are. Furthermore,  $E \subseteq A^*$  and  $F \subseteq B^*$ .

To see that  $E$  and  $F$  reduce  $A^*$  and  $B^*$ , fix  $x \in A^* \cup B^*$ . Then at least one of  $A_x$  and  $B_x$  is all of  $\omega^\omega$ . In the former case,  $A'_x = \omega^\omega$ , and in the latter case  $B'_x = \omega^\omega$ . Then for each  $z \in \omega^\omega$ ,  $(x, z)$  is either in  $C'$  or  $D'$ , and one of the sets  $\{[z]_{\text{Tu}} : (x, z) \in C'\}$  and  $\{[z]_{\text{Tu}} : (x, z) \in D'\}$  contains a Turing cone. Then player  $I$  has a winning strategy in one of the games  $\mathcal{G}_A(x)$  and  $\mathcal{G}_B(x)$ .

For the second part (redirecting our labels) we find a reduction for  $A^* = \{x : \exists y \in \omega^\omega (x, y) \in A\}$  and  $B^* = \{x : \exists y \in \omega^\omega (x, y) \in B\}$ . Let

$$A' = \{(x, z) \in (\omega^\omega)^2 : \exists y \leq_{\text{Tu}} z (x, y) \in A\}$$

and let

$$B' = \{(x, z) \in (\omega^\omega)^2 : \exists y \leq_{\text{Tu}} z (x, y) \in B\}.$$

Since  $\Gamma$  is closed under countable unions,  $A'$  and  $B'$  are in  $\Gamma$ , so there exist  $C$  and  $D$  in  $\Gamma$  reducing them. Let

$$C' = \{(x, z) \in (\omega^\omega)^2 : \exists z' \equiv_{\text{Tu}} z (x, z') \in C\}$$

and let

$$D' = \{(x, z) \in (\omega^\omega)^2 : \forall z' \equiv_{\text{Tu}} z (x, z') \in D\}.$$

Then  $C'$  and  $D'$  are in  $\Gamma$ , since  $\Gamma$  is closed under countable intersections and countable unions, and they reduce  $A'$  and  $B'$ .

Let  $\mathcal{G}_A(x)$  (for  $x \in \omega^\omega$ ) be the game where players  $I$  and  $II$  collaborate to build  $z \in \omega^\omega$ , where  $I$  wins if and only if  $(x, z) \in C'$ , and let  $E$  be the set of  $x$  for which  $I$  has a winning strategy. Let  $\mathcal{G}_B(x)$  (for  $x \in \omega^\omega$ ) be the game where players  $I$  and  $II$  collaborate to build  $z \in \omega^\omega$ , where  $I$  wins if and only if  $(x, z) \in D'$ , and let  $F$  be the set of  $x$  for which  $I$  has a winning strategy. Then  $E$  and  $F$  are in  $\exists^\omega \Gamma$  (via the assertion that player  $I$  has a winning strategy, and the assumption that  $\forall^\omega \Gamma \subseteq \Gamma$ ), and they are disjoint. Again,  $E \subseteq A^*$  and  $F \subseteq B^*$ .

To see that  $E$  and  $F$  reduce  $A^*$  and  $B^*$ , fix  $x \in A^* \cup B^*$ . Then at least one of  $A_x$  and  $B_x$  is nonempty. Then again one of the sets  $\{[z]_{\text{Tu}} : (x, z) \in C'\}$  and  $\{[z]_{\text{Tu}} : (x, z) \in D'\}$  contains a Turing cone. Then player  $I$  has a winning strategy in one of the games  $\mathcal{G}_A(x)$  and  $\mathcal{G}_B(x)$ .  $\square$

## 4.2 The prewellordering property

Given a subset  $P$  of a space in  $\mathcal{X}$ , a *norm* on  $P$  is a function from  $P$  to the ordinals. A norm is *regular* if its range is an ordinal. We write  $o(\varphi)$  for the ordertype of the range of a norm  $\varphi$ , and call this the *length* or the *rank* of  $\varphi$ .

A *prewellordering* on a set  $P$  is a binary relation  $\leq$  on which is reflexive, transitive, total (so for all  $x, y \in X$ , at least one of  $x \leq y$  and  $y \leq x$  holds) and wellfounded. Equivalently (via the canonical rank function for a wellfounded preorder), a prewellordering on a set  $P$  is a set of the form

$$\{(x, y) \in P \times P : f(x) \leq f(y)\},$$

where  $f$  is a (regular) norm on  $P$ . Similarly, a *strict prewellordering* of  $P$  is a set of the form  $\{(x, y) \in P \times P : f(x) < f(y)\}$ , for some norm  $f$  on  $P$ . The *length* of a (strict) prewellordering then is the ordertype of the range of any such  $f$ . Given a pointclass  $\Gamma$ , we let  $\delta(\Gamma)$  be the supremum of the lengths of the prewellorderings in  $\Gamma \cap \check{\Gamma}$ .

The following is the central definition of this section.

**4.2.1 Definition.** Given a pointclass  $\Gamma$ , a space  $X$  in  $\mathcal{X}$  and a set  $P \subseteq X$ , a  $\Gamma$ -*norm* on  $P$  is norm  $\varphi$  on  $P$  for which each of the following sets are in  $\Gamma$ :

- $\leq_\varphi^*$ , the set of pairs  $(x, y) \in X \times X$  such that  $x \in P$  and either  $y \notin P$  or  $\varphi(x) \leq \varphi(y)$ ;
- $<_\varphi^*$ , the set of pairs  $(x, y) \in X \times X$  such that  $x \in P$  and either  $y \notin P$  or  $\varphi(x) < \varphi(y)$ .

**4.2.2 Remark.** Suppose that  $\Gamma$  is a boldface pointclass, and  $\varphi$  is a  $\Gamma$ -norm on a subset  $P$  of a space  $X$  in  $\mathcal{X}$ . Let  $Y$  be a space in  $\mathcal{X}$  and let  $f: Y \rightarrow X$  be a continuous function. Then  $\varphi \circ f$  is a  $\Gamma$ -norm on  $f^{-1}[P]$ . In particular, if  $P$  is  $\Gamma$ -complete, and  $P$  is the domain of a  $\Gamma$ -norm, then every member of  $\Gamma$  is the domain of some  $\Gamma$ -norm.

**4.2.3 Remark.** Let  $\Gamma$  be a pointclass. If  $\Gamma$  is selfdual and closed under intersections, then, for each  $A \in \Gamma$ , any constant function from  $A$  to the ordinals is a  $\Gamma$ -norm. If  $\Gamma$  is a lightface pointclass, and for each  $A \in \Gamma$  there is a constant  $\Gamma$ -norm on  $A$ , then  $\Gamma$  is selfdual.

**4.2.4 Remark.** Let  $\Gamma$  be a pointclass, let  $P$  be an element of  $\Gamma$  contained in a space  $X$  in  $\mathcal{X}$ . Let  $\varphi$  be a  $\Gamma$ -norm on  $P$ . For points  $x, y$  in  $X$ , the negation of  $x \leq_\varphi^* y$  is equivalent to the statement

$$x \notin P \vee (x, y \in P \wedge \varphi(y) < \varphi(x))$$

and the negation of  $x <_\varphi^* y$  is equivalent to the statement

$$x \notin P \vee (x, y \in P \wedge \varphi(y) \leq \varphi(x)).$$

**4.2.5 Remark.** It follows from Remark 4.2.4 that if  $\Gamma$  is a boldface pointclass closed under intersections and unions,  $\varphi$  is a  $\Gamma$ -norm on a set  $P$  and  $z \in P$ , then  $\{(x, y) \in P^2 : \varphi(x) \leq \varphi(y) < \varphi(z)\}$  is a prewellordering in  $\Gamma \cap \check{\Gamma}$ .

A pointclass  $\Gamma$  is said to have the *prewellordering property* if every member of  $\Gamma$  has a  $\Gamma$ -norm. We write  $\text{PWO}(\Gamma)$  to mean that  $\Gamma$  has the prewellordering property. The two following results derive boundedness and reduction properties from the prewellordering property.

**Proposition 4.2.6.** *Suppose that  $\Gamma$  is a  $\forall^{\omega^\omega}$ -closed boldface pointclass closed under unions, and that  $\varphi$  is a  $\Gamma$ -norm on a set  $P \in \Gamma \setminus \check{\Gamma}$ . Let  $Q$  be a subset of  $P$  in  $\check{\Gamma}$ . Then there exists a  $y \in P$  such that  $\varphi(x) < \varphi(y)$  for all  $x \in Q$ .*

*Proof.* Supposing otherwise,  $P \in \check{\Gamma}$ , as  $y \in P$  if and only if there exists an  $x \in Q$  such that  $x <_\varphi^* y$  fails.  $\square$

**Theorem 4.2.7.** *If  $\Gamma$  is a boldface pointclass closed under intersections and unions and satisfying the prewellordering property, then  $\text{Red}(\Gamma)$  holds.*

*Proof.* Let  $A$  and  $B$  be subsets of  $\omega^\omega$  in  $\Gamma$ . For each  $i \in 2$ , let  $f_i: \omega^\omega \rightarrow \omega^\omega$  be such that, for each  $x \in \omega^\omega$ ,  $f_i(x)(0) = i$  and  $f_i(x)(n+1) = x(n)$  for all  $n \in \omega$ . Let  $\varphi$  be a  $\Gamma$ -norm on  $f_0[A] \cup f_1[B]$ . Let  $A' = \{x \in \omega^\omega : f_0(x) \leq_\varphi^* f_1(x)\}$  and let  $B' = \{x \in \omega^\omega : f_1(x) <_\varphi^* f_0(x)\}$ . Then  $A'$  and  $B'$  reduce  $A$  and  $B$ .  $\square$

**4.2.8 Remark.** The pointclass  $\Pi_1^1$  has the prewellordering property (see 4B.2 of [32]). Adapting an argument of Novikov, Moschovakis showed that if  $\Gamma$  is a  $\forall^{\omega^\omega}$ -closed boldface pointclass and  $\text{PWO}(\Gamma)$  holds, then  $\text{PWO}(\exists^{\omega^\omega} \Gamma)$  also holds. The First Periodicity Theorem (due to Martin and Moschovakis) says that, assuming  $\text{AD} + \text{DC}_\mathbb{R}$ , if  $\Gamma$  is an  $\exists^{\omega^\omega}$ -closed boldface pointclass closed under intersections and  $\text{PWO}(\Gamma)$  holds, then  $\text{PWO}(\forall^{\omega^\omega} \Gamma)$  also holds. See 4B.3 and 6B.1 of [32].

**4.2.9 Remark.** Suppose that  $\Gamma$  is a boldface pointclass, and  $P$  is an element of  $\Gamma \setminus \check{\Gamma}$ , and  $\varphi$  is a  $\Gamma$ -norm on  $P$  with range some limit ordinal  $\gamma$ . By Remark 4.2.5,  $\varphi$  gives rise to a  $\gamma$ -sequence of sets in  $\Gamma \cap \check{\Gamma}$  whose union is  $P$ . In particular,  $\Gamma$  is not closed under  $\gamma$ -unions.

Given a set  $A \subseteq \omega^\omega$  and  $n \in \omega$ , a  $\Sigma_1^2(A)$  subset of  $(\omega^\omega)^n$  is a set of the form

$$\{(x_1, \dots, x_n) \in (\omega^\omega)^n : \exists B \subseteq \omega^\omega \varphi(A, B, x_1, \dots, x_n, y)\},$$

where the quantifiers in  $\varphi$  range over the hereditarily countable sets, and  $y$  is an element of  $\omega^\omega$ . If  $\varphi(A, B, x_1, \dots, x_n, y)$  holds for suitable  $A, B, x_1, \dots, x_n$  and  $y$ , we say that the formula  $\exists B \subseteq \omega^\omega \varphi(A, B, x_1, \dots, x_n, y)$  is *witnessed* by  $B$ . The pointclass  $\Sigma_1^2(A)$  is (by definition) the smallest boldface pointclass containing each  $\Sigma_1^2(A)$  subset of  $\omega^\omega$ .

**4.2.10 Remark.** It is straightforward to verify that each pointclass  $\Sigma_1^2(A)$  contains every subset of  $\omega^\omega$  projective in  $A$  and is closed under  $\exists^{\omega^\omega}$  and intersections. We do not know whether each such pointclass is closed under  $\forall^{\omega^\omega}$ , or even countable intersections, although they are in many natural models of AD. If the cardinal  $\Theta$  is regular (i.e., there is no cofinal function from  $\omega^\omega$  to  $\Theta$ ), then the  $\forall^{\omega^\omega}$ -closure of  $\Sigma_1^2(A)$  can be shown by taking a witness  $B$  of sufficiently high Wadge rank.

The pointclasses  $\Sigma_1^2(A)$  play an important role in Chapter 5. We introduce them here to give another example of a class of pointclasses with the prewellordering property (see Remark 4.2.12).

**4.2.11 Remark.** A set  $C \subseteq \omega^\omega$  is  $\Sigma_1^2(A)$  if, for some first order formula  $\varphi$  in the language of set theory, and some  $y \in \omega^\omega$ ,  $C$  is the set of  $x \in \omega^\omega$  for which there is an  $\omega$ -structure  $(M, E)$  containing  $\omega^\omega \cup \{A\}$  (where  $E$  is the  $\in$ -relation of  $M$ ) such that  $(M, E) \models \varphi(A, y, x)$ . Modifying this formulation one can define a universal  $\Sigma_1^2(A)$  set from the set of  $(n, x, y) \in \omega \times \omega^\omega \times \omega^\omega$  for which there exists an  $\omega$ -structure  $(M, E)$  containing  $\omega^\omega \cup \{A\}$  and a set  $T$  (naturally coded by a set of reals) such that  $T$  is the theory of  $M$  (in parameters from  $M$ , using the Gödel numbering for first order formulas) and  $n$  is the Gödel number of a formula  $\varphi$  such that  $(M, E) \models \varphi(A, x, y)$ . It follows from Proposition 2.4.2 that  $\Sigma_1^2(A)$  is nonselfdual.

We write  $\Pi_1^2(A)$  for  $\check{\Sigma}_1^2(A)$  (the class of complements of members of  $\Sigma_1^2(A)$ ), and  $\Delta_1^2$  for  $\Sigma_1^2(A) \cap \Pi_1^2(A)$ . We write  $\delta_1^2(A)$  for  $\delta(\Sigma_1^2(A))$ . Proposition 2.5.8 shows that, for each  $A \subseteq \omega^\omega$  whose Wadge rank is defined,  $\delta_1^2(A)$  is at least the Wadge rank of  $A$ .

**4.2.12 Remark.** Assuming  $\text{AD} + \text{DC}_\mathbb{R}$ , the pointclass  $\Sigma_1^2(A)$  has the prewellordering property for each  $A \subseteq \omega^\omega$ . To see this, fix a  $\Sigma_1^2(A)$  set  $X$  of the form

$$\{x \in \omega^\omega : \exists B \subseteq \omega^\omega \psi(A, B, x, z)\}.$$

Orders  $\leq^*$  and  $<^*$  witnessing the prewellordering property for  $X$  can be defined using the Wadge rank of a witness  $B$ . That is, define  $x \leq^* y$  to hold if there exists a  $B \subseteq \omega^\omega$  such that  $\psi(A, B, x, z)$  holds, and for every continuous function  $f: \omega^\omega \rightarrow \omega^\omega$  such that  $f^{-1}[B] <_W A$ ,  $\psi(A, f^{-1}[B], y, z)$  fails. Similarly, define  $x <^* y$  to hold if there exists a  $B \subseteq \omega^\omega$  such that  $\psi(A, B, x, z)$  holds, and for every continuous function  $f: \omega^\omega \rightarrow \omega^\omega$ ,  $\psi(A, f^{-1}[B], y, z)$  fails.

Selfdual boldface pointclasses which are closed under unions but not wellordered unions give rise to pointclasses with the prewellordering property.

**Theorem 4.2.13** (ZF; Martin). *Let  $\Delta$  be a selfdual boldface pointclass closed under unions. Suppose that  $\Delta$  is not closed under wellordered unions, and let  $\rho$  be the least ordinal  $\gamma$  such that  $\bigcup_\gamma \Delta \not\subseteq \Delta$ . Then  $\bigcup_\rho \Delta$  has the prewellordering property. Moreover, each element of  $\bigcup_\rho \Delta \setminus \Delta$ , has a prewellordering of length  $\rho$  whose proper initial segments are all in  $\Delta$ .*

*Proof.* Since  $\Delta$  is closed under unions,  $\rho$  is an infinite regular cardinal. By Remark 4.2.2, it suffices to find a  $\bigcup_\rho \Delta$ -norm as desired for an arbitrary  $A$  in  $\mathcal{P}(\omega^\omega) \cap \bigcup_\rho \Delta \setminus \Delta$ . Fix such an  $A$ . By the minimality of  $\rho$  and the assumptions on  $\Delta$ , there exists a function  $f: \rho \rightarrow \Delta \setminus \{\emptyset\}$  such that

- for all  $\alpha < \beta < \rho$ ,  $f(\alpha) \cap f(\beta) = \emptyset$ ;
- $A = \bigcup_{\alpha < \rho} f(\alpha)$ .

Define  $\varphi: A \rightarrow \rho$  by setting  $\varphi(x)$  to be the unique  $\alpha < \rho$  such that  $x \in f(\alpha)$ . Then  $\varphi$  is a  $\bigcup_\rho \Delta$ -norm on  $A$ . To see this, note first that, for all  $x, y$  in  $\omega^\omega$ ,  $x <_\varphi^* y$  if and only if there exists an  $\alpha < \rho$  such that  $x \in f(\alpha)$  and  $y \notin \bigcup_{\beta \leq \alpha} f(\beta)$ , and this set is in  $\bigcup_\rho \Delta$ . Similarly,  $x \leq_\varphi^* y$  if and only if there exists an  $\alpha < \rho$  such that  $x \in f(\alpha + 1)$  and  $y \notin \bigcup_{\beta \leq \alpha} f(\beta)$ , and this set is also in  $\bigcup_\rho \Delta$ . Finally, for each  $\gamma < \rho$ , the set  $\{(x, y) \in A : \varphi(x) \leq \varphi(y) < \gamma\}$  is a member of  $\Delta$ , being a union of fewer than  $\rho$  many sets from  $\Delta$ .  $\square$

**4.2.14 Remark.** Theorem 4.2.13 implies that if  $\Delta$  is a selfdual boldface pointclass containing a countable Wadge-cofinal subset but no Wadge-maximal set, then  $\bigcup_\omega \Delta$  has the prewellordering property. This holds in particular whenever  $\Delta$  is the pointclass of sets projective in a fixed subset of  $\omega^\omega$ . This shows moreover that if  $\Delta$  is a selfdual boldface pointclass closed under countable unions, and every element of  $\Delta$  is a member of subset of  $\Delta$  with a countable Wadge-cofinal subset but no Wadge-maximal set, then  $\Delta$  has the prewellordering property.

Along with Proposition 2.3.3, the following theorem shows, under the additional assumptions of Lipschitz Determinacy and Baire( $\mathcal{P}(\omega^\omega)$ ), that at most one member of any complementary pair of nonselfdual boldface pointclasses can have the prewellordering property. The proof is an adaptation of the proof of Theorem 4.1.4.

**Theorem 4.2.15.** *If  $\Gamma$  is a boldface pointclass which is closed under intersections and has a universal set, then  $\Gamma$  and  $\check{\Gamma}$  do not both have the prewellordering property.*

*Proof.* Let  $U \subseteq \omega^\omega \times \omega^\omega$  be a universal  $\Gamma$ -set, and let  $\varphi$  be a  $\Gamma$ -norm on  $U$ . Let  $\pi: \omega^\omega \rightarrow \omega^\omega \times \omega^\omega$  be a continuous bijection, and let  $\pi_0$  and  $\pi_1$  be such that  $\pi(x) = (\pi_0(x), \pi_1(x))$  for all  $x \in \omega^\omega$ . Let  $B$  be the set of  $(x, y)$  such that

$$(\pi_0(x), y) <_\varphi^* (\pi_1(x), y),$$

and let  $C$  be the set of  $(x, y)$  such that

$$(\pi_1(x), y) <_{\varphi}^* (\pi_0(x), y).$$

Then  $B$  and  $C$  are disjoint and in  $\Gamma$ .

Let  $\rho: \omega \times 2 \rightarrow \omega$  be a continuous bijection. For each  $i \in 2$ , let  $\rho^*$  be the function on  $\omega^\omega \times \omega^\omega$  defined by setting  $\rho_i^*(x, y)$  to be  $(\langle \rho(x(0), i), x(1), x(2), \dots \rangle, y)$ . Let  $P$  be the set

$$\rho_0^*[(\omega^\omega \times \omega^\omega) \setminus B] \cup \rho_1^*[(\omega^\omega \times \omega^\omega) \setminus C].$$

Then  $P$  is in  $\check{\Gamma}$ . Let  $\psi$  be a  $\check{\Gamma}$ -norm on  $P$ . Let  $E$  be the set of  $(x, y) \in \omega^\omega \times \omega^\omega$  such that

$$\rho_0^*(x, y) <_{\psi}^* \rho_1^*(x, y).$$

Since  $B \cap C = \emptyset$ ,  $E$  is also the set of  $(x, y)$  such that

$$\neg(\rho_1^*(x, y) \leq_{\psi}^* \rho_0^*(x, y)),$$

so  $E$  is in  $\Delta$ .

We derive a contradiction to Theorem 2.4.2 by showing that  $E$  is universal for  $\Delta$ . Fixing  $D \in \Delta$ , let  $x \in \omega^\omega$  be such that  $\omega^\omega \setminus D = U_{\pi_0(x)}$  and  $D = U_{\pi_1(x)}$ . Then  $\omega^\omega \setminus D = B_x$  and  $D = C_x$ , since, for all  $y \in \omega^\omega$ ,

$$y \in D \Leftrightarrow \pi_1(x), y \in U \Leftrightarrow \pi_0(x), y \notin U.$$

Finally, for each  $y \in \omega^\omega$ ,

$$(x, y) \in C \Rightarrow (\rho_0^*(x, y) \in P \wedge \rho_1^*(x, y) \notin P) \Rightarrow (x, y) \in E$$

and

$$(x, y) \in B \Rightarrow (\rho_0^*(x, y) \notin P \wedge \rho_1^*(x, y) \in P) \Rightarrow (x, y) \notin E.$$

It follows that  $E_x = D$ . □

Finally, the following theorem shows that for certain pointclasses, every witness to the prewellordering property has the same length. Part of the proof of Theorem 4.2.16 is reused for Theorem 4.3.3.

**Theorem 4.2.16** (Moschovakis). *Suppose that  $\Gamma$  is a  $\forall^{\omega^\omega}$ -closed boldface pointclass closed under unions, and that  $\langle U_n : n \in \omega \setminus \{0\} \rangle$  is a sequence of  $\Gamma$ -universal sets with the recursion property. Then for every  $n \in \omega \setminus \{0\}$  and every regular  $\Gamma$ -norm  $\varphi$  on  $U_n$ ,  $o(\varphi) = \delta(\Gamma)$ .*

*Proof.* Fix  $n \in \omega \setminus \{0\}$ , and let  $\varphi$  be a  $\Gamma$ -norm on  $U_n$ . By Remark 4.2.5, and the fact that  $\Gamma$  is closed under unions, we get that  $o(\varphi) \leq \delta(\Gamma)$ .

For the reverse inequality, let  $\prec$  be a strict prewellordering in  $\check{\Gamma}$  on a set  $Y \subseteq (\omega^\omega)^n$  (this is implied by the corresponding order  $\preceq$  being in  $\Gamma \cap \check{\Gamma}$ ). Since any two spaces in  $\mathcal{X}$  are homeomorphic, it suffices to show that the rank of  $\prec$  is at most  $o(\varphi)$ . Let  $Q$  be the set of  $(x, y) \in \omega^\omega \times (\omega^\omega)^n$  such that for all  $z \in (\omega^\omega)^n$ ,  $z \prec y$  implies  $(x, z) <_{\varphi}^* (x, y)$ . Then  $Q \in \Gamma$ . Applying the recursion property for  $U_n$ , we get an  $x^* \in \omega^\omega$  such that  $U_{n, x^*} = Q_{x^*}$ .

It suffices to see that the two following facts hold for all  $y \in Y$  :

- $y \in U_{n,x^*}$ ;
- for all  $z \in (\omega^\omega)^n$ ,  $z \prec y$  implies  $\varphi(x^*, z) < \varphi(x^*, y)$ .

This we do by induction on the  $\prec$ -rank of  $y$ . Fix  $y \in Y$ , and suppose that for all  $z \in (\omega^\omega)^n$  with  $z \prec y$ ,  $z$  is in  $U_{n,x^*}$  and, for all  $w \in (\omega^\omega)^n$ , if  $w \prec z$  then  $\varphi(x^*, w) < \varphi(x^*, z)$ . To see that  $y$  is in  $U_{n,x^*}$ , we show it is in  $Q_{x^*}$ , i.e., that for all  $z \in (\omega^\omega)^n$ , if  $z \prec y$  then  $(x^*, z) <_\varphi^* (x^*, y)$ . Fixing  $z \in (\omega^\omega)^n$  such that  $z \prec y$ , have that the failure of  $(x^*, z) <_\varphi^* (x^*, y)$  implies that either  $(x^*, z) \notin U_n$  or  $(x^*, y) \in U_n$  and  $\varphi(x^*, y) \leq \varphi(x^*, z)$ . Since  $(x^*, z)$  is in  $U_n$ , we have that  $(x^*, y)$  is in  $U_n$ . Finally, fix once again a  $z \in (\omega^\omega)^n$  such that  $z \prec y$ . Since  $(x^*, y) \in U_n$ , we have that  $(x^*, z) <_\varphi^* (x^*, y)$ , and (again applying the fact that  $(x^*, y)$  is in  $U_n$ ) this means that  $\varphi(x^*, z) < \varphi(x^*, y)$ , as desired.  $\square$

**4.2.17 Remark.** Suppose that  $\Gamma$  is a nonselfdual boldface pointclass closed under unions and intersections. Then  $\delta(\Gamma)$  is a limit ordinal, and (by Theorem 2.4.3),  $\Gamma$  has a universal set. Suppose that in addition Lipschitz Determinacy and Baire( $\mathcal{P}(\omega)$ ) hold, and that  $\Gamma$  is  $\forall^{\omega^\omega}$ -closed and has the prewellordering property. Theorems 2.4.9 and 4.2.16 and Remark 4.2.5 together then imply that  $\check{\Gamma}$  is not closed under unions of length  $\delta(\Gamma)$ .

### 4.3 Prewellorderings and wellfounded relations

Theorems 4.3.1 and 4.3.2 connect the length of prewellorderings in a boldface pointclass  $\Gamma$  with the closure of  $\Gamma$  under wellfounded unions. The other three theorems in this section relate the prewellordering property for a boldface nonselfdual pointclass with the supremum of the ranks of the wellfounded relations in associated pointclasses. Aside from Theorem 4.3.5, this section is based on material from [6]

**Theorem 4.3.1** (ZF + AD; Martin). *Let  $\Gamma$  be a nonselfdual boldface pointclass closed under  $\exists^{\omega^\omega}$  and intersections. Let  $\rho$  be the length of some strict prewellordering in  $\Gamma$ . Then  $\bigcup_\rho \Gamma \subseteq \Gamma$ .*

*Proof.* Fix  $f: \rho \rightarrow \Gamma \cap \mathcal{P}(\omega^\omega)$ , and let  $F = \bigcup_{\alpha < \rho} f(\alpha)$ . We want to see that  $F$  is in  $\Gamma$ . Let  $\prec$  be a strict prewellordering in  $\Gamma$  of length  $\rho$ . Let  $X$  be the domain of  $\prec$ , and for each  $x \in X$  let  $\alpha_x$  denote the  $\prec$ -rank of  $x$ . Applying Theorem 2.4.3, let  $U \subseteq (\omega^\omega)^2$  be a universal  $\Gamma$ -set, and let  $Z$  be  $\{(x, y) \in X \times \omega^\omega : U_y = f(\alpha_x)\}$ .

By the Coding Lemma (Theorem 3.0.1), there is a set  $A \subseteq Z$  in  $\text{pos-}\Sigma_1^1(\prec)$  (which is contained in  $\Gamma$ , since  $\Gamma$  is  $\exists^{\omega^\omega}$ -closed and closed under intersections) such that for all  $x \in X$ , if  $Z_x \neq \emptyset$  then there is an  $x' \in X$  with  $\alpha_x = \alpha_{x'}$  and  $A_{x'} \neq \emptyset$ . Then for all  $z \in \omega^\omega$ ,  $z \in F$  if and only if there exists a pair  $(x, y) \in A$  with  $(y, z) \in U$ .  $\square$

**Theorem 4.3.2** (ZF + AD; Martin). *Let  $\Gamma$  be a nonselfdual pointclass closed under  $\forall^{\omega^\omega}$  and unions, and assume that  $\Gamma$  has the prewellordering property. Let  $\Delta = \Gamma \cap \check{\Gamma}$ . Then  $\Delta$  is closed under wellordered unions and intersections of length less than  $\delta(\Gamma)$ .*

*Proof.* Since  $\Delta$  is closed under complements, it suffices to see that it is closed under unions of length less than  $\delta(\Gamma)$ . Supposing that it is not, let  $\rho < \delta(\Gamma)$  be minimal such that there exists a function  $f: \rho \rightarrow \Delta$  with  $\bigcup_{\alpha < \rho} f(\alpha) \notin \Delta$ . Since  $\Delta$  is closed under unions,  $\rho$  is a limit ordinal. By Theorem 4.2.13, the pointclass  $\bigcup_\rho \Delta$  has the prewellordering property. By Theorems 2.4.9 and 4.2.16, and Remark 4.2.5, there is a prewellordering in  $\Delta$  of length  $\rho$ . Since  $\Delta$  is closed under complements and intersections, the corresponding strict prewellordering is in  $\Delta$  and therefore in  $\check{\Gamma}$ . By Theorem 4.3.1,  $\bigcup_\rho \Delta \subseteq \check{\Gamma}$ . Since  $\bigcup_\rho \Delta$  is not contained in  $\Delta$ , it must be (by Lipschitz Determinacy) that  $\bigcup_\rho \Delta = \check{\Gamma}$ , so  $\text{PWO}(\check{\Gamma})$  holds. We then have a contradiction to Theorem 4.2.15.  $\square$

The proof of the following theorem is similar to that of Theorem 4.2.16, and a similar fact makes up part of Theorem 4.3.5.

**Theorem 4.3.3** (Moschovakis). *Suppose that Lipschitz Determinacy holds, and that every subset of  $\omega^\omega$  has the property of Baire. Let  $\Gamma$  be a nonselfdual boldface pointclass closed under  $\forall^{\omega^\omega}$  and unions, and suppose that  $\text{PWO}(\Gamma)$  holds. Then  $\delta(\Gamma)$  is the supremum of the ranks of the wellfounded relations in  $\check{\Gamma}$ .*

*Proof.* That  $\delta(\Gamma)$  is at most the supremum of the ranks of the wellfounded relations in  $\check{\Gamma}$  follows immediately from its definition. We show that every wellfounded relation in  $\check{\Gamma}$  has length less than  $\delta(\Gamma)$ .

Let  $\prec$  be a strict wellfounded relation in  $\check{\Gamma}$  on a set  $X \in \check{\Gamma} \cap \mathcal{P}(\omega^\omega)$ . We will show that the rank of  $\prec$  is at most  $\delta(\Gamma)$ . This will suffice, since there are strict wellfounded relations in  $\check{\Gamma}$  whose rank is more than that of  $\prec$  (one more, for instance), and the argument will apply to these relations also.

Applying Theorem 2.4.9, let  $\bar{U} = \langle U_n : n < \omega \setminus \{0\} \rangle$  be a sequence of universal sets for  $\Gamma$  with the recursion property. Let  $\varphi$  be a regular  $\Gamma$ -norm on  $U_2$ . By Theorem 4.2.16,  $o(\varphi) = \delta(\Gamma)$ .

Let  $A$  be the set of  $(x, y) \in \omega^\omega \times \omega^\omega$  such that, for all  $z \in X$ , if  $z \prec y$  then  $(x, z) \prec_\varphi^* (x, y)$ . By the recursion property of  $U_1$ , there is an  $x \in \omega^\omega$  such that  $A_x = U_{1,x}$ .

We verify first by induction on the  $\prec$ -rank of  $y \in \omega^\omega$  that  $(x, y) \in U_1$  for all  $y \in X$ . First note that if  $(x, y) \notin U_1$  and  $(x, z)$  is in  $U_1$  for all  $z \prec y$ , then  $(x, y)$  is in  $A$ , which implies that in fact  $(x, y)$  is in  $U_1$ . It follows then that for all  $y, z \in X$ , if  $z \prec y$  then  $\varphi(x, z) < \varphi(x, y)$ , as desired.  $\square$

The following theorem can be used in conjunction with Theorem 4.3.5 to show (assuming  $\text{AD} + \text{DC}_{\mathbb{R}}$ ) that, for certain selfdual boldface pointclasses  $\Delta$ , the supremum of the Wadge ranks of the members of  $\Delta$  is a regular cardinal, .

**Theorem 4.3.4** (ZF + AD). *If  $\Gamma$  is a nonselfdual boldface pointclass satisfying the prewellordering property and closed under  $\forall^{\omega^\omega}$  and unions then the supremum of the ranks of the wellfounded relations in  $\check{\Gamma}$  is a regular cardinal.*

*Proof.* Let  $\kappa$  be the supremum of the ranks of the wellfounded relations in  $\check{\Gamma}$ . Since  $\check{\Gamma}$  is closed under unions,  $\kappa$  is a limit ordinal, and all wellfounded relations in  $\check{\Gamma}$  have rank less than  $\kappa$ . Let  $\rho$  be the cofinality of  $\kappa$ , suppose that



$\rho < \kappa$ , and let  $f: \rho \rightarrow \kappa$  be cofinal. Applying Theorem 4.3.3, let  $\prec$  be a strict prewellordering in  $\Gamma \cap \check{\Gamma}$  of length  $\rho$ . Let  $X$  be the domain of  $\prec$ , and, for each  $x \in X$ , let  $\alpha_x$  be the  $\prec$ -rank of  $x$ . Applying Theorem 2.4.3, let  $U \subseteq (\omega^\omega)^3$  be a universal  $\check{\Gamma}$  set, and let  $Z$  be the set of  $(x, y) \in X \times \omega^\omega$  for which  $U_y$  is a wellfounded relation of length at least  $f(\alpha_x)$ . By the Coding Lemma (Theorem 3.0.1), there exists a pos- $\Sigma_1^1(\prec)$ -set  $A \subseteq Z$  such that, for all  $x \in X$ , if  $Z_x \neq \emptyset$  then  $A_x \neq \emptyset$ . Then  $A$  is in  $\check{\Gamma}$ . Now define the relation  $<$  on  $(\omega^\omega)^3$  by setting  $(x, y, z) < (a, b, c)$  if and only if  $(x, y) \in A$ ,  $(x, y) = (a, b)$  and  $(y, z, c) \in U$ . Then  $<$  is a wellfounded relation in  $\Gamma$  of rank  $\kappa$ , giving a contradiction.  $\square$

A *projective algebra* is a selfdual boldface pointclass containing all closed subsets of spaces in  $\mathcal{X}$  and closed under real quantification (some authors, e.g. [18], include closure under wellordered unions in the definition of projective algebra). Given a pointclass  $\Gamma$ , we write  $o(\Gamma)$  for the set of Wadge ranks of elements of  $\Gamma \cap \mathcal{P}(\omega^\omega)$ . The following is a weak version of a theorem from [18].

**Theorem 4.3.5** (ZF + AD). *Let  $\Gamma$  be a nonselfdual boldface pointclass closed under  $\forall^{\omega^\omega}$  and unions, and suppose that  $\text{PWO}(\Gamma)$  holds. Let  $\Delta = \Gamma \cap \check{\Gamma}$ , and suppose that  $\Delta$  is a projective algebra. Then  $\delta(\Gamma)$  is equal to the supremum of the ranks of the wellfounded relations in  $\Delta$ . If in addition the Wadge rank of each element of  $\Delta$  exists, then  $\delta(\Gamma) = o(\Delta)$ .*

*Proof.* Let  $\kappa$  be the supremum of the ranks of the wellfounded relations in  $\Delta$ . That  $\delta(\Gamma) \leq \kappa$  follows immediately; that  $\delta(\Gamma) = \kappa$  follows from Theorem 4.3.3. That  $o(\Delta) \leq \delta(\Gamma)$  follows from Proposition 2.5.8.

Assume now that the Wadge rank of each element of  $\Delta$  exists. To see that  $\delta(\Gamma) \leq o(\Delta)$ , note first by Theorem 4.3.2 that  $\Delta$  is closed under unions of length less than  $\delta(\Gamma)$ . One can then construct sets in  $\Delta$  of Wadge ranks cofinal in  $\delta(\Gamma)$  by following Solovay's construction, as in Theorem 2.5.9 and Remark 2.5.10.  $\square$

## 4.4 Closure under wellordered unions

This section gives applications of the material in Section 4.2 and 4.3, showing that certain pointclasses are closed under wellordered unions. These results will be used in Chapters 6 and 8. Theorem 4.4.1 is Theorem 2.1 of [7].

**Theorem 4.4.1** (ZF+AD). *Let  $\Gamma$  be a boldface nonselfdual pointclass containing  $\Sigma_1^1$  which is closed under  $\exists^{\omega^\omega}$  and  $\forall^{\omega^\omega}$ . Then at least one of  $\Gamma$  and  $\check{\Gamma}$  is closed under wellordered unions.*

*Proof.* Supposing otherwise, let  $\kappa$  be the least ordinal  $\gamma$  such that  $\Gamma$  is not closed under  $\gamma$ -unions, and let  $\lambda$  be the least ordinal  $\gamma$  such that  $\check{\Gamma}$  is not closed under  $\gamma$ -unions. Then  $\kappa$  and  $\lambda$  are regular cardinals. Without loss of generality,  $\lambda \leq \kappa$ . We will find a strict prewellordering in  $\check{\Gamma}$  of length  $\kappa$ . By Theorem 4.3.1, this will contradict the choice of  $\lambda$ .

By Corollary 2.3.2,  $\check{\Gamma} \subseteq \bigcup_\kappa \Gamma$ . Fix  $A \subseteq \omega^\omega$  in  $\check{\Gamma} \setminus \Gamma$ , and let  $\langle A_\alpha : \alpha < \kappa \rangle \in \Gamma^\kappa$  be such that  $A = \bigcup_{\alpha < \kappa} A_\alpha$ . Applying Theorem 2.4.3, let  $U \subseteq (\omega^\omega)^2$  be a

universal  $\Gamma$  set, and let  $B = \{x \in \omega^\omega : U_x \subseteq A\}$ . Since  $\check{\Gamma}$  is  $\forall^{\omega^\omega}$ -closed,  $B \in \check{\Gamma}$ . There exists then a sequence  $\langle B_\alpha : \alpha < \kappa \rangle \in \Gamma^\kappa$  such that  $B = \bigcup_{\alpha < \kappa} B_\alpha$ . For each  $\alpha < \kappa$ , let  $C_\alpha = \bigcup_{x \in B_\alpha} U_x$ . Then  $A = \bigcup_{\alpha < \kappa} C_\alpha$ , and every subset of  $A$  in  $\Gamma$  is contained in  $C_\alpha$  for some  $\alpha < \kappa$ . Since each  $C_\alpha$  is in  $\Gamma$ ,  $A$  is not contained in  $\bigcup_{\alpha < \gamma} C_\alpha$  for any  $\gamma < \kappa$ . For each  $x \in A$ , let  $\alpha_x$  be the least  $\alpha$  with  $x \in A_\alpha$ .

Let  $\mathcal{G}$  be the game where  $I$  plays  $x \in \omega^\omega$ ,  $II$  plays  $y$  and  $z$ , and  $II$  wins if and only if  $x \notin A$  or

$$\exists \beta > \alpha_x (U_y = \bigcup_{\alpha \leq \beta} C_\alpha \wedge z \in C_\beta \setminus \bigcup_{\alpha < \beta} C_\alpha).$$

Since every subset of  $A$  in  $\Gamma$  is contained in some  $C_\alpha$ ,  $I$  cannot have a winning strategy in  $\mathcal{G}$ . Let  $\tau$  be a winning strategy for  $II$ . Given  $x \in \omega^\omega$ , let  $y_\tau(x)$  and  $z_\tau(x)$  denote, respectively, the  $y$  and  $z$  parts of the response to  $x$  (played by  $I$ ) given by  $\tau$ , and, when  $x \in A$ , let  $\beta_\tau(x)$  be such that  $U_{y_\tau(x)} = \bigcup_{\alpha \leq \beta_\tau(x)} C_\alpha$ . Define the binary relation  $\prec$  on  $A$  by setting  $x_0 \prec x_1$  to hold if and only if  $z_\tau(x_1) \notin U_{y_\tau(x_0)}$ . Then for all  $x_0, x_1$  in  $A$ ,  $x_0 \prec x_1$  holds if and only if  $\beta_\tau(x_0) < \beta_\tau(x_1)$ . It follows that  $\prec$  is a strict prewellordering in  $\check{\Gamma}$  of length  $\kappa$ .  $\square$

**4.4.2 Remark.** Remark 4.2.17 and Theorem 4.4.1 together imply that if AD holds and  $\Gamma$  is a boldface nonselfdual pointclass containing  $\Sigma_1^1$  and closed under  $\exists^{\omega^\omega}$  and  $\forall^{\omega^\omega}$ , and PWO( $\Gamma$ ) holds, then  $\Gamma$  is closed under wellordered unions.

Lemma 4.4.3 and Theorem 4.4.5 below concern pointclasses which are closed under  $\exists^{\omega^\omega}$  but not  $\forall^{\omega^\omega}$ . The proofs of Lemma 4.4.3 and Theorem 4.4.4 are extracted from the proof of Lemma 2.18 of [6]. The proof of the first part of Lemma 4.4.3 is very similar to the proof of Theorem 4.4.1.

**Lemma 4.4.3** (ZF + AD). *Suppose that  $\Gamma$  is a nonselfdual boldface pointclass containing  $\Sigma_1^1$  and closed under  $\exists^{\omega^\omega}$  and finite intersections, but not  $\forall^{\omega^\omega}$  or wellordered unions.*

1. *Let  $\Gamma_1$  be  $\exists^{\omega^\omega} \check{\Gamma}$  and let  $\kappa$  be the least ordinal  $\gamma$  such that  $\bigcup_\gamma \Gamma \not\subseteq \Gamma$ . Then  $\bigcup_\kappa \Gamma_1 = \Gamma_1 = \bigcup_\kappa \Gamma$  and  $\Gamma_1$  is nonselfdual.*
2. *Let  $\Delta_1 = \Gamma_1 \cap \check{\Gamma}_1$ . Then  $\Delta_1$  is not closed under wellorded unions. Letting  $\rho$  be the least ordinal  $\gamma$  such that  $\bigcup_\gamma \Delta_1 \not\subseteq \Delta_1$ ,  $\rho \leq \kappa$ ,  $\Gamma_1 = \bigcup_\rho \Delta_1$  and  $\Gamma_1$  has the prewellordering property.*

*Proof.* Note that  $\kappa$  is a regular cardinal. Since  $\Gamma$  is  $\exists^{\omega^\omega}$ -closed, but not  $\forall^{\omega^\omega}$ -closed,  $\Gamma$  is nonselfdual. It follows by Theorem 2.4.3 and Remark 2.4.4 that  $\Gamma_1$  is also nonselfdual. By Corollary 2.3.2,  $\check{\Gamma} \subseteq \bigcup_\kappa \Gamma$  and  $\Gamma \subseteq \Gamma_1$ . Since  $\bigcup_\kappa \Gamma$  is closed under  $\exists^{\omega^\omega}$ ,  $\Gamma_1$  is also contained in  $\bigcup_\kappa \Gamma$ . We will show first that there is a strict prewellordering of length  $\kappa$  in  $\Gamma_1$ . From this and Theorem 4.3.1 it will follow that  $\bigcup_\kappa \Gamma_1 \subseteq \Gamma_1$ , from which the first part of the lemma follows.

Let  $A \subseteq \omega^\omega$  be any set in  $\check{\Gamma} \setminus \Gamma$ . Let  $S \subseteq (\omega^\omega)^2$  be a universal  $\Sigma_1^1$  set, and let  $B = \{x \in \omega^\omega : S_x \subseteq A\}$ . Since  $\check{\Gamma}$  is closed under  $\forall^{\omega^\omega}$  and unions,  $B \in \check{\Gamma}$ ,

so  $B$  can be written as  $\bigcup_{\alpha < \kappa} B_\alpha$ , where each  $B_\alpha$  is in  $\Gamma$ . For each  $\alpha < \kappa$ , let  $A_\alpha = \bigcup_{x \in B_\alpha} S_x$ . Then  $A = \bigcup_{\alpha < \kappa} A_\alpha$ , and every  $\Sigma_1^1$  subset of  $A$  (being  $S_x$  for some  $x \in \omega^\omega$ ) is contained in some  $A_\alpha$ . Since each  $A_\alpha$  is in  $\Gamma$ ,  $A$  is not contained in  $\bigcup_{\alpha < \gamma} A_\alpha$  for any  $\gamma < \kappa$ . For each  $x \in A$ , let  $\alpha_x$  be the least  $\alpha$  with  $x \in A_\alpha$ .

Let  $U \subseteq (\omega^\omega)^2$  be a universal  $\Gamma$  set. Let  $\mathcal{G}$  be the game where  $I$  plays  $x \in \omega^\omega$ ,  $II$  plays  $y$  and  $z$ , and  $II$  wins if and only if  $x \notin A$  or

$$\exists \beta > \alpha_x (U_y = \bigcup_{\alpha \leq \beta} A_\alpha \wedge z \in A_\beta \setminus \bigcup_{\alpha < \beta} A_\alpha).$$

Since every  $\Sigma_1^1$  subset of  $A$  is contained in some  $A_\alpha$ ,  $I$  cannot have a winning strategy in  $\mathcal{G}$ . Let  $\tau$  be a winning strategy for  $II$ . Given  $x \in \omega^\omega$ , let  $y_\tau(x)$  and  $z_\tau(x)$  denote, respectively, the  $y$  and  $z$  parts of the response to  $x$  (played by  $I$ ) given by  $\tau$ , and, when  $x \in A$ , let  $\beta_\tau(x)$  be such that  $U_{y_\tau(x)} = \bigcup_{\alpha \leq \beta_\tau(x)} A_\alpha$ . Define the binary relation  $\prec$  on  $A$  by setting  $x_0 \prec x_1$  to hold if and only if  $z_\tau(x_1) \notin U_{y_\tau(x_0)}$ . Then for all  $x_0, x_1$  in  $A$ ,  $x_0 \prec x_1$  holds if and only if  $\beta_\tau(x_0) < \beta_\tau(x_1)$ . It follows that  $\prec$  is a strict prewellordering in  $\check{\Gamma}$  of length  $\kappa$ .

For the second part of the lemma, since  $\Gamma \subseteq \Delta_1$  and  $\bigcup_\kappa \Gamma$  is nonselfdual,  $\Delta_1$  is not closed under wellordered unions, and  $\rho \leq \kappa$ . Since  $\bigcup_\rho \Delta_1$  properly contains  $\Delta_1$  and is contained in  $\bigcup_\kappa \Gamma_1 = \Gamma_1$ ,  $\Gamma_1 = \bigcup_\rho \Delta_1$ , by Proposition 2.3.3. It follows from Theorem 4.2.13 that  $\Gamma_1$  has the prewellordering property.  $\square$

The last paragraph of the proof of the following theorem is taken from the proof of Lemma 2.18 of [6].

**Theorem 4.4.4** (ZF + AD + DC $_{\mathbb{R}}$ ). *Suppose that  $\Delta$  is a projective algebra such that  $\bigcup_\omega \Delta \not\subseteq \Delta$ . Then  $\bigcup_\omega \Delta$  is nonselfdual, has the prewellordering property and is closed under wellordered unions.*

*Proof.* Since  $\Delta$  is a projective algebra and  $\bigcup_\omega \Delta \not\subseteq \Delta$ , there exists a sequence  $\langle B_i : i \in \omega \rangle \in (\mathcal{P}(\omega^\omega) \setminus \omega^\omega)^\omega$  whose range is Wadge cofinal in  $\Delta$ . Let  $\pi : \omega^\omega \rightarrow (\omega^\omega)^\omega$  be a continuous bijection, and let  $\pi_i$  ( $i \in \omega$ ) be the functions on  $\omega^\omega$  such that  $\pi(x) = \langle \pi_i(x) : i \in \omega \rangle$  for all  $x \in \omega^\omega$ . Recalling our coding in Remark 2.5.2 of continuous functions  $f_x^c$  by  $x \in \mathcal{F}^c \subseteq \omega^\omega$ , for each  $i \in \omega$  let  $C_i$  be the set of  $(x, y) \in (\omega^\omega)^2$  such that  $\pi_i(x) \in \mathcal{F}^c$  and  $f_{\pi_i(x)}^c(y) \in B_i$ . Since  $\Delta$  is a projective algebra, each  $C_i$  is in  $\Delta$ . Let  $C = \bigcup_{i \in \omega} C_i$ .

We claim that  $C$  is universal for the pointclass  $\bigcup_\omega \Delta$ . To verify this, consider a set  $\bigcup_{i \in \omega} A_i$ , where each  $A_i$  is in  $\Delta$ . Applying CC $_{\mathbb{R}}$ , we may find  $E_i$  and  $x \in \omega^\omega$  such that  $\{\pi_i(x) : i \in \omega\} \subseteq \mathcal{F}^c$ , each  $E_i$  is  $(f_{x_i}^c)^{-1}[B_i]$  and  $\bigcup_{i \in \omega} A_i = \bigcup_{i \in \omega} E_i$ . Then  $\bigcup_{i \in \omega} E_i = C_x$ . Since  $\bigcup_\omega \Delta$  has a universal set, it is nonselfdual, by Proposition 2.4.2.

Let  $\Gamma = \bigcup_\omega \Delta$  and let  $\Gamma_1 = \exists^{\omega^\omega} \check{\Gamma}$ . Then  $\Gamma$  is closed under  $\exists^{\omega^\omega}$  and intersections. It follows by Theorem 4.2.13 that  $\Gamma$  has the prewellordering property. By the First Periodicity Theorem (see Remark 4.2.8),  $\check{\Gamma}_1 = \forall^{\omega^\omega} \Gamma$  does as well. It follows by Theorem 4.2.15 that  $\Gamma_1$  does not have the prewellordering property. It follows then by the second part of Lemma 4.4.3 that  $\Gamma$  must be  $\forall^{\omega^\omega}$ -closed or closed under wellordered unions. However, it follows by Remark 4.4.2 that  $\Gamma$  is closed under wellordered unions even if it is  $\forall^{\omega^\omega}$ -closed.  $\square$

The following is Lemma 2.4 of [18]. It implies Corollary 4.4.6 below, which we will use in Chapter 6 to analyze pointclasses of Suslin sets.

**Theorem 4.4.5** (ZF + AD). *Let  $\Gamma$  be an  $\exists^{\omega^\omega}$ -closed boldface pointclass containing  $\Sigma_1^1$  which is closed under countable unions and intersections but not  $\forall^{\omega^\omega}$ -closed. If  $\text{Red}(\Gamma)$  holds, then  $\Gamma$  is closed under wellordered unions.*

*Proof.* Since  $\Gamma$  is  $\exists^{\omega^\omega}$ -closed and not  $\forall^{\omega^\omega}$ -closed,  $\Gamma$  is nonselfdual. Working toward a contradiction, let  $\kappa$  be the least ordinal  $\gamma$  such that for some  $f: \gamma \rightarrow \Gamma$ ,  $\bigcup_{\alpha < \gamma} f(\alpha) \notin \Gamma$ . Then  $\kappa$  is a regular uncountable cardinal. Fix such a function  $f$ , and let  $F = \bigcup_{\alpha < \kappa} f(\alpha)$ . By Theorem 2.1.4, the boldface pointclass  $\bigcup_\kappa \Gamma$  contains both  $\Gamma$  and  $\check{\Gamma}$ . Letting  $\Gamma_1 = \exists^{\omega^\omega} \check{\Gamma}$  and  $\Delta_1 = \Gamma_1 \cap \check{\Gamma}_1$ , we have the following by Lemma 4.4.3:

- $\Gamma \subseteq \Gamma_1 = \bigcup_\kappa \Gamma_1 = \bigcup_\kappa \Gamma$ ;
- $\Gamma_1$  is nonselfdual;
- $\Gamma \subseteq \Delta_1$ , which is not closed under wellordered unions;
- letting  $\rho$  be the least ordinal  $\gamma$  such that  $\bigcup_\rho \Delta_1 \not\subseteq \Delta_1$ ,  $\rho \leq \kappa$ ,  $\Gamma_1 = \bigcup_\rho \Delta_1$ ;
- $\Gamma_1$  has the prewellordering property, and thus the reduction property, by Theorem 4.2.7.

By Theorem 4.1.8 and the assumption that  $\Gamma$  has the reduction property,  $\check{\Gamma}_1$  has the reduction property. We now have a contradiction, by Remark 4.1.5.  $\square$

**Corollary 4.4.6** (ZF + AD). *If  $\Delta$  is a projective algebra then every member of  $\Delta$  is an element of a nonselfdual  $\exists^{\omega^\omega}$ -closed pointclass contained in  $\Delta$  and closed under wellordered unions.*

*Proof.* Fix  $A \in \Delta \cap \mathcal{P}(\omega^\omega)$ . The pointclasses  $\Sigma_1^1(A)$  and  $\Sigma_2^1(A)$  have universal sets, so they are nonselfdual and properly contained in  $\Delta$ . Moreover, they are closed under countable unions and intersections but not  $\forall^{\omega^\omega}$ -closed (since if they were they would contain their complement classes). By Remark 4.1.5 and Theorem 4.1.8, one of these two pointclasses satisfies the reduction property and (by Theorem 4.4.5) is closed under wellordered unions.  $\square$

## Chapter 5

# Strong partition cardinals

In this chapter we introduce strong partition cardinals and prove that under AD they are cofinal below  $\Theta$ . Again, the material presented here is a small part of a much larger story.

Given an ordinal  $\alpha$  and a set  $X$  of ordinals,  $[X]^\alpha$  denotes the collection of subsets of  $X$  of ordertype  $\alpha$ . For notational convenience, we identify each element of  $[X]^\alpha$  with the corresponding order-preserving function on  $\alpha$  which enumerates it. Given ordinals  $\alpha, \beta, \delta$  and  $\gamma$  the formula  $\alpha \rightarrow (\beta)_\gamma^\delta$  asserts that for every partition of  $[\alpha]^\delta$  into  $\gamma$  many pieces, there exists an  $X \in [\alpha]^\beta$  such that  $[X]^\delta$  is contained in one piece. We write  $\alpha \rightarrow (\beta)^\delta$  when  $\gamma = 2$ . A *strong partition cardinal* is a regular uncountable cardinal  $\kappa$  such that  $\kappa \rightarrow (\kappa)_\mu^\kappa$  holds for all  $\mu < \kappa$ .

**5.0.1 Remark.** Some authors define strong partition cardinals using the ostensibly weaker property  $\kappa \rightarrow (\kappa)^\kappa$ . We do not know whether ZF implies that these two definitions are equivalent. However, if  $\kappa \rightarrow (\kappa)^\kappa$  holds, then  $\kappa$  is a regular cardinal and  $\kappa \rightarrow (\kappa)_\mu^\mu$  holds for all  $\mu < \kappa$ . The latter claim can be seen by considering, given an  $f: [\kappa]^\mu \rightarrow \mu$ , the function  $f': [\kappa]^\kappa \rightarrow 2$  defined by setting  $f'(X) = 0$  if and only if  $f(a) = f(b)$ , where  $a$  is the initial segment of  $X$  of ordertype  $\mu$ , and  $b$  is the initial segment of  $X \setminus a$  of ordertype  $\mu$ . The property  $\forall \mu < \kappa (\kappa \rightarrow (\kappa)_\mu^\mu)$  (in particular its consequence given in Theorem 5.0.3 below) is enough for our application of strong partition cardinals in Theorem 7.0.1.

**5.0.2 Remark.** A theorem of Kleinberg (Theorem 28.10 of [10]) shows that if  $\kappa \rightarrow (\kappa)_2^{\omega_1 + \omega_1}$ , then  $\kappa$  is measurable.

The following notation will be used in Theorem 5.0.3, and again in the proof of Theorem 7.0.1. Given an ordinal-valued function  $g$  on  $\omega \cdot \alpha$ , for some ordinal  $\alpha$ , let  $g^*$  be the function on  $\alpha$  defined by setting  $g^*(\xi)$  to be the supremum of  $\{g(\omega \cdot \xi + n) : n \in \omega\}$ . For each set  $X$  of ordinals, let  $X^*(\alpha)$  be  $\{g^* : g \in [X]^{\omega \cdot \alpha}\}$ . Given an ordinal  $\delta$ , let  $\mu_{\delta, \alpha}$  be the set of  $A \subseteq [\delta]^\alpha$  for which there exists a club  $C \subseteq \delta$  with  $C^*(\alpha) \subseteq A$ .

**Theorem 5.0.3.** *Let  $\delta$  be a regular uncountable cardinal and let  $\alpha < \delta$  be such that  $\delta \rightarrow (\delta)_\rho^{\omega \cdot \alpha}$  holds for all  $\rho < \delta$ . Then  $\mu_{\delta, \alpha}$  is a  $\delta$ -complete ultrafilter on  $[\delta]^\alpha$ .*

*Proof.* Fix  $\rho < \delta$  and suppose that  $[\delta]^\alpha$  is the union of sets  $A_\beta$  ( $\beta < \rho$ ). For each  $\beta < \rho$ , let  $B_\beta$  be  $\{g \in [\delta]^{\omega \cdot \alpha} : g^* \in A_\beta\}$ . Since  $\delta \rightarrow (\delta)_\rho^{\omega \cdot \alpha}$  holds, there exist an  $H \in [\delta]^\delta$  and a  $\beta < \rho$  such that  $[H]^{\omega \cdot \alpha} \subseteq B_\beta$ . Let  $C$  be the set of limit points of  $H$ . Then  $C$  is a club subset of  $\delta$ . Given  $h \in [C]^{\omega \cdot \alpha}$ , define  $g \in [H]^{\omega \cdot \alpha}$  by setting  $g(\gamma)$  to be  $\min(H \setminus h(\gamma))$ . Then  $h^* = g^*$ , which is in  $A_\beta$ . It follows that  $C^*(\alpha) \subseteq A_\beta$ .  $\square$

Our aim in this chapter is to prove the following theorem, which is due to Kechris, Kleinberg, Moschovakis and Woodin [14].

**Theorem 5.0.4.** *If AD holds then the strong partition cardinals are cofinal below  $\Theta$ .*

The proof uses the pointclasses  $\Sigma_1^2(A)$  and their associated ordinals  $\delta_1^2(A)$  as defined in Section 4.2. We do not know whether AD implies that cardinals of the form  $\delta_1^2(A)$  are strong partition cardinals. One problem is that we do not know how to show that the pointclass  $\Sigma_1^2(A)$  has the prewellordering property without some additional assumption. In Remark 4.2.12 we show this using the additional assumption of  $\text{DC}_\mathbb{R}$ , which implies that the Wadge hierarchy is wellfounded. A second problem is that we do not know whether AD implies that  $\Sigma_1^2(A)$  is  $\forall^{\omega^\omega}$ -closed, although it is in many natural models of AD. Adding these two properties as hypotheses, we get the following.

**Theorem 5.0.5** ( $\text{ZF} + \text{AD} + \text{DC}_\mathbb{R}$ ). *For each  $A \subseteq \omega^\omega$  for which  $\Sigma_1^2(A)$  is  $\forall^{\omega^\omega}$ -closed,  $\delta_1^2(A)$  is a strong partition cardinal.*

To get around these issues, we relativize  $\Sigma_1^2(A)$  to the inner model  $L(A, \mathbb{R})$ . We write  $\text{loc-}\Sigma_1^2(A)$  for

$$\Sigma_1^2(A)^{L(A, \mathbb{R})},$$

$\text{loc-}\Delta_1^2(A)$  for  $\Delta_1^2(A)^{L(A, \mathbb{R})}$  and  $\text{loc-}\delta_1^2(A)$  for  $\delta_1^2(A)^{L(A, \mathbb{R})}$ . As with  $\Sigma_1^2(A)$ , each pointclass  $\text{loc-}\Sigma_1^2(A)$  is nonselfdual, has a universal set and closed under  $\exists^{\omega^\omega}$  and intersections. Since  $\Theta$  is regular in models of the form  $L(A, \mathbb{R})$ , pointclasses of the form  $\text{loc-}\Sigma_1^2(A)$  are also  $\forall^{\omega^\omega}$ -closed.

The Solovay Basis Theorem (Theorem 5.0.6 below) says that for each  $A \subseteq \omega^\omega$ ,  $\text{loc-}\Delta_1^2(A)$  is  $\Sigma_1$ -elementary in  $\mathcal{P}(\omega^\omega) \cap L(A, \mathbb{R})$ . The proof of the theorem, which we leave to the reader, follows from the fact that in models of the form  $L(A, \mathbb{R})$ , every set is ordinal definable from  $A$  and a member of  $\omega^\omega$ , which implies that every set has an elementary submodel which is the surjective image of  $\omega^\omega$ . This means that there is a minimal (relative to a parameter from  $\omega^\omega$ ) witness to each true  $\Sigma_1^2(A)$  statement, and that that minimal witness is in  $\text{loc-}\Delta_1^2(A)^{L(A, \mathbb{R})}$ . In particular, membership in  $\text{loc-}\Sigma_1^2(A)$  sets is witnessed by  $\text{loc-}\Delta_1^2(A)$  sets. This gives another proof that pointclasses of the form  $\text{loc-}\Sigma_1^2(A)$  are  $\forall^{\omega^\omega}$ -closed.

**Theorem 5.0.6** (Solovay Basis Theorem). *Let  $A$  be a subset of  $\omega^\omega$  and let  $x$  and  $y$  be elements of  $\omega^\omega$ . Let  $\varphi$  be a formula whose quantifiers range over the hereditarily countable sets. If there exists a set  $B$  in  $\mathcal{P}(\omega^\omega) \cap L(A, \mathbb{R})$  such that  $\varphi(A, B, x, y)$  holds, then there exists such a set  $B$  in  $\text{loc-}\Delta_1^2(A)$ .*

**5.0.7 Remark.** To see that the pointclasses  $\text{loc-}\Sigma_1^2(A)$  satisfy the prewellordering property (without assuming  $\text{DC}_{\mathbb{R}}$ ), fix a  $\Sigma_1^2(A)$  set  $X$  of the form

$$\{x \in \omega^\omega : L(A, \omega^\omega) \models \exists B \subseteq \omega^\omega \psi(A, B, x, y)\}.$$

Orders  $\leq^*$  and  $<^*$  witnessing that  $X$  has the prewellordering property can be defined using the least ordinal  $\alpha$  such that  $L_\alpha(A, \omega^\omega)$  contains a set  $B$  as in the definition of  $X$ . Elementary submodel arguments similar to the proof of the Solovay Basis Theorem show that this gives a  $\text{loc-}\Sigma_1^2(A)$ -norm on  $X$ .

The following gives Theorem 5.0.4.

**Theorem 5.0.8.** *If AD holds then  $\text{loc-}\delta_1^2(A)$  is a strong partition cardinal, for each  $A \subseteq \omega^\omega$ .*

Our proofs of Theorems 5.0.5 and 5.0.8 diverge at just one point, establishing that each of the relevant pointclasses satisfies the prewellordering property (see the discussion just before the statement of Theorem 5.0.11). We complete the proof of Theorem 5.0.8 (so also Theorem 5.0.4) by presenting an argument of Martin connecting closure properties of pointclasses with partition properties of their associated ordinals. The following definition (a slightly modified version of Definition 2.33 on page 1783 of [6]) is due to Martin.

**5.0.9 Definition.** Let  $\lambda \leq \kappa$  be ordinals. We say that  $\kappa$  is  $\lambda$ -reasonable if there exist a nonselfdual  $\forall^{\omega^\omega}$ -closed boldface pointclass  $\Gamma$  and a function

$$\varphi: \omega^\omega \rightarrow \mathcal{P}(\lambda \times \kappa)$$

such that, letting  $\Delta = \Gamma \cap \check{\Gamma}$ ,

1. for each function  $F: \lambda \rightarrow \kappa$  there is an  $x \in \omega^\omega$  such that  $\varphi(x) = F$ ;
2. for each pair  $(\beta, \gamma) \in \lambda \times \kappa$ , the set

$$R_{\beta, \gamma} = \{x \in \omega^\omega : \{\eta < \kappa : (\beta, \eta) \in \varphi(x)\} = \{\gamma\}\}$$

is in  $\Delta$ ;

3. for each  $\beta < \lambda$  and each  $A$  in  $\exists^{\omega^\omega} \Delta \cap \mathcal{P}(R_\beta)$  there exists a  $\gamma_0 < \kappa$  such that

$$A \subseteq \bigcup_{\gamma < \gamma_0} R_{\beta, \gamma},$$

where  $R_\beta = \bigcup \{R_{\beta, \gamma} : \gamma < \kappa\}$ .

Given a function  $\varphi$ , sets  $R_\beta$  ( $\beta < \lambda$ ) as in Definition 5.0.9 and an  $x$  in  $\bigcap_{\beta < \lambda} R_\beta$ ,  $\varphi(x)$  is a function from  $\lambda$  to  $\kappa$ .

We first show how reasonableness gives partition properties, and then how the established properties of the pointclasses  $\text{loc-}\Sigma_1^2(A)$  give reasonableness. Given a set  $C$  of ordinals, we let  $C(\omega)$  denote the set of  $\gamma \in C$  for which the ordertype of  $C \cap \gamma$  is  $\eta + \omega$ , for some ordinal  $\eta$ .

**Theorem 5.0.10** (ZF + AD; Martin). *Let  $\Gamma$  be a nonselfdual boldface pointclass such that  $\delta(\Gamma)$  is a regular cardinal and  $\Gamma \cap \check{\Gamma}$  is a projective algebra closed under intersections and unions of length less than  $\delta(\Gamma)$ . Let  $\lambda$  be an ordinal such that  $\omega \cdot \lambda \leq \delta(\Gamma)$  and  $\Gamma$  witnesses that  $\delta(\Gamma)$  is  $\omega \cdot \lambda$ -reasonable. Then for each function  $P: [\kappa]^\lambda \rightarrow 2$  there is a club  $C_0 \subseteq \kappa$  such that, for any cofinal  $C \subseteq C_0$ ,  $P$  is constant on  $[C(\omega)]^\lambda$ .*

*Proof.* Let  $\kappa$  be  $\delta(\Gamma)$  and let  $\Delta$  be  $\Gamma \cap \check{\Gamma}$ . Let  $\varphi, R_{\beta, \gamma}$  ( $\beta < \omega \cdot \lambda, \gamma < \kappa$ ) and  $R_\beta$  ( $\beta < \omega \cdot \lambda$ ) be as in Definition 5.0.9, with respect to  $\omega \cdot \lambda, \kappa$  and  $\Gamma$ . Fix  $P: [\kappa]^\lambda \rightarrow 2$ .

Consider the following game in which player  $I$  plays the values of  $x \in \omega^\omega$  and player  $II$  plays the values of  $y \in \omega^\omega$ .

I	$x(0)$	$x(1)$	$x(2)$	$\dots$
II	$y(0)$	$y(1)$	$\dots$	

The winner of the game is determined as follows. If there is a least ordinal  $\beta < \omega \cdot \lambda$  such that  $\{x, y\} \not\subseteq R_\beta$ , then  $II$  if and only if  $x \notin R_\beta$ . Otherwise, letting  $f_{x, y}: \lambda \rightarrow \kappa$  be defined by setting  $f_{x, y}(\beta)$  to be

$$\sup\{\max(\varphi(x)(\eta), \varphi(y)(\eta)) : \eta < \omega \cdot (\beta + 1)\},$$

$II$  wins if and only if  $P(f_{x, y}) = 1$ .

Suppose first that  $\tau$  is a winning strategy for player  $II$ . Fix for this paragraph a pair  $(\beta, \gamma)$  in  $(\omega \cdot \lambda) \times \kappa$ . Let

$$S_{\beta, \gamma} = \bigcap_{\delta \leq \beta} \bigcup_{\eta \leq \gamma} R_{\delta, \eta}.$$

Then  $S_{\beta, \gamma}$  is in  $\Delta$ , since  $\Delta$  is closed under intersections and unions of length less than  $\kappa$ . Therefore, recalling that  $x \circ \tau$  denotes the sequence of moves made by player  $II$  when player  $I$  plays  $x$  and player  $II$  plays according to  $\tau$ , the set

$$S_{\beta, \gamma} \circ \tau = \{x \circ \tau : x \in S_{\beta, \gamma}\}$$

is in  $\Delta$ , as  $\Delta$  is a projective algebra. We have that  $S_{\beta, \gamma} \subseteq \bigcap_{\delta \leq \beta} R_\delta$ , which, since  $\tau$  is a winning strategy for  $II$ , implies that  $S_{\beta, \gamma} \circ \tau \subseteq R_\beta$ . Letting  $\theta(\beta, \gamma)$  be

$$\sup\{\varphi(y)(\beta) : y \in S_{\beta, \gamma} \circ \tau\},$$



we have by part (3) of Definition 5.0.9 that  $\theta(\beta, \gamma) < \kappa$ .

Let  $C_0$  be the set of  $\eta < \kappa$  such that  $\theta(\beta, \gamma) < \eta$  for all  $\beta \in \eta \cap (\omega \times \lambda)$  and  $\gamma < \eta$ . Since  $\kappa$  is regular,  $C_0$  is club in  $\kappa$ . Let  $C$  be any cofinal subset of  $C_0$ . Suppose now that  $F: \lambda \rightarrow C(\omega)$  is increasing. We want to see that  $P(F) = 1$ . Applying part (1) of Definition 5.0.9, let  $x$  be such that  $\varphi(x)$  is an increasing function, with range contained in  $C$ , such that

$$F(\beta) = \sup\{\varphi(x)(\delta) : \delta < \omega \cdot (\beta + 1)\}$$

for all  $\beta < \omega \cdot \lambda$ . Let  $y = x \circ \tau$ . Then  $y$  is in  $\bigcap_{\beta < \omega \cdot \lambda} R_\beta$ . It suffices to see that  $\varphi(y)(\beta) \leq \varphi(x)(\beta + 1)$  for all  $\beta < \omega \cdot \lambda$ , since then  $F = f_{x,y}$ , so  $P(F) = 1$ .

Fix  $\beta < \omega \cdot \lambda$ . We have that  $x$  is in  $S_{\beta, \varphi(x)(\beta)}$ , so  $y$  is in  $S_{\beta, \varphi(x)(\beta)} \circ \tau$ . Then

$$\varphi(y)(\beta) \leq \theta(\beta, \varphi(x)(\beta)) < \varphi(x)(\beta + 1),$$

since  $\varphi(x)$  is increasing and  $\varphi(x)(\beta + 1)$ , being in  $C$ , is closed under  $\theta$ .

A winning strategy for player  $I$  can be used as a winning strategy for player  $II$  in the game where the values of  $P$  are reversed. It follows that if there is a winning strategy for player  $I$ , then there is homogeneous set for value 0.  $\square$

Remarks 4.2.10, 4.2.11 and 5.0.7 show that, assuming AD, the pointclasses  $\text{loc-}\Sigma_1^2(A)$  are nonselfdual boldface pointclass with the prewellordering property, closed under  $\forall^{\omega^\omega}$  and unions, and that the corresponding selfdual pointclasses  $\Delta$  are projective algebras. In addition, Remark 4.2.12 shows that (aside from closure under  $\forall^{\omega^\omega}$ ) the same holds the pointclasses  $\Sigma_1^2(A)$ , if in addition the Wadge hierarchy is assumed to be wellfounded. Theorems 4.3.4 and 4.3.5 together show that the ordinals  $\text{loc-}\delta_1^2(A)$  are regular cardinals (and the same holds for the ordinals  $\delta_1^2(A)$ , if  $\Sigma_1^2$  is  $\forall^{\omega^\omega}$ -closed and  $\Sigma_1^2(A)$  has the prewellordering property). Theorem 5.0.11 then gives the versions of Theorems 5.0.5 and 5.0.8 for the weaker notion of strong partition cardinal discussed in Remark 5.0.1.

**Theorem 5.0.11** (ZF + AD). *Let  $\Gamma$  be a nonselfdual boldface pointclass with the prewellordering property, closed under  $\forall^{\omega^\omega}$  and unions, such that  $\Gamma \cap \check{\Gamma}$  is a projective algebra. Let  $\kappa$  be  $\delta(\Gamma)$ , and suppose that  $\kappa$  is a regular cardinal. Then  $\kappa \rightarrow (\kappa)^\kappa$ .*

*Proof.* Let  $\Delta$  be  $\Gamma \cap \check{\Gamma}$ . By Theorem 4.3.2,  $\Delta$  is closed under intersections and unions of length less than  $\delta(\Gamma)$ . By Theorem 5.0.10, it suffices to see that  $\kappa$  is  $\omega \cdot \kappa$ -reasonable. We will find a function  $\varphi: \omega^\omega \rightarrow \mathcal{P}(\kappa \times \kappa)$  such that  $\Gamma$  and  $\varphi$  witness this. By Theorems 2.4.9 and 4.2.16 there exist a set  $X \subseteq \omega^\omega$  in  $\Gamma \setminus \check{\Gamma}$  and a surjective  $\Gamma$ -norm  $f: X \rightarrow \kappa$ . By Remark 4.2.5, for each  $x \in X$ , the set

$$\{y \in X : f(y) \leq f(x)\}$$

is in  $\Delta$ . Let, for each  $x \in X$ ,

$$[x]_f = \{y \in X : f(y) = f(x)\},$$

$$[<x]_f = \{y \in X : f(y) < f(x)\}$$

and

$$C_x = \omega^\omega \setminus ([x]_f \cup [ < x ]_f).$$

Since  $\Delta$  is a projective algebra closed under unions and intersections of length less than  $\kappa$ , each of these sets is in  $\Delta$ .

Applying Theorem 2.4.9, fix a sequence  $\bar{U} = \langle U_n : n < \omega \setminus \{0\} \rangle$  of  $\Sigma_1^1$ -universal sets with the s-m-n property. Define  $\varphi: \omega^\omega \rightarrow \mathcal{P}(\kappa \times \kappa)$  by setting, for each  $z \in \omega^\omega$  and  $(\beta, \gamma) \in \kappa \times \kappa$ ,  $(\beta, \gamma) \in \varphi(z)$  if and only if there exist  $x, y$  in  $X$  with

- $f(x) = \beta$ ,
- $f(y) = \gamma$  and
- $(x, y) \in U_{2,z}([x]_f, C_x)$ .
- for all  $(v, w) \in U_{2,z}([x]_f, C_x)$ ,  $w \in X$  and  $f(w) = \gamma$ .

The Uniform Coding Lemma (Theorem 3.0.3) implies that part (1) of Definition 5.0.9 is satisfied by  $\Gamma$  and  $\varphi$ . To see this, given  $F: \kappa \rightarrow \kappa$ , let  $Z_F$  be

$$\{(x, y) \in X \times X : F(f(x)) = f(y)\}.$$

By the Uniform Coding Lemma, there is a  $z \in \omega^\omega$  such that, for all  $x \in X$ ,

1.  $U_{2,z}([x]_f, C_x) \subseteq Z_F \cap ([x]_f \times \omega^\omega)$ ,
2.  $U_{2,z}([x]_f, C_x) \neq \emptyset$  if and only if  $Z_F \cap ([x]_f \times \omega^\omega) \neq \emptyset$ .

It follows from this that  $\varphi(z) = F$ .

To see that part (2) of Definition 5.0.9 is satisfied, fix  $(\beta, \gamma) \in \kappa \times \kappa$ . By the definition of  $R_{\beta, \gamma}$ , for each  $z \in \omega^\omega$ ,  $z \in R_{\beta, \gamma}$  implies  $(\beta, \gamma) \in \varphi(z)$ . The reverse implication follows from last condition on the definition of  $\varphi$ . It suffices then to see that  $\{z \in \omega^\omega : (\beta, \gamma) \in \varphi(z)\}$  is in  $\Delta$ . This can be seen by fixing  $x_*, y_* \in X$  such that  $f(x_*) = \beta$  and  $f(y_*) = \gamma$ , and applying the fact that  $\Delta$  is a projective algebra.

For part (3), fix  $\beta < \kappa$  and  $A \subseteq R_\beta$  with  $A$  in  $\exists^\omega \Delta$  (which is the same as  $\Delta$ ). Fix  $x_* \in X$  with  $f(x_*) = \beta$ . Let  $D$  be the set of  $y \in \omega^\omega$  for which there exist  $z \in A$  and  $x \in [x_*]_f$  with  $(x, y) \in U_{2,z}([x_*]_f, C_{x_*})$ . Then  $D \subseteq X$  and  $D \in \Delta$ , so, by Proposition 4.2.6, there is a  $\gamma_0 < \kappa$  such that  $f[D] \subseteq \gamma_0$ . Fixing  $z \in A$  then, we have that  $z \in R_{\beta, \gamma}$  for some  $\gamma < \kappa$ . Then  $(\beta, \gamma) \in \varphi(z)$ . If  $x$  and  $y$  witness this, then  $x \in [x_*]_f$  and  $y \in D$ , which implies that  $\gamma < \gamma_0$ .  $\square$

Given ordinals  $\mu, \kappa$ , we define  $(\mu, \kappa)$ -club directedness to be the statement that if  $\langle \mathcal{C}_\alpha : \alpha < \mu \rangle$  is a sequence of nonempty sets of club subsets of  $\kappa$ , then there is a club subset of  $\kappa$  contained in at least one member of each  $\mathcal{C}_\alpha$ . Given  $\mu < \kappa$  and a function  $P: [\kappa]^\kappa \rightarrow \mu$ , for each  $\alpha < \mu$ , let  $P_\alpha: [\kappa]^\kappa \rightarrow 2$  be defined by letting  $P_\alpha(f)$  be 0 if  $P(f) = \alpha$  and 1 otherwise. Applying Theorem 5.0.10, we get for each  $\alpha < \mu$  a nonempty set  $\mathcal{C}_\alpha$  of club subsets of  $\kappa$  with the properties

of  $C_0$  as stated there, with respect to  $P_\alpha$ . If  $C$  is a club subset of  $\kappa$  contained in at least one member of each  $\mathcal{C}_\alpha$ , then  $P$  is constant on  $[C(\omega)]^\kappa$ .

To finish the proofs of Theorems 5.0.5 and 5.0.8 then it suffices to show that established properties of the pointclasses  $\text{loc-}\Sigma_1^2(A)$  and  $\Sigma_1^2(A)$  imply  $(\mu, \kappa)$ -club directedness, where  $\kappa$  is  $\delta(\Gamma)$  for the given pointclass  $\Gamma$ , and  $\mu < \kappa$ . We do this using Definition 5.0.9 and the Moschovakis Coding Lemma.

**Theorem 5.0.12** (ZF + AD). *Let  $\Gamma$  be a nonselfdual boldface pointclass closed under  $\forall^{\omega^\omega}$  and unions, and assume that  $\Gamma$  has the prewellordering property. Let  $\kappa$  a regular uncountable cardinal such that the pointclass  $\Gamma \cap \check{\Gamma}$  is closed under wellordered unions and intersections of length less than  $\kappa$ . If  $\Gamma$  witnesses that  $\kappa$  is  $\omega \cdot \kappa$ -reasonable, then, for all  $\mu < \kappa$ ,  $(\mu, \kappa)$ -club directedness holds.*

*Proof.* Let  $\varphi, \Delta, R_{\beta, \gamma}$  ( $\beta < \lambda, \gamma < \kappa$ ) and  $R_\beta$  ( $\beta < \lambda$ ) be as in Definition 5.0.9, with respect to  $\omega \cdot \kappa, \kappa$  and  $\Gamma$ . Fix  $\mu < \kappa$ , and let  $\langle \mathcal{C}_\alpha : \alpha < \mu \rangle$  be a sequence of nonempty sets of club subsets of  $\kappa$ . Let  $X$  be  $\bigcup_{\alpha < \mu} R_{0, \alpha}$ , and let  $f : X \rightarrow \mu$  be such that  $x \in R_{0, f(x)}$  for all  $x \in X$ . Then  $f$  is onto. For each  $x \in X$ , let  $[x]_f$  be  $\{z \in X : f(x) = f(z)\}$ . Let  $<_f$  be the set  $\{(x, y) \in X \times X : f(x) < f(y)\}$ . Since  $\Delta$  is closed under wellordered unions and intersections of length less than  $\kappa$ ,  $<_f$  is in  $\Delta$ . Let  $Z$  be the set of  $(x, y) \in X \times \omega^\omega$  such that  $\varphi(y)$  is an increasing function whose range is in  $\mathcal{C}_{f(x)}$ . Let  $A$  be as in the conclusion of the Moschovakis Coding Lemma (Theorem 3.0.1), with respect to  $X, Z$ , and  $f$ . So  $A$  is a  $\text{pos-}\Sigma_1^1(<_f)$  subset of  $Z$  such that for all  $x \in X$ , if  $Z \cap ([x]_f \times \omega^\omega)$  is nonempty then so is  $A \cap ([x]_f \times \omega^\omega)$ . Then  $A$  is in  $\exists^{\omega^\omega} \Delta$ , and so is  $A_1 = \{y : \exists x \in X (x, y) \in A\}$ . Furthermore,  $A_1 \subseteq \bigcap_{\beta < \kappa} R_\beta$ . It follows from condition (3) of Definition 5.0.9 that there is a function  $g : \kappa \rightarrow \kappa$  such that

$$A_1 \subseteq \bigcap_{\beta < \kappa} \bigcup_{\gamma < g(\beta)} R_{\beta, \gamma}.$$

Let  $D$  be the set of closure points of  $g$ ; then  $D$  is a club subset of  $\kappa$ .

We want to see that  $D$  is contained in at least one member of each  $\mathcal{C}_\alpha$ . To see that this is the case, fix  $\alpha < \mu$ ,  $x \in X$  such that  $f(x) = \alpha$  and a  $y \in \omega^\omega$  such that  $\varphi(y)$  is an increasing function whose range is contained in some element of  $\mathcal{C}_\alpha$ . Then  $(x, y)$  is in  $Z$ , so there exists an  $(x', y') \in A$  such that  $f(x) = f(x')$ . Then  $y'$  is in  $A_1$ , so the range of  $y'$  is an element of  $\mathcal{C}_\alpha$  containing  $D$ .  $\square$

**5.0.13 Remark.** The version of Martin's theorem presented in this section may not be optimal. In Remark 2.32 of [6] it is stated that if  $\Gamma$  and  $\varphi$  witness that  $\kappa$  is  $\lambda$ -reasonable, then  $\Gamma \cap \check{\Gamma}$  is closed under unions of length less than  $\kappa$ ,  $\Gamma$  is closed under intersections and countable unions, and  $\text{PWO}(\check{\Gamma})$  holds.

It is an open question whether AD implies that there are no strong partition cardinals greater than (or equal to)  $\Theta$ . Remark 8.1.15 discusses a connection between this question and the question of whether  $\text{AD} + \text{DC}_\mathbb{R}$  implies that all sets of reals are  $\infty$ -Borel.



## Chapter 6

# Suslin sets and Uniformization

### 6.1 Suslin sets

Recall that a *tree*  $T$  on a set  $X$  is a set of finite sequences from  $X$  which is closed under initial segments. We write  $[T]$  for the set of infinite sequences whose finite initial segments are all in  $T$ . A tree is *wellfounded* if  $[T] = \emptyset$ ; otherwise, it is *illfounded*.

If  $T$  is a tree on a product of two sets  $X \times Y$ , and  $s$  is a finite sequence of elements of  $X$ , then  $T_s$  denotes the set of  $t \in Y^{<\omega}$  for which  $(s, t) \in T$ . For each  $f \in X^\omega$ , we write  $T_f$  for  $\bigcup_{n \in \omega} T_{f \upharpoonright n}$ . The *projection* of  $T$ ,  $p[T]$ , is the set of  $f \in X^\omega$  for which  $T_f$  is illfounded, i.e., for which there exists a  $g \in Y^\omega$  such that  $(f, g) \in [T]$  (identifying, for notational simplicity, pairs of sequences with sequences of pairs).

If  $T$  is an illfounded tree on a set  $X$ , and  $\leq$  is a wellordering of  $X$ , the *leftmost branch* of  $T$  relative to  $\leq$  ( $\text{lb}_\leq(T)$ ) is the unique  $f \in X^\omega$  such that, for each  $n \in \omega$ ,  $f(n)$  is the  $\leq$ -least  $x \in X$  for which the tree

$$\{\sigma \in X^{<\omega} : (f \upharpoonright n) \frown \langle x \rangle \frown \sigma \in T\}$$

is illfounded. When  $X$  is an ordinal or a finite product of ordinals we use the ordinal ordering or the corresponding lexicographical ordering for  $\leq$  and write  $\text{lb}(T)$ .

**6.1.1 Definition.** Given an ordinal  $\gamma$  and a set  $X$ , a set  $A \subseteq X^\omega$  is  $\gamma$ -*Suslin* if there exists a tree  $T \subseteq (X \times \gamma)^{<\omega}$  such that  $A = p[T]$ . A set  $A \subseteq X^\omega$  is *Suslin* if it is  $\gamma$ -Suslin for some ordinal  $\gamma$ .

**6.1.2 Remark.** Every Suslin subset of  $\omega^\omega$  is  $\gamma$ -Suslin for some  $\gamma < \Theta$ . To see this, suppose that  $\alpha$  is an ordinal and  $T$  is tree on  $\omega \times \alpha$ , and observe that the set  $\{\text{lb}(T_x)(n) : x \in p[T], n \in \omega\}$  is a surjective image of  $\omega^\omega$ , and therefore has ordertype less than  $\Theta$ . Alternately one can use the fact that (assuming only ZF) if  $\beta$  is an ordinal with  $T \in L_\beta(T, \mathbb{R})$ , then  $L_\beta(T, \mathbb{R})$  has an elementary submodel which contains  $\omega^\omega$  and is a surjective image of  $\omega^\omega$ .

Given an ordinal  $\gamma$ , we write  $\mathcal{S}_\gamma$  for the smallest boldface pointclass containing each  $\gamma$ -Suslin subset of  $\omega^\omega$  and  $\mathcal{S}_{<\gamma}$  for  $\bigcup_{\alpha<\gamma} \mathcal{S}_\alpha$ . Then  $\mathcal{S}_\gamma$  and  $\mathcal{S}_{<\gamma}$  are boldface pointclasses closed under  $\exists^{\omega^\omega}$ , intersections and unions.

**6.1.3 Remark.** If  $\gamma$  is less than  $\Theta$ , and AD holds, then, by the Coding Lemma,  $\text{CC}_{\mathcal{P}(\gamma)}$  holds, so  $\mathcal{S}_\gamma$  is closed under countable intersections and countable unions.

The following definition is due to Kechris.

**6.1.4 Definition.** A cardinal  $\kappa$  is *Suslin* if it is infinite and there exists  $A \subseteq \omega^\omega$  which is  $\kappa$ -Suslin but not  $\gamma$ -Suslin for any  $\gamma < \kappa$ .

In symbols, an infinite cardinal  $\kappa$  is Suslin if and only if  $\mathcal{S}_\kappa \setminus \mathcal{S}_{<\kappa} \neq \emptyset$ . By Remark 6.1.2, every Suslin cardinal is less than  $\Theta$ . By Remark 6.1.3, AD implies that a limit of Suslin cardinals of countable cofinality is a Suslin cardinal if it is below  $\Theta$  (more is true, as we shall see in Section 11.4).

**6.1.5 Remark.** If Lipschitz Determinacy holds, and  $\kappa < \lambda$  are Suslin cardinals, then, by Theorem 2.1.4,  $\mathcal{S}_\kappa \subseteq \mathcal{S}_\lambda$ . It follows (under the same assumption) that if  $\kappa$  is a limit of Suslin cardinals, then  $\mathcal{S}_{<\kappa}$  is a projective algebra. If AD holds,  $\lambda$  is a Suslin cardinal and  $\Delta$  is a projective algebra contained in  $\mathcal{S}_\lambda$ , then  $\mathcal{S}_\lambda$  contains  $\bigcup_\omega \Delta$  (see Remark 6.1.3). If in addition  $\text{DC}_\mathbb{R}$  holds and  $\Delta$  is not closed under countable unions then, by Theorem 4.4.4, every wellordered union of sets in  $\Delta$  is in  $\mathcal{S}_\lambda$ .

**6.1.6 Remark.** Moschovakis's Second Periodicity Theorem (see Section 6C of [32]) implies that under  $\text{AD} + \text{DC}_\mathbb{R}$ , the pointclass of Suslin sets is closed under  $\forall^{\omega^\omega}$  and  $\exists^{\omega^\omega}$ . If  $\kappa$  is the largest Suslin cardinal (which can happen), it follows that  $\mathcal{S}_\kappa$  is closed under  $\forall^{\omega^\omega}$  and  $\exists^{\omega^\omega}$ .

**6.1.7 Remark.** By the Moschovakis Coding Lemma, if AD holds then for each  $\kappa < \Theta$  there is a subset of  $\omega^\omega$  which is not  $\kappa$ -Suslin. It follows that if every subset of  $\omega^\omega$  is Suslin, then the Suslin cardinals are cofinal in  $\Theta$ . The converse is shown in Corollary 6.1.19 below.

The proof of the following theorem uses the Coding Lemma to code trees projecting to sets of reals. The theorem will be used in the proof of Theorem 6.1.10 below.

**Theorem 6.1.8 (ZF + AD).** *Let  $\Gamma$  be a boldface pointclass closed under  $\exists^{\omega^\omega}$ , let  $\kappa$  be an infinite cardinal and let  $\preceq$  be a prewellordering of a subset of  $\omega^\omega$ . If  $\preceq$  has length  $\kappa$  and both  $\preceq$  and its corresponding strict prewellordering are in  $\Gamma$ , then  $\mathcal{S}_\kappa \subseteq \Gamma$ .*

*Proof.* Fix  $A \in \mathcal{S}_\kappa \cap \mathcal{P}(\omega^\omega)$ . Let  $T \subseteq (\omega \times \kappa)^{<\omega}$  be a tree of cardinality  $\kappa$  projecting to  $A$ . Let  $\preceq$  be as in the statement of the theorem, let  $\prec$  be the strict part of  $\preceq$  and let  $X$  be the domain of  $\preceq$ . Let  $r: X \rightarrow \kappa$  be the function which sends each member of  $X$  to its  $\preceq$ -rank, and let  $b: \kappa \rightarrow T$  be a bijection. Let  $\pi: \omega^\omega \rightarrow (\omega^{<\omega} \times (\omega^\omega)^{<\omega})$  be a recursive bijection. Let  $Z$  be the set of

$(x, y) \in X \times \omega^\omega$  such that  $\pi(y)$  is a tuple  $(s, z_0, \dots, z_{|s|})$  such that  $s$  is the first coordinate of  $b(r(x))$  and for all  $k \leq |s|$ ,  $b(r(z_k))$  is the initial segment of  $b(r(x))$  of length  $k$ .

By the Coding Lemma, there is a set  $R \subseteq Z$  in  $\Gamma$  such that for all  $x$ , if there exists a  $y$  with  $(x, y) \in Z$ , then there exist  $(x', y') \in R$  with  $r(x) = r(x')$ . Then  $A$  is the set of  $w \in \omega^\omega$  for which there exist  $\langle q_i : i \in \omega \rangle \in (\omega^\omega)^\omega$  such that for all  $n \in \omega$  there exist  $(x, y) \in R$  such that, letting  $\pi(y) = (s, z_0, \dots, z_{|s|})$ ,  $s = w \upharpoonright n$  and, for each  $m \leq n$ ,  $r(z_m) = r(q_m)$ . This shows that  $A$  is in  $\Gamma$ .  $\square$

Theorem 6.1.10 below shows that if  $\text{AD} + \text{DC}_\mathbb{R}$  holds and  $\kappa$  is a Suslin cardinal, then  $\mathcal{S}_\kappa$  is nonselfdual. The proof uses the following definition.

**6.1.9 Definition.** Let  $\gamma$  be an ordinal, let  $X$  be a set and let  $T$  be a tree on  $X \times \gamma$ . The *minimization* of  $T$  is the tree

$$\{(x \upharpoonright n, \text{lb}(T_x) \upharpoonright n) : x \in p[T], n \in \omega\}.$$

We say that  $T$  is *minimal* if it is equal to its minimization.

Note that the minimization of a tree  $T$  is minimal and has the same projection as  $T$ .

**Theorem 6.1.10** ( $\text{ZF} + \text{AD} + \text{DC}_\mathbb{R}$ ; Kechris [11]). *If  $\kappa$  is a Suslin cardinal, then  $\mathcal{S}_\kappa$  is nonselfdual.*

*Proof.* Assume toward a contradiction that  $\mathcal{S}_\kappa$  is closed under complements. Since  $\mathcal{S}_\kappa$  is (always)  $\exists^{\omega^\omega}$ -closed, it follows from this assumption that it is also  $\forall^{\omega^\omega}$ -closed. Since  $\mathcal{S}_\kappa$  is closed under countable unions, it follows from Corollary 4.4.6 that every member of  $\mathcal{S}_\kappa$  is an element of a nonselfdual  $\exists^{\omega^\omega}$ -closed pointclass contained in  $\mathcal{S}_\kappa$  and closed under wellordered unions. This shows that for any  $A \in \mathcal{S}_\kappa$ , any wellordered union of subsets of  $\omega^\omega$  Wadge-reducible to  $A$  is in  $\mathcal{S}_\kappa$ . In combination with Theorem 6.1.8 it also shows that  $\mathcal{S}_\kappa$  does not contain a prewellordering of length  $\kappa$ .

Let  $A$  be a subset of  $\omega^\omega$  in  $\mathcal{S}_\kappa \setminus \mathcal{S}_{<\kappa}$ . Let  $\Gamma$  be a nonselfdual pointclass contained in  $\mathcal{S}_\kappa$  with  $A \in \Gamma$  such that  $\Gamma$  is closed under wellordered unions. Then every element of  $\mathcal{S}_{<\kappa}$  is in  $\Gamma$ . If the cofinality of  $\kappa$  were uncountable, it would follow that  $\mathcal{S}_\kappa = \Gamma$ , giving a contradiction. To see this, let  $T$  be a tree on  $\omega \times \kappa$ , and for each  $\alpha < \kappa$ , let  $T \upharpoonright \alpha$  denote  $T \cap (\omega \times \alpha)^{<\omega}$ . If  $\kappa$  has uncountable cofinality, then  $p[T] = \bigcup_{\alpha < \kappa} p[T \upharpoonright \alpha]$ , and the latter is a wellordered union of sets in  $\mathcal{S}_{<\kappa}$ .

To complete the proof, we show that  $\kappa$  has uncountable cofinality. Let  $\delta$  be the supremum of the lengths of the prewellorderings in  $\mathcal{S}_\kappa$ . Since  $\mathcal{S}_\kappa$  is a projective algebra,  $o(\mathcal{S}_\kappa) \leq \delta$ , by Proposition 2.5.8. Since  $\mathcal{S}_\kappa$  is closed under countable unions,  $\delta$  has uncountable cofinality. We show that  $\kappa = \delta$ . As noted in the first paragraph of this proof, there is no prewellordering in  $\mathcal{S}_\kappa$  of length  $\kappa$ , so  $\delta \leq \kappa$ . To show that  $\kappa \leq \delta$ , let  $T$  be a minimal tree on  $\omega \times \kappa$  projecting to  $A$ . Then  $T$  has cardinality  $\kappa$ . For each  $\sigma \in T$ , let  $A_\sigma$  be the set of  $x \in A$  for which  $\sigma$  is an initial segment of  $(x, \text{lb}(T_x))$ . Under our assumption that  $\mathcal{S}_\kappa$  is closed under complements, each  $A_\sigma$  is in  $\mathcal{S}_\kappa$ .

By the minimality of  $T$ , each  $A_\sigma$  is nonempty, and for distinct  $\sigma$  and  $\tau$  in  $T$  of the same length,  $A_\sigma$  and  $A_\tau$  are disjoint. For each  $\rho < o(\mathcal{S}_\kappa)$ , let  $T \restriction \rho$  be the set of  $\sigma \in T$  for which the Wadge rank of  $A_\sigma$  is less than  $\rho$ . For each  $n \in \omega$ , let  $<_{n,\rho}$  be the lexicographical order on the members of  $T \restriction \rho$  of length  $n$ . We claim that each  $<_{n,\rho}$  has ordertype less than  $\delta$ . The claim completes the proof of the theorem, since it implies that  $T$  is a union of  $|o(\mathcal{S}_\kappa)|$  many sets of cardinality at most  $|\delta|$ . To prove the claim, fix  $n \in \omega$  and  $\rho < o(\mathcal{S}_\kappa)$ . The union of all the sets  $A_\tau \times A_\sigma$  for  $\tau <_{n,\rho} \sigma$  is a wellordered union of sets of Wadge rank bounded below  $o(\mathcal{S}_\kappa)$ , and is therefore in  $\mathcal{S}_\kappa$ . This is a prewellordering with the same length as  $<_{n,\rho}$ , which must therefore be less than  $\delta$ .  $\square$

**6.1.11 Remark.** Remark 6.1.6 and Theorems 6.1.8 and 6.1.10 imply that if  $\text{AD} + \text{DC}_\mathbb{R}$  holds and  $\kappa$  is the largest Suslin cardinal, then all prewellorderings in  $\mathcal{S}_\kappa \cap \dot{\mathcal{S}}_\kappa$  have length less than  $\kappa$ .

Lemma 6.1.14 below shows that if  $\lambda$  is a Suslin cardinal, then there exists a strictly  $\subseteq$ -increasing  $\lambda$ -sequence of Suslin sets, induced by a single tree on  $\omega \times \lambda$ .

Let  $<_\ell^2$  be the following version of the lexicographical order on finite sequences of pairs of ordinals :  $(s_0, t_0) <_\ell^2 (s_1, t_1)$  if and only if there exists an  $i \in \omega$  such that the following hold:

- $(s_0 \restriction i, t_0 \restriction i) = (s_1 \restriction i, t_1 \restriction i)$ ;
- $s_0(i) < s_1(i) \vee (s_0(i) = s_1(i) \wedge t_0(i) < t_1(i))$ .

If  $A$  is a set of pairwise incompatible sequences of pairs (i.e., such that no sequence in the set extends any other one), and  $n \in \omega$  is such that each member of  $A$  has length at most  $n$ , then the restriction of  $<_\ell^2$  to  $A$  wellorders  $A$ .

**6.1.12 Definition.** Let  $\eta \leq \gamma$  be ordinals, and let  $T$  be a tree on  $\omega \times \gamma$ . We say that  $T$  is  $\eta$ -full if

$$\{\langle (0, \alpha) \rangle : \alpha < \eta\} \subseteq T.$$

**Lemma 6.1.13.** *Let  $\gamma$  and  $\eta$  be ordinals, let  $T$  be a minimal tree on  $\omega \times \gamma$  such that every member of  $p[T]$  takes value 0 at 0, and let  $A \subseteq T \setminus \{\emptyset\}$  be an antichain such that*

- *every element of  $[T]$  intersects  $A$ ;*
- *the restriction of  $<_\ell^2$  to  $A$  has ordertype  $\eta$ .*

*Then there is a minimal  $\eta$ -full tree  $T'$  on  $\omega \times \max\{\gamma, \eta\}$ , definable from  $A$  and  $T$ , such that  $p[T] = p[T']$ .*

*Proof.* Enumerate  $A$  in  $<_\ell^2$ -order as  $\langle (s_\alpha, t_\alpha) : \alpha < \eta \rangle$ . For each  $\alpha < \eta$ , let

- $T_\alpha$  be the set of  $(s, t) \in T$  which are compatible with  $(s_\alpha, t_\alpha)$ ;
- $f_\alpha : \gamma^{<\omega} \rightarrow \max\{\gamma, \eta\}^{<\omega}$  be defined by setting
  - $|f_\alpha(t)| = |t|$ ;



- for all  $i \in |t| \cap |t_\alpha|$ ,  $f_\alpha(t)(i) = \alpha$ ;
- for all  $i \in |t| \setminus |t_\alpha|$ ,  $f_\alpha(t)(i) = t(i)$ ;
- $T'_\alpha$  be  $\{(s, f_\alpha(t)) : (s, t) \in T_\alpha\}$ .

Let  $T' = \bigcup_{\alpha < \eta} T'_\alpha$ . Then  $T'$  is  $\eta$ -full, and since  $p[T_\alpha] = p[T'_\alpha]$  for all  $\alpha < \eta$ , and every element of  $[T]$  intersects  $A$ ,  $p[T'] = p[T]$ . To see that  $T'$  is minimal, fix  $(s', t') \in T'$  and  $\alpha < \eta$  such that  $(s', t') \in T'_\alpha$ . We want to find an  $x \in p[T']$  such that  $(s', t')$  is an initial segment of  $(x, \text{lb}(T'_x))$ . Replacing  $(s', t')$  with  $(s_\alpha, t_\alpha)$  in the case  $|s'| < |s_\alpha|$ , it suffices to consider the case where  $|s'| \geq |s_\alpha|$ . Let  $(s, t) \in T_\alpha$  be such that  $s' = s$  and  $t' = f_\alpha(t)$ . Since  $T$  is minimal, there exists an  $x \in p[T_\alpha]$  such that  $(s, t)$  is an initial segment of  $(x, \text{lb}(T_x))$ . Applying the definition of  $<_\ell^2$  we have that for all  $\beta < \alpha$ ,  $x \notin p[T_\beta]$ , so  $x \notin p[T'_\beta]$ . Then  $\text{lb}(T'_x) = \text{lb}(T'_{\alpha,x})$ , and  $t'$  is an initial segment of  $\text{lb}(T'_x)$  as desired.  $\square$

**Lemma 6.1.14.** *If  $\lambda$  is a Suslin cardinal then there is a minimal  $\lambda$ -full tree on  $\omega \times \lambda$ .*

*Proof.* Let  $T_0$  be a tree on  $\omega \times \lambda$  projecting to a subset of  $\omega^\omega$  which is not  $\gamma$ -Suslin for any  $\gamma < \lambda$ , and such that each member of  $p[T_0]$  takes value 0 at 0. Let  $T_1$  be the minimization of  $T_0$ . Since  $T_0$  witnesses that  $\lambda$  is Suslin,  $|T_1| = \lambda$ .

If some  $T_1$ -antichain has  $<_\ell^2$ -ordertype  $\lambda$ , let  $A$  be such an antichain and let  $T'_1$  be the set of nodes of  $T_1$  compatible with a member of  $A$ . Then  $T'_1$  is minimal, and the lemma follows by applying Lemma 6.1.13 with  $T'_1$  and  $A$ .

Suppose then that  $T_1$  does not contain such an antichain. For each  $n \in \omega$ , let  $\kappa_n$  be the cardinality of the set of nodes of  $T_1$  of length  $n + 1$ . Then each  $\kappa_n$  is less than  $\lambda$ , and  $\sup_{n \in \omega} \kappa_n = \lambda$ . We can then complete the proof by using (for each  $n \in \omega$ ) the  $(n + 1)$ st level of  $T_1$  to produce a minimal  $\kappa_n$ -full tree on  $\omega \times \kappa_n$ , modifying the trees to make their projections disjoint, and combining these trees to make a minimal  $\lambda$ -full tree on  $\omega \times \lambda$ . We give the details below.

For each  $n \in \omega$ , let  $A_n$  be the length- $\kappa_n$   $<_\ell^2$ -initial segment of the  $(n + 1)$ st level of  $T_1$ , and let  $T_{1,n}$  be the set of nodes of  $T_1$  compatible with a member of  $A_n$ . Then  $T_{1,n}$  is minimal; let  $T_{2,n}$  be the result of applying Lemma 6.1.13 to  $T_{1,n}$  and  $A_n$ .

We now modify the trees  $T_{2,n}$ , achieving minimality (via the function  $\pi_{0,n}^*$  defined below) by making the members of their left coordinates disjoint (aside from 0) and fullness (via  $\pi_{1,n}^*$ ). Let  $\pi: \omega \times \omega \rightarrow \omega$  be an injection, and for each  $n \in \omega$  let  $\pi_{0,n}^*$  be the length-preserving function on  $\omega^{<\omega}$  defined by setting  $\pi_{0,n}^*(s)(0)$  to be  $s(0)$  (in the case where  $s$  is nonempty) and  $\pi_{0,n}^*(s)(i) = \pi(n, i)$  for all nonzero  $i \in \text{dom}(s)$ . Again for each  $n \in \omega$ , let  $\gamma_n = \sum_{m < n} \kappa_m$  (using ordinal addition) and let  $\pi_{1,n}^*$  be the length-preserving function on  $\gamma^{<\omega}$  defined by setting  $\pi_{1,n}^*(t)(0)$  to be  $\gamma_n + t(0)$  (in the case where  $t$  is nonempty) and  $\pi_{1,n}^*(t)(i) = t(i)$  for all nonzero  $i \in \text{dom}(t)$ . Finally, let  $T_2$  be

$$\{(\pi_{0,n}^*(s), \pi_{1,n}^*(t)) : n \in \omega, (s, t) \in T_{2,n}\}.$$

Then  $T_2$  is as desired.  $\square$

**Corollary 6.1.15.** *Let  $\lambda$  is a Suslin cardinal and let  $\Delta$  be the smallest pointclass containing  $\mathcal{S}_\lambda \cup \dot{\mathcal{S}}_\lambda$ . Then there is a prewellordering of  $\omega^\omega$  of length  $\lambda$  in  $\bigcup_\lambda \Delta$ .*

*Proof.* Let  $T$  be a minimal  $\lambda$ -full tree on  $\omega \times \lambda$ . Define  $\leq$  on  $\omega^\omega$  by setting  $x \leq y$  to hold if either  $x \notin p[T]$  or  $x, y \in p[T]$  and  $\text{lb}(T_x)(0) \leq \text{lb}(T_y)(0)$ . Then  $\leq$  is a prewellordering of length  $\lambda$ .  $\square$

Every  $\Sigma_1^1$  prewellordering of  $\omega^\omega$  has countable length (see [32, 20], for instance). Applying this fact in a forcing extension by  $\text{Col}(\omega, \kappa)$ , we get the following.

**Theorem 6.1.16** (Kunen-Martin). *If  $\kappa$  is an infinite cardinal and  $\leq$  is a  $\kappa$ -Suslin prewellordering of  $\omega^\omega$ , then  $\leq$  has length less than  $\kappa^+$ .*

**6.1.17 Remark.** The Kunen-Martin theorem can be restated as saying that for any uncountable cardinal  $\kappa$ , all prewellorderings in  $\mathcal{S}_{<\kappa}$  have length less than  $\kappa$ .

**Theorem 6.1.18** (ZF+AD). *If  $\kappa$  is a limit of Suslin cardinals, then  $\delta(\mathcal{S}_{<\kappa}) = \kappa$ .*

*Proof.* That  $\delta(\mathcal{S}_{<\kappa}) \leq \kappa$  follows from both Theorem 6.1.16 and Remark 6.1.17. The reverse inequality follows from Corollaries 4.4.6 and 6.1.15, noting that  $\mathcal{S}_{<\kappa}$  is a projective algebra, by Remark 6.1.5.  $\square$

Finally we have the equivalence, under AD, of the Suslin cardinals being cofinal in  $\Theta$  with the statement every subset of  $\omega^\omega$  is Suslin.

**Corollary 6.1.19** (ZF + AD). *The Suslin cardinals are cofinal in  $\Theta$  if and only if every subset of  $\omega^\omega$  is Suslin.*

*Proof.* For the forward direction, Lipschitz Determinacy implies that if  $A \subseteq \omega^\omega$  is not Suslin, then every Suslin prewellordering of a subset of  $\omega^\omega$  is a continuous preimage of  $A$ . Theorem 6.1.18 implies that if AD holds and the Suslin cardinals are cofinal in  $\Theta$  then  $\Theta$  is the supremum of the lengths of the Suslin prewellorderings. The failure of the forward direction of the corollary would then imply the existence of a surjection from  $\omega^\omega$  onto  $\Theta$ , giving a contradiction. The reverse direction follows from the Moschovakis Coding Lemma, as discussed in Remark 6.1.7.  $\square$

Given a cardinal  $\kappa$ , we say that a subset of topological space  $X$  is *weakly  $\kappa$ -Borel* if it is in the smallest collection of subsets of  $X$  containing the closed sets and closed under complements and unions and intersections of cardinality at most  $\kappa$ . For instance, the weakly  $\aleph_0$ -Borel sets are just the Borel sets in the usual sense. A set is *weakly  $<\kappa$ -Borel* if it is in the smallest collection of subsets of  $X$  containing the closed sets and closed under complements and unions and intersections of cardinality less than  $\kappa$ . A set is *weakly  $\infty$ -Borel* if it is weakly  $\kappa$ -Borel for some cardinal  $\kappa$ . A standard elementary submodel argument shows that if a subset of  $\omega^\omega$  is weakly  $\infty$ -Borel then it is weakly  $\kappa$ -Borel for some  $\kappa < \Theta$ .

Some authors use “ $\infty$ -Borel” for what we are calling “weakly  $\infty$ -Borel” and use “effectively  $\infty$ -Borel” for the notion of  $\infty$ -Borel defined in Chapter 8. Example 8.1.6 shows that the two notions do not always coincide.

**6.1.20 Remark.** Suppose that  $\text{AD} + \text{DC}_{\mathbb{R}}$  holds, and that  $\Delta$  is a projective algebra which is not closed under countable unions. Then the pointclass  $\bigcup_{\omega} \Delta$  is  $\exists^{\omega^{\omega}}$ -closed. By Theorem 4.4.4,  $\bigcup_{\omega} \Delta$  is nonselfdual and has the prewellordering property. By Theorem 2.4.3 it has a universal set, which implies that the pointclass  $\forall^{\omega^{\omega}} \bigcup_{\omega} \Delta$  is nonselfdual, since it also has a universal set. By the First Periodicity Theorem (see Remark 4.2.8),  $\forall^{\omega^{\omega}} \bigcup_{\omega} \Delta$  has the prewellordering property. Let  $\Gamma = \forall^{\omega^{\omega}} \bigcup_{\omega} \Delta$  and let  $\Lambda = \Gamma \cap \tilde{\Gamma}$ . By Theorem 4.3.2, every weakly  $<\delta(\Gamma)$ -Borel subset of  $\omega^{\omega}$  is in  $\Lambda$ . For any  $A \subseteq \omega^{\omega}$ , the pointclass  $\Delta_{\omega}(A)$  of sets projective in  $A$  is a projective algebra containing  $A$  which is not closed under countable unions. Since the pointclass  $\Lambda$  defined above is a proper subset of  $\Gamma$ , this shows that (assuming  $\text{AD} + \text{DC}_{\mathbb{R}}$ ) there cannot be a  $\kappa < \Theta$  such that every subset of  $\omega^{\omega}$  is weakly  $\kappa$ -Borel.

The converse of the following theorem is also true (see Theorem 8.1.13).

**Theorem 6.1.21** ( $\text{ZF} + \text{AD} + \text{DC}_{\mathbb{R}}$ ). *Suppose that  $\kappa$  is a limit of Suslin cardinals and that  $\kappa$  has uncountable cofinality. Then every subset of  $\omega^{\omega}$  which is weakly  $<\kappa$ -Borel is also  $<\kappa$ -Suslin.*

*Proof.* We make use of Remark 6.1.3 and the comments immediately before. Let  $\gamma < \kappa$  be a limit of Suslin cardinals of countable cofinality. By Theorem 6.1.18,  $\delta(\mathcal{S}_{<\gamma}) = \gamma$ . By Remark 6.1.5,  $\mathcal{S}_{<\gamma}$  is a projective algebra which is not closed under countable unions. Letting  $\Gamma$  be the pointclass  $\forall^{\omega^{\omega}} \bigcup_{\omega} \mathcal{S}_{<\gamma}$  and  $\Lambda$  be  $\Gamma \cap \tilde{\Gamma}$ , we have by Remark 6.1.20 that every  $<\gamma$ -Borel set is in  $\Lambda$ . Since  $\mathcal{S}_{\gamma}$  is closed under countable unions, it contains  $\bigcup_{\omega} \mathcal{S}_{<\gamma}$ . If  $\lambda > \gamma$  is a Suslin cardinal greater than  $\gamma$ , then  $\mathcal{S}_{\lambda}$  is  $\exists^{\omega^{\omega}}$ -closed and contains the complement of every set in  $\mathcal{S}_{\gamma}$ , so  $\Lambda \subseteq \tilde{\Gamma} \subseteq \mathcal{S}_{\lambda}$ .  $\square$

The following theorem will be used to show that if  $\text{AD}$  holds then  $\text{AD}^+$  holds in  $L(\mathbb{R})$  (see Remark 7.0.4 and Corollary 8.2.5).

**Theorem 6.1.22** (Martin-Steel [30]). *Assuming  $\text{AD} + V=L(\mathbb{R})$ , the Suslin sets are exactly the  $\Sigma_1^2$  sets.*

## 6.2 Uniformization

Given sets  $X, Y$  and  $A \subseteq X \times Y$ , we say that a function  $f: X \rightarrow Y$  *uniformizes*  $A$  if  $\text{dom}(f) = \{x \in X \mid A_x \neq \emptyset\}$ , and  $(x, f(x)) \in A$  for all  $x \in \text{dom}(f)$ . We say then that  $A$  is *uniformized*. If  $A \subseteq \alpha^{\omega} \times \beta^{\omega}$  is the projection of a tree  $T$  on  $\alpha \times \beta \times \gamma$ , for some ordinals  $\alpha, \beta$  and  $\gamma$ , then a uniformizing function for  $A$  can be constructed from  $T$  by considering  $T$  as a tree on  $\alpha \times (\beta \times \gamma)$  and using the function  $x \mapsto \text{lb}(T_x)$  for  $x \in p[T]$ . This gives the following fact.

**Theorem 6.2.1.** *Every Suslin subset of  $(\omega^{\omega})^2$  is uniformized.*

We let **Uniformization** be the statement that every subset of  $(\omega^{\omega})^2$  has a uniformizing function.

**6.2.2 Remark.** Using a coding a finite sequences of reals by individual reals, it is easy to see that Uniformization implies  $\text{DC}_{\mathbb{R}}$ . Similarly,  $\text{AD}_{\mathbb{R}}$  implies Uniformization, via the one-round game where player  $I$  plays an  $x \in \omega^\omega$ , and player  $II$  wins by playing a  $y \in \omega^\omega$  such that the pair  $(x, y)$  is in the given payoff set.

**6.2.3 Definition.** Given a set  $X$ , a set  $A \subseteq X^\omega$  is *quasi-determined* (as a subset of  $X^\omega$ ) if there is a function  $\pi: X^{<\omega} \rightarrow \mathcal{P}(X) \setminus \{\emptyset\}$  such that one of the two following statements holds.

1. For every  $x \in X^\omega$ , if  $x(2n) \in \pi(x \upharpoonright 2n)$  holds for all  $n \in \omega$ , then  $x \in A$ .
2. For every  $x \in X^\omega$ , if  $x(2n+1) \in \pi(x \upharpoonright (2n+1))$  hold for all  $n \in \omega$ , then  $x \notin A$ .

A function  $\pi: X^{<\omega} \rightarrow \mathcal{P}(X) \setminus \{\emptyset\}$  is then said to be a *quasi-strategy* in the game  $G_A$ . Such a function  $\pi$  as in case (1) above is a *winning quasi-strategy* for player  $I$ ; in case (2) it is a winning quasi-strategy for player  $II$ . We let  $\text{quasi-AD}_X$  denote the statement that every subset of  $X^\omega$  is quasi-determined, as a subset of  $X^\omega$ .

If  $X$  is wellorderable, then  $\text{quasi-AD}_X$  and  $\text{AD}_X$  are equivalent. Remark 6.2.4 shows that a form of uniformization suffices to prove this.

**6.2.4 Remark.** If  $X$  is a set, and every subset of  $X^{<\omega} \times X$  is uniformized, then every quasi-determined game on  $X$  is determined. As  $\text{quasi-AD}_{\mathbb{R}}$  implies Uniformization (via a game where  $I$  plays a real, and  $II$  plays the infinite string of coordinates of a partner real),  $\text{quasi-AD}_{\mathbb{R}}$  and  $\text{AD}_{\mathbb{R}}$  are equivalent.

Given a set  $X$ , we let  $\text{OD}_X$  denote the class of sets which are definable from an ordinal and finite sequence from  $X$  (i.e., which are ordinal definable from a finite sequence from  $X$ ; see page 194 of [8] or page 145 of [24]). If there is a wellordering of  $X$  in  $\text{OD}_X$ , then there is a class-length wellordering of  $\text{OD}_X$  which is ordinal definable from a finite sequence from  $X$ .

The following theorem is the fundamental tool for showing the failure of Uniformization in certain models of AD. In particular, it can be used to show that, in  $L(\mathbb{R})$ , there are  $\prod_1^2$  sets which cannot be uniformized, and are therefore not Suslin.

**Theorem 6.2.5** (Kechris-Solovay). *Suppose that there is a set  $Z$  such that every subset of  $\omega^\omega$  is ordinal definable from  $Z$  and a member of  $\omega^\omega$ , and that for all  $x \in \omega^\omega$ ,  $\omega^\omega \not\subseteq \text{OD}_{\{Z, x\}}$ . Then the set  $\{(x, y) \in (\omega^\omega)^2 \mid y \notin \text{OD}_{\{Z, x\}}\}$  is not uniformizable.*

*Proof.* Any uniformizing function would be ordinal definable from  $Z$  and some  $x \in \omega^\omega$ , which would mean that  $f(x) \in \text{OD}_{\{Z, x\}}$ .  $\square$

## 6.3 The Solovay sequence

The *Solovay sequence* [36] is the unique continuous sequence  $\langle \theta_\alpha : \alpha \leq \beta \rangle$  such that

- $\theta_0$  is the least ordinal which is not the surjective image of  $\omega^\omega$  under an OD function;
- for every ordinal  $\alpha$  such that  $\alpha + 1 \leq \beta$ ,  $\theta_{\alpha+1}$  is the least ordinal which is not the surjective image of  $\omega^\omega$  under an  $\text{OD}_{\{A\}}$  function for any  $A \subseteq \omega^\omega$  of Wadge rank  $\theta_\alpha$ ;
- $\theta_\beta = \Theta$ .

We call  $\beta$  the *length* of the Solovay sequence. In  $L(\mathbb{R})$ ,  $\theta_0 = \Theta$ , so  $\beta = 0$ . For each  $\alpha \leq \beta$ , if we let  $\Gamma_\alpha$  be the set subsets of  $\omega^\omega$  of Wadge rank less than  $\theta_\alpha$ , then  $\text{HOD}_{\Gamma_\alpha}$  is a model of ZF containing  $\omega^\omega$  whose subsets of  $\omega^\omega$  are exactly the members of  $\Gamma_\alpha$ .

**6.3.1 Remark.** If  $\text{AD}_{\mathbb{R}}$  holds, and the length of the Solovay sequence has uncountable cofinality, then  $L(\Gamma_\alpha) \models \text{AD}_{\mathbb{R}}$  for club many  $\alpha$  on the Solovay sequence. To see this, suppose that  $\text{AD}_{\mathbb{R}}$  holds, and let  $\{A_x : x \in \omega^\omega\}$  be a set of subsets of  $(\omega^\omega)^\omega$ . Consider the real game in which player *I* plays  $x$  and the two players play the real game with payoff set  $A_x$ , with player *II* choosing first which player to be in the game with payoff set  $A_x$ . Player *I* cannot have a winning strategy. This shows for instance that, under  $\text{AD}_{\mathbb{R}}$ , for each  $A \subseteq \omega^\omega$  there is a  $B \subseteq \omega^\omega$  such that every real game with payoff set Wadge below  $A$  has a winning strategy with payoff set below  $B$ .

Theorems 2.5.9 and 6.2.5 give the following. Recall from Remark 2.1.8 that Lipschitz Determinacy implies that  $\aleph_1 \not\leq 2^{\aleph_0}$ . By Martin's theorem on the wellfoundedness of the Wadge hierarchy (Corollary 2.1.10), the assumption on Wadge ranks would follow from the additional assumption that every subset of  $\omega^\omega$  satisfies the Baire property.

**Theorem 6.3.2.** *Suppose that Lipschitz Determinacy and Uniformization hold, and that the Wadge rank of each subset of  $\omega^\omega$  exists. Then the length of the Solovay sequence is a limit ordinal, and there does not exist a  $\Theta$ -sequence of distinct elements of  $\mathcal{P}(\omega^\omega)$ .*

*Proof.* Suppose first that the length of the Solovay sequence is not a limit ordinal. Then there is a set  $A \subseteq \omega^\omega$  such that every ordinal below  $\Theta$  is a surjective image of  $\omega^\omega$  via a function which is ordinal definable from  $A$ . By Theorem 2.5.9, for each  $\gamma < \Theta$  there is a  $\gamma$ -sequence of subsets of  $\omega^\omega$  of increasing Wadge rank, ordinal definable from  $A$ . Since every set of reals has Wadge rank less than  $\Theta$ , it follows from Lipschitz Determinacy that every set of reals is definable from  $A$ , a real and an ordinal. On the other hand, since Uniformization and  $\aleph_1 \not\leq 2^{\aleph_0}$  hold (see Remark 2.1.8), Theorem 6.2.5 implies that there is a set  $B \subseteq \omega^\omega$  which is not ordinal definable from  $A$  and a real.

For the second part of the theorem, suppose that  $f: \Theta \rightarrow \mathcal{P}(\omega^\omega)$  is an injection. Then  $\{\text{WR}(f(\alpha)) : \alpha < \Theta\}$  is cofinal in  $\Theta$ , since otherwise there would be a surjection from  $\omega^\omega$  to  $\Theta$ . Then every set of reals is ordinal definable from  $f$  and a real, again contradicting Uniformization.  $\square$

**6.3.3 Remark.** Under  $\text{AD}^+$ , nonlimit members of the Solovay sequence below  $\Theta$  have cofinality  $\omega$ , by Theorem 13.1.7. On the other hand, if  $\text{AD} + \text{DC}_{\mathbb{R}}$  holds and the length of the Solovay sequence is not a limit ordinal, then  $\Theta$  is regular, since in this case there is an  $A \subset \omega^\omega$  such that every element of  $\mathcal{P}(\omega^\omega)$  is definable from  $A$ , an element of  $\omega^\omega$  and an ordinal, as shown in the proof of Theorem 6.3.2. A cofinal function  $c: \lambda \rightarrow \Theta$  and a surjection  $s: \omega^\omega \rightarrow \lambda$  could then be combined to build a set of reals of Wadge rank at least  $\Theta$  (using the definability order to pick, for each  $(x, y) \in (\omega^\omega)^2$  a set in  $\text{OD}_{\{y\}}$  of Wadge rank at least  $c(s(x))$ , if one exists), which is impossible.

A similar argument shows (from  $\text{AD} + \text{DC}_{\mathbb{R}}$ ) that if the length of the Solovay sequence is not a limit ordinal, then there is a function choosing for each  $\alpha < \Theta$  a set of reals of Wadge rank at least  $\alpha$ . If  $\text{AD} + \text{DC}_{\mathbb{R}}$  holds,  $V = \text{HOD}_{\mathcal{P}(\omega^\omega)}$  and the Solovay sequence has limit length then there is no  $\Theta$ -sequence of distinct elements of  $\mathcal{P}(\omega^\omega)$ , since any such sequence would have to be ordinal definable from a set of reals.

Solovay [36] showed that, assuming  $\text{AD}_{\mathbb{R}} + V = L(\mathcal{P}(\omega^\omega))$ ,  $\text{DC}$  holds if and only if the length of the Solovay sequence has uncountable cofinality. For the easier direction, note that  $\text{Lipschitz Determinacy} + \text{DC}_{\mathbb{R}} + \text{CC}_{\mathcal{P}(\omega^\omega)}$  implies that  $\text{cof}(\Theta)$  is uncountable, since otherwise there exists an  $\omega$ -sequence of sets of reals whose Wadge ranks are unbounded in  $\Theta$ , and thereby a set of reals of Wadge rank at least  $\Theta$ , contradicting Corollary 2.5.11. The following theorem gives the harder direction. Part of the proof is reused in the proof of Theorem 9.0.2, which shows that  $\text{DC}_{\mathbb{R}}$  follows from the assumption that the ultrapower of the ordinals by the Turing measure is wellfounded.

A ranking function on a tree  $T$  is a function  $\rho: T \rightarrow \text{Ord}$  such that  $\rho(s) > \rho(t)$  whenever  $t$  is a proper extension of  $s$ . The existence of a ranking function for  $T$  implies that  $T$  is wellfounded; the reverse implication follows from  $\text{DC}_T$ . The canonical ranking function  $\text{rank}_T$  of a wellfounded tree  $T$  sends each node  $t$  of  $T$  to the least ordinal greater than all the values  $\text{rank}_T(s)$  for  $s$  a proper extension of  $t$  in  $T$ .

**Theorem 6.3.4** (Solovay [36]). *If  $\text{AD} + \text{DC}_{\mathbb{R}}$  holds,  $V = \text{HOD}_{\mathcal{P}(\omega^\omega)}$  and  $\text{cof}(\Theta)$  is uncountable, then  $\text{DC}$  holds.*

*Proof.* By Remark 0.4.2, it suffices to show that  $\text{DC}_{\mathcal{P}(\omega^\omega)}$  holds. Let  $T$  be a tree on  $\mathcal{P}(\omega^\omega)$  without terminal nodes. We want to see that  $T$  has an infinite path. For each  $\xi < \Theta$ , let  $T_\xi$  be the set of  $t \in T$  such that every element of the range of  $t$  has Wadge rank less than  $\xi$ . Then  $T_\xi$  is a surjective image of  $\omega^\omega$ , and, since  $\text{DC}_{\mathbb{R}}$  holds, either  $T_\xi$  has an infinite path or  $T_\xi$  has a ranking function. We assume that the latter case holds for all  $\xi$ , since otherwise we are done.

For each  $t \in T$ , let  $w(t)$  be the least Wadge rank of a set  $A \subseteq \omega^\omega$  such that  $t \cap \langle A \rangle$  is in  $T$ . Suppose that for each  $\xi < \Theta$ ,  $w[T_\xi]$  is bounded in  $\Theta$ . Define

$s: \Theta \rightarrow \Theta$  by setting  $s(\xi)$  to be the supremum of  $w[T_\xi]$ . Then since  $\text{cof}(\Theta) > \aleph_0$  there is a  $\zeta < \Theta$  closed under  $w$ . Then  $T_\zeta$  is a subtree of  $T$  without terminal nodes, contradicting the assumption from the previous paragraph.

We continue the proof under the assumption that for some  $\xi_* < \Theta$ ,  $w[T_{\xi_*}]$  is cofinal in  $\Theta$ . Since  $T_{\xi_*}$  is a surjective image of  $\omega^\omega$ , the ordertype of  $w[T_{\xi_*}]$  must be less than  $\Theta$ . It follows that  $\Theta$  is singular. Let  $\lambda$  be its cofinality (necessarily a regular uncountable cardinal) and let  $c: \lambda \rightarrow \Theta$  be increasing, continuous and cofinal.

Toward a contradiction, suppose that  $T$  does not contain an infinite path. Then each  $T_\xi$  is also wellfounded. Let  $s: \omega^\omega \rightarrow \lambda$  be a surjection. For each  $x \in \omega^\omega$ , let  $s'(x) = \sup\{s(y) : y \in \omega^\omega, y \leq_{\text{TU}} x\}$ . Then  $s'$  is Turing invariant (i.e., Turing-equivalent inputs give the same output). Since  $\lambda > \aleph_0$ ,  $s'(x) < \lambda$  for each  $x \in \omega^\omega$ . For each  $t \in T$ , let  $r_t$  be the function on  $\omega^\omega$  defined by setting  $r_t(x)$  to be  $\text{rank}_{T_{s'(x)}}(t)$  if  $t \in T_{s'(x)}$  and 0 otherwise. Then each  $r_t$  is Turing invariant, and  $r_t(x) < r_{t'}(x)$  for a Turing cone of  $x$  whenever  $t < t'$  in  $T$ . Let  $R = \{r_t : t \in T\}$  and for each  $\alpha < \lambda$  let  $R_\alpha = \{r_t : t \in T_{c(\alpha)}\}$ .

Define the preorder  $\preceq$  on  $R$  by setting  $r_t \preceq r_{t'}$  to hold if  $r_t(x) \leq r_{t'}(x)$  holds for a Turing cone of  $x$ . Since every element of  $T$  has a proper extension in  $T$ ,  $\preceq$  has no minimal element. Since  $\text{DC}_\mathbb{R}$  holds and each  $T_\xi$  is wellfounded (i.e.,  $\text{rank}_{T_\xi}(t)$  exists for each  $t \in T_\xi$ ), the restriction of  $\preceq$  to each set  $R_\alpha$  is wellfounded, and in particular has a minimal element. Let  $\alpha_0 = 0$ , and for each  $n \in \omega$  let  $\alpha_{n+1}$  be the least  $\alpha < \lambda$  such that  $R_{\alpha_{n+1}}$  contains a proper  $\preceq$  lower bound for  $R_{\alpha_n}$ . Letting  $\xi = \sup\{\alpha_n : n \in \omega\}$ ,  $\xi < \lambda$  and  $(R_\xi, \preceq)$  is illfounded, giving a contradiction.  $\square$





## Part II

AD<sup>+</sup>



# Chapter 7

## $<\Theta$ -Determinacy

In this chapter we introduce  $<\Theta$ -Determinacy, which is one of the axioms of  $\text{AD}^+$ . In conjunction with the results of Chapter 5, Theorem 7.0.1 shows that  $\text{AD}$  implies the determinacy of games on ordinals below  $\Theta$  whose payoff sets are Suslin and co-Suslin. The axiom  $<\Theta$ -Determinacy asserts the determinacy of a set of ordinal games whose payoff sets are coded by a subset of  $\omega^\omega$  via a continuous function. As discussed in Remark 7.0.4,  $\text{AD}$  implies that  $<\Theta$ -Determinacy holds in  $L(\mathbb{R})$ .

The *Lusin-Sierpiński order* (sometimes called the Brouwer-Kleene order) is the ordering on finite sequences of ordinals defined by setting  $s <_{\text{LS}} t$  if either  $s$  properly extends  $t$  or  $s(n) < t(n)$  for the least  $n$  such that  $s(n) \neq t(n)$  (identifying a finite sequence  $s$  with the corresponding function with domain  $|s|$ ). Then  $<_{\text{LS}}$  is a (class-sized) strict linear order. Given a tree  $T$  consisting of finite sequences of ordinals,  $<_{\text{LS}}$  is wellfounded if and only if  $T$  is (i.e., if and only if  $T$  has no infinite branches). To see this, note that any node of  $T$  such that the restriction of  $<_{\text{LS}}$  is illfounded must have a proper extension with the same property.

As noted in Remark 5.0.1, strong partition cardinals satisfy the conditions on  $\delta$  in the statement of Theorem 7.0.1. The game  $\mathcal{G}_A$  in the statement of the theorem is as defined in Section 1.1 : players *I* and *II* alternate playing members of  $\lambda$ , with player *I* winning if and only if the resulting sequence is in  $A$ .

**Theorem 7.0.1** ( $\text{ZF} + \text{DC}_{\mathbb{R}}$ ; Moschovakis, Woodin). *Suppose that  $\delta$  is regular cardinal such that*

$$\forall \mu < \delta (\delta \rightarrow (\delta)_{\mu}^{\mu})$$

*holds. Let  $\lambda < \delta$  be an ordinal, and let  $A \subseteq \lambda^\omega$  be such that  $A$  and  $\lambda^\omega \setminus A$  are both  $\lambda$ -Suslin. Then  $\mathcal{G}_A$  is determined.*

*Proof.* Fix trees  $T$  and  $S$  on  $\lambda \times \lambda$  such that  $A = p[T]$  and  $\lambda^\omega \setminus A = p[S]$ .

For each  $u \in \lambda^{<\omega}$ , recalling that  $T_u$  denotes the set  $\{s : (u, s) \in T\}$ , let

$$T_u^* = \bigcup \{T_{u \restriction n} : n \leq |u|\}$$

and let

$$S_u^* = \bigcup \{S_{u \upharpoonright n} : n \leq |u|\}.$$

For each such  $u$ ,  $<_{\text{LS}} \upharpoonright T_u^*$  and  $<_{\text{LS}} \upharpoonright S_u^*$  are wellorderings; let  $\zeta_u$  and  $\xi_u$  denote their respective ordertypes. Then  $\max\{\zeta_u, \xi_u\} < \delta$ . For each  $x \in \lambda^\omega$ ,  $x \in A$  if and only if the restriction of  $<_{\text{LS}}$  to

$$T_x = \left( \bigcup_{n \in \omega} T_{x \upharpoonright n} \right)$$

is illfounded, and  $x \notin A$  if and only if the restriction of  $<_{\text{LS}}$  to

$$S_x = \left( \bigcup_{n \in \omega} S_{x \upharpoonright n} \right)$$

is illfounded.

Let  $OP_u^T$  denote the set of functions from  $T_u^*$  to  $\delta$  for which, for all  $s, t \in T_u^*$ , if  $s <_{\text{LS}} t$  then  $f(s) < f(t)$ . Let  $OP_u^S$  denote the set of functions from  $S_u^*$  to  $\delta$  for which, for all  $s, t \in S_u^*$ , if  $s <_{\text{LS}} t$  then  $f(s) < f(t)$ .

We define two games of length  $\omega$ ,  $\mathcal{G}^T$  and  $\mathcal{G}^S$ . In  $\mathcal{G}^T$ , player  $I$  plays ordinals  $\alpha_i \in \lambda$  (for  $i \in \omega$  even) and player  $II$  plays pairs  $(\alpha_i, f_i)$  (for  $i \in \omega$  odd) such that each  $\alpha_i$  is in  $\lambda$  and each  $f_i$  is a function in  $OP_{\langle \alpha_0, \dots, \alpha_i \rangle}^T$ . Player  $II$  wins a run  $\mathcal{G}^T$  if and only if  $f_i \subseteq f_j$  for all  $i < j$  in  $\omega$ .

I	$\alpha_0$	$\alpha_2$	$\alpha_4$	$\dots$
II	$\alpha_1, f_1$	$\alpha_3, f_3$	$\dots$	

The game  $\mathcal{G}^T$

In  $\mathcal{G}^S$ , player  $II$  plays ordinals  $\alpha_i \in \lambda$  (for  $i \in \omega$  odd) and player  $I$  plays pairs  $(\alpha_i, f_i)$  (for  $i \in \omega$  even) such that each  $\alpha_i$  is in  $\lambda$  and each  $f_i$  is a function in  $OP_{\langle \alpha_0, \dots, \alpha_i \rangle}^S$ . Player  $I$  wins a run of  $\mathcal{G}^S$  if and only if  $f_i \subseteq f_j$  for all  $i < j$  in  $\omega$ .

I	$\alpha_0, f_0$	$\alpha_2, f_2$	$\alpha_4, f_4$	$\dots$
II	$\alpha_1$	$\alpha_3$	$\dots$	

The game  $\mathcal{G}^S$

The games  $\mathcal{G}^T$  and  $\mathcal{G}^S$  are each closed. By the Gale-Stewart theorem on the determinacy of closed games [3], either player  $I$  has a winning strategy for  $\mathcal{G}^T$  or player  $II$  has a winning quasi-strategy (not necessarily a strategy, as the set of possible moves for  $II$  may not be wellorderable). Similarly, in  $\mathcal{G}^S$ , either player  $II$  has a winning strategy, or player  $I$  has a winning quasi-strategy. It cannot

be, however, that player  $II$  has a winning quasi-strategy  $\tau_{II}$  in  $\mathcal{G}^T$  and player  $I$  has a winning quasi-strategy  $\tau_I$  in  $\mathcal{G}^S$ . Supposing that such quasi-strategies existed, using  $\text{DC}_{\mathbb{R}}$ , the fact that  $\delta < \Theta$ , the Moschovakis Coding Lemma (to code subsets of  $\delta$  by reals), and the fact that each function  $f_i$  as above can be coded by a subset of  $\delta$ , we could find  $\langle (\alpha_i, f_i) : i \in \omega \rangle$  such that

$$\langle \alpha_0, (\alpha_1, f_1), \alpha_2, (\alpha_3, f_3), \dots \rangle$$

is according to  $\tau_{II}$  and

$$\langle (\alpha_0, f_0), \alpha_1, (\alpha_2, f_2), \alpha_3, \dots \rangle$$

is according to  $\tau_I$ . Then  $\bigcup_{i \in \omega} f_{2i+1}$  would witness that  $\langle \alpha_i : i \in \omega \rangle \notin A$ , and  $\bigcup_{i \in \omega} f_{2i}$  would witness that  $\langle \alpha_i : i \in \omega \rangle \in A$ , giving a contradiction.

The proof is then completed by proving the following claims.

**Claim 1.** *If  $I$  has a winning strategy in  $\mathcal{G}^T$ , then  $I$  has a winning strategy in  $\mathcal{G}_A$ .*

**Claim 2.** *If  $II$  has a winning strategy in  $\mathcal{G}^S$ , then  $II$  has a winning strategy in  $\mathcal{G}_A$ .*

Recall that for an ordinal  $\alpha$  and a set of ordinals  $X$ ,  $[X]^\alpha$  denotes the collection of subsets of  $X$  of ordertype  $\alpha$ , and that we identify each element of  $[X]^\alpha$  with the corresponding order-preserving function on  $\alpha$  which enumerates it. Recalling notation from Chapter 5, given an ordinal-valued function  $g$  on  $\omega \cdot \alpha$ , for some ordinal  $\alpha$ , let  $g^*$  be the function on  $\alpha$  defined by setting  $g^*(\xi)$  to be the supremum of  $\{g(\omega \cdot \xi + n) : n \in \omega\}$ . For each set  $X$  of ordinals, let  $X^*(\alpha)$  be  $\{g^* : g \in [X]^{\omega \cdot \alpha}\}$ . Let  $\mu_\alpha$  be the set of  $A \subseteq [\delta]^\alpha$  for which there exists a club  $C \subseteq \delta$  with  $C^*(\alpha) \subseteq A$ . By Theorem 5.0.3, each  $\mu_\alpha$  is a  $\delta$ -complete (so  $\lambda$ -complete) measure on the corresponding set  $[\delta]^\alpha$ .

Proof of Claim 1 : For each  $u \in \omega^{<\omega}$ , let  $Q_u$  be the bijection from  $[\delta]^{\zeta_u}$  to  $OP_u^T$  with the property that  $Q_u(g)$  and  $g$  have the same range, for all  $g \in [\delta]^{\zeta_u}$ . Let  $\nu_u$  be  $\{Q_u[A] : A \in \mu_{\zeta_u}\}$ . Then  $\nu_u$  is a  $\delta$ -complete ultrafilter on  $OP_u^T$ .

Fix a winning strategy  $\Sigma$  for  $I$  in  $\mathcal{G}^T$ . We define a strategy  $\sigma$  for  $I$  in  $\mathcal{G}_A$ . We let  $\sigma(\langle \rangle) = \Sigma(\langle \rangle)$ . For each positive  $n \in \omega$ , we let  $\sigma(\langle \alpha_0, \dots, \alpha_{2n-1} \rangle)$  be the unique value  $\alpha$  such that, letting  $u = \langle \alpha_0, \dots, \alpha_{2n-1} \rangle$ , for  $\nu_u$ -many  $f \in OP_u^T$ ,

$$\Sigma(\alpha_0, (\alpha_1, f \restriction T_{u \restriction 2}^*), \alpha_2, (\alpha_3, f \restriction T_{u \restriction 4}^*), \dots, \alpha_{2n-2}, (\alpha_{2n-1}, f)) = \alpha.$$

Now suppose that  $\langle \alpha_i : i \in \omega \rangle$  is the result of a run of  $\mathcal{G}_A$  according to  $\sigma$ , and that  $I$  has lost, so that  $\langle \alpha_i : i \in \omega \rangle$  is not in  $A$ . Let  $x = \langle \alpha_i : i \in \omega \rangle$ . Then  $x \notin p[T]$ , so  ${}_{<LS} \restriction T_x$  is wellfounded and there exists a function  $f : T_x \rightarrow \delta$  preserving  ${}_{<LS}$ , i.e., such that  $f \restriction T_{x \restriction n}^* \in OP_{x \restriction n}^T$  for all  $n \in \omega$ .

For each  $n \in \omega$ , let  $A_{2n+2}$  be the set in  $\nu_{x \restriction 2n+2}$  used to choose  $\alpha_{2n+2}$ . Using  $\text{CC}_{\mathbb{R}}$ , the Moschovakis Coding Lemma and the fact that  $\delta < \Theta$ , we can find a sequence  $\langle C_{2n+2} : n \in \omega \rangle$  of club subsets of  $\delta$  such that each  $C_{2n+2}$  witnesses

(via the corresponding function  $Q_{x \restriction 2n+2}$ ) that the corresponding set  $A_{2n+2}$  is in  $\nu_{x \restriction 2n+2}$  (so, for all  $g \in [C_{2n+2}]^{\omega \cdot \zeta_{x \restriction 2n+2}}$ ,  $Q_{x \restriction 2n+2}(g^*)$  is in  $A_{2n+2}$ ). Let  $C$  be  $\bigcap_{n \in \omega} C_{2n+2}$ , and let  $C'$  be the set of  $\beta \in C$  such that the ordertype of  $C \cap \beta$  is  $\gamma + \omega$ , for some ordinal  $\gamma$ .

Let  $g$  be an order-preserving function from the range of  $f$  into  $C'$ , and for each  $n \in \omega$ , let  $f_{2n+1}$  be  $(g \circ f) \restriction T_{x \restriction 2n+2}^*$ . Then  $Q_{x \restriction 2n+2}^{-1}(f_{2n+1})$  is equal to  $h^*$  for some  $h \in [C]^{\omega \cdot \zeta_{x \restriction 2n+2}}$ , so  $f_{2n+1}$  is in  $A_{2n+2}$ . We have then that the run of  $\mathcal{G}^T$  given by

$$\alpha_0, (\alpha_1, f_1), \alpha_2, (\alpha_3, f_3), \dots$$

is according to  $\Sigma$  yet losing for player  $I$ , giving a contradiction.

Proof of Claim 2: For each  $u \in \omega^{<\omega}$ , let  $Q_u$  be the bijection from  $[\delta]^{\xi_u}$  to  $OP_u^S$  with the property that  $Q_u(g)$  and  $g$  have the same range, for all  $g \in [\delta]^{\xi_u}$ . Let  $\nu_u$  be  $\{Q_u[A] : A \in \mu_{\xi_u}\}$ . Then  $\nu_u$  is a  $\delta$ -complete ultrafilter on  $OP_u^S$ .

Fix a winning strategy  $\Sigma$  for  $II$  in  $\mathcal{G}^S$ . We define a strategy  $\sigma$  for  $II$  in  $\mathcal{G}_A$ . For each  $n \in \omega$ , we let  $\sigma(\langle \alpha_0, \dots, \alpha_{2n} \rangle)$  be the unique value  $\alpha$  such that, letting  $u = \langle \alpha_0, \dots, \alpha_{2n} \rangle$ , for  $\nu_u$ -many  $f \in OP_u^S$ ,

$$\Sigma((\alpha_0, f \restriction S_{u \restriction 1}^*), \alpha_1, (\alpha_2, f \restriction S_{u \restriction 1}^*), \dots, \alpha_{2n-1}, (\alpha_{2n}, f)) = \alpha.$$

Now suppose that  $\langle \alpha_i : i \in \omega \rangle$  is the result of a run of  $\mathcal{G}_A$  according to  $\sigma$ , and that  $II$  has lost, so that  $\langle \alpha_i : i \in \omega \rangle$  is in  $A$ . Let  $x = \langle \alpha_i : i \in \omega \rangle$ . Then  $x \notin p[S]$ , so  $<_{LS} \restriction S_x$  is wellfounded and there exists a function  $f : S_x \rightarrow \delta$  preserving  $<_{LS}$ , i.e., such that  $f \restriction S_{x \restriction n}^* \in OP_{x \restriction n}^S$  for all  $n \in \omega$ .

For each  $n \in \omega$ , let  $A_{2n+1}$  be the set in  $\nu_{x \restriction 2n+1}$  used to choose  $\alpha_{2n+1}$ . Using  $CC_{\mathbb{R}}$ , the Moschovakis Coding Lemma and the fact that  $\delta < \Theta$ , we can find a sequence  $\langle C_{2n+1} : n \in \omega \rangle$  of club subsets of  $\delta$  such that each  $C_{2n+1}$  witnesses (via the corresponding function  $Q_{x \restriction 2n+1}$ ) that the corresponding set  $A_{2n+1}$  is in  $\nu_{x \restriction 2n+1}$  (so, for all  $g \in [C_{2n+1}]^{\omega \cdot \xi_{x \restriction 2n+1}}$ ,  $Q_{x \restriction 2n+1}(g^*)$  is in  $A_{2n+1}$ ). Let  $C$  be  $\bigcap_{n \in \omega} C_{2n+1}$ , and let  $C'$  be the set of  $\beta \in C$  such that the ordertype of  $C \cap \beta$  is  $\gamma + \omega$ , for some ordinal  $\gamma$ .

Let  $g$  be an order-preserving function from the range of  $f$  into  $C'$ , and for each  $n \in \omega$ , let  $f_{2n}$  be  $(g \circ f) \restriction S_{x \restriction 2n+1}^*$ . Then  $Q_{x \restriction 2n+1}^{-1}(f_{2n})$  is equal to  $h^*$  for some  $h \in [C]^{\omega \cdot \xi_{x \restriction 2n+1}}$ , so  $f_{2n}$  is in  $A_{2n+1}$ . We have then that the run of  $\mathcal{G}^S$  given by

$$(\alpha_0, f_0), \alpha_1, (\alpha_2, f_2), \alpha_3, \dots$$

is according to  $\Sigma$  yet losing for player  $II$ , giving a contradiction.  $\square$

**7.0.2 Remark.** Consider the game of length  $\omega$  in which player  $I$  plays  $\alpha \in \omega_1$  (and then makes no other moves) and then player  $II$  plays (digit by digit) an  $x \in \omega^\omega$ , with  $II$  winning if and only if  $x$  HC-codes (in the sense given in Section 0.2) a wellordering of  $\omega$  of ordertype  $\alpha$ . Player  $I$  can never have a winning strategy in this game, and if the Axiom of Choice holds, then player  $II$  has a winning strategy. By part (2) of Remark 1.1.2, if AD holds then there is no injection from  $\omega_1$  into  $\omega^\omega$ , so this game is not determined. The statement that  $\aleph_1 \not\leq 2^{\aleph_0}$  implies that the payoff set for this game is not Suslin.

Modifying this example slightly, suppose that  $\pi: \omega^\omega \rightarrow \omega_1$  is a surjection, and consider the real game where players  $I$  and  $II$  collaborate to build a sequence  $\langle x_i : i \in \omega \rangle \in (\omega^\omega)^\omega$ , with player  $II$  winning if  $\{\pi(x_i) : i < \omega\}$  is an ordinal. If Uniformization holds, player  $II$  has a winning strategy. However, if  $\aleph_1 \not\leq 2^{\aleph_0}$ , then player  $II$  does not have a winning strategy in the corresponding ordinal game. Moreover, player  $II$  doesn't have a winning strategy for the game in which players  $I$  and  $II$  collaborate to build a sequence  $\langle \alpha_i : i \in \omega \rangle \in \omega_1^\omega$ , with player  $II$  winning if the ordertype of  $\{\alpha_i : i < \omega\}$  is at least  $\alpha_0$ .

Suppose now that  $\lambda$  is an ordinal, and that  $f: \lambda^\omega \rightarrow \omega^\omega$  is a continuous function, with respect to the discrete topology on  $\lambda$  and the corresponding product topology on  $\lambda^\omega$ . Given  $A \subseteq \omega^\omega$ , let  $\mathcal{G}_{f,A}$  be the game in which players  $I$  and  $II$  alternate choosing ordinals  $\alpha_i \in \lambda$  ( $i \in \omega$ ), where  $I$  wins if and only if  $f(\langle \alpha_i : i \in \omega \rangle)$  is in  $A$ .

I	$\alpha_0$	$\alpha_2$	$\alpha_4$	$\dots$
II		$\alpha_1$	$\alpha_3$	$\dots$

The game  $\mathcal{G}_{f,A}$

We let  $\lambda$ -Determinacy denote the statement that  $\mathcal{G}_{f,A}$  is determined for all continuous  $f: \lambda^\omega \rightarrow \omega^\omega$  and  $A \subseteq \omega^\omega$ . Given an ordinal  $\gamma$ , we let  $<\gamma$ -Determinacy denote the statement that  $\lambda$ -Determinacy holds for all  $\lambda < \gamma$ .

If  $f: \lambda^\omega \rightarrow \omega^\omega$  and  $A \subseteq \omega^\omega$  is Suslin and co-Suslin, then so is the payoff set  $f^{-1}[A]$  for  $\mathcal{G}_{f,A}$ . Putting together Theorems 5.0.4 and 7.0.1 then, we get the following.

**Corollary 7.0.3** (ZF + AD). *For every  $\lambda < \Theta$  and every continuous function  $f: \lambda^\omega \rightarrow \omega^\omega$ , if  $A \subseteq \omega^\omega$  is Suslin and co-Suslin, then the game  $\mathcal{G}_{f,A}$  is determined.*

**7.0.4 Remark.** Suppose that  $f: \lambda^\omega \rightarrow \omega^\omega$  is a continuous function, for some ordinal  $\lambda < \Theta$ , and that  $A \subseteq \omega^\omega$ . Let  $\prec$  be a strict prewellordering of a subset of  $\omega^\omega$  of length  $\lambda$ . By the Moschovakis Coding Lemma, every strategy in  $\mathcal{G}_{f,A}$  is coded by a subset of  $\omega^\omega$  in  $\text{pos-}\Sigma_1^1(\prec)$ , as is  $f$ . In particular, if  $\mathcal{G}_{f,A}$  is determined, then it is determined in  $L(A, \prec, \mathbb{R})$ . Similarly, since the negation of  $<\Theta$ -Determinacy is a  $\Sigma_1^2$  statement (using the Coding Lemma), Theorem 6.1.22 implies that, assuming  $\text{AD} + V = L(\mathbb{R})$ , the failure of  $<\Theta$ -Determinacy would imply the nondeterminacy of a game of the form  $\mathcal{G}_{f,A}$  for some Suslin, co-Suslin set  $A$ , contradicting Corollary 7.0.3.

Using the fact that, under AD, every subset of  $\omega_1$  is constructible from a real, and the sharp of each real exists, it is not hard to see that AD implies  $\omega_1$ -Determinacy (this observation is apparently due to the author). It is apparently an open question whether AD implies  $\omega_2$ -Determinacy.

We will consider yet another form of ordinal determinacy in Chapter 12.





# Chapter 8

## $\infty$ -Borel sets

In this chapter we complete the definition of  $\text{AD}^+$  by introducing the notion of  $\infty$ -Borel set. In Section 8.1 we introduce two equivalent notions of  $\infty$ -Borel code. In Section 8.2 we show that every  $\infty$ -Borel set  $A$  has an  $\infty$ -Borel code which is projective in  $A$  (Theorem 8.2.3). In conjunction with Remark 7.0.4, this shows that the implication  $\text{AD} \Rightarrow \text{AD}^+$  holds in the inner model  $L(\mathbb{R})$ .

### 8.1 $\infty$ -Borel codes

Given infinite ordinals  $\gamma$  and  $\delta$ , we let  $\mathcal{L}_{\gamma,\delta}^0$  be the language with 0-ary predicates  $P_\alpha$  ( $\alpha \in \delta$ ), closed under negation as well as conjunctions and disjunctions indexed by members of  $\gamma$ . We write  $\mathcal{L}_\gamma^0$  for  $\mathcal{L}_{\gamma,\omega}^0$ , which is the primary case of interest.

We associate to each sentence in  $\mathcal{L}_{\gamma,\delta}^0$  a unique code (in  $\mathcal{P}(\gamma)$ , if  $\delta \leq \gamma$  and  $\gamma$  is a cardinal) via a definable (injective) pairing function  $\prec \cdot, \cdot \succ$  on the ordinals (as defined in Section 0.2), and write  $\varphi_S$  for the sentence coded by the set  $S$ . Fixing one such coding, we let

- $\varphi_{\{ \prec 0, \alpha \succ \}}$  be  $P_\alpha$  for each  $\alpha \in \delta$ ,
- $\varphi_{\{ \prec 1, \alpha \succ : \alpha \in S \}}$  be  $\neg \varphi_S$ , and
- $\varphi_{\{ \prec 2 + \alpha, \beta \succ : \alpha < \eta, \beta \in S_\alpha \}}$  be  $\bigwedge_{\alpha < \eta} \varphi_{S_\alpha}$ .

We let  $\mathcal{L}_{\infty,\delta}^0$  be the class of sentences which are in  $\mathcal{L}_{\gamma,\delta}^0$  for some infinite ordinal  $\gamma$ . Each  $x \in 2^\delta$  can be thought of as an  $\mathcal{L}_{\infty,\delta}^0$ -structure, where, for each  $\alpha \in \delta$ ,  $P_\alpha$  is interpreted as true if and only if  $x(\alpha) = 1$ . For a sentence  $\varphi$  of  $\mathcal{L}_{\infty,\delta}^0$ , we let  $A_\varphi$  be the set of  $x \in 2^\delta$  such that  $x \models \varphi$ , and say that  $\varphi$  is an  $\infty$ -Borel code for  $A_\varphi$ , or a  $\gamma$ -Borel code, if  $\varphi$  is in  $\mathcal{L}_{\gamma,\delta}^0$ . We also say that a set of ordinals  $S$  is an  $\infty$ -Borel code or  $\gamma$ -Borel code if the corresponding formula  $\varphi_S$  is.

**8.1.1 Definition.** Given an infinite ordinal  $\gamma$ , a subset of  $2^\delta$  is  $\gamma$ -Borel if it is equal to  $A_\varphi$  for some  $\varphi \in \mathcal{L}_{\gamma,\delta}^0$ , and  $<\gamma$ -Borel if it is  $\eta$ -Borel for some infinite ordinal  $\eta < \gamma$ . A subset of  $2^\delta$  is  $\infty$ -Borel if it is  $\gamma$ -Borel for some infinite ordinal  $\gamma$ .

**8.1.2 Remark.** For each infinite cardinal  $\kappa$ , the class of  $\kappa$ -Borel subsets of  $2^\omega$  is closed under continuous preimages, and thus Wadge reducibility. To see this, fix  $\varphi \in \mathcal{L}_{\kappa,\omega}^0$  and a continuous function  $f: 2^\omega \rightarrow 2^\omega$ . To express “ $f(x) \models \varphi$ ” it suffices to find, for each  $n \in \omega$ , an expression for “ $f(x)(n) = 1$ ”, since then one can form a  $\psi$  with  $A_\psi = f^{-1}[A_\varphi]$  by replacing each instance of each  $P_n$  in  $\varphi$  with the corresponding expression. The existence of such expressions follows from the continuity of  $f$ .

The following remark shows that Suslin subsets of  $2^\delta$  (for any ordinal  $\delta$ ) are  $\infty$ -Borel. This also follows easily from item (2) of Theorem 8.1.7 below. Given a tree  $T$  and a node  $t \in T$ , we let  $T_t$  (for this remark) be the set of  $s$  such that the concatenation  $t \smallfrown s$  is in  $T$ . We define  $\text{rank}(T)$  to be 0 if  $T$  is empty or  $T = \{\emptyset\}$ . Otherwise,  $\text{rank}(T)$  is the strict supremum of the set  $\{\text{rank}(T_t) : t \in T, |t| = 1\}$ .

**8.1.3 Remark.** Let  $T$  be a tree on  $\delta \times \gamma$ , for some ordinals  $\delta \leq \gamma$ , with  $\gamma$  infinite. Given a node  $(s, t) \in T$  and an  $x \in 2^\delta$ , let  $T_{x,s,t}$  denote the set of  $(s', t') \in \delta^{<\omega} \times \gamma^{<\omega}$  for which  $s \smallfrown s'$  is an initial segment of  $x$  and  $(s \smallfrown s', t \smallfrown t')$  is in  $T$ . One can define recursively on  $\alpha \leq |\gamma|^+$  formulas  $\varphi_{(s,t)}^\alpha \in \mathcal{L}_{\gamma,\delta}^0$ , simultaneously for each  $(s, t) \in \delta^{<\omega} \times \gamma^{<\omega}$ , such that  $A_{\varphi_{(s,t)}^\alpha}$  is the set of  $x \in 2^\omega$  for which  $\text{rank}(T_{x,s,t}) = \alpha$ . The projection of  $T$  is then defined by  $\bigwedge_{\alpha < |\gamma|^+} \neg \varphi_{(\emptyset, \emptyset)}^\alpha$ .

The following is a variation of Remark 8.1.3 showing that there exists an  $\omega_1$ -sequence of  $\infty$ -Borel codes for disjoint subsets of  $2^\omega$ .

**8.1.4 Example.** Fixing a tree  $T$  on  $\delta \times \gamma$ , for some ordinals  $\delta$  and  $\gamma$ , one can define, by recursion on  $\alpha$ ,  $\mathcal{L}_{\infty,\omega}^0$ -sentences  $\varphi_{T,\alpha}$  in such a way that each  $A_{\varphi_{T,\alpha}}$  is the set of  $x \in 2^\delta$  for which the tree  $T_x = \{t \in \gamma^{<\omega} : (x \restriction |t|, t) \in T\}$  has rank at most  $\alpha$ . Starting with a natural tree on  $\omega \times \omega_1$  whose projection is the set of  $x \in 2^\omega$  which HC-code wellorders of  $\omega$ , this operation gives a sequence of  $\mathcal{L}_{\omega_1}^0$ -sentences  $\langle \varphi_\alpha : \alpha < \omega_1 \rangle$  such that each  $A_{\varphi_\alpha}$  is the set of HC-codes for wellorders of  $\omega$  of ordertype  $\alpha$ .

**8.1.5 Remark.** There is in general no method for picking an  $\infty$ -Borel code for each  $\infty$ -Borel set. To see this, let  $\mathbb{E}_0$  be the equivalence relation of mod-finite agreement on  $2^\omega$ , and suppose that every subset of  $2^\omega$  has the property of Baire. If  $F$  were a function picking for each  $\mathbb{E}_0$ -class a code for the class, then by the Kuratowski-Ulam theorem (which among other things implies that if every set of reals has the property of Baire then every wellordered union of meager sets is meager) comeagerly many reals would have the same code for their class.

**8.1.6 Remark.** The collection of  $\infty$ -Borel subsets of  $2^\omega$  is not necessarily closed under wellordered unions, even assuming DC. For example, let  $\{x_\alpha : \alpha < \omega_1\}$  be mutually generic Cohen reals over  $L$ , and consider the model  $L(\omega^\omega, F)$ , where  $\omega^\omega$

is the set of all reals in the forcing extension, and  $F$  is the function taking  $\alpha < \omega_1$  to the  $\mathbb{E}_0$ -class of  $x_\alpha$ . This model satisfies  $\text{DC}_{\mathbb{R}}$  since the forcing extension satisfies AC. It then satisfies DC, as in Remark 0.4.2. The set  $\bigcup_{\alpha < \omega_1} F(\alpha)$  cannot be  $\infty$ -Borel in this model, even though it is a wellordered union of countable sets. To see this, note first that any  $\infty$ -Borel code for this set would be ordinal definable from  $F$  and finitely many reals. These finitely many reals would appear in  $L[\{x_\alpha : \alpha < \beta\}]$  for some  $\beta < \omega_1$ . The homogeneity of the rest of the forcing would then imply that the  $\infty$ -Borel code for the union would be in this model. This is impossible, since for instance if  $\omega_1^L$  is countable it is possible to build two generic filters for the remainder of the forcing for which the resulting model  $L(\omega^\omega, F)$  is the same, and such that some real  $x$  is equal to  $x_\beta$  in one extension and not a member of any  $F(\alpha)$  in the other. For instance, starting with one such generic, replace  $x_\beta$  with

$$\{(2n, x_\beta(n) : n \in \omega\} \cup \{(2n+1, x_{\beta+1}(n) : n \in \omega\}$$

and  $x_{\beta+1+\alpha}$  with  $x_{\beta+2+\alpha}$  for each  $\alpha < \omega_1$ . (This example was constructed by the author in collaboration with Trevor Wilson, although it was very likely noticed earlier. A similar example can be constructed adding countably many Cohen reals; in this case  $\text{DC}_{\mathbb{R}}$  fails.)

Theorem 8.1.7 gives a number of convenient reformulations of the notion of  $\infty$ -Borel set. Item (2) below is especially useful.

Given ordinals  $\alpha$  and  $\delta$ ,  $x \in 2^\delta$  and  $C \subseteq \alpha^\omega \times 2^\delta$ , let  $G(\alpha, x, C)$  be the length- $\omega$  game on  $\alpha$  such that player  $I$  wins if  $(\vec{\beta}, x) \in C$ , where  $\vec{\beta}$  is the sequence produced by the run of the game. The notion of closure in parts (3) and (4) below refers to the product topology on both  $\alpha^\omega$  and  $2^\delta$ .

**Theorem 8.1.7** (ZF). *Let  $\delta$  be an ordinal. The following are equivalent, for a set  $A \subseteq 2^\delta$ .*

1.  $A$  is  $\infty$ -Borel.
2. For some set of ordinals  $S$  and some first-order formula  $\theta$ ,

$$A = \{x \in 2^\delta : L[S, x] \models \theta(S, x)\}.$$

3. For some ordinal  $\alpha$  and some clopen  $C \subseteq \alpha^\omega \times 2^\delta$ ,  $A$  is the set of  $x \in 2^\delta$  for which player  $I$  has a winning strategy in  $G(\alpha, x, C)$ .
4. For some ordinal  $\alpha$  and some closed  $C \subseteq \alpha^\omega \times 2^\delta$ ,  $A$  is the set of  $x \in 2^\delta$  for which player  $I$  has a winning strategy in  $G(\alpha, x, C)$ .

*Proof.* The implication from (1) to (2) follows from the fact that sentences in  $\mathcal{L}_{\infty, \delta}^0$  can be coded by sets of ordinals, and the definability of the relation  $x \models \varphi$ . For the reverse direction, for each ordinal  $\alpha$ , let

- $\mathcal{L}_\alpha^*$  be the extension of the first-order language of set theory given by adding predicate symbols  $\dot{S}, \dot{x}$  and constant symbols  $\dot{\beta}$  for each  $\beta \in \alpha$  and

- $T_\alpha$  denote the collection of  $\mathcal{L}_\alpha^*$  sentences of the form

$$\forall x_1 \dots \forall x_n ((x_1 = X_{\varphi_1}^{\beta_1} \wedge \dots \wedge x_n = X_{\varphi_n}^{\beta_n}) \rightarrow \psi)$$

where

- each  $\beta_i$  is in  $\alpha$ ;
- each  $\varphi_i$  is a unary formula in the corresponding  $\mathcal{L}_{\beta_i}^*$ ;
- each expression of the form  $x_i = X_{\varphi_i}^{\beta_i}$  denotes the formula saying that  $x_i$  is the set of sets satisfying  $\varphi_i$  in the structure

$$(L_{\dot{\beta}_i}[\dot{S}, \dot{x}]; \dot{S}, \dot{x}, \dot{\delta} (\delta \in d_i), \in),$$

where  $d_i$  is the set of symbols from  $\{\dot{\delta} : \delta < \beta_i\}$  appearing in  $\varphi_i$ ;

- $\psi$  is a  $\mathcal{L}_\alpha^*$  formula with free variables  $x_1, \dots, x_n$  (so each sentence of  $\mathcal{L}_\alpha^*$  is itself an element of  $T_\alpha$ ).

Since each expression of the form  $x_i = X_{\varphi_i}$  above asserts that  $x_i$  is defined by  $\varphi_i$ , each sentence in  $T_\alpha$  holds in the same structures as the corresponding sentence

$$\exists x_1 \dots \exists x_n (x_1 = X_{\varphi_1}^{\beta_1} \wedge \dots \wedge x_n = X_{\varphi_n}^{\beta_n} \wedge \psi).$$

We refer to this below as the *existential form* of the sentence.

Working (in  $L[S]$ ) by recursion on  $\alpha$ , we describe a procedure which associates to each sentence  $\theta \in T_\alpha$  a sentence  $\rho_{\theta, \alpha}$  of  $\mathcal{L}_{\omega \cup \alpha, \delta}^0$  such that for all  $x \in 2^\delta$ ,  $L_\alpha[S, x] \models \theta$  (with the symbols  $\dot{\beta}$  ( $\beta < \alpha$ ),  $\dot{S}$  and  $\dot{x}$  given their natural interpretations) if and only if  $x \models \rho_{\theta, \alpha}$ . For finite  $\alpha$  this follows from the fact that the  $\mathcal{L}_\alpha^*$ -theory of  $(L_\alpha[S, x]; S, x, \beta (\beta \in \alpha), \in)$  depends only on  $x \upharpoonright \alpha$  (as  $S$  is fixed) so the desired  $\rho_{\theta, \alpha}$  can consist of a disjunction of conjunctions describing  $x \upharpoonright \alpha$  exactly (and since there is a canonical finite set of such sentences our procedure can pick the least one in a suitable ordering).

For the limit and successor cases, we induct on the complexity of  $\psi$  in the representation of the members of  $T_\alpha$  given above. For limit  $\alpha$ , the induction hypothesis gives us the desired sentences  $\rho_{\theta, \alpha}$  when  $\psi$  is a  $\Delta_0$  formula, and the steps corresponding to  $\wedge$ ,  $\vee$  and  $\neg$  are handled by combining the formulas  $\rho_{\theta, \alpha}$  in the same manner (to see that this works in the case of  $\vee$  and  $\neg$ , use the fact that each sentence in  $T_\alpha$  has an equivalent existential form). If  $\psi$  has the form  $Qx\psi'$ , for  $Q$  either  $\forall$  or  $\exists$ , we let  $\rho_{\theta, \alpha}$  be the conjunction (when  $Q = \forall$ ) or the disjunction (when  $Q = \exists$ ) of all the sentences  $\rho_{\theta_\varphi^\beta, \alpha}$ , where  $\beta < \alpha$ ,  $\varphi$  is a unary formula in  $\mathcal{L}_\beta^*$  and  $\theta_\varphi^\beta$  is formed from  $\theta$  by adding  $\forall x$  to the beginning of  $\theta$  and  $x = X_\varphi^\beta$  at the appropriate place. The point here is that  $\alpha \times \mathcal{L}_\alpha^*$  is wellorderable, and each set in the range of the quantifier  $Q$  is defined by some formula in  $\bigcup_{\beta < \alpha} \mathcal{L}_\beta^*$ .

The successor step from  $\alpha$  to  $\alpha + 1$  is similar, except that, given a sentence  $\theta \in T_{\alpha+1}$  as above, with  $\psi$  a  $\Delta_0$  formula, we must deal with the possibility that some of the formulas  $\varphi_i$  define subsets of  $L_\alpha[S, x]$ , so that we cannot simply

apply the induction hypothesis. Each such  $T_{\alpha+1}$  sentence  $\theta$  is equivalent to a sentence  $\theta'$  in  $T_\alpha$ , formed by moving the introduction of the corresponding variables and their definitions into the  $\psi$  part of  $\theta'$ . For instance, if  $x_i = X_{\varphi_i}^\alpha$  appears in  $\theta$ , then it does not appear in  $\theta'$ , and  $\psi'$  (the  $\psi$  part of  $\theta'$ ) asserts about the set of things satisfying  $\varphi_i$  in

$$(L_\alpha[\dot{S}, \dot{x}]; \dot{S}, \dot{x}, \dot{\delta} (\delta \in d_i), \in)$$

what  $\psi$  says about  $x_i$ . This involves rewriting the atomic formulas of  $\psi$  which include  $x_i$ . Rewriting formulas of the form  $y = x_i$  or  $y \in x_i$  can be done in a straightforward fashion. The formula  $x_i \in y$  can be expressed by saying that there exists a set  $z$  containing everything satisfying the defining formula  $\varphi_i$ , and consisting only of things satisfying this formula, and that  $z$  is in  $y$ . The formula  $\psi$  may also need to be rewritten to deal with the predicates for  $S$  and  $x$ , resulting (if  $\alpha < \delta$ ) in formulas  $\psi_0$  and  $\psi_1$  to accommodate the two possible values of  $x(\alpha)$ . Letting  $\theta_0$  and  $\theta_1$  be the two rewritten forms of  $\theta$ , we can let  $\rho_{\theta, \alpha+1}$  be  $(P_\alpha \wedge \rho_{\theta_0, \alpha}) \vee ((\neg P_\alpha) \wedge \rho_{\theta_1, \alpha})$ . The rest of this step is the same as the limit case.

To complete the proof, fix an ordinal  $\gamma > \delta \cup \sup(S)$  such that for all  $x \in 2^\delta$ ,  $L_\gamma[S, x] \models \theta(S, x)$  if and only if  $L[S, x] \models \theta(S, x)$ , and note that there is an element of  $T_\gamma$  which, for each  $x \in 2^\delta$ , holds in the expanded version of  $L_\gamma[S, x]$  if and only if  $L[S, x] \models \theta(S, x)$ .

That (3) implies (4) is immediate. The implication from (4) to (2) follows from the absoluteness of the existence of winning strategies in closed games (via a ranking function, as in [3]). To get from (1) to (3), consider a witness  $\varphi$  to (1) as a tree. Players  $I$  and  $II$  choose a path from the root of the tree to a terminal node, with  $I$  choosing at disjunctions,  $II$  choosing at conjunctions, and  $I$  winning if and only if  $x(\gamma) = 1$ , for  $P_\gamma$  the terminal node reached by the run of the game. If  $x \models \varphi$ , then  $I$  can play to maintain that  $x$  satisfies each sub-sentence visited during the run of the game, and conversely for  $II$ .  $\square$

**8.1.8 Remark.** The construction of the  $\mathcal{L}_{\infty, \delta}^*$ -sentence in the implication from (2) to (1) in the proof of Theorem 8.1.7 is carried out in  $L[S]$ , aside from the choice of  $\gamma$ . The construction is also sufficiently uniform to produce, for any set of witnesses to (2) a corresponding collection of witnesses to (1).

**8.1.9 Remark.** Part (2) of Theorem 8.1.7 makes sense for subsets of spaces of the form  $\omega^\delta$  and their products (or sets of subsets of any set in  $L$ ), so we will say that a set  $A \subseteq (\omega^\delta)^n$  (for some  $n \in \omega$  and some ordinal  $\delta$ ) is  $\infty$ -Borel if there exists a set  $S$  of ordinals such that  $A = \{x \in (\omega^\delta)^n : L[S, x] \models \varphi(S, x)\}$ , for some binary formula  $\varphi$ . One could also revise the infinitary language  $\mathcal{L}_{\infty, \delta}^0$  above to describe (for instance) subsets of  $2^{\delta_1 \times \dots \times \delta_n}$ , starting with variables  $P_s$  ( $s \in \delta_1 \times \dots \times \delta_n$ ), interpreted so that  $x \models P_s$  if and only if  $x(s) = 1$ . Under this formulation everything goes through as above (trivially, as we are merely relabeling the set of predicates). One can then identify elements of  $\omega^\omega$ , for instance, with elements of  $2^{\omega \times \omega}$ . However, this identification is not surjective, so, for instance,

a  $\mathcal{L}_{\omega \times \omega}^0$ -sentence defining a subset of  $\omega^\omega$  should include conjuncts expressing  $\bigvee_{m \in \omega} P_{(n,m)}$  for each  $n \in \omega$ .

**8.1.10 Definition.** Given a set of ordinals  $S$  and a binary formula  $\varphi$  from the first order language of set theory, we say that the pair  $(S, \varphi)$  is an  $\infty$ -Borel\* code for the set

$$\{x \in (\omega^\delta)^n : L[S, x] \models \varphi(S, x)\}.$$

If in addition  $S$  is a subset of an ordinal  $\gamma$ , we say that the pair  $(S, \varphi)$  is a  $\gamma$ -Borel\* code for the set

$$\{x \in (\omega^\delta)^n : L_\gamma[S, x] \models \varphi(S, x)\}.$$

**8.1.11 Remark.** For any infinite cardinal  $\gamma$ , and any infinite ordinal  $\delta$ , every  $\gamma$ -Borel subset of  $2^\delta$  has a  $\gamma$ -Borel\* code, using the coding of sentences in  $\mathcal{L}_{\gamma, \delta}^0$  by sets of ordinals introduced at the beginning of this section. The proof of (1) from (2) in Theorem 8.1.7 contains a reflection argument, so it does not immediately give a bound in the reverse direction. However, if  $S$  is a set of ordinals, then for all  $x, y \in \omega^\omega$ ,  $y \in L[S, x]$  if and only if  $y \in L_\gamma[S, x]$ , where  $\gamma$  is the least uncountable cardinal (of  $L(S, \mathbb{R})$ ) greater than the supremum of  $S$ . It follows that  $\{(x, y) : y \in L[S, x]\}$  and its complement are  $\gamma$ -Borel. We will use this observation in Chapter 11.

**8.1.12 Remark.** If  $\delta < \Theta$  and  $A \subseteq 2^\delta$  is  $\infty$ -Borel, then this is witnessed by an element of  $\mathcal{L}_{\gamma, \delta}^0$  for some  $\gamma < \Theta$ . This can be shown by taking a continuum-sized elementary submodel (containing  $\omega^\omega$ ) of a suitable model of the form  $L_\alpha(S, \mathbb{R})$ , where  $S$  is a set of ordinals such that  $\varphi_S$  is an  $\infty$ -Borel code for  $A$ . Since every element of  $L_\alpha(S, \mathbb{R})$  is definable in  $L_\alpha(S, \mathbb{R})$  from  $S$ , an ordinal and an element of  $\omega^\omega$ , such a submodel can be built assuming only ZF.

If  $A \subseteq \omega^\omega$  is  $\infty$ -Borel, and  $\aleph_1 \not\leq 2^{\aleph_0}$  holds, then  $A$  has the property of Baire and is Lebesgue measurable. To see this for the Baire property, let  $\mathbb{P}$  be Cohen forcing and let  $\dot{c}$  be the canonical  $\mathbb{P}$ -name for the generic Cohen real. Suppose that  $(S, \varphi)$  is an  $\infty$ -Borel\* code for  $A$ . Since  $\aleph_1 \not\leq 2^{\aleph_0}$  and  $L[S] \models \text{AC}$ ,  $\omega^\omega \cap L[S]$  is countable, so the set of reals which are Cohen-generic over  $L[S]$  is comeager. For each such real  $x$ , let  $g_x$  be the filter for which  $\dot{c}_{g_x} = x$ . Let  $A'$  be the set of  $x$  which are Cohen-generic over  $L[S]$  for which there is a condition  $p \in g_x$  forcing that  $L[S, \dot{c}] \models \varphi(S, \dot{c})$ . Then  $A'$  is Borel, and the symmetric difference  $A \triangle A'$  is contained in the set of reals which are not Cohen-generic over  $L[S]$ . For Lebesgue measurability the argument is the same, using random forcing instead of Cohen forcing. One can use the same argument with Mathias forcing to show that  $\infty$ -Borel subsets of  $\mathcal{P}(\omega)$  satisfies the Ramsey property. Unlike the cases of the Baire property and Lebesgue measurability, it is still unknown whether AD alone implies that every subset of  $\mathcal{P}(\omega)$  has the Ramsey property.

As shown in Remark 8.1.3 and Theorem 8.1.7, Suslin subsets of  $2^\delta$  are  $\infty$ -Borel, for any ordinal  $\delta$ . Theorem 6.2.1 says that Suslin sets can be uniformized. The proof of that theorem shows that if  $\gamma$  is an ordinal and  $T$  is a tree on  $\delta \times \gamma$  such that  $p[T]$  is nonempty, then  $p[T]$  has a member in  $L[T]$ . However, it is

possible to have a  $\varphi \in \mathcal{L}_\infty^0$  such that  $A_\varphi$  is nonempty in  $V$  but empty in  $L[\varphi]$ . For instance, there is a  $\varphi$  in  $L$  such that  $A_\varphi$  is the set of Cohen reals over  $L$ . Another way to see this is to note that consistently there are  $\infty$ -Borel subsets of  $(\omega^\omega)^2$  which are not uniformized (for instance, in  $L(\mathbb{R})$ , assuming AD, by Corollary 8.2.5 and Theorem 6.2.5) whereas if  $S$  were an  $\infty$ -Borel code for a set  $A \subseteq (\omega^\omega)^2$  and  $L[S, x] \cap A$  were nonempty for each  $x$ , then a uniformizing function could be found by picking the constructibly least member of each set  $L[S, x] \cap A$  relative to  $S$  and  $x$ .

We will make use of the following lemma in Section 11.4. Recall that the weakly  $\kappa$ -Borel sets were defined near the end of Section 6.1.

**Theorem 8.1.13** (ZF+AD+DC $_{\mathbb{R}}$ ). *Suppose that  $\kappa$  is a limit of Suslin cardinals and that  $\kappa$  has uncountable cofinality. Then the following are equivalent for each  $A \subseteq \omega^\omega$ .*

- $A$  is weakly  $<\kappa$ -Borel.
- $A$  is  $<\kappa$ -Borel.
- $A$  is  $<\kappa$ -Suslin.

*Proof.* For any infinite cardinal  $\lambda$ ,  $\lambda$ -Borel sets are weakly  $\lambda$ -Borel, and, by either Remark 8.1.3 or Theorem 8.1.7,  $\lambda$ -Suslin sets are  $\lambda^+$ -Borel. Theorem 6.1.21 says that, under AD+DC $_{\mathbb{R}}$ , if  $\kappa$  is a limit of Suslin cardinals of uncountable cofinality, then all weakly  $<\kappa$ -Borel sets are  $<\kappa$ -Suslin.  $\square$

In conjunction with Remark 2.5.9 (on Solovay's diagonalization method for building sets of large Wadge rank), Theorems 6.1.18 and 6.1.21 give the following.

**Theorem 8.1.14** (ZF + AD + DC $_{\mathbb{R}}$ ). *If  $\kappa$  is a limit of Suslin cardinals of uncountable cofinality, then  $o(\mathcal{S}_{<\kappa}) = \delta(\mathcal{S}_{<\kappa}) = \kappa$ .*

*Proof.* That  $\delta(\mathcal{S}_{<\kappa}) = \kappa$  follows from Theorem 6.1.18. Since  $\mathcal{S}_{<\kappa}$  is a projective algebra,  $o(\mathcal{S}_{<\kappa}) \leq \delta(\mathcal{S}_{<\kappa})$ , by Proposition 2.5.8. Since in addition (by Theorem 8.1.13) it is closed under unions and intersections of length less than  $\kappa$ , Remark 2.5.9 gives that  $o(\mathcal{S}_{<\kappa}) \geq \kappa$ .  $\square$

**8.1.15 Remark.** Woodin has shown that, assuming AD+DC $_{\mathbb{R}}$ , for each ordinal  $\gamma < \Theta$ , all sets of reals in  $L(\mathcal{P}(\gamma))$  are  $\infty$ -Borel. While we do not use or prove this result in this book, we note the following consequence. Suppose that AD+DC $_{\mathbb{R}}$  holds, and that  $\lambda$  is a strong partition cardinal above the Wadge rank of some set which is not  $\infty$ -Borel. Then, in  $L(\mathcal{P}(\lambda))$ , AD+DC $_{\mathbb{R}}$  holds and  $\lambda$  is a strong partition cardinal greater than  $\Theta$ . It is currently unknown whether AD+DC $_{\mathbb{R}}$  is consistent with the existence of a strong partition cardinal greater than or equal to  $\Theta$ .

Part (1) of Corollary 10.3.7 shows that, assuming AD+DC $_{\mathbb{R}}$ , every set of reals in  $L(Y, \mathbb{R})$  is  $\infty$ -Borel, whenever  $Y$  is a set of ordinals.

## 8.2 Local $\infty$ -Borel codes

Our goal in this section is to show that, assuming Lipschitz Determinacy, if  $A \subseteq \omega^\omega$  is  $\infty$ -Borel, then  $A$  has an  $\infty$ -Borel code in  $L(A, \mathbb{R})$ . Given  $A \subseteq \omega^\omega$ , let  $\mathcal{P}_A$  be the set of prewellorderings  $\omega^\omega$  Wadge-reducible to either  $A$  or its complement (i.e., those prewellorderings  $P$  such that either  $P \leq_{\text{Wa}} A$  or  $P \leq_{\text{Wa}} \omega^\omega \setminus A$ ) and let  $\delta_A$  denote the supremum of the lengths of the members of  $\mathcal{P}_A$ .

**Lemma 8.2.1.** *For any  $A \subseteq \omega^\omega$ ,  $\delta_A < \Theta^{L(A, \mathbb{R})}$ , and no member of  $\mathcal{P}_A$  has length  $\delta_A$ .*

*Proof.* For the first part, one can naturally form a wellfounded relation isomorphic to the disjoint union of the members of  $\mathcal{P}_A$ . For instance, recalling from Remark 2.5.2 that  $\mathcal{F}^{c,2,1}$  denotes the set of  $x \in \omega^\omega$  such that  $f_x^{c,2,1}$  is a continuous function from  $(\omega^\omega)^2$  to  $\omega^\omega$  (relative to a fixed universal closed subset of  $(\omega^\omega)^4$ ), let  $\leq_A$  be the relation on (a subset of)  $(\omega^\omega)^2$  defined by setting  $(x_0, y_0) \leq_A (x_1, y_1)$  to hold if  $x_0 = x_1 \in \mathcal{F}^{c,2,1}$ ,  $(f_{x_0}^{c,2,1})^{-1}[A]$  is a prewellordering and

$$f_{x_0}^{c,2,1}(y_0, y_1) \in [A].$$

Then  $\leq_A$  is a wellfounded preorder projective in  $A$ , and the rank of  $\leq_A$  is  $\delta_A$ . The map sending each  $(x, y)$  in the domain of  $\leq_A$  to its  $\leq_A$ -rank shows that  $\delta_A$  is less than  $\Theta^{L(A, \mathbb{R})}$ .

For the second part, each prewellordering in  $\mathcal{P}_A$  is isomorphic to one whose domain is not all of  $\omega^\omega$ , and such a prewellordering can be extended by one point to another one in  $\mathcal{P}_A$ .  $\square$

In this section we show (assuming  $\text{AD} + \text{DC}_{\mathbb{R}}$ ) that if  $A$  is  $\infty$ -Borel then it is  $\delta_A$ -Borel.

We define the *rank*  $\text{rk}(\varphi)$  of a sentence in  $\mathcal{L}_{\infty, \delta}^0$  (for some ordinal  $\delta$ ) by recursively setting

- $\text{rk}(P_\alpha)$  to be 0 for each predicate  $P_\alpha$ ;
- $\text{rk}(\neg\varphi)$  to be  $\text{rk}(\varphi)$  for all  $\varphi \in \mathcal{L}_{\infty, \delta}^0$ ;
- $\text{rk}(\bigvee_{\alpha \in Y} \varphi_\alpha)$  and  $\text{rk}(\bigwedge_{\alpha \in Y} \varphi_\alpha)$  to be  $\sup\{\text{rk}(\varphi_\alpha) + 1 : \alpha \in Y\}$  for all sets  $Y \subseteq \text{Ord}$  and all formulas  $\varphi_\alpha$  ( $\alpha \in Y$ ) in  $\mathcal{L}_{\infty, \delta}^0$ ,

We now define a procedure that removes redundant terms from an  $\infty$ -Borel code. For each  $\varphi \in \mathcal{L}_{\infty, \delta}^0$  we define the formula  $\varphi^* \in \mathcal{L}_{\infty, \delta}^0$  recursively as follows:

- $P_\alpha^*$  is  $P_\alpha$ , for each predicate  $P_\alpha$ ;
- $(\neg\varphi)^*$  is  $\neg\varphi^*$ , for all  $\varphi \in \mathcal{L}_{\infty, \delta}^0$ ;
- for all sets  $Y \subseteq \text{Ord}$ , and all  $\mathcal{L}_{\infty, \delta}^0$ -sentences  $\varphi_\alpha$  ( $\alpha \in Y$ ),



- $(\bigvee_{\alpha \in Y} \varphi_\alpha)^*$  is  $\bigvee_{\alpha < \gamma} \varphi_{\pi(\alpha)}^*$ , where  $\gamma$  is the ordertype of the set

$$X = \{\alpha \in Y : A_{\varphi_\alpha} \not\subseteq \bigcup_{\beta \in \alpha \cap Y} A_{\varphi_\beta}\}$$

and  $\pi: \gamma \rightarrow X$  is an order-preserving bijection (i.e., the inverse of the Mostowski collapse of  $X$ );

- $(\bigwedge_{\alpha \in Y} \varphi_\alpha)^*$  is  $\bigwedge_{\alpha < \gamma} \varphi_{\pi(\alpha)}^*$ , where  $\gamma$  is the ordertype of the set

$$X = \{\alpha \in Y : A_{\varphi_\alpha} \not\supseteq \bigcap_{\beta \in \alpha \cap Y} A_{\varphi_\beta}\}$$

and  $\pi: \gamma \rightarrow X$  is an order-preserving bijection.

Then for all  $\varphi \in \mathcal{L}_{\infty, \delta}^0$ ,  $A_\varphi = A_{\varphi^*}$ ,  $\text{rk}(\varphi) \geq \text{rk}(\varphi^*)$  and  $\varphi^{**} = \varphi^*$ . Note that the operation  $\varphi \mapsto \varphi^*$  merely removes from each conjunction and disjunction in  $\varphi$  those terms which have no effect on the corresponding intersection or union, and reindexes the conjuncts or disjuncts.

We define the following ordinals, all of which are at most  $\Theta$ .

- $\chi_B$ , the least  $\chi$  such that all  $\infty$ -Borel subsets of  $2^\omega$  are  $\chi$ -Borel;
- $\kappa_B$ , the supremum of the lengths of wellordered sequences of  $\infty$ -Borel codes for nonempty disjoint subsets of  $2^\omega$ ;
- $\lambda_B$ , the supremum of the lengths of the  $\infty$ -Borel prewellorderings of  $2^\omega$ ;
- $\rho_B$ , the supremum of the Wadge ranks of  $\infty$ -Borel subsets of  $2^\omega$ .

The existence of analytic (and therefore  $\infty$ -Borel) sets which are not Borel shows that  $\chi_B \geq \aleph_1$ . Corollary 10.3.8 shows that the four ordinals have the same value, assuming  $\text{AD} + \text{DC}_{\mathbb{R}}$ .

We start with a lemma relating  $\chi_B$ ,  $\kappa_B$  and  $\lambda_B$ , and return to these ordinals at the end of the section.

**Lemma 8.2.2.**  $\chi_B \leq \kappa_B \leq \lambda_B$

*Proof.* To see that  $\chi_B \leq \kappa_B$ , let  $\varphi$  be any sentence in  $\mathcal{L}_\infty^0$ . For any disjunction  $\bigvee_{\alpha < \gamma} \psi_\alpha$  in  $\varphi^*$  the sentences  $\psi_\alpha \wedge \neg(\bigvee_{\beta < \alpha} \psi_\beta)$  are  $\infty$ -Borel codes for nonempty disjoint sets. A similar remark applies to the conjunctions. It follows that  $\varphi^*$  is in  $\mathcal{L}_{\kappa_B}^0$ .

To see that  $\kappa_B \leq \lambda_B$ , note that one can convert a sequence of  $\infty$ -Borel codes for nonempty disjoint sets to an  $\infty$ -Borel code for a prewellordering in a uniform fashion by means of uniformly coding the product. Consider  $\langle \varphi_\alpha : \alpha < \gamma \rangle$  with the sets  $A_{\varphi_\alpha}$  disjoint and nonempty. The set

$$\bigcup_{\alpha \leq \beta < \gamma} A_{\varphi_\alpha} \times A_{\varphi_\beta}$$

is a prewellordering of length  $\gamma$ . If each  $\varphi_\alpha$  is in  $\mathcal{L}_\xi^0$  (for some ordinal  $\xi$ ), then the resulting code is in  $\mathcal{L}_{\xi \cup \gamma}^0$ .  $\square$

The main theorem in this section is the following.

**Theorem 8.2.3** (ZF + Lipschitz Determinacy). *If  $A \subseteq 2^\omega$  is  $\infty$ -Borel then  $A$  is  $\delta_A$ -Borel.*

*Proof.* If  $\delta_A \geq \chi_B$  we are done. Supposing otherwise, using a sentence  $\varphi$  in  $\mathcal{L}_\infty^0$  of minimal rank such that  $\varphi^* \notin \mathcal{L}_{\delta_A}^0$ , we may find a sequence  $\langle \varphi_\alpha : \alpha \leq \delta_A \rangle$ , consisting of elements of  $\mathcal{L}_{\delta_A}^0$ , such that the corresponding sets  $A_{\varphi_\alpha}$  are nonempty and pairwise disjoint. Then the prewellordering (call it  $\leq$ ) built as in the second half of the proof of Lemma 8.2.2 is  $\delta_A$ -Borel, and, as it has length  $\delta_A$ , it is not in  $\mathcal{P}_A$ . By Lipschitz Determinacy, then,  $A$  is Wadge below  $\leq$ , which means that  $A$  is  $\delta_A$ -Borel.  $\square$

Suppose now that Lipschitz Determinacy +  $\text{DC}_\mathbb{R}$  holds, and let  $A \subseteq 2^\omega$  be  $\infty$ -Borel. By Proposition 2.5.8, there is a prewellordering of  $\mathcal{F}^c$  of length  $\delta_A$  which is projective in  $A$ . By the Moschovakis Coding Lemma, every subset of  $\delta_A$  is coded (using this prewellordering) by a set of reals projective in  $A$ . Putting this all together with Theorem 8.2.3 gives the following corollaries.

**Corollary 8.2.4** (ZF + Lipschitz Determinacy +  $\text{DC}_\mathbb{R}$ ). *If  $A \subseteq 2^\omega$  is  $\infty$ -Borel then  $A = A_\varphi$  for some  $\varphi \in \mathcal{L}_{\delta_A}^0 \cap L(A, \mathbb{R})$ .*

Corollary 10.3.7 below is a stronger version of the following corollary, modulo the additional assumption of  $\text{DC}_\mathbb{R}$ .

**Corollary 8.2.5** (ZF + AD). *In  $L(\mathbb{R})$ , every subset of  $2^\omega$  is  $\infty$ -Borel.*

*Proof.* We work in  $L(\mathbb{R})$ , noting that AD implies that  $\text{DC}_\mathbb{R}$  holds there by Theorem 0.4.1. By the Moschovakis Coding Lemma, the assertion that there is a subset of  $2^\omega$  which is not  $\infty$ -Borel is  $\Sigma_1^2$ . By the Solovay Basis Theorem (Theorem 5.0.6), this assertion, if true, is witnessed by a  $\Delta_1^2$  set. By Theorem 6.1.22, in  $L(\mathbb{R})$ , all  $\Delta_1^2$  sets are Suslin, and therefore  $\infty$ -Borel.  $\square$

Combining Theorem 0.4.1 (for  $\text{DC}_\mathbb{R}$ ) and Remark 7.0.4 (for  $<\Theta$ -Determinacy) with Corollary 8.2.5, we get that the implication from AD to  $\text{AD}^+$  holds in  $L(\mathbb{R})$ .

**Corollary 8.2.6** (ZF + AD).  $L(\mathbb{R}) \models \text{AD}^+$ .

A similar argument shows that  $\text{AD}^+$  follows from the conjunction of AD with  $\Sigma_1^2$ -reflection into the Suslin, co-Suslin sets. Woodin has shown that the converse holds.

**Corollary 8.2.7** (ZF + AD). *If every true  $\Sigma_1^2$  sentence is witnessed by a set which is Suslin and co-Suslin, then  $\text{AD}^+$  holds.*

*Proof.* No counterexample to any of the three parts of  $\text{AD}^+$  can be witnessed by a set which is Suslin and co-Suslin. For  $\text{DC}_\mathbb{R}$  this follows from the fact that Suslin sets can be uniformized, and for  $<\Theta$ -Determinacy it follows from Theorem 7.0.1. For the  $\infty$ -Borel property it follows from each of Remark 8.1.3 and Theorem 8.1.7. The assertion that any of the three parts of  $\text{AD}^+$  fails is a  $\Sigma_1^2$  sentence.

For  $\text{DC}_{\mathbb{R}}$  this is immediate. For  $<\Theta$ -Determinacy this follows from the Coding Lemma. For the  $\infty$ -Borel property this follows from the Coding Lemma and Theorem 8.2.3.  $\square$

Remark 7.0.4 and Theorem 8.2.3 show that  $\text{AD}^+$  reflects down to inner models containing the reals.

**Theorem 8.2.8** ( $\text{ZF} + \text{AD}^+$ ). *If  $M$  is an inner model of  $\text{ZF}$  containing  $\omega^\omega$ , then  $M \models \text{AD}^+$ .*

*Proof.* Since  $\omega^\omega \subseteq M$ ,  $M \models \text{AD} + \text{DC}_{\mathbb{R}}$ . By Remark 7.0.4,  $M$  satisfies  $<\Theta$ -Determinacy. By Theorem 8.2.3 (noting that the ordinals  $\delta_A$  have the same value whether computed in  $M$  or  $V$ ), every subset of  $\omega^\omega$  in  $M$  is  $\infty$ -Borel in  $M$ .  $\square$

The rest of this section concerns the four ordinals defined before Lemma 8.2.2. Since the collection of  $\infty$ -Borel subsets of  $\omega^\omega$  is closed under continuous preimages, if Lipschitz Determinacy +  $\text{DC}_{\mathbb{R}}$  holds then  $\rho_B = \Theta$  if and only if all subsets of  $\omega^\omega$  are  $\infty$ -Borel. The same is true for  $\lambda_B$  (and, as well shall see,  $\chi_B$  and  $\kappa_B$ ).

**Theorem 8.2.9** ( $\text{ZF} + \text{Lipschitz Determinacy}$ ). *The equation  $\lambda_B = \Theta$  holds if and only if every subset of  $\omega^\omega$  is  $\infty$ -Borel.*

*Proof.* By definition,  $\Theta$  is the supremum of the lengths of the prewellorderings of  $\omega^\omega$ , so if every subset of  $\omega^\omega$  is  $\infty$ -Borel then  $\lambda_B = \Theta$ . For the other direction, suppose that  $A \subseteq \omega^\omega$  is not  $\infty$ -Borel and that  $P$  is an  $\infty$ -Borel prewellordering. Then  $A \not\leq_W P$  as the  $\infty$ -Borel sets form an initial segment of the Wadge hierarchy, so  $P \in \mathcal{P}_A$ . By Lemma 8.2.1, the length of  $P$  is less than  $\delta_A$ , which is less than  $\Theta$ .  $\square$

**Theorem 8.2.10** ( $\text{ZF} + \text{AD} + \text{DC}_{\mathbb{R}}$ ). *The following statements are equivalent.*

1. *All subsets of  $\omega^\omega$  are  $\infty$ -Borel.*
2. *At least one of  $\chi_B$ ,  $\kappa_B$ ,  $\lambda_B$  and  $\rho_B$  is  $\Theta$ .*
3.  *$\chi_B = \kappa_B = \lambda_B = \rho_B = \Theta$*

*Proof.* Statement (2) follows immediately from statement (3). Theorem 8.2.9 implies that (1) implies (2) and (3) implies (1). By Lemma 8.2.2 and Theorem 8.2.9, and the remark on  $\rho_B$  before the statement of Theorem 8.2.9, (2) implies (1).

By Lemma 8.2.2, all that remains to be shown is that if all subsets of  $\omega^\omega$  are  $\infty$ -Borel then  $\chi_B = \Theta$ . This follows from the Moschovakis Coding Lemma, which implies that for each  $\chi < \Theta$  there is a surjection from  $\omega^\omega$  to  $\mathcal{P}(\chi)$ , and therefore (if  $\chi$  is a cardinal, which  $\chi_B$  is) one from  $\omega^\omega$  to the  $\chi$ -Borel sets.  $\square$



## Chapter 9

# Cone measure ultrapowers

In this chapter we analyze ultraproducts by the cone measures introduced in Section 1.2. The results of this analysis will be used throughout the rest of the book.

Given an equivalence relation  $E$  on a set  $X$ , a function  $f$  with domain  $X$  is said to be  $E$ -invariant if  $f(x) = f(y)$  whenever  $xEy$ . Given an ordered equivalence relation  $(E, \leq_E)$  (on some set  $X$ ) such that the cone measure  $\mu_E$  is an ultrafilter on the set  $\mathcal{C}_E$  consisting of the  $E$ -equivalence classes, we let  $\text{WF}_E$  denote the assertion that the ultrapower  $\text{Ord}^{\mathcal{C}_E}/\mu_E$  (formed by taking all  $E$ -invariant functions on  $X$ , modulo  $\mu_E$ ) is wellfounded. Note that in this situation  $\text{Ord}^{\mathcal{C}_E}/\mu_E$  is wellfounded if and only if  $V^{\mathcal{C}_E}/\mu_E$  is. Given a function  $f$  with domain  $\mathcal{C}_E$ , or an  $E$ -invariant function  $f$  with domain  $X$ , we write  $[f]_{\mu_E}$  for the element of  $V^{\mathcal{C}_E}/\mu_E$  represented by  $f$ . Given two ordered equivalence relations  $(E, \leq_E)$  and  $(F, \leq_F)$  on the same set  $X$ , if  $\leq_F \subseteq \leq_E$  then  $\text{Ord}^{\mathcal{C}_E}/\mu_E$  embeds into  $\text{Ord}^{\mathcal{C}_F}/\mu_F$ , so  $\text{WF}_F$  implies  $\text{WF}_E$ .

The wellfoundedness of the ultrapower  $\text{Ord}^{\mathcal{C}_E}/\mu_E$  follows from DC plus the assumption that  $\mu_E$  is a countably complete ultrafilter on  $\mathcal{C}_E$ . Recall that DC holds in models of the form  $L(A, \mathbb{R})$  satisfying  $\text{DC}_{\mathbb{R}}$ , if  $A$  is a set contained in  $L(\mathbb{R})$ .

Woodin has shown that  $\text{AD}^+$  implies  $\text{WF}_{\text{Tu}}$ , where Tu is Turing equivalence. We will not prove or use this result in this book, however, and instead will assume statements of the form  $\text{WF}_E$  when they are needed.

**9.0.1 Remark.** By Remark 1.2.7, if  $(E, \leq_E)$  is an ordered equivalence relation such that all downward cones are countable and the corresponding cone measure  $\mu_E$  is a countably complete ultrafilter, then  $E$  induces a countably complete ultrafilter on  $\omega_1$ , which gives the statement  $\aleph_1 \not\leq 2^{\aleph_0}$ . Suppose that  $M_x$  ( $x \in \omega^\omega$ ) is an  $E$ -invariant association of elements of  $\omega^\omega$  to models of ZFC such that the corresponding  $\mu_E$ -ultrapower  $M$  is wellfounded. Since  $\aleph_1 \not\leq 2^{\aleph_0}$ ,  $\omega_1^V$  is strongly inaccessible in  $M$ . It follows then that for each ordinal  $\gamma < \omega_1^V$ ,  $V_\gamma^{M_x} = V_\gamma^M$  for an  $E$ -cone of  $x$ .

Theorem 9.0.2 below shows that  $\text{DC}_{\mathbb{R}}$  follows from certain instances of  $\text{WF}_E$

(and in fact one needs only that the ultrapower of  $\omega_1$  is wellfounded). We say that an ordered equivalence relation  $(E, \leq_E)$  (or the corresponding order  $\leq_E$ ) is *locally countable* if for all  $x \in \omega^\omega$ ,  $\bigcup \mathcal{D}_E(x)$  is countable.

**Theorem 9.0.2** (Solovay [36]). *Let  $(E, \leq_E)$  be a locally countable ordered equivalence relation on  $\omega^\omega$ . If the  $E$ -cone measure is an ultrafilter, and the image of  $\omega_1$  in the corresponding ultrapower is wellfounded, then  $\text{DC}_{\mathbb{R}}$  holds.*

*Proof.* Let  $T$  be a tree of finite sequences from  $\omega^\omega$  without an infinite path. We will find a ranking function for  $T$ . For each  $\bar{x} \in T$  and each  $z \in \omega^\omega$ , let  $f_{\bar{x}}([z]_E)$  be 0 if the members of  $\bar{x}$  are not all in  $\mathcal{D}_E(z)$ , and the rank of  $T \restriction \mathcal{D}_E(z)$  below  $\bar{x}$  otherwise (which exists, since, as  $\mathcal{D}_E(z)$  is countable,  $\text{DC}_{\mathcal{D}_E(z)}$  holds). Then if  $\bar{x}$  properly extends  $\bar{y}$  in  $T$ ,  $f_{\bar{x}}([z]_E) < f_{\bar{y}}([z]_E)$  for all  $[z]_E$  containing both  $\bar{x}$  and  $\bar{y}$ . The map  $\bar{x} \mapsto [f_{\bar{x}}]_{\mu_E}$  then ranks  $T$ .  $\square$

A similar argument gives Theorem 9.0.3 below, which is similar to Theorem 4.3.5 and will be used in Section 9.2. Recall that, given a set  $A \subseteq \omega^\omega$ ,  $\Delta_\omega(A)$  is the smallest projective algebra with  $A$  as an element, and we say that the members of  $\Delta_\omega(A)$  are *projective in  $A$* . We let  $\delta_\omega(A)$  be the supremum of the lengths of the prewellorderings in  $\Delta_\omega(A)$ .

**Theorem 9.0.3.** *Suppose that there exists a locally countable ordered equivalence relation  $(E, \leq_E)$  on  $\omega^\omega$  such that  $\mu_E$  is an ultrafilter and  $\text{WF}_E$  holds. Then for each  $A \subseteq \omega^\omega$ ,  $\delta_\omega(A)$  is the supremum of the ranks of the wellfounded preorders on  $\omega^\omega$  in  $\Delta_\omega(A)$ .*

*Proof.* It suffices to fix a wellfounded transitive relation  $R$  on  $\omega^\omega$  in  $\Delta_\omega(A)$  and show that its rank is less than  $\delta_\omega(A)$ . Given  $x \in \omega^\omega$ , define a function  $\rho_x: \mathcal{C}_E \rightarrow \omega_1$  by setting  $\rho_x(e)$  to be the rank of  $R_x \restriction \{x \in \omega^\omega : \exists y \in e \ x \leq_E y\}$ , where  $R_x$  is

$$\{(y, z) \in \omega^\omega \times \omega^\omega : x R y R z\}.$$

Define the relation  $\leq^*$  on  $\omega^\omega$  by setting  $x \leq^* y$  if and only if

$$\{e \in \mathcal{C}_E : \rho_x(e) \leq \rho_y(e)\} \in \mu_E.$$

Since  $\mu_E$  is an ultrafilter and  $\text{WF}_E$  holds,  $\leq^*$  is a prewellordering. Furthermore,  $\leq^*$  is projective in  $A$ . Since  $R$  is transitive, if  $x R y$ , then for all  $e \in \mathcal{C}_E$ ,  $\rho_y(e) \leq \rho_x(e)$ . It follows that the rank of  $R$  is less than or equal to the length of  $\leq^*$ .  $\square$

## 9.1 $S$ -cones

We are interested in a special class of ordered equivalence relations, for which we establish the following notation.

**9.1.1 Definition.** Let  $S$  be a set of ordinals.

- We write  $\leq_S$  for the binary relation on  $\omega^\omega$  defined by setting  $x \leq_S y$  to hold if and only if  $x \in L[S, y]$ .

- We write  $\equiv_S$  for the equivalence relation  $\leq_S \cap \geq_S$  on  $\omega^\omega$ .
- For each  $x \in \omega^\omega$ ,  $[x]_S$  denotes the  $\equiv_S$ -equivalence class of  $x$ .
- We write  $\mathcal{D}_S$  for the set of  $\equiv_S$ -classes.
- An  $S$ -cone is either a set of the form  $\{y \in \omega^\omega : x \in L[S, y]\}$  or a set of the form  $\{[y]_S : y \in \omega^\omega, x \in L[S, y]\}$ , for some  $x \in \omega^\omega$ .
- We write  $\mu_S$  for the set of subsets of  $\mathcal{D}_S$  which contain an  $S$ -cone.
- A function  $f$  on  $\omega^\omega$  is  $S$ -invariant if  $f(x) = f(y)$  whenever  $x \equiv_S y$ .
- Given an  $S$ -invariant function  $f$  with domain  $\omega^\omega$ ,  $[f]_{\mu_S}$  denotes the element of  $V^{\mathcal{D}_S}/\mu_S$  represented by  $f$ .

Recall from Corollary 1.2.6 that TD implies that  $\mu_S$  is a countably complete ultrafilter on  $\mathcal{D}_S$ , for each set  $S \subseteq \text{Ord}$ . We will be primarily interested in ultrapowers of the form  $V^{\mathcal{D}_S}/\mu_S$ , for  $S$  a set of ordinals. Elements of this ultrapower are represented by  $S$ -invariant functions on  $\omega^\omega$ . When  $f$  is such a function, we sometimes write  $\prod f(x)/\mu_S$  for  $[f]_{\mu_S}$ . We also write  $\prod V/\mu_S$  for  $V^{\mathcal{D}_S}/\mu_S$  and  $\prod \text{Ord}/\mu_S$  for  $\text{Ord}^{\mathcal{D}_S}/\mu_S$ . We start by showing that the function  $f(x) = \omega_1^{L[S, x]}$  represents  $\omega_1$  (this fact appears as Lemma 3.3 of [21]).

**Theorem 9.1.2** (ZF + AD). *If  $S$  is a set of ordinals then  $\prod \omega_1^{L[S, x]}/\mu_S = \omega_1$ .*

*Proof.* Let  $f$  be the function on  $\omega^\omega$  defined by setting  $f(x)$  to be  $\omega_1^{L[S, x]}$ . Since  $\mu_S$  is a countably complete ultrafilter (by Corollary 1.2.6), each  $\alpha < \omega_1$  is represented by the function with constant value  $\alpha$ . For each  $\alpha < \omega_1$ ,

$$\{[x]_S : x \in \omega^\omega, \omega_1^{L[S, x]} > \alpha\}$$

is in  $\mu_S$ , so  $[f]_{\mu_S} > \alpha$ .

Suppose now that  $g$  is an  $S$ -invariant function on  $\omega^\omega$  such that

$$\{[x]_S : x \in \omega^\omega, \omega_1^{L[S, x]} > g(x)\} \in \mu_S.$$

We want to see that there is an  $\alpha < \omega_1$  such that  $\{[x]_S : x \in \omega^\omega, g(x) = \alpha\} \in \mu_S$ . By the countable completeness of  $\mu_S$ , this follows from the existence of an  $\alpha < \omega_1$  such that  $\{[x]_S : x \in \omega^\omega, g(x) < \alpha\} \in \mu_S$ , which we will show.

Let WO be the set of wellorderings of  $\omega$ , and for each  $z \in \text{WO}$ , let  $|z|$  denote the ordinal ordertype of  $z$ . Let  $\mathcal{G}$  be the game in which player  $I$  produces  $x \in \omega^\omega$ ,  $II$  produces  $y \in \omega^\omega$  and  $z \subseteq \omega \times \omega$  (all bit-by-bit), and  $II$  wins if and only if  $x \leq_S y$ ,  $z \in \text{WO}$  and  $|z| > g(y)$ .

Let  $x_0 \in \omega^\omega$  be such that  $\omega_1^{L[S, x]} > g(x)$  for all  $x \in \omega^\omega$  with  $x_0 \in L[S, x]$ . Given a strategy  $\sigma$  for player  $I$ , let  $y \in \omega^\omega$  be such that  $x_0$  and  $\sigma$  are in  $L[S, y]$ , and let  $z \in \text{WO} \cap L[S, y]$  be such that  $|z| > g(y)$  (such a  $z$  exists since  $g(y) < \omega_1^{L[S, y]}$ ). If  $x$  is the response given by  $\sigma$  to  $y$  and  $z$  then  $x \in L[S, y]$  and

$g(y) < |z|$ , so  $II$  wins the corresponding run of the game. So  $I$  cannot have a winning strategy.

Then AD implies that there is a winning strategy  $\sigma$  for  $II$ . Denote by  $y_\sigma(x)$  and  $z_\sigma(x)$  the values  $y$  and  $z$  respectively given by  $\sigma$  in response to a play  $x$  for  $I$ . Then  $\{z_\sigma(x) : x \in \omega^\omega\}$  is an analytic subset of WO, so by  $\Sigma_1^1$ -boundedness (mentioned just before and generalized by Theorem 6.1.16) there is a countable ordinal  $\alpha$  greater than all the members of  $\{|z_\sigma(x)| : x \in \omega^\omega\}$ . Then for all  $x \in \omega^\omega$  with  $\sigma \in L[S, x]$ ,  $y_\sigma(x) \equiv_S x$ , so  $\alpha > |z_\sigma(x)| > g(y_\sigma(x)) = g(x)$ .  $\square$

Theorem 9.1.3 shows that if  $\mu_S$  is an ultrafilter then for an  $S$ -cone of  $y \in \omega^\omega$ ,  $L[S, y]$  satisfies GCH below  $\omega_1^V$ . This fact was first proved by Steel in [39] assuming AD and later by Woodin from the hypothesis below. It will be used in the proof of Theorem 11.2.1. The (standard) generalization of the theorem from CH from  $\diamond_{\omega_1}$  was observed by the author.

**Theorem 9.1.3** (ZF + CC $_{\mathbb{R}}$ ). *Let  $S$  be a set of ordinals such that  $\leq_S$  is locally countable and  $\mu_S$  is an ultrafilter on  $\mathcal{D}_S$ . Then there is an  $x \in \omega^\omega$  such that for all  $y \in \omega^\omega$  with  $x \leq_S y$ , and all  $\gamma < \omega_1^V$  which are infinite cardinals in  $L[S, y]$ ,*

$$L[S, y] \models \diamond_{\gamma+}.$$

*Proof.* By Remark 1.2.7,  $\mu_S$  being an ultrafilter implies that  $\omega_1^V$  is measurable, which in turn implies that  $\aleph_1 \not\leq 2^{\aleph_0}$ . Since  $\mu_S$  is an ultrafilter, there is an  $x_0 \in \omega^\omega$  such that either

- $L[S, y] \models \diamond_{\omega_1}$  for all  $y \in \omega^\omega$  with  $x_0 \in L[S, y]$  or
- $L[S, y] \models \neg \diamond_{\omega_1}$  for all  $y \in \omega^\omega$  with  $x_0 \in L[S, y]$ .

We show that the former holds, by finding one such  $y \geq_S x_0$ .

Let  $P$  be the set of ordered pairs from  $\omega^{<\omega}$ , and let  $\pi : P \rightarrow \omega$  be a recursive bijection. Let

$$p : \omega^\omega \times \omega^\omega \rightarrow \mathcal{P}(\omega)$$

be defined by setting  $p(x, y)$  to be  $\pi[\{(x \upharpoonright n, y \upharpoonright n) : n \in \omega\}]$ . Then  $p$  sends each pair from  $\omega^\omega$  to an infinite subset of  $\omega$ , and distinct pairs from  $\omega^\omega$  to sets having finite intersection.

Recall that the partial order  $\text{Col}^*(\omega_1, \omega^\omega)$  consists of injections from countable ordinals to  $\omega^\omega$ , ordered by extension, and adds a bijection  $g : \omega_1 \rightarrow \omega^\omega$ . Let  $\dot{X}$  be a  $\text{Col}^*(\omega_1, \omega^\omega)$ -name for the  $p$ -image of the set of  $(x, y) \in \omega^\omega \times \omega^\omega$  such that  $g^{-1}(x) < g^{-1}(y)$ . Let  $Q_{\dot{X}}$  be a  $\text{Col}^*(\omega_1, \omega^\omega)$ -name for the Jensen-Solovay almost disjoint coding forcing for the realization of  $\dot{X}$  (see Section 0.5 for more on both of these partial orders).

Suppose that  $M$  is a transitive model of ZFC, and that  $(g, z)$  is  $M$ -generic for  $(\text{Col}^*(\omega_1, \omega^\omega) * Q_{\dot{X}})^M$ . Then  $z$  is a subset of  $\omega$  such that, for each pair  $(x, y)$  from  $(\omega^\omega)^V$ ,  $z \cap p(x, y)$  is finite if and only if  $g^{-1}(x) < g^{-1}(y)$ . It follows that  $g$  is an element of  $M[z]$  (see Remark A.0.5 of [27] for a discussion of



why  $M[z]$  is a generic extension of  $M$ ). A standard genericity argument shows that  $M[g] \models \Diamond_{\omega_1}$ . Since  $Q_{\dot{X}_g}$  is c.c.c. and has cardinality  $\aleph_1$  in  $M[g]$ , another standard argument shows that  $\Diamond_{\omega_1}$  holds in  $M[g][z]$  (which is the same as  $M[z]$ ).

Since  $\omega_1^V$  is measurable, and since  $L[S, x_0] \models \text{AC}$ , the set

$$\mathcal{P}(\text{Col}^*(\omega_1, \omega^\omega) * Q_{\dot{X}})^{L[S, x_0]}$$

is countable, so there exists a pair  $(g, z)$  which is  $L[S, x_0]$ -generic for this partial order. The argument just given shows that  $L[S, x_0, z] \models \Diamond_{\omega_1}$ , so any real in  $L[S, x_0, z]$  coding the pair  $(x_0, z)$  works as our desired  $y$ .

Again using the measurability of  $\omega_1^V$ ,  $\mathcal{P}(\alpha) \cap L[S, x_0]$  is countable for each  $\alpha < \omega_1^V$  so there exist  $L[S, x_0]$ -generic filters for each corresponding partial order  $\text{Col}(\omega, \alpha)$  (see Section 0.5 again). Since these extensions must satisfy  $\Diamond_{\omega_1}$ , we have that  $L[S, x_0] \models \Diamond_{\gamma^+}$  for all  $\gamma < \omega_1^V$  which are infinite cardinals in  $L[S, x_0]$ . To see this, fix such a  $\gamma$ , and recall that  $\Diamond_{\gamma^+}$  is equivalent to the version in which the witnessing sequence is allowed  $\gamma$  many guesses at each  $\alpha < \gamma^+$  (see [23], for instance). Working in  $L[S, x_0]$ , suppose that  $\langle \sigma_\alpha : \alpha < \gamma^+ \rangle$  is such that each  $\sigma_\alpha$  is a  $\text{Col}(\omega, \gamma)$ -name for a subset of  $\alpha$ , and the realizations of the  $\sigma_\alpha$ 's are forced to be witness to  $\Diamond_{\omega_1}$ . For each  $\alpha < \gamma^+$ , let  $A_\alpha$  be the set of  $a \in \mathcal{P}(\alpha) \cap L[S, x_0]$  such that the statement  $\dot{a} = \sigma_\alpha$  is forced by some condition in  $\text{Col}(\omega, \gamma)$ . Since  $\text{Col}(\omega, \gamma)$  has cardinality  $\gamma$ , each  $A_\alpha$  has cardinality at most  $\gamma$ . Applying the definition of  $\Diamond_{\omega_1}$  to the elements of  $\mathcal{P}(\gamma^+) \cap L[S, x_0]$ , we see that  $\langle A_\alpha : \alpha < \gamma^+ \rangle$  witnesses the generalized form of  $\Diamond_{\gamma^+}$  in  $L[S, x_0]$ . The same considerations apply to each  $y \in \omega^\omega$  such that  $y \geq_S x_0$ .  $\square$

## 9.2 Measurable cardinals from cone measures

We give here an application of Theorem 9.0.3 demonstrating the existence of measurable cardinals. Our only use of the theorem is in the proof of Theorem 10.1.8, for which Remark 5.0.2 would also suffice.

**Theorem 9.2.1** (ZF + AD + DC $_{\mathbb{R}}$ ). *For each  $A \subseteq \omega^\omega$ ,  $\delta_\omega(A)$  is a limit of measurable cardinals.*

*Proof.* Since  $\delta_\omega(A) < \Theta$ , there is a set  $C \subseteq \omega^\omega$  of Wadge rank greater than  $\delta_\omega(A)$ . By the Moschovakis Coding Lemma, it suffices to establish the theorem in  $L(C, \mathbb{R})$ , where DC holds and therefore the ultrapower of the ordinals by the Turing measure is wellfounded, so Theorem 9.0.3 applies. Given  $B \subseteq \omega^\omega$ , let  $\rho_B$  denote the supremum of the ranks of the wellfounded relations in  $\Sigma_1^1(B)$ , and let  $\delta_B^1$  be the supremum of the ranks of the  $\Delta_1^1(B)$  prewellorderings. By Theorem 9.0.3,  $\delta_\omega(A) = \sup\{\rho_B : B \in \Delta_\omega(A)\}$ .

Let  $B \subseteq \omega^\omega$  be projective in  $A$ , let  $U \subseteq (\omega^\omega)^3$  be a complete  $\Sigma_1^1(B)$ -set, and let  $B^*$  be the set of  $x$  for which  $U_x$  is a wellfounded preorder. Define a prewellordering  $\leq^*$  on  $B^*$  by setting  $x \leq^* y$  if and only if  $\text{rk}(U_x) \leq \text{rk}(U_y)$ , where  $\text{rk}(U_x)$  denotes the rank of  $U_x$ .

**Claim 1.** *If  $Z \subseteq B^*$  is in  $\Sigma_1^1(B)$ , then  $\{\text{rk}(U_x) : x \in Z\}$  is bounded in  $\rho_B$ .*

*Proof.* The supremum of this set is the rank of the transitive wellfounded relation  $\{((x, y), (x, z)) : x \in Z, (x, y, z) \in U\}$ , which is in  $\Sigma_1^1(B)$ .  $\square$

We adapt Solovay's original proof of the measurability of  $\omega_1$  under AD to find a measure on  $\rho_B$ . For each  $X \subseteq \rho_B$  we consider the game  $\mathcal{G}_X$ , where player  $I$  builds  $x_i \in \omega^\omega$  for even  $i \in \omega$  and player  $II$  builds  $x_i \in \omega^\omega$  for odd  $i \in \omega$ . We use the way of doing this illustrated in the following diagram.

I	$x_0(0)$	$x_0(1), x_2(0)$	$x_0(2), x_2(1), x_4(0)$	$\dots$
II		$x_1(0)$	$x_1(1), x_3(0)$	$\dots$

The game  $\mathcal{G}_X$ .

Given  $x_i$  ( $i \in \omega$ ) produced by a run of this game, the winner is decided as follows.

- If there is an  $i \in \omega$  such that  $x_i \notin B^*$ , then  $I$  wins if the least such  $i$  is odd, and  $II$  wins if the least such  $i$  is even.
- If  $x_i \in B^*$  for all  $i \in \omega$ , and there is an  $i \in \omega$  such that  $\neg(x_i \leq^* x_{i+1})$ , then  $I$  wins if the least such  $i$  is even, and  $II$  wins if the least such  $i$  is odd.
- If  $x_i \in B^*$  and  $x_i \leq^* x_{i+1}$  holds for all  $i \in \omega$ , then  $I$  wins if and only if  $\sup\{\text{rk}(U_{x_i}) : i \in \omega\} \in X$ .

A winning strategy for player  $II$  in  $\mathcal{G}_X$  (for some  $X \subseteq \rho_B$ ) can be converted into a winning strategy for player  $I$  in  $\mathcal{G}_{\rho_B \setminus X}$ , playing any fixed member of  $B^*$  for  $x_0$ . It follows that, for each  $X \subseteq \rho_B$ , player  $I$  has a winning strategy in at least one of the games  $\mathcal{G}_X$  and  $\mathcal{G}_{\rho_B \setminus X}$ . Let  $W$  be the set of  $X \subseteq \rho_B$  for which  $I$  has a winning strategy.

Claim 2 shows that  $W$  is a countably complete ultrafilter. Its completeness, which by the claim is at least  $\delta_B^1$ , is a measurable cardinal. The theorem then follows from Claim 2 and the fact that  $\{\delta_B^1 : B \in \Delta_\omega(A)\}$  is cofinal in  $\delta_\omega(A)$  (by the definition of  $\delta_\omega(A)$ ).

**Claim 2.**  $W$  is closed under intersections of cardinal less than  $\delta_B^1$ .

We finish by proving the claim, using the Moschovakis Coding Lemma plus the boundedness fact established in Claim 1. Fix  $\gamma < \delta_B^1$  and a prewellordering  $\leq_R$  in  $\Delta_1^1(B)$  of a set  $R \subseteq \omega^\omega$  such that  $\leq_R$  has length  $\gamma$ . Let  $=_R$  be  $\leq_R \cap \geq_R$ , and let  $<_R$  be  $\leq_R \setminus =_R$ . For each  $a \in R$ , let  $\text{rk}_R(a)$  denote the rank of  $a$  in  $\leq_R$ . Suppose that  $\langle X_\alpha : \alpha < \gamma \rangle$  is a sequence of sets in  $W$ . It suffices to show that the intersection  $\bigcap \{X_\alpha : \alpha < \gamma\}$  is nonempty. Let  $Y$  be the set of pairs  $(a, b)$  such that  $a \in R$  and  $b$  is a winning strategy for player  $I$  in  $\mathcal{G}_{X_{\text{rk}_R(a)}}$  (more formally,  $b$  is a code for a strategy using some fixed bijection between  $\omega$  and  $\omega^{<\omega}$ ). By the

Coding Lemma, there is a  $\text{pos-}\Sigma_1^1(<_R)$  set  $Y^* \subseteq Y$  such that for each  $\alpha < \gamma$  there is an  $(a, b) \in Y^*$  with  $\text{rk}_R(a) = \alpha$ . Let  $Z = \{b : \exists a (a, b) \in Y^*\}$ . Then  $Z$  is a collection of (codes for) winning strategies and  $Z$  is in  $\Sigma_1^1(B)$ .

Let  $\Sigma$  be the set of finite sequences  $\langle y_i : i < j_* \rangle$  from  $B^*$  such that, whenever

- $\langle x_i : i < \omega \rangle$  is a run of  $\mathcal{G}_\emptyset$  (the payoff set is irrelevant here) where  $I$  plays according to a strategy from  $Z$ , and
- $x_{2j+1} = y_j$  for all  $j < j_*$ ,

$\text{rk}(U_{x_{2j}}) < \text{rk}(U_{x_{2j+1}})$  for all  $j < j_*$ . By Claim 1, each sequence in  $\Sigma$  has a proper extension in  $\Sigma$ . By  $\text{DC}_{\mathbb{R}}$ , there is a sequence  $\langle y_i : i \in \omega \rangle$  whose finite initial segments are all in  $\Sigma$ . Since  $\langle y_j : j < \omega \rangle$  forms a losing play for player  $II$  against any of the strategies from  $Z$ , it follows that  $\sup\{\text{rk}(U_{y_j}) : j \in \omega\}$  is in  $X_\alpha$  for all  $\alpha < \gamma$ , as desired.  $\square$

### 9.3 The degree order on sets of ordinals

This section introduces an ordering on sets of ordinals in terms of their induced notions of relative constructibility on the reals. This ordering is related to the notion of  $\infty$ -Borel code (introduced in Chapter 8) and connected to Uniformization (see Corollary 10.2.7). Theorem 9.3.8 gives a summary of consequences of this relation in the context of Turing Determinacy. Theorem 9.6.6 gives a consequence of the existence of a maximal set in this order, in the context of AD.

Given  $S, T \subseteq \text{Ord}$ , we write  $S \leq_{\mathcal{D}} T$  ( $\mathcal{D}$  for “degree”) to mean that

$$\{[x]_{\text{Tu}} : \omega^\omega \cap L[S, x] \subseteq L[T, x]\} \in \mu_{\text{Tu}}.$$

We write  $S \equiv_{\mathcal{D}} T$  for

$$(S \leq_{\mathcal{D}} T) \wedge (T \leq_{\mathcal{D}} S)$$

and  $S <_{\mathcal{D}} T$  for

$$(S \leq_{\mathcal{D}} T) \wedge \neg(T \leq_{\mathcal{D}} S).$$

**9.3.1 Remark.** A reflection argument shows that for any set  $S$  of ordinals there is a bounded  $T \subseteq \Theta$  with  $S \leq_{\mathcal{D}} T$ . Ultimately we will be concerned only with the restriction of  $\leq_{\mathcal{D}}$  to bounded subsets of  $\Theta$ .

Theorem 9.3.2 shows that  $\leq_{\mathcal{D}}$  is a total order when  $\mu_{\text{Tu}}$  is a countably complete ultrafilter. As with Theorem 9.1.3, Theorem 9.3.2 was first proved by Steel in [39] assuming AD and later by Woodin from  $\text{TD} + \text{CC}_{\mathbb{R}}$ . Recall that TD was later shown to imply  $\text{CC}_{\mathbb{R}}$  [33].

**Theorem 9.3.2 (ZF+TD).** *If  $S$  and  $T$  are sets of ordinals such that  $\neg(T \leq_{\mathcal{D}} S)$  then for a Turing cone of  $x \in \omega^\omega$ , for all  $\gamma < \omega_1^V$ ,  $\mathcal{P}(\gamma) \cap L[S, x]$  is a set of cardinality  $|\gamma|^{L[T, x]}$  in  $L[T, x]$ .*

To prove Theorem 9.3.2, we use a variation of Mathias forcing relative to a nonprincipal ultrafilter  $U$  on  $\omega$ . Our partial order is a slight simplification of the one used originally by Woodin, and is forcing-equivalent to it. Let  $\mathcal{S}$  be the set of nonempty finite subsets of  $\omega$  and let  $\mathcal{F}$  be the set of functions from  $\omega$  to  $U$ . We let  $\max(s)$  denote the largest element of a set  $s$  in  $\mathcal{S}$ . The domain of  $\mathbb{P}_U$  is  $\mathcal{S} \times \mathcal{F}$ . The order on  $\mathbb{P}_U$  is defined as follows :  $(s', F') \leq (s, F)$  if the following hold:

- $s' \cap (\max(s) + 1) = s$ ;
- for each  $k \in s' \setminus s$ ,  $k \in F(\max(s \cap k))$ ;
- for each  $i \in \omega$ ,  $F'(i) \subseteq F(i)$ .

The second condition above distinguishes  $\mathbb{P}_U$  from the usual Mathias forcing relative to an ultrafilter. This condition enables Lemmas 9.3.3 and 9.3.4 below. The corresponding facts hold for Mathias forcing, but only for Ramsey ultrafilters; the ultrafilters we use below are not Ramsey.

Note that if  $(s', F') \leq (s, F)$  and  $i$  is the least member of  $s' \setminus s$ , then  $(s', F') \leq (s \cup \{i\}, F) \leq (s, F)$ . That is,  $(s', F')$  can be reached from  $(s, F)$  by one-point extensions.

Let us say that a function  $F \in \mathcal{F}$  is *refining* if  $F(i) \subseteq F(j) \subseteq (\omega \setminus (j + 1))$  whenever  $j < i \in \omega$ . For densely many  $\mathbb{P}_U$ -conditions  $(s, F)$ ,  $F$  is refining. Given  $F \in \mathcal{F}$ , say that an infinite  $a \subseteq \omega$  is *F-fast* if, for all  $i \in a$ ,  $a \setminus (i + 1) \subseteq F(i)$ . Every infinite subset of an  $F$ -fast set is  $F$ -fast. Every infinite  $a \subseteq \omega$  (in the ground model or any outer model) generates a filter  $G_a \subseteq \mathbb{P}_U$  consisting of those  $(s, F) \in \mathbb{P}_U$  such that  $s$  is an initial segment of  $a$  and  $a \setminus (\max(s) + 1)$  is  $F$ -fast.

The following lemma gives a characterization of  $\mathbb{P}_U$ -generic sets.

**Lemma 9.3.3.** *Assume that  $\text{CC}_{\mathbb{R}}$  holds. Let  $M$  be a transitive model of ZF, and let  $U$  be a nonprincipal ultrafilter on  $\omega$  in  $M$ . Let  $a$  be an infinite subset of  $\omega$ . Then  $G_a$  is  $M$ -generic for  $\mathbb{P}_U$  if and only if, for each  $F \in \mathcal{F}$  in  $M$ , there is an  $i \in a$  such that  $a \setminus i$  is  $F$ -fast.*

*Proof.* The forward direction follows immediately from the fact that for every  $s \in \mathcal{S}$ , and any two functions  $F, F' \in \mathcal{F}$ ,  $(s, F)$  and  $(s, F')$  are compatible. For the reverse direction, fix  $a$  with the given property and let  $D$  be a dense open subset of  $\mathbb{P}_U$  in  $M$ . Working in  $M$ , we recursively define a ranking function  $\rho_D$  on  $\mathcal{S}$ , letting  $\rho_D(s)$  be

- 0 if there exists an  $F \in \mathcal{F}$  such that  $(s, F) \in D$ ;
- $\alpha$  if  $\neg(\rho_D(s) < \alpha)$  and  $\{i \in \omega : \rho_D(s \cup \{i\}) < \alpha\} \in U$ ;
- $\omega_1$  if  $\neg(\rho_D(s) < \omega_1)$ .

Observe that, for each  $s \in \mathcal{S}$ , if  $\{i \in \omega : \rho_D(s \cup \{i\}) < \omega_1\} \in U$ , then  $\rho_D(s)$  is at most

$$\sup\{\rho_D(s \cup \{i\}) : i \in \omega, \rho_D(s \cup \{i\}) < \omega_1\},$$

which is less than  $\omega_1$ . It follows that, by  $\text{CC}_{\mathbb{R}}$ , we can choose, for each  $s \in \mathcal{S}$  a function  $F_s: \omega \rightarrow U$  such that

- if  $\rho_D(s) = 0$ , then  $(s, F_s) \in D$ ;
- if  $\rho_D(s) < \omega_1$ , then for all  $i \in F_s(\max(s))$ ,  $i > \max(s)$  and

$$\rho_D(s \cup \{i\}) < \rho_D(s);$$

- if  $\rho_D(s) = \omega_1$ , then for all  $i \in F_s(\max(s))$ ,  $i > \max(s)$  and

$$\rho_D(s \cup \{i\}) = \omega_1.$$

Define  $F_*: \omega \rightarrow U$  by setting  $F_*(i)$  to be  $\bigcap \{F_s(i) : s \in \mathcal{S} \cap \mathcal{P}(i+1)\}$ .

We claim that  $\rho_D(s) < \omega_1$  for all  $s \in \mathcal{S}$ . To see this, suppose that  $s \in \mathcal{S}$  is such that  $\rho_D(s) = \omega_1$ . By the choice of  $F_s$ , and the definition of  $F_*$ , we have (passing from  $(s, F_*)$  to  $(s', F')$  by one-point extensions) that  $\rho_D(s') = \omega_1$  whenever  $(s', F') \in \mathbb{P}_U$  is such that  $(s', F') \leq (s, F_*)$ . This contradicts our assumption that  $D$  is dense.

Finally, fix an  $i \in a$  such that  $a \setminus i$  is  $F^*$ -fast. Then the sequence of values  $\langle \rho_D(a \cap (j+1)) : j \in a \setminus i \rangle$  must decrease until reaching 0. If  $j \in a \setminus i$  is such that  $\rho_D(a \cap (j+1)) = 0$ , then  $(a \cap (j+1), F_*)$  is in  $G_a \cap D$ .  $\square$

Lemma 9.3.3 implies that infinite subsets of  $\mathbb{P}_U$ -generic subsets of  $\omega$  are  $\mathbb{P}_U$ -generic.

**Lemma 9.3.4.** *Let  $M$  be a transitive model of ZF, let  $U$  be a nonprincipal ultrafilter on  $\omega$  in  $U$ , and suppose that  $a \in [\omega]^\omega$  is such that  $G_a$  is an  $M$ -generic filter for  $\mathbb{P}_U$ . Then for any infinite  $a' \subseteq a$ ,  $G_{a'}$  is an  $M$ -generic filter for  $\mathbb{P}_U$ .*

Recall that a subset of HF is *recursive* if it is  $\Delta_1$  over HF. Say that a function  $\pi: 2^{<\omega} \rightarrow [\omega]^{<\omega}$  is *diffuse* if, letting  $\pi_*(x)$  denote  $\bigcup \{\pi(t|i) : i < \omega\}$  for each  $x \in 2^\omega$ , for all distinct  $x_0, \dots, x_{n-1} \in 2^\omega$  and  $a_0, \dots, a_{n-1} \in \mathcal{P}(\omega)$ , if  $a_i \in \{\pi_*(x_i), \omega \setminus \pi_*(x_i)\}$  for each  $i < n$  then  $\bigcap_{i < n} a_i$  is infinite. We will use a fixed recursive diffuse function below. One such function can be built by assigning disjoint infinite subsets  $a_{\bar{s}, f}$  of  $\omega$  to each pair  $(\bar{s}, f)$  such that, for some  $n, k \in \omega$ ,  $\bar{s} = \langle s_i : i < n \rangle$  is an  $n$ -tuple of distinct functions from  $k$  to 2 and  $f$  is a function from  $n$  to 2. Then for each  $t \in 2^{<\omega}$ , let  $\pi(t)$  be the set of  $\ell$  such that there exists such a pair  $(\bar{s}, f)$  with  $\ell \in a_{\bar{s}, f}$ , and some  $s_i$  with  $f(i) = 1$  is an initial segment of  $t$ . Then, given  $x_0, \dots, x_{n-1}$  and  $a_0, \dots, a_{n-1}$  as above, let  $k$  be large enough so that the sequence  $\bar{s} = \langle x_0|k, \dots, x_{n-1}|k \rangle$  is injective, and let  $f: n \rightarrow 2$  be such that  $f(i) = 1$  if and only if  $a_i = \pi_*(x_i)$ . Then  $\bigcap_{i < n} a_i$  contains  $a_{\bar{s}, f} \setminus k$ .

Given a function  $\pi: 2^{<\omega} \rightarrow [\omega]^{<\omega}$ , say that a set  $b \subseteq \omega$  is  $\pi$ -weak if there exist infinitely many  $x \in 2^\omega$  such that at least one of  $b \cap \pi_*(x)$  and  $b \setminus \pi_*(x)$  is finite.

**Lemma 9.3.5.** *If  $\pi: 2^{<\omega} \rightarrow [\omega]^{<\omega}$  is diffuse,  $n$  is a positive integer and  $b_i$  ( $i < n$ ) are  $\pi$ -weak subsets of  $\omega$ , then  $\bigcup_{i < n} b_i \neq \omega$ .*

*Proof.* Since each  $b_i$  is  $\pi$ -weak, there exist distinct  $x_i \in 2^\omega$  ( $i < n$ ) and  $a_i \in \{\pi_*(x_i), \omega \setminus \pi_*(x_i)\}$  ( $i < n$ ) such that, for each  $i < n$ ,  $b_i \cap a_i$  is finite. No member of the set

$$\left(\bigcap_{i < n} a_i\right) \setminus \bigcup_{i < n} (b_i \cap a_i)$$

is in  $\bigcup_{i < n} b_i$ . Since  $\pi$  is diffuse, this set is infinite.  $\square$

We separate out one lemma which will be used again in the proof of Theorem 10.2.6.

**Lemma 9.3.6.** *Let  $\pi: 2^{<\omega} \rightarrow [\omega]^{<\omega}$  be a recursive diffuse function in a model  $M$  of ZF. Let  $U$  be, in  $M$ , an ultrafilter on  $\omega$  disjoint from the collection of  $\pi$ -weak sets, and suppose that  $\mathcal{P}(\mathbb{P}_U)^M$  is countable. Let  $y$  be in  $2^\omega \setminus M$ , and let  $a$  be a subset of  $\omega$ . Then there exists a  $c \subseteq \omega$  which is  $M$ -generic for  $\mathbb{P}_U^M$  such that  $a \in L[c, y]$ .*

*Proof.* By  $\Sigma_1^1$  absoluteness, for each  $b \in U$ , every  $z \in 2^\omega$  such that at least one of  $b \cap \pi^*(z)$  and  $b \setminus \pi^*(z)$  is finite is in  $M$ . It follows that  $\pi^*(y)$  and  $\omega \setminus \pi^*(y)$  have infinite intersection with each member of  $U$ .

By Lemma 9.3.3, there exists an  $M$ -generic  $b \subseteq \omega$  such that  $b \cap \pi^*(y)$  and  $b \setminus \pi^*(y)$  are both infinite. To see this, enumerate all functions  $F: \omega \rightarrow U$  in  $M$  as  $\langle F_i : i \in \omega \rangle$ . Let  $b$  be  $\{n_i : i \in \omega\}$  where, for each  $i \in \omega$ ,  $n_i < n_{i+1}$ ,  $n_i \in \bigcap_{j < i} F_j(\{n_j\})$  and  $n_i \in \pi^*(y)$  if and only if  $i$  is even. Let  $c$  be an infinite subset of  $b$ , enumerated in increasing order by  $\langle m_i : i < \omega \rangle$ , such that, for each  $i \in \omega$ ,  $m_i \in \pi^*(y)$  if and only if  $i \in a$ . Then  $c$  is  $M$ -generic, by Lemma 9.3.4. Since  $\pi$  is recursive,  $\pi^*(y) \in L[y]$ , so  $a \in L[c, y]$ .  $\square$

The end of the proof of Theorem 9.3.2 uses following theorem of Solovay, which will be used again in Section 9.6. We will use the theorem only for the case where  $x$  is a set of ordinals, in which case the proof is slightly easier.

**Theorem 9.3.7** (Solovay). *Suppose that  $M \subseteq N$  are transitive models of ZF,  $\mathbb{P}$  is a partial order in  $M$  and  $\mathcal{P}(\mathbb{P} \times \mathbb{P}) \cap N$  is countable. Suppose that  $x \in M[G]$  for every  $N$ -generic  $G \subseteq \mathbb{P}$ . Then  $x \in M$ .*

*Proof.* It suffices to prove the theorem in the case where  $x$  is transitive, replacing  $x$  with  $\text{TC}(\{x\})$  if necessary. Since  $\mathcal{P}(\mathbb{P} \times \mathbb{P}) \cap N$  is countable, there exists an  $N$ -generic  $(G, H)$  for  $\mathbb{P} \times \mathbb{P}$ . Arguing by induction on the rank of  $x$ , we may assume that the members of  $x$  are all in  $M$ . Then if  $\tau$  and  $\sigma$  are  $\mathbb{P}$ -names in  $M$  such that  $\tau_G = \sigma_H = x$ , it follows by genericity that there is some condition  $(p, q) \in G \times H$  such that, for all  $y \in M$   $p$  decides the statement  $\check{y} \in \tau$  and  $q$  decides the statement  $\check{y} \in \sigma$ , from which it follows that  $x \in M$ .  $\square$

*Proof of Theorem 9.3.2.* Let  $x \in \omega^\omega$ ,  $y \in 2^\omega$  be such that  $y \in L[T, x] \setminus L[S, x]$ . Let  $\pi: 2^{<\omega} \rightarrow [\omega]^{<\omega}$  be a recursive diffuse function. Let  $U$  be, in  $L[S, x]$ , an ultrafilter on  $\omega$  not containing any  $\pi$ -weak subset of  $\omega$ , and let  $\mathbb{P}$  be  $\mathbb{P}_U^{L[S, x]}$ . Let  $a \subseteq \omega$  HC-code an enumeration in ordertype  $\omega$  of the  $\mathbb{P}$ -names in  $L[S, x]$  for

elements of  $\omega^\omega$ . By Lemma 9.3.6, there is an  $L[S, x]$ -generic set  $c$  for  $\mathbb{P}$  such that  $a \in L[c, y]$ . Then  $\omega^\omega \cap L[S, x][c]$  is a countable set in  $L[T, x, c]$ .

It follows by Turing Determinacy that there is an  $x_* \in \omega^\omega$  such that, for all  $y \geq_{\text{Tu}} x_*$ ,  $\omega^\omega \cap L[S, y]$  is a countable set in  $L[T, y]$ . Applying this fact and Theorem 9.3.7 (with  $M = N = L[T, y]$ ) to the  $\text{Col}(\omega, \gamma)$ -extension of any such model  $L[T, y]$ , we see that for all such  $y$ ,  $\mathcal{P}(\gamma) \cap L[S, y]$  is a set of cardinality  $|\gamma|^{L[T, y]}$  in  $L[T, y]$ .  $\square$

Theorem 9.3.8 lists some consequences of Theorem 9.3.2. Note that neither part of the theorem assumes (or implies) that  $S$  is in  $L[T, y]$ . A natural application of part (1b) is when  $S$  is an  $\infty$ -Borel code, as defined in Chapter 8.

**Theorem 9.3.8** (ZF + TD). *Let  $S$  and  $T$  be sets of ordinals.*

1. *If  $x$  is a base of a Turing cone witnessing that  $S \leq_{\mathcal{D}} T$ , then the following hold for all  $y \geq_{\text{Tu}} x$ .*

- (a) *For every  $\gamma < \omega_1^V$ ,  $\mathcal{P}(\gamma) \cap L[S, y] \subseteq L[T, y]$ .*
- (b) *For every formula  $\varphi$ ,*

$$\{z \in \omega^\omega : L[S, z] \models \varphi(S, z)\} \cap L[T, y] \in L[T, y].$$

2. *If  $\neg(T \leq_{\mathcal{D}} S)$ , then the following hold for a Turing cone of  $y \in \omega^\omega$ .*

- (a) *For every  $\gamma < \omega_1^V$ ,*

$$\mathcal{P}(\gamma) \cap L[S, y] \in H(|\gamma|^+)^{L[T, y]}.$$

- (b) *Every regular cardinal of  $L[T, y]$  below  $\omega_1^V$  is a strongly inaccessible cardinal in  $L[S, y]$ .*

*Proof.* Part (1a) follows a collapse argument using Theorem 9.3.7, as in the last paragraph of the proof of Theorem 9.3.2. For part (1b), it is enough by Theorem 9.3.7 to see that the set  $A = \{z \in \omega^\omega : L[S, z] \models \varphi(S, z)\} \cap L[T, y]$  is in  $L[T, y][g]$  whenever  $g$  is  $L[T, y]$ -generic for the partial order  $\text{Col}(\omega, \mathbb{R})$  (as defined in  $L[T, y]$ ). Fixing such a  $g$ , we have that  $A$  is in  $L[B]$ , for some bounded subset  $B$  of  $\omega_1^V$  in  $L[S, y, g]$ . It follows from part (1a) then that  $B$  and  $A$  are in  $L[T, y]$ . Part (2a) is Theorem 9.3.2, and part (2b) follows directly from part (2a).  $\square$

## 9.4 Pointed trees

In this section we prove a result due to Martin which will be used in Sections 9.5 and 9.6. We continue to use the notation from Definition 9.1.1, and add two additional terms in Definition 9.4.1. Recall that a tree is perfect if each node has an incompatible pair of extensions, and, given a tree  $a$ , that  $[a]$  denotes the set of infinite branches through  $a$ .

**9.4.1 Definition.** Given a set of ordinals  $S$ , we say that

- a set  $A \subseteq \omega^\omega$  is  *$S$ -positive* if it intersects every  $S$ -cone;
- a tree  $a \subseteq \omega^{<\omega}$  is  *$S$ -pointed* if  $a$  is perfect and, for every  $x \in [a]$ ,  $a \in L[S, x]$ ;

The following theorem, which appears in [13], shows that, assuming AD, the sets of the form  $[a]$ , for  $a$  an  $S$ -pointed tree, are dense in the containment order on the  $S$ -positive subsets of  $\omega^\omega$ .

**Theorem 9.4.2** (ZF + AD; Martin). *Let  $S$  be a set of ordinals and let  $A$  be a subset of  $\omega^\omega$ . Then  $A \subseteq \omega^\omega$  is  $S$ -positive if and only if there is an  $S$ -pointed perfect tree  $a$  such that  $[a] \subseteq A$ .*

*Proof.* The reverse implication (which does not use the assumption of AD) follows from the fact that for any perfect tree  $a \subseteq \omega^{<\omega}$  and any  $x \in \omega^\omega$  there is a  $y \in [a]$  such that  $x \in L[a, y]$ .

For the forward implication, consider the game  $\mathcal{G}$  where  $I$  and  $II$  respectively build  $x$  and  $y$  in  $\omega^\omega$ , and  $II$  wins if and only if  $x \leq_S y$  and  $y \in A$ . If  $\sigma$  is a strategy for  $I$  and  $y \in A$  is such that  $\sigma$  is in  $L[S, y]$ , then  $y$  is a winning play for  $II$  against  $\sigma$ . This shows that if  $A$  is  $S$ -positive then  $I$  cannot have a winning strategy. Suppose then that  $\sigma$  is a winning strategy for  $II$ . For each  $s \in \omega^{<\omega}$ , let  $\sigma(s)$  denote the sequence of ( $|s|$ -many) moves made according to  $\sigma$  in response to  $s$ . Let  $z \in 2^\omega$  be such that  $\sigma \in L[S, z]$ , and let  $Z$  be the set of  $t \in \omega^{<\omega} \cup \omega^\omega$  such that  $t(2n) = z(n)$  whenever  $n \in \omega$  and  $2n \in \text{dom}(t)$ . Working in  $L[S, z]$  we can choose for each  $s \in 2^{<\omega}$ , recursively in  $|s|$ , a  $t_s$  in  $Z \cap \omega^{<\omega}$ , in such a way that, for all  $s, s' \in 2^{<\omega}$ ,

- if  $|s| = |s'|$  then  $|t_s| = |t_{s'}|$ ;
- if  $s \perp s'$  then  $\sigma(t_s) \neq \sigma(t_{s'})$ .

The achievability of the second condition above follows from the fact that  $\sigma$  is a winning strategy for  $II$ , which implies that, for any finite  $t \in Z$ , the set

$$\{x \circ \sigma : x \in \omega^\omega \cap Z, t \subseteq x\}$$

is  $S$ -positive, and therefore has size at least 2. Let  $a$  be the tree of initial segments of the members of  $\{\sigma(t_s) : s \in 2^{<\omega}\}$ . We claim that  $a$  is as desired. To see this, fix  $y \in [a]$  and  $x \in \omega^\omega \cap Z$  such that  $y = x \circ \sigma$ . Since  $\sigma$  is a winning strategy for  $II$ ,  $y$  is in  $A$ , and  $x$  is in  $L[S, y]$ . This implies that  $z$  and  $\sigma$  are in  $L[S, y]$  and therefore that  $a$  is in  $L[S, y]$  as desired.  $\square$

## 9.5 Coding ultrapowers

This section continues to use the notation from Definitions 9.1.1 and 9.4.1, for a set of ordinals  $S$ . We work under the assumption that  $\mu_S$  is an ultrafilter (which AD implies, by Corollary 1.2.6) on  $\mathcal{D}_S$ , and let  $\delta_S^\infty$  be  $\prod \omega_2^{L[S, x]} / \mu_S$ , assuming



that this ultrapower is wellfounded (as it is if DC holds). In Remark 9.6.7 we will show (assuming AD plus the wellfoundedness of the ultrapower  $\text{Ord}^{\mathcal{D}_S}/\mu_S$ ) that  $\delta_S^\infty \leq \Theta$ . In this section we will show that  $\delta_S^\infty < \Theta$ , under the additional assumption that, for some set of ordinals  $T$ ,  $S <_{\mathcal{D}} T$ .

Given a set of ordinals  $T$ , we define the relation  $<_T^c$  on  $(\omega^\omega)^2$  by setting  $(x, y) <_T^c (z, w)$  to hold if and only if

- $x = z$ ;
- $y$  and  $w$  are in  $L[T, x]$ ;
- $y$  comes before  $w$  in the constructibility order in  $L[T, x]$  using  $T$  and  $x$  as predicates.

In the statement of Theorem 9.5.1 the set  $\leq_S \times <_T^c$  is used for convenience as a set coding both  $\leq_S$  and  $<_T^c$ .

**Theorem 9.5.1** (ZF+AD). *Let  $S$  and  $T$  be sets of ordinals. Suppose that  $f$  is an  $S$ -invariant function on  $\omega^\omega$  such that, for an  $S$ -cone of  $x$ ,  $f(x)$  is a transitive set in  $H(\aleph_1)^{L[T, x]}$ . Then  $\prod f(x)/\mu_S$  is isomorphic to a relation which is projective in  $\leq_S \times <_T^c$ . If in addition  $\text{DC}_{\mathbb{R}}$  holds then  $\prod f(x)/\mu_S$  is wellfounded.*

*Proof.* Let  $R_f$  be the set of  $(x, y)$  such that  $y \in \omega^\omega \cap L[T, x]$  and  $y$  HC-codes  $f(x)$ , and let  $R_f^*$  be the set of  $<_T^c$ -minimal pairs in  $R_f$ . Then  $R_f^*$  is a uniformizing function for  $R_f$ . Note however that  $R_f^*$  need not be  $S$ -invariant.

Let  $E$  be the set of  $y \in \omega^\omega$  which HC-code a transitive set. Then  $E$  is coanalytic. If  $y \in E$  HC-codes a transitive set  $M$ , then, since  $M$  is rigid, for each  $k \in \omega$  there is a unique  $p$  such that there is an isomorphism between

$$(\omega, \{(n, m) \in \omega \times \omega : y(2^n 3^m) = 0\})$$

and  $(\text{TC}(\{M\}), \in)$  sending  $k$  to  $p$ . We call this element  $p_{y,k}$ . The following sets are projective:

- $\{(y, y', k, k') \in E^2 \times \omega^2 : p_{y,k} = p_{y',k'}\}$ ;
- $\{(y, y', k, k') \in E^2 \times \omega^2 : p_{y,k} \in p_{y',k'}\}$ ;

Let  $H$  be the set of  $S$ -invariant functions  $h$  such that  $h(x) \in f(x)$  for an  $S$ -cone of  $x \in \omega^\omega$ . For each  $h \in H$  there is a  $k \in \omega$  such that the set  $A_{h,k} = \{x \in \omega^\omega : p_{R_f^*(x),k} = h(x)\}$  is  $S$ -positive. Since  $R_f^*$  may not be  $S$ -invariant, there may be more than one such  $k$ . Any such pair  $(k, A_{h,k})$  codes  $h$  on an  $S$ -cone, however, in the sense that, for an  $S$ -cone of  $z \in \omega^\omega$ ,

$$h(z) = h(x) = p_{R_f^*(x),k}$$

for any  $x \in [z]_S \cap A_{h,k}$ .

Let  $\langle \sigma_i : i < \omega \rangle$  be a recursive listing of  $\omega^{<\omega}$ , and let  $C_f$  be the set of pairs  $(k, b) \in \omega \times 2^\omega$  such that  $a_b = \{\sigma_i : b(i) = 1\}$  is an  $S$ -pointed perfect tree and, for all  $x, y \in [a_b]$ , if  $x \equiv_S y$  then  $p_{R_f^*(x),k} = p_{R_f^*(y),k}$ . By Theorem 9.4.2, every

$S$ -positive set contains the ( $S$ -positive) set of paths through some  $S$ -pointed tree. It follows that for each  $h \in H$ , given  $k \in \omega$  such that  $A_{h,k}$  is  $S$ -positive, there is a  $b \in 2^\omega$  such that  $[a_b] \subseteq A_{h,k}$  and  $(k, b) \in C_f$ .

For each  $(k, b) \in C_f$  and  $z \in \omega^\omega$ , set  $h_{(k,b)}([z]_S)$  to be the common value of  $p_{R_f^*(x),k}$  for all  $x \in [z]_S \cap [a_b]$  (if there is such a value, and  $\emptyset$  otherwise). By the two previous paragraphs, for each  $h \in H$  there exists a  $(k, b) \in C_f$  such that  $h(z) = h_{(k,b)}([z]_S)$  for an  $S$ -cone of  $z \in \omega^\omega$ . Define an equivalence relation  $\sim$  on  $C_f$  by setting

$$(k, b) \sim (k', b')$$

if  $h_{(k,b)}$  and  $h_{(k',b')}$  agree on an  $S$ -cone, and let  $[k, b]_f$  denote the  $\sim$ -class of  $(k, b)$ . Define  $[k, b]_f \in_f [k', b']_f$  to hold whenever

$$h_{(k,b)}([z]_S) \in h_{(k',b')}([z]_S)$$

for an  $S$ -cone of  $z \in \omega^\omega$ . Then  $(C_f / \sim, \in_f)$  is isomorphic to  $\prod f(x) / \mu_S$ . Since  $C_f$ ,  $\sim$  and the relation on  $C_f$  induced by  $\in_f$  are projective in  $\leq_S \times <_T^c$  the first conclusion of the theorem follows. The second conclusion follows then from the fact that  $C_f$  is a surjective image of  $\omega^\omega$ .  $\square$

## 9.6 Forcing with positive sets

This section applies the results of the previous three sections to the analysis of ultrapowers of the form  $\prod V / \mu_S$ . Given a set  $S$  consisting of ordinals, let  $\mathbb{P}_S$  be the partial order of  $S$ -positive subsets of  $\omega^\omega$  and let  $\mathbb{S}_S$  be the partial order of  $S$ -pointed perfect trees, each ordered by containment. By Theorem 9.4.2, each  $\mathbb{P}_S$  condition contains the set of infinite paths through some  $S$ -pointed tree, which is in turn  $S$ -positive. It follows that a  $V$ -generic filter for either partial order generates one for the other.

A  $V$ -generic filter  $G \subseteq \mathbb{P}_S$  induces an ultrapower  $\prod_{x \in \omega^\omega} V / G$ , whose elements are represented by (not necessarily  $S$ -invariant) functions in  $V$  with domain  $\omega^\omega$ . For each hereditarily countable set  $a$ , the constant function from  $\omega^\omega$  to  $\{a\}$  represents  $a$  in this ultrapower. The identity function represents the generic  $x_G \in \omega^\omega$  added by  $\mathbb{P}_S$ . The following theorem shows that the corresponding ultrapower of any set of ordinals, in particular the ultrapower of  $S$  itself, is the same in the ultrapowers with respect to  $\mu_S$  and  $G$ , where the  $\mu_S$  ultrapower is computed in  $V$  and the  $G$  ultrapower is computed in the  $\mathbb{P}_S$ -extension.

**Theorem 9.6.1.** *Let  $S$  be a set of ordinals such that  $\leq_S$  is locally countable and  $\mu_S$  is a countably complete ultrafilter on  $\mathcal{D}_S$ . Let  $G \subseteq \mathbb{P}_S$  be a  $V$ -generic filter. Let  $k$  be the map sending  $[f]_{\mu_S}$  to  $[f]_G$ , for each  $S$ -invariant function  $f$  in  $V$  from  $\omega^\omega$  to  $\text{Ord}$ . Then  $k$  maps  $(\prod \text{Ord} / \mu_S)^V$  isomorphically to  $(\prod_{x \in \omega^\omega} \text{Ord} / G)^{V[G]}$ .*

*Proof.* Since each  $S$ -positive set has  $S$ -positive intersection with each  $S$ -cone, and  $\mu_S$  is an ultrafilter,  $k$  is injective and order preserving. We show that it is onto. Let  $g: A \rightarrow \text{Ord}$  be a function in  $V$  with  $S$ -positive domain. Define  $f$  on

$$\bigcup \{[x]_S : x \in A\}$$

by letting  $f(x)$  be  $\min(g[[x]_S \cap A])$ . Then  $f$  is  $S$ -invariant. Let  $C$  be

$$\{x \in A : f(x) = g(x)\}.$$

It suffices (by the genericity of  $G$ ) to show that  $C$  is  $S$ -positive, since  $C$  forces that  $[g]_G = [f]_G = k([f]_{\mu_S})$ . Supposing otherwise, let  $x \in \omega^\omega$  be such that  $\{y \in \omega^\omega : y \geq_S x\}$  is disjoint from  $C$ . Since  $A$  is  $S$ -positive, there is a  $y \geq_S x$  in  $A$ . Then there is a  $z \in [y]_S \cap A$  such that  $g(z) = f(y)$ . Then  $z \in C$  and  $z \geq_S x$ , giving a contradiction.  $\square$

We let  $j_G$  denote the embedding of  $V$  into  $\prod_{x \in \omega^\omega} V/G$  sending each element of  $V$  to the class represented by its corresponding constant function, assuming that  $G$  is a  $V$ -generic filter for  $\mathbb{P}_S$ . We let  $j_S$  be the corresponding embedding of  $V$  into  $\prod V/\mu_S$ . The following corollary of Theorem 9.6.1 says that the two embeddings agree on sets of ordinals.

**Corollary 9.6.2.** *Let  $S$  be a set of ordinals such that  $\leq_S$  is locally countable and  $\mu_S$  is a countably complete ultrafilter on  $\mathcal{D}_S$ . Let  $G \subseteq \mathbb{P}_S$  be a  $V$ -generic filter. Then for every set  $T$  of ordinals,  $j_G(T) = j_S(T)$ .*

*Proof.* By Theorem 9.6.1, for each ordinal  $\gamma$  there is an  $S$ -invariant function  $f: \omega^\omega \rightarrow \text{Ord}$  representing  $\gamma$  in both ultrapowers. Given  $f$ , either  $f(x) \in T$  for an  $S$ -cone of  $x$  or  $f(x) \notin T$  for an  $S$ -cone of  $x$ . In either case the same must hold on a set in  $G$ .  $\square$

**9.6.3 Remark.** We are primarily interested here in the ultraproduct

$$\prod_{x \in \omega^\omega} L[S, x]/G,$$

which by Theorem 9.6.1 and Corollary 9.6.2 is isomorphic to  $\prod L[S, x]/\mu_S$ . Since the functions used to form  $\prod_{x \in \omega^\omega} L[S, x]/G$  are not required to be  $S$ -invariant, the usual proof of Łoś's Theorem goes through (using the constructibility order on  $L[S, x]$ ), showing that whenever  $f_1, \dots, f_n$  are functions on  $\omega^\omega$  in  $V$  such that each value  $f_i(x)$  is in the corresponding model  $L[S, x]$ , and  $\varphi$  is an  $n$ -ary formula,

$$\prod_{x \in \omega^\omega} L[S, x]/G \models \varphi([f_1]_G, \dots, [f_n]_G)$$

if and only if the set of  $x$  for which  $L[S, x] \models \varphi(f_1(x), \dots, f_n(x))$  is in  $G$ . In particular, for each  $n \in \omega$  the function  $x \mapsto \omega_n^{L[S, x]}$  represents the value of  $\omega_n$  in the ultrapower.

Assuming that the ultrapower  $\text{Ord}^{\mathcal{D}_S}/\mu_S$  is wellfounded, we let  $S^\infty$  be the set of ordinals represented by the function with constant value  $S$ . Given  $A \subseteq \omega^\omega$ , we say that  $S$  *computes*  $A$  on a cone if  $A \cap L[S, x] \in L[S, x]$  for an  $S$ -cone of  $x \in \omega^\omega$ . Part (1b) of Theorem 9.3.8 shows that if  $S$  and  $T$  are sets of ordinals such that  $S \leq_{\mathcal{D}} T$ , then  $T$  computes  $\{x \in \omega^\omega : L[S, x] \models \varphi(S, x)\}$  on a cone, for any binary formula  $\varphi$ .

The following is a summary of information collected so far.

**Corollary 9.6.4.** *Let  $S$  be a set of ordinals such that  $\leq_S$  is locally countable,  $\mu_S$  is a countably complete ultrafilter on  $\mathcal{D}_S$  and  $\prod \text{Ord}/\mu_S$  is wellfounded. Let  $G \subseteq \mathbb{P}_S$  be a  $V$ -generic filter and let  $x_G$  be the corresponding generic element of  $\omega^\omega$ . Then the following hold.*

1. *The ultraproduct  $\prod_{x \in \omega^\omega} L[S, x]/G$  is isomorphic to  $L[S^\infty, x_G]$ .*
2. *For all  $\gamma < j_G(\omega_1^V)$  which are infinite cardinals of  $L[S^\infty, x_G]$ ,*

$$L[S^\infty, x_G] \models \diamond_{\gamma^+}.$$

$$3. \omega_1^V = \omega_1^{L[S^\infty, x_G]}$$

$$4. \delta_S^\infty = \omega_2^{L[S^\infty, x_G]}$$

$$5. (\omega^\omega)^V \subseteq L[S^\infty, x_G]$$

6. *If  $A \subseteq \omega^\omega$  is such that  $S$  computes  $A$  on a cone, then the function sending each  $x \in \omega^\omega$  to  $A \cap L[S, x]$  represents a set  $A^* \subseteq (\omega^\omega)^{V[G]}$  in  $L[S^\infty, x_G]$  such that  $A^* \cap (\omega^\omega)^V = A$ .*

*Proof.* For part (1), the fact that the identity function on  $\omega^\omega$  represents  $x_G$  shows that  $\prod_{x \in \omega^\omega} L[S, x]/G$  is isomorphic to  $L[j_G(S), x_G]$ . Corollary 9.6.2 implies that this is the same as  $L[S^\infty, x_G]$ . Part (2) follows from Remark 9.6.3 and Theorem 9.1.3. Part (3) follows from Theorems 9.6.1 and 9.1.2, and Remark 9.6.3. Part (4) follows from Theorem 9.6.1 and Remark 9.6.3. Parts (5) and (6) follow from part (1) and Theorem 9.6.1.  $\square$

**9.6.5 Remark.** Let  $S$  be a set of ordinals such that  $\mu_S$  is a countably complete ultrafilter on  $\mathcal{D}_S$  and  $\text{Ord}^{\mathcal{D}_S}/\mu_S$  is wellfounded. For all  $x, y \in \omega^\omega$ ,  $y \in L[S^\infty, x]$  if and only if  $y \in L[S, x]$  for an  $S$ -cone of  $z$ , i.e., if and only if  $y \in L[S, x]$ . It follows that  $\leq_S$  and  $\leq_{S^\infty}$  are the same relation.

The following theorem will be used in Sections 11.2 and 11.4. The last part of the theorem uses the notion of  $\infty$ -Borel code as defined in Section 8.1.

**Theorem 9.6.6 (ZF + AD).** *Let  $S$  be a set of ordinals such that  $\text{Ord}^{\mathcal{D}_S}/\mu_S$  is wellfounded. Suppose that  $A \subseteq \omega^\omega$  is such that  $S$  computes  $A$  on a cone. Then  $A \in L(S^\infty, \mathbb{R})$ . In particular, if  $S$  is a  $\leq_{\mathcal{D}}$ -maximal set of ordinals, then every  $\infty$ -Borel set is in  $L(S^\infty, \mathbb{R})$ .*

*Proof.* We show that  $A \in L(S^\infty, \mathbb{R})[x_G]$  whenever  $x_G$  is  $V$ -generic for  $\mathbb{S}_{S^\infty}$ . The theorem then follows by Theorem 9.3.7. By Remark 9.6.5, the partial orders  $\mathbb{S}_S$  and  $\mathbb{S}_{S^\infty}$  are the same, and by Theorem 9.4.2 they are forcing-equivalent to  $\mathbb{P}_S$ . It follows that if  $x_G$  is  $V$ -generic for  $\mathbb{S}_{S^\infty}$  then it is  $V$ -generic for  $\mathbb{P}_S$ , and therefore by part (6) of Corollary 9.6.4 that there is an  $A^* \in L[S^\infty, x_G]$  such that  $A^* \cap (\omega^\omega)^V = A$ . Since  $(\omega^\omega)^V \in L(S^\infty, \mathbb{R})[x_G]$  we are done. The second part of the theorem follows from the first part, and part (1b) of Theorem 9.3.8.  $\square$

By Theorems 8.2.3 and 9.6.6, the  $\infty$ -Borel sets are all  $\infty$ -Borel in  $L(S^\infty, \mathbb{R})$ , assuming that  $S$  is a  $\leq_{\mathcal{D}}$ -maximal set of ordinals, along with the other the hypotheses of Theorem 9.6.6.

Similarly, if AD holds and  $A \subseteq \omega^\omega$  is not  $\infty$ -Borel, then every  $\infty$ -Borel set is  $\infty$ -Borel in  $L(A, \mathbb{R})$  by Theorem 8.2.3. If  $S$  is a set of ordinals, one can apply Theorem 9.6.6 in  $L(S, A, \mathbb{R})$  assuming (in addition) only  $\text{DC}_{\mathbb{R}}$  (i.e., without assuming that  $\prod \text{Ord}/\mu_S$  is wellfounded in  $V$ ). In this case, however, the  $S^\infty$  in the statement of the theorem is as computed in  $L(S, A, \mathbb{R})$ .

**Corollary 9.6.7** (ZF + AD). *Let  $S$  be a set of ordinals such that  $\prod \text{Ord}/\mu_S$  is wellfounded. Then  $\Theta \geq \delta_S^\infty$ .*

*Proof.* Let  $x_G$  be  $V$ -generic for  $\mathbb{S}_S$ . The partial order  $\mathbb{S}_S$  is a surjective image of  $\omega^\omega$  (in  $V$ ), so forcing with  $\mathbb{S}_S$  preserves  $\Theta^V$  as a cardinal. Since  $L[S^\infty, x_G]$  is an inner model of  $V[G]$ ,  $\Theta^V$  is a cardinal in  $L[S^\infty, x_G]$ . Since  $\omega_1^{L[S^\infty, x_G]} = \omega_1^V < \Theta^V$ , it must be that  $\Theta^V \geq \omega_2^{L[S^\infty, x_G]}$ , which is equal to  $\delta_S^\infty$ , by Corollary 9.6.4.  $\square$

We say that  $S$  is *strongly maximal* if it computes each subset of  $\omega^\omega$  on a cone. By part (1b) of Theorem 9.3.8, every  $\leq_{\mathcal{D}}$ -maximal set of ordinals is strongly maximal, if every set of reals is  $\infty$ -Borel.

**Corollary 9.6.8.** *Let  $S$  be a strongly maximal set of ordinals such that  $\leq_S$  is locally countable,  $\mu_S$  is a countably complete ultrafilter on  $\mathcal{D}_S$  and  $\prod \text{Ord}/\mu_S$  is wellfounded. Then  $\delta_S^\infty \geq \Theta$ .*

*Proof.* It follows from Corollary 9.6.4 that each prewellordering of  $\omega^\omega$  in  $V$  is a subset of a prewellordering of  $(\omega^\omega)^{L[S^\infty, x_G]}$  in  $L[S^\infty, x_G]$ . Since  $L[S^\infty, x_G]$  satisfies CH, and  $\delta_S^\infty = \omega_2^{L[S^\infty, x_G]}$ , this implies that  $\delta_S^\infty \geq \Theta$ .  $\square$



# Chapter 10

## Vopěnka algebras

This chapter presents a version of Vopěnka's theorem (Theorem 10.1.2) - which implies among other things that every set of ordinals is generic over the inner model  $\text{HOD}$  (see page 249 of [8]) - and various applications and extensions due to Woodin. One application is Theorem 10.1.8, which shows, assuming  $\text{AD}$ , that for any  $A \subseteq \omega^\omega$ , either  $\mathcal{P}(\omega^\omega) \subseteq \text{L}(A, \mathbb{R})$  or  $A^\#$  exists. In Section 10.2 we show, assuming  $\text{TD} + \text{DC}_{\mathbb{R}}$ , that the  $\infty$ -Borel sets are closed under projections (Theorem 10.2.3). Corollary 10.2.7 shows, under the same assumption, that **Uniformization** is equivalent to the assertion that  $\leq_{\mathcal{D}}$  has no maximal element. In Section 10.3 we introduce a variant of the Vopěnka algebra using  $\infty$ -Borel codes. An application, Corollary 10.3.7 shows, again assuming  $\text{TD} + \text{DC}_{\mathbb{R}}$ , that whenever  $Y$  is a set of ordinals, every subset of  $\omega^\omega$  in  $L(Y, \mathbb{R})$  is  $\infty$ -Borel.

The material in this chapter uses various relativizations of the inner model  $\text{HOD}$ . Recall that for a set  $X$ ,  $\text{OD}_X$  is the class of all sets which are definable from a finite sequence from  $X \cup \text{Ord}$ . We let  $\text{HOD}_X$  denote the class of all sets in  $\text{OD}_X$  whose transitive closures are contained in  $\text{OD}_X$ .

A proof of Theorem 10.0.1 below is given in Section 5.2 of [35]. Note that the members of  $X$  are not necessarily in  $\text{HOD}_{\{X\}}$ . This can happen for instance if  $X$  is an uncountable subset of  $\omega^\omega$ .

**Theorem 10.0.1** (Gödel). *If  $X$  is a set such that  $X \in \text{OD}_X$ , then  $\text{HOD}_X$  is a model of ZF. If in addition there is a wellordering of  $X$  in  $\text{OD}_X$ , then  $\text{HOD}_X$  is a model of ZFC.*

### 10.1 The Vopěnka algebra

We start by fixing an indexing of the subsets of  $\mathcal{P}(Y)$  in  $\text{OD}_{\bar{x}}$ , where  $\bar{x}$  is a finite sequence and  $Y$  is a set in  $\text{OD}_{\bar{x}}$ . Let  $\kappa$  be the least cardinal  $\lambda$  such that  $\bar{x} \in V_\lambda$  and every subset of  $\mathcal{P}(Y)$  in  $\text{OD}_{\bar{x}}$  is in  $\text{OD}_{\bar{x}}^{V_\lambda}$ . Let  $P_{\bar{x}, Y}$  be the set of pairs  $(n, \bar{\alpha})$  such that

- $n$  is the Gödel number of a ternary formula  $\varphi$ ,

- $\bar{\alpha}$  is a finite sequence of elements of  $\kappa$  and
- the set  $B_{n,\bar{\alpha}} = \{C \subseteq Y : V_\kappa \models \varphi(C, \bar{x}, \bar{\alpha})\}$  is nonempty.

Then  $P_{\bar{x},Y}$  is in  $\text{HOD}_{\bar{x}}$ , as is the reflexive and transitive ordering  $\leq_{\bar{x},Y}$  on  $P_{\bar{x},Y}$  defined by setting  $(n, \bar{\alpha}) \leq_{\bar{x},Y} (m, \bar{\beta})$  to hold when  $B_{n,\bar{\alpha}} \subseteq B_{m,\bar{\beta}}$ . We let  $\equiv_{\bar{x},Y}$  be the induced equivalence relation on  $P_{\bar{x},Y}$ , and for each  $(n, \bar{\alpha}) \in P_{\bar{x},Y}$ , let  $[n, \bar{\alpha}]_{\bar{x},Y}$  denote the corresponding equivalence class. We let  $\mathbb{V}_{\bar{x},Y}$  (the *Vopěnka algebra* for  $\mathcal{P}(Y)$  relative to  $\bar{x}$ ) be the partial order whose domain is the set of  $\equiv_{\bar{x},Y}$ -classes of  $P_{\bar{x},Y}$ , with the order inherited from  $\leq_{\bar{x},Y}$ .

**10.1.1 Remark.** There are natural modifications of the definition of  $P_{\bar{x},Y}$  and  $\mathbb{V}_{\bar{x},Y}$  just given to allow for the case where  $X$  is an arbitrary (possibly infinite) element of  $\text{HOD}_X$ . This variation does not generalize the version under consideration here, since we are not requiring  $\bar{x}$  to be in  $\text{HOD}_{\bar{x}}$ . In this case the members of  $P_{X,Y}$  can be triples  $(n, \bar{x}, \bar{\alpha})$  with  $\bar{x}$  a finite sequence from  $X$ , and the sets  $B_{n,\bar{\alpha}}$  are replaced with sets  $B_{n,\bar{x},\bar{\alpha}}$  of the form  $\{C \subseteq Z : V_\kappa \models \varphi(C, \bar{x}, \bar{\alpha})\}$ . The analysis then goes through largely as we have here, except that the last paragraph of the proof of Theorem 10.1.2 is complicated slightly by the fact that the corresponding model  $\text{HOD}_{X \cup \{E\}}$  may not satisfy Choice. This form of the Vopěnka algebra can be used to force over models of the form  $\text{HOD}_\Gamma$ , where  $\Gamma$  is an initial segment of the Wadge hierarchy.

**Theorem 10.1.2.** *Let  $\bar{x}$  be a finite sequence, and let  $Y$  be a set in  $\text{HOD}_{\bar{x}}$ . For each  $E \subseteq Y$ , there is a  $\text{HOD}_{\bar{x}}$ -generic filter  $G \subseteq \mathbb{V}_{\bar{x},Y}$  such that*

$$\text{HOD}_{\bar{x} \smallfrown \langle E \rangle} = \text{HOD}_{\bar{x}}[G].$$

*Proof.* Fix  $E \subseteq Y$ . If  $D$  is a subset of  $P_{\bar{x},Y}$  and an element of  $\text{HOD}_{\bar{x}}$ , then

$$\bigcup \{B_{n,\bar{\alpha}} : (n, \bar{\alpha}) \in D\}$$

is in  $\text{OD}_{\bar{x}}$ . It follows that if this union is not all of  $\mathcal{P}(Y)$ , then there is a pair  $(m, \bar{\beta})$  in  $P_{\bar{x},Y}$  such that

$$B_{m,\bar{\beta}} = \mathcal{P}(Y) \setminus \bigcup \{B_{n,\bar{\alpha}} : (n, \bar{\alpha}) \in D\}.$$

From this it follows that the set of equivalence classes of pairs  $(n, \bar{\alpha}) \in P_{\bar{x},Y}$  for which  $E \in B_{n,\bar{\alpha}}$  is a  $\text{HOD}_{\bar{x}}$ -generic filter for  $\mathbb{V}_{\bar{x},Y}$ . Let  $G_E$  denote this filter. The definition just given shows that  $G_E$  is in  $\text{HOD}_{\bar{x} \smallfrown \langle E \rangle}$  and therefore that  $\text{HOD}_{\bar{x} \smallfrown \langle G_E \rangle}$  is contained in  $\text{HOD}_{\bar{x} \smallfrown \langle E \rangle}$ .

The set  $K$  consisting of those pairs  $(y, (n, \bar{\alpha})) \in Y \times P_{\bar{x},Y}$  for which

$$B_{n,\bar{\alpha}} = \{C \subseteq Y : y \in C\}$$

is also a member of  $\text{HOD}_{\bar{x}}$ . Since  $E$  is equal to

$$\{y \in Y : \exists [n, \bar{\alpha}]_{\bar{x},Y} \in G_E (y, (n, \bar{\alpha})) \in K\},$$



it follows that  $E$  is in  $\text{HOD}_{\bar{x}}[G_E]$ . Since  $\text{HOD}_{\bar{x}}[G_E] \subseteq \text{HOD}_{\bar{x} \cap \langle G_E \rangle}$ , it follows also that  $E$  is in  $\text{HOD}_{\bar{x} \cap \langle G_E \rangle}$ ,  $\text{HOD}_{\bar{x} \cap \langle E \rangle} = \text{HOD}_{\bar{x} \cap \langle G_E \rangle}$  and that each of these models contains  $\text{HOD}_{\bar{x}}[G_E]$ .

To see that  $\text{HOD}_{\bar{x} \cap \langle E \rangle} \subseteq \text{HOD}_{\bar{x}}[G_E]$ , it suffices to see that every set of ordinals in  $\text{HOD}_{\bar{x} \cap \langle E \rangle}$  is in  $\text{HOD}_{\bar{x}}[G_E]$ , since  $\text{HOD}_{\bar{x} \cap \langle E \rangle}$  satisfies Choice. Fix then an ordinal  $\zeta$  and a set  $Q \subseteq \zeta$  in  $\text{HOD}_{\bar{x} \cap \langle E \rangle}$ . Fix a quaternary formula  $\varphi$  and a finite set of ordinals  $\bar{\alpha}$  such that  $Q = \{\xi \in \zeta : \varphi(\xi, \bar{x}, \bar{\alpha}, E)\}$ . Let  $T$  be

$$\{(\xi, (n, \bar{\beta})) \in \zeta \times P_{\bar{x}, Y} : B_{n, \bar{\beta}} = \{C \subseteq Y : \varphi(\xi, \bar{x}, \bar{\alpha}, C)\}\}.$$

Then  $T$  is in  $\text{HOD}_{\bar{x}}$ , and  $Q$ , which is

$$\{\xi \in \zeta : \exists [n, \bar{\beta}]_{\bar{x}, Y} \in G_E (\xi, (n, \bar{\beta})) \in T\},$$

is in  $\text{HOD}_{\bar{x}}[G_E]$ . □

**10.1.3 Remark.** Theorem 10.1.2 does not imply that  $G_E$  is in  $\text{HOD}_{\bar{x}}[E]$ . Consider the case where  $Y = \omega$  and  $\bar{x} = \langle A \rangle$ , for some  $A \subseteq 2^\omega$ . By Theorem 10.1.5 below,  $\text{HOD}_{\langle A \rangle}^{L(A, \mathbb{R})}$  is equal to  $L[B]$ , for some set of ordinals  $B$ . The set  $S$  of subsets of  $\omega$  whose characteristic functions are in  $A$  is in  $\text{HOD}_{\langle A \rangle}^{L(A, \mathbb{R})}$ . If  $G_E$  were uniformly definable from  $E \subseteq \omega$  in  $\text{HOD}_{\langle A \rangle}^{L(A, \mathbb{R})}[E]$ , then this definition would give an  $\infty$ -Borel\* code for  $A$ , along with  $B$  and the ordinal index of  $S$  in the constructibility order on  $L[B]$  relative to  $B$ . Remark 8.1.6 shows that  $\text{ZF} + \text{DC}$  does not prove that every set of reals is  $\infty$ -Borel.

**10.1.4 Remark.** Theorem 9.1.3 shows that, assuming  $\text{CC}_{\mathbb{R}}$  and that  $S$  is a set of ordinals for which  $\leq_S$  is locally countable and  $\mu_S$  is an ultrafilter on  $\mathcal{D}_S$ ,

$$L[S, x] \models \text{CH} + 2^{\aleph_1} = \aleph_2$$

for an  $S$ -cone of  $x \in \omega^\omega$ . Fix such an  $S$  and  $x$  and work in  $L[S, x]$  for the rest of this paragraph. Since  $\text{CH}$  and  $2^{\aleph_1} = \aleph_2$  hold, we have that  $\mathbb{V}_{\langle S \rangle, \omega}$  has cardinality at most  $\aleph_2$ . One can show that the cardinality of  $\mathbb{V}_{\langle S \rangle, \omega}$  is exactly  $\aleph_2$ , using sets of the form  $\{(w, y, z) \in (\omega^\omega)^3 : \pi_{\alpha, w}(y) < \pi_{\alpha, w}(z)\}$ , where  $\pi_{\alpha, w}$  is the least bijection between  $\omega^\omega$  and  $\alpha$  ordinal definable from  $S$  and  $w$ , in the natural definability order. For each  $w \in \mathcal{P}(\omega^\omega)$ , let  $<_w$  be the constructibility order on  $\mathcal{P}(\omega^\omega) \cap L[S, w]$  using  $S$  and  $w$  and parameters, and for each  $\alpha < \omega_2$ , let  $A_{\alpha, w}$  be the collection of subsets of  $\omega^\omega$  in  $L[S, w]$  of  $<_w$ -rank less than  $\alpha$ . Since  $\text{CH}$  holds, each set of the form  $\bigcup \{A_{\alpha, w} : w \in \omega^\omega\}$  has cardinality at most  $\aleph_1$ . It follows that in  $\text{HOD}_{\langle S \rangle}$ ,  $\mathbb{V}_{\langle S \rangle, \omega}$  is a union of  $\omega_2^{L[S, x]}$  many sets of cardinality less than  $\omega_2^{L[S, x]}$ , so the cardinality of  $\mathbb{V}_{\langle S \rangle, \omega}$  (again, as computed in  $\text{HOD}_{\langle S \rangle}$ ) is exactly  $\omega_2^{L[S, x]}$ .

Since  $x$  is generic over  $\text{HOD}_{\langle S \rangle}^{L[S, x]}$  via  $\mathbb{V}_{\langle S \rangle, \omega}^{L[S, x]}$  by Theorem 10.1.2, it follows that the models  $\text{HOD}_{\langle S \rangle}^{L[S, x]}$  and  $L[S, x]$  have the same cardinals above and including  $\omega_2^{L[S, x]}$ .

The following theorem, due to Woodin, shows that certain models of the form  $\text{HOD}_{\bar{x}}$  have the form  $L[B]$ , for  $B$  a set of ordinals. The case where  $F = \omega^\omega$  and  $S \subseteq \text{Ord}$  was done first by Kechris and Woodin. The proof the theorem uses parts of the proof of Theorem 10.1.2.

**Theorem 10.1.5.** *Suppose that  $V = L(F)$ , for some transitive set  $F$ , and let  $S$  be any set in  $L(F)$ . Then  $\text{HOD}_{\langle S \rangle} = L[B]$ , for some set  $B$  of ordinals.*

Before proving Theorem 10.1.5, we prove a lemma (a variation of Theorem 9.3.7), and then prove the theorem in the case  $V = L[F]$ , for  $F$  a set of ordinals.

**Lemma 10.1.6.** *Let  $M_1$  and  $M_2$  be transitive models of ZFC, with  $M_1 \subseteq M_2$ . Suppose that  $\mathbb{P}$  is a partial order in  $M_1$ ,  $G \subseteq \mathbb{P}$  is an  $M_2$ -generic filter and  $M_2 \subseteq M_1[G]$ . Then  $M_1 = M_2$ .*

*Proof.* Since  $M_1$  and  $M_2$  are models of ZFC, it is enough to see that every set of ordinals in  $M_2$  is in  $M_1$ . Letting  $X$  be a set of ordinals in  $M_2$ , we have that  $X = \tau_G$ , for some  $\mathbb{P}$ -name  $\tau$  in  $M_1$ . Since  $G$  is  $M_2$ -generic, this means that some condition in  $\mathbb{P}$  decides all of  $\tau$ , so  $X$  is in  $M_1$ .  $\square$

Now, suppose that  $V = L[F]$ , where  $F$  is a subset of an ordinal  $\gamma$ . Since Choice holds in  $\text{HOD}_{\langle S \rangle}$ , there exist in  $\text{HOD}_{\langle S \rangle}$  an ordinal  $\eta$  and bijection  $\pi: \mathbb{V}_{\langle S \rangle, \gamma} \rightarrow \eta$ . Let  $\leq_\gamma$  be the partial order on  $\eta$  induced by  $\pi$ . Let  $K$  be the set of pairs

$$(\delta, (n, \bar{\alpha})) \in \gamma \times P_{\langle S \rangle, \gamma}$$

such that  $B_{n, \bar{\alpha}} = \{C \subseteq \gamma : \delta \in C\}$  (i.e., the corresponding set  $K$  from the proof of Theorem 10.1.2 with  $\langle S \rangle$  as  $\bar{x}$  and  $\gamma$  as  $Y$ ), and let  $K_\gamma$  be the  $\pi$ -image of  $K$ , that is,

$$\{(\delta, \pi([(n, \bar{\alpha})]_{\langle S \rangle, \gamma})) : (\delta, (n, \bar{\alpha})) \in K\}.$$

Then  $\leq_\gamma$  and  $K_\gamma$  are in  $\text{HOD}_{\langle S \rangle}$ . By Theorem 10.1.2,  $F$  is  $\mathbb{V}_{\langle S \rangle, \gamma}$ -generic over  $\text{HOD}_{\langle S \rangle}$ , via the generic filter  $G_F$ . Then  $\pi[G_F]$  is  $\text{HOD}_{\langle S \rangle}$ -generic for  $\leq_\gamma$ , and  $F$  is in  $L[\leq_\gamma, K_\gamma][G_F]$ . Then

$$V = L[F] = \text{HOD}_{\langle S \rangle}[G_F] = L[\leq_\gamma, K_\gamma][\pi[G_F]],$$

which implies that  $\text{HOD}_{\langle S \rangle} = L[\leq_\gamma, K_\gamma]$ , by Lemma 10.1.6, with  $L[\leq_\gamma, K_\gamma]$  as  $M_1$  and  $\text{HOD}_{\langle S \rangle}$  as  $M_2$ .

We introduce a new partial order to deal with the general case. Recall that, given an infinite set  $Y$ ,  $\text{Col}^*(\omega, Y)$  is the partial order of finite partial injections from  $\omega$  to  $Y$  (each with domain some  $n \in \omega$ ), ordered by extension. Given a filter  $G \subseteq \text{Col}^*(\omega, Y)$ , let  $a_G = \{(i, j) \in \omega \times \omega : g(i) \in g(j)\}$ , where  $g = \bigcup G$ . Then  $a_G$  is a subset of  $\omega \times \omega$ , and, if  $Y$  is transitive,  $V[G] = V[a_G]$ .

As above, suppose that  $\bar{x}$  is a finite sequence and  $Y$  is an (infinite) set such that  $Y$  is in  $\text{OD}_{\bar{x}}$ . Let  $\kappa$  be the least cardinal  $\lambda$  such that  $\bar{x} \in V_\lambda$ ,  $Y$  is in  $\text{OD}_{\bar{x}}^{V_\lambda}$  and every subset of  $\mathcal{P}(\text{Col}^*(\omega, Y))$  in  $\text{OD}_{\bar{x}}$  is in  $\text{OD}_{\bar{x}}^{V_\lambda}$ . Let  $P_{\bar{x}, Y}^\omega$  be the set of triples  $(n, m, \bar{\alpha})$  such that

- $n \in \omega$ ,
- $m$  is the Gödel number of a ternary formula  $\varphi$ ,
- $\bar{\alpha}$  is a finite sequence of elements of  $\kappa$  and
- the set  $B_{n,m,\bar{\alpha}} = \{p \in \text{Col}^*(\omega, Y) : \text{dom}(p) = n, V_\kappa \models \varphi(p, \bar{x}, \bar{\alpha})\}$  is nonempty.

Given  $(n, m, \bar{\alpha}) \in P_{\bar{x}, Y}^\omega$  and  $k \leq n$ , let  $B_{n,m,\bar{\alpha}} \upharpoonright k$  denote the set

$$\{p \upharpoonright k : p \in B_{n,m,\bar{\alpha}}\}.$$

This set is evidently also in  $\text{HOD}_{\bar{x}}$ . The set  $P_{\bar{x}, Y}^\omega$  is in  $\text{HOD}_{\bar{x}}$ , as is the reflexive and transitive ordering  $\leq_{\bar{x}, Y}^\omega$  on  $P_{\bar{x}, Y}^\omega$  defined by setting

$$(n, m, \bar{\alpha}) \leq_{\bar{x}, Y}^\omega (k, j, \bar{\beta})$$

to hold if  $n \geq k$  and

$$B_{n,m,\bar{\alpha}} \upharpoonright k \subseteq B_{k,j,\bar{\beta}}.$$

Let  $\equiv_{\bar{x}, Y}^\omega$  be the induced equivalence relation on  $P_{\bar{x}, Y}^\omega$ , and let  $[n, m, \bar{\alpha}]_{\bar{x}, Y}^\omega$  denote the  $\equiv_{\bar{x}, Y}^\omega$ -class of a tuple  $(n, m, \bar{\alpha})$ . Let  $\mathbb{V}_{\bar{x}, Y}^\omega$  be the partial order whose domain is the set of  $\equiv_{\bar{x}, Y}^\omega$ -classes of  $P_{\bar{x}, Y}^\omega$ , with the order inherited from  $\leq_{\bar{x}, Y}^\omega$ . Each injection  $f: \omega \rightarrow Y$  defines a filter  $H_f$  on  $\mathbb{V}_{\bar{x}, Y}^\omega$ , consisting of the  $\equiv_{\bar{x}, Y}^\omega$ -classes of those  $(n, m, \bar{\alpha}) \in P_{\bar{x}, Y}^\omega$  for which  $f \upharpoonright n$  is in  $B_{n,m,\bar{\alpha}}$ .

In the following lemma we require only that  $Y \in \text{OD}_{\bar{x}}$ , whereas the hypotheses of Theorem 10.1.2 have  $Y \in \text{HOD}_{\bar{x}}$ . This distinction is essential for the proofs of Theorems 10.1.5 and 10.1.8.

**Lemma 10.1.7.** *For any set finite sequence  $\bar{x}$  with  $\bar{x} \in \text{OD}_{\bar{x}}$ , and any infinite set  $Y$  in  $\text{OD}_{\bar{x}}$ , if  $G \subseteq \text{Col}^*(\omega, Y)$  is a  $V$ -generic filter, and  $g = \bigcup G$ , then  $H_g$  is  $\mathbb{V}_{\bar{x}, Y}^\omega$ -generic over  $\text{HOD}_{\bar{x}}$ , and  $a_g$  is in  $\text{HOD}_{\bar{x}}[H_g]$ .*

*Proof.* To see that  $H_g$  is  $\mathbb{V}_{\bar{x}, Y}^\omega$ -generic over  $\text{HOD}_{\bar{x}}$ , let  $D$  be a dense open subset of  $\mathbb{V}_{\bar{x}, Y}^\omega$  in  $\text{HOD}_{\bar{x}}$ , and let  $p$  be a condition in  $\text{Col}^*(\omega, Y)$ . Let  $D^0$  be the set of  $(n, m, \bar{\alpha}) \in P_{\bar{x}, Y}^\omega$  with  $[(n, m, \bar{\alpha})]_{\bar{x}, Y}^\omega \in D$ . Applying the genericity of  $G$ , it suffices to find a condition  $p' \leq p$  and a tuple  $(n, m, \bar{\alpha})$  in  $D^0$  such that  $p'$  is in  $B_{n,m,\bar{\alpha}}$ . Let  $n_p$  be the domain of  $p$ . The set

$$\bigcup \{B_{n,m,\bar{\alpha}} \upharpoonright n_p : (n, m, \bar{\alpha}) \in D^0, n \geq n_p\},$$

being in  $\text{OD}_{\bar{x}}$ , must be the set of injections from  $n_p$  to  $Y$ , since otherwise the complement of this set has the form  $B_{n_p,j,\bar{\beta}}$  for some  $j$  and  $\bar{\beta}$ , and we get a contradiction by considering a  $(n, m, \bar{\alpha}) \in D^0$  with  $B_{n,m,\bar{\alpha}} \upharpoonright n_p \subseteq B_{n_p,j,\bar{\beta}}$ . It follows that there exist  $p' \leq p$  and  $(n, m, \bar{\alpha})$  as desired.

As in the proof of Theorem 10.1.2, let  $K$  be the set of pairs

$$((i, j), (n, m, \bar{\alpha})) \in (\omega \times \omega) \times P_{\bar{x}, Y}^\omega$$

such that  $\{i, j\} \subseteq n$  and  $p(i) \in p(j)$  for all  $p \in B_{n,m,\bar{\alpha}}$ . Then again  $K$  is in  $\text{HOD}_{\bar{x}}$ , and, as

$$a_g = \{(i, j) \in \omega \times \omega : \exists [n, m, \bar{\alpha}]_{\bar{x}, Y}^\omega \in H_g((i, j), (n, m, \bar{\alpha})) \in K\},$$

$a_g$  is in  $\text{HOD}_{\bar{x}}[H_g]$ .  $\square$

Theorem 10.1.5 now follows.

*Proof of Theorem 10.1.5.* We may assume that  $F = V_\alpha$ , for some infinite ordinal  $\alpha$  (so that  $F$  is ordinal definable). In  $\text{HOD}_{\langle S \rangle}$ , there exist an ordinal  $\gamma$  and bijection  $\pi: \mathbb{V}_{\langle S \rangle, F}^\omega \rightarrow \gamma$ . Let  $\leq_\gamma$  be the partial order on  $\gamma$  induced by  $\pi$ . Let  $K$  be the set of pairs

$$((i, j), (n, m, \bar{\alpha})) \in (\omega \times \omega) \times P_{\langle S \rangle, F}^\omega$$

such that  $\{i, j\} \subseteq n$  and  $p(i) \in p(j)$  for all  $p \in B_{n,m,\bar{\alpha}}$  (i.e., the set  $K$  from the proof of Lemma 10.1.7 with  $F$  as  $Y$  and  $\langle S \rangle$  as  $\bar{x}$ ), and let  $K_\gamma$  be the  $\pi$ -image of  $K$ , that is,

$$\{((i, j), \pi([n, m, \bar{\alpha}]_{\langle S \rangle, F}^\omega)) : ((i, j), (n, m, \bar{\alpha})) \in K\}.$$

Let  $G_0 \subseteq \text{Col}^*(\omega, F)$  be a  $V$ -generic filter and let  $g = \bigcup G_0$ . Then  $\pi[H_g]$  is a  $\text{HOD}_{\langle S \rangle}$ -generic filter for  $\leq_\gamma$ . As in the proof of Lemma 10.1.7,  $a_g$  is in the model  $L[\leq_\gamma, K_\gamma][\pi[H_g]]$ , and therefore so is  $F$ . Applying Lemma 10.1.6 with  $L[\leq_\gamma, K_\gamma]$  as  $M_1$ ,  $\text{HOD}_{\langle S \rangle}$  as  $M_2$ ,  $\leq_\gamma$  as  $\mathbb{P}$  and  $\pi[H_g]$  as  $G$ , we get that  $L[\leq_\gamma, K_\gamma] = \text{HOD}_{\langle S \rangle}$ . As the model  $L[\leq_\gamma, K_\gamma]$  has the form  $L[B]$ , for some set  $B$  of ordinals, we are done.  $\square$

The proof of the following theorem uses the following standard facts about sharps: (1) if  $\kappa$  is a measurable cardinal then the sharp of each bounded subset of  $\kappa$  exists; (2) if  $X$  is a set such that  $X^\#$  exists, and  $Y$  exists in a set-generic forcing extension of  $L[X]$ , then  $Y^\#$  exists. Unfortunately, we do not know of a reference for basic properties of sharps of arbitrary sets. The second fact just listed follows from the fact that, for an arbitrary set  $X$ ,  $X^\#$  exists if and only if, for arbitrarily large ordinals  $\gamma$ ,  $\gamma$  is the critical point of an elementary embedding from  $L(X)$  to itself.

**Theorem 10.1.8** (ZF + AD). *Suppose that  $A \subseteq \omega^\omega$  and  $\mathcal{P}(\omega^\omega) \not\subseteq L(A, \mathbb{R})$ . Then  $A^\#$  exists.*

*Proof.* Every  $\text{OD}_{\langle A \rangle}$  subset of  $\omega^\omega$  in  $L(A, \mathbb{R})$  is definable in  $L(A, \mathbb{R})$  from  $A$  and an element of  $\Theta$ . It follows that the partial order  $\mathbb{V}_{\langle A \rangle, H(\aleph_1) \cup \{A\}}^\omega$  as computed in the model  $L(A, \mathbb{R})$  has cardinality at most  $\Theta^{L(A, \mathbb{R})}$  in  $L(A, \mathbb{R})$ , and moreover from the proof of Theorem 10.1.5 that  $\text{HOD}_{\langle A \rangle}^{L(A, \mathbb{R})}$  is equal to  $L[B]$  for some  $B \subseteq \Theta^{L(A, \mathbb{R})}$  such that (by Lemma 10.1.7)  $A$  exists in a extension of  $L[B]$  via a generic filter for  $\mathbb{V}_{\langle A \rangle, H(\aleph_1) \cup \{A\}}^\omega$  induced by any  $V$ -generic filter for  $\text{Col}^*(\omega, H(\aleph_1) \cup \{A\})$ . Since  $\Theta$  is both greater than  $\Theta^{L(A, \mathbb{R})}$  and a limit of measurable cardinals (by Remark 5.0.2 or Theorem 9.2.1, for instance),  $B^\#$  exists. It follows that  $A^\#$  exists in any forcing extension of  $V$  by  $\text{Col}^*(\omega, H(\aleph_1) \cup \{A\})$ . Since it must exist as the same set in any such extension,  $A^\#$  exists already in  $V$ .  $\square$

## 10.2 Codes for projections, and Uniformization

In this section we prove that, assuming  $\text{TD} + \text{DC}_{\mathbb{R}}$ , the set of  $\infty$ -Borel sets is closed under projections (Theorem 10.2.3), and we derive some consequences for Uniformization (Theorem 10.2.6, for instance). We use the results of Chapter 9 and the notation  $\leq_S$ ,  $\mu_S$ ,  $\mathcal{D}_S$ , etc. introduced in Definition 9.1.1. Theorem 1.2.2 shows that  $\text{AD}$  implies that, for each set of ordinals  $S$ ,  $\mu_S$  is an ultrafilter on the corresponding set  $\mathcal{D}_S$ ; if  $\text{CC}_{\mathbb{R}}$  holds (as it does under  $\text{AD}$ ) then  $\mu_S$  is countably complete. If there exists any set of ordinals  $S$  for which  $\mu_S$  is a countably complete ultrafilter on  $\mathcal{D}_S$ , then  $\omega_1^V$  is measurable (see Remark 1.2.7), so  $\aleph_1 \not\leq 2^{\aleph_0}$ .

Since Lemma 10.2.1 below seems to hold only for  $\infty$ -Borel subsets of  $\omega^\omega$  (as opposed to  $\omega^\delta$  for larger  $\delta$ ), this section concentrates on  $\infty$ -Borel sets of reals. For each set  $S$  of ordinals, and each  $x \in \omega^\omega$ , let  $Q_x^S$  denote the definably least poset on the ordinals isomorphic in  $\text{HOD}_S^{L[S,x]}$  to the poset  $\mathbb{V}_{\langle S \rangle, \omega}^{L[S,x]}$  (via the definability order on  $\text{HOD}_S^{L[S,x]}$ , say), and let  $K_x^S$  (also a set of pairs of ordinals) denote the corresponding version of the set  $K$  from the proof of Theorem 10.1.2 (i.e., relabeled as in the proof of Theorem 10.1.5 to refer to  $Q_x^S$ ). Note that  $Q_x^S$  depends only on  $[x]_S$ . We will use this notation throughout this section.

Theorem 10.2.2 below shows that if  $\text{DC}_{\mathbb{R}}$  holds and  $\mu_S$  is an ultrafilter for each set of ordinals  $S$  then the collection of  $\infty$ -Borel sets of reals is projectively closed. We start with a lemma whose hypothesis is weaker (and appeal to the fact that  $\aleph_1 \not\leq 2^{\aleph_0}$  is equivalent to the assertion that  $\omega_1^V$  is strongly inaccessible in every inner model satisfying Choice). Background material on some of the forcing claims (e.g., involving regular embeddings) made in the last paragraph of the proof can be found in the appendix to [27].

**Lemma 10.2.1** ( $\text{ZF} + \aleph_1 \not\leq 2^{\aleph_0}$ ). *Let  $S$  be a set of ordinals, let  $\varphi$  be a ternary formula and let  $B$  denote the set*

$$\{x \in \omega^\omega : \exists y \in \omega^\omega L[S, x, y] \models \varphi(S, x, y)\}.$$

*Then the following are equivalent, for each  $x \in \omega^\omega$ .*

1.  $x \in B$
2. *For some  $w \in \omega^\omega$ , for all  $z \geq_S w$ ,  $L[S, Q_z^S, K_z^S, x] \models$  “there is some  $p \in \text{Col}(\omega, 2^{Q_z^S})$  forcing that there exists a  $y \in \omega^\omega$  such that  $L[S, x, y] \models \varphi(S, x, y)$ .”*
3. *For some  $z \in \omega^\omega$ ,  $L[S, Q_z^S, K_z^S, x] \models$  “there is some  $p \in \text{Col}(\omega, 2^{Q_z^S})$  forcing that there exists a  $y \in \omega^\omega$  such that  $L[S, x, y] \models \varphi(S, x, y)$ .”*

*Proof.* That (2) implies (3) is immediate. To see that (3)  $\Rightarrow$  (1), fix one such  $z$ . The cardinality of  $Q_z^S$  in  $L[S, z]$  is at most  $(2^{2^{\aleph_0}})^{L[S, z]}$ , which is below  $\omega_1^V$ , as  $\omega_1^V$  is strongly inaccessible in  $L[S, z]$ . Since  $\omega_1^V$  is strongly inaccessible in  $L[S, Q_z^S, K_z^S, x]$ ,

$$\mathcal{P}(Q_z^S) \cap L[S, Q_z^S, K_z^S, x]$$

is countable, and  $L[S, Q_z^S, K_z^S, x]$ -generic filters for  $Q_z^S$  exist below each condition. Since forcing with  $\text{Col}(\omega, 2^{Q_z^S})$  adds a generic filter for  $Q_z^S$ , statement (1) follows.

For (1)  $\Rightarrow$  (2), fix  $y_0 \in \omega^\omega$  such that  $L[S, x, y_0] \models \varphi(S, x, y_0)$ , and  $z \in \omega^\omega$  such that  $z \geq_S y_0$  and  $z \geq_S x$ . Letting  $G_z \subseteq Q_z^S$  be the filter induced by  $z$  as in the proof of Theorem 10.1.2,  $G_z$  is  $\text{HOD}_{\langle S \rangle}^{L[S, z]}$ -generic. It follows that  $G_z$  is  $Q_z^S$ -generic over  $L[S, Q_z^S, K_z^S]$  and (using  $K_z^S$ ) that  $z$  is a member of  $L[S, Q_z^S, K_z^S][G_z]$ . Since  $x$  is in  $L[S, Q_z^S, K_z^S][G_z]$ ,  $L[S, Q_z^S, K_z^S, x]$  is a generic extension of  $L[S, Q_z^S, K_z^S]$ , and  $L[S, Q_z^S, K_z^S][G_z]$  is a generic extension of  $L[S, Q_z^S, K_z^S, x]$  by a partial order of cardinality at most  $(2^{|Q_z^S|})^{L[S, Q_z^S, K_z^S]}$  in  $L[S, Q_z^S, K_z^S, x]$ . This partial order regularly embeds into  $\text{Col}(\omega, 2^{Q_z^S})$  in  $L[S, Q_z^S, K_z^S, x]$ . This gives (2).  $\square$

If DC holds in  $L(S, \mathbb{R})$  (as it does if  $\text{DC}_{\mathbb{R}}$  holds), then for each set  $S$  of ordinals the ultrapower

$$V^{\mathcal{D}_S} / \mu_S$$

(formed using all  $S$ -invariant functions in  $L(S, \mathbb{R})$ ) is wellfounded in  $L(S, \mathbb{R})$ . We let  $S^\infty$ ,  $Q_S^\infty$  and  $K_S^\infty$  be the sets in  $\prod_{z \in \omega^\omega} V / \mu_S$  represented by the functions  $z \mapsto S$  and  $z \mapsto Q_z^S$  and  $z \mapsto K_z^S$ , respectively.

**Theorem 10.2.2** (ZF +  $\text{DC}_{\mathbb{R}}$ ). *Let  $S$  be a set of ordinals such that  $\leq_S$  is locally countable and  $\mu_S$  is an ultrafilter. Let  $\varphi$  be a ternary formula, let  $S$  be a set of ordinals, and let  $B$  be*

$$\{x \in \omega^\omega : \exists y \in \omega^\omega L[S, x, y] \models \varphi(S, x, y)\}.$$

*Then there exist a set of ordinals  $T$  in  $\text{OD}_{\langle S \rangle}$  and a binary formula  $\psi$  such that*

$$B = \{x \in \omega^\omega : L[T, x] \models \psi(T, x)\}.$$

*Proof.* By  $\text{DC}_{\mathbb{R}}$ ,  $\mu_S$  is countably complete. Since  $\mu_S$  is an ultrafilter,  $\omega_1^V$  is a measurable cardinal, which implies that  $\aleph_1 \not\leq 2^{\aleph_0}$  (as in Remark 1.2.7). We work in  $L(S, \mathbb{R})$ , which satisfies DC by the assumption that  $V$  satisfies  $\text{DC}_{\mathbb{R}}$ .

Since DC holds in  $L(S, \mathbb{R})$ , for each  $x \in \omega^\omega$  the ultraproduct

$$\prod_{z \in \omega^\omega} L[S, Q_z^S, K_z^S, x] / \mu_S$$

is wellfounded in  $L(S, \mathbb{R})$ . Furthermore, for each  $x \in \omega^\omega$ ,

$$L[S^\infty, Q_S^\infty, K_S^\infty, x] = \prod_{z \in \omega^\omega} L[S, Q_z^S, K_z^S, x] / \mu_S$$

and, by Lemma 10.2.1,  $x$  is in  $B$  if and only if  $L[S^\infty, Q_S^\infty, K_S^\infty, x]$  satisfies the statement “there is some  $p \in \text{Col}(\omega, 2^{Q_S^\infty})$  forcing that

$$\exists y \in \omega^\omega L[S^\infty, x, y] \models \varphi(S^\infty, x, y).”$$

Then we can let  $T$  be set of the ordinals coding the triple  $(S^\infty, Q_S^\infty, K_S^\infty)$  under some fixed coding in  $L$  of triples of ordinals by ordinals, and let  $\psi$  be the corresponding version of the statement just given.  $\square$

The following is an immediate consequence of Theorem 10.2.2 and Corollary 1.2.6, which implies that, assuming TD, the assumptions on  $S$  in the hypotheses of Theorem 10.2.2 hold for any set  $S$  of ordinals.

**Theorem 10.2.3** (ZF + TD +  $\text{DC}_\mathbb{R}$ ). *The set of  $\infty$ -Borel subsets of  $(\omega^\omega)^{<\omega}$  is projectively-closed.*

**10.2.4 Remark.** Since  $T$  can be taken to be in  $\text{OD}_{\langle S \rangle}$  in the conclusion of Theorem 10.2.2, TD +  $\text{DC}_\mathbb{R}$  implies that there is an ordinal definable class-sized function (minimizing in the ordinal definability order from  $S$ ) taking in  $\infty$ -Borel\* codes  $(S, \varphi)$  for subsets of  $(\omega^\omega)^2$  and returning the corresponding  $\infty$ -Borel\* codes for  $\{x \in \omega^\omega : \exists y \in \omega^\omega L[S, x, y] \models \varphi(S, x, y)\}$ .

Theorem 10.2.6 below is a uniformization result derived from Lemma 10.2.1 and Theorem 10.2.2. Corollary 10.2.7 connects Uniformization with the nonexistence of a  $\leq_{\mathcal{D}}$ -maximal set of ordinals. Recall that for a set  $A \subseteq (\omega^\omega)^2$  and  $y \in \omega^\omega$ ,  $A_y$  denotes the cross-section  $\{z \in \omega^\omega : (y, z) \in A\}$ . As defined in Section 6.2, such a set  $A$  can be uniformized if there is a function  $f: \{y \in \omega^\omega : A_y \neq \emptyset\} \rightarrow \omega^\omega$  such that  $(y, f(y)) \in A$  for all  $y \in \text{dom}(f)$ . Recall also that, for  $x, y \in \omega^\omega$ ,  $x \oplus y$  denotes the element  $z$  of  $\omega^\omega$  such that, for all  $n \in \omega$ ,  $z(2n) = x(n)$  and  $z(2n+1) = y(n)$ .

The proof of Theorem 10.2.6 uses the following observation.

**Lemma 10.2.5.** *Suppose that  $A$  and  $B$  are sets, with  $A \subseteq (\omega^\omega)^2$ . Suppose that for a Turing cone of  $x \in \omega^\omega$ , for all  $y \in L[x]$ , if  $A_y \neq \emptyset$  then  $A_y \cap \text{OD}_{\{B, x\}} \neq \emptyset$ . Then  $A$  can be uniformized.*

*Proof.* Let  $x_0$  be a base for a cone witnessing the relevant assumption of the lemma. Given  $y$  such that  $A_y \neq \emptyset$ , define  $f(y)$  to be the  $\text{OD}_{\{B, x_0, y\}}$ -least  $z \in \text{OD}_{\{B, x_0 \oplus y\}}$  with  $(y, z) \in A$ .  $\square$

**Theorem 10.2.6** (ZF + TD +  $\text{DC}_\mathbb{R}$ ). *Suppose that  $(S, \varphi)$  is an  $\infty$ -Borel\* code for a set  $A \subseteq (\omega^\omega)^2$ . Let  $B$  be a set such that  $S \in \text{OD}_{\{B\}}$  and suppose that the  $\mu_S$ -ultrapower of the ordinals is wellfounded. Suppose that, for a Turing cone of  $x \in \omega^\omega$ ,  $\text{OD}_{\{B, x\}} \cap \omega^\omega \not\subseteq L[S^\infty, Q_S^\infty, K_S^\infty, x]$ . Then  $A$  can be uniformized.*

*Proof.* Again, TD implies that  $\aleph_1 \not\leq 2^{\aleph_0}$ . Let  $A^*$  be the set of  $y \in \omega^\omega$  with  $A_y \neq \emptyset$ . We prove that for a Turing cone of  $x \in \omega^\omega$ , for all  $y \in L[x] \cap A^*$ ,  $A_y \cap \text{OD}_{\{B, x\}} \neq \emptyset$ . The theorem then follows from Lemma 10.2.5. Let  $x_0 \in \omega^\omega$  be such that for all  $x \geq_T x_0$ ,

$$\text{OD}_{\{B, x\}} \cap \omega^\omega \not\subseteq L[S^\infty, Q_S^\infty, K_S^\infty, x].$$

Applying TD, suppose toward a contradiction that for a Turing cone of  $x$  there exists a  $y \in L[x] \cap A^*$  with  $A_y \cap \text{OD}_{\{B, x\}} = \emptyset$ . Let  $x_1 \geq_{\text{Tu}} x_0$  be a base for a Turing cone witnessing this.

By Lemma 10.2.1 (and the homogeneity of  $\text{Col}(\omega, Q_S^\infty)$  to remove the reliance on the condition  $p$  given there), for every  $y \in A^*$ ,

$$L[S^\infty, Q_S^\infty, K_S^\infty, y] \models \text{"}V^{\text{Col}(\omega, Q_S^\infty)} \models \exists z L[S, y, z] \models \varphi(S, y, z)\text{"}.$$

Applying Lemma 9.3.5 and the discussion just before it, let  $\pi: 2^{<\omega} \rightarrow [\omega]^{<\omega}$  be a recursive diffuse function and let  $U$  be, in the model  $L[S^\infty, Q_S^\infty, K_S^\infty, x_1]$ , an ultrafilter on  $\omega$  not containing any  $\pi$ -weak set. Let  $\mathbb{P}$  denote the partial order  $\mathbb{P}_U^{L[S^\infty, Q_S^\infty, K_S^\infty, x_1]}$ , again as in Section 9.3. Fix

$$t \in \text{OD}_{\{B, x_1\}} \cap \omega^\omega \setminus L[S^\infty, Q_S^\infty, K_S^\infty, x_1].$$

For any  $L[S^\infty, Q_S^\infty, K_S^\infty, x_1]$ -generic

$$g \subseteq \mathbb{P},$$

and any

$$y \in L[S^\infty, Q_S^\infty, K_S^\infty, x_1][g] \cap A^*,$$

$$L[S^\infty, Q_S^\infty, K_S^\infty, x_1][g] \models \text{"}V^{\text{Col}(\omega, Q_S^\infty)} \models \exists z \in \omega^\omega L[S, y, z] \models \varphi(S, y, z)\text{"},$$

since  $L[S^\infty, Q_S^\infty, K_S^\infty, x_1][g]$  is an outer model of  $L[S^\infty, Q_S^\infty, K_S^\infty, y]$ .

We have that

$$L[S^\infty, Q_S^\infty, K_S^\infty, x_1] = \prod L[S, Q_x^S, K_x^S, x_1] / \mu_S.$$

Since  $\aleph_1 \not\leq 2^{\aleph_0}$ ,  $L[S^\infty, Q_S^\infty, K_S^\infty, x_1] \cap \mathcal{P}(\mathcal{P}(\omega^\omega))$  is countable, and (by Remark 9.0.1) there is an  $x_2 \geq_{\text{Tu}} x_1$  in  $\omega^\omega$  such that, for all  $x \geq_{\text{Tu}} x_2$ ,

1.  $L[S^\infty, Q_S^\infty, K_S^\infty, x_1] \cap \mathcal{P}(\mathcal{P}(\omega^\omega)) = L[S, Q_x^S, K_x^S, x_1] \cap \mathcal{P}(\mathcal{P}(\omega^\omega))$ ;
2.  $\mathbb{P} \in L[S, Q_x^S, K_x^S, x_1]$ ;
3. every nice  $\mathbb{P}$ -name in  $L[S^\infty, Q_S^\infty, K_S^\infty, x_1]$  for an element of  $\omega^\omega$  is in the model  $L[S, Q_x^S, K_x^S, x_1]$ ;
4. for any  $L[S, Q_x^S, K_x^S, x_1]$ -generic  $g \subseteq \mathbb{P}$  and any

$$y \in L[S, Q_x^S, K_x^S, x_1][g] \cap A^*,$$

$$L[S, Q_x^S, K_x^S, x_1][g] \models \text{"}V^{\text{Col}(\omega, Q_x^S)} \models \exists z \in \omega^\omega L[S, y, z] \models \varphi(S, y, z)\text{"}.$$

By Lemma 9.3.6, there exists an  $L[S^\infty, Q_S^\infty, K_S^\infty, x_1]$ -generic filter  $g^* \subseteq \mathbb{P}$ , with corresponding generic  $a^* \subseteq \omega$ , such that there is a  $w \in \omega^\omega \cap L[a^*, t]$  which HC-codes an  $\omega$ -sequence listing all the members of  $\mathcal{P}(\mathbb{P} \times Q_{x_2}^S \times \omega)$  in  $L[S, Q_{x_2}^S, K_{x_2}^S, x_1]$ .

Let  $y \in A^*$  be an element of

$$L[x_1 \oplus a^*].$$

It suffices to show that  $A_y \cap \text{OD}_{\{B, x_1 \oplus a^*\}} \neq \emptyset$ , which we now do.



Since  $t \in \text{OD}_{\{B, x_1 \oplus a^*\}}$  and  $w \in L[a^*, t]$ , there exists an  $L[S, Q_{x_2}^S, K_{x_2}^S, x_1][a^*]$ -generic filter  $G \subseteq \text{Col}(\omega, Q_{x_2}^S)$  in  $\text{OD}_{\{B, x_1, a^*\}}$ . By item (4) from the choice of  $x_2$ , there is a

$$z \in \omega^\omega \cap L[S, Q_{x_2}^S, K_{x_2}^S, x_1][a^*][G]$$

such that  $L[S, y, z] \models \varphi(S, y, z)$ . Since

$$\omega^\omega \cap L[S, Q_{x_2}^S, K_{x_2}^S, x_1][a^*][G] \subseteq \text{OD}_{\{B, x_1 \oplus a^*\}},$$

we are done.  $\square$

In Section 9.3 we defined, for sets  $S, T$  of ordinals,  $S \leq_{\mathcal{D}} T$  to mean that  $\omega^\omega \cap L[S, x] \subseteq L[T, x]$  for a Turing cone of  $x$ . Theorem 10.2.6 shows that if  $\text{TD} + \text{DC}_{\mathbb{R}}$  holds, and  $A$  is an  $\infty$ -Borel subset of  $(\omega^\omega)^2$  with an  $\infty$ -Borel code  $S$  such that some set of ordinals constructing the tuple  $(S, S^\infty, Q_S^\infty, K_S^\infty)$  is not  $\leq_{\mathcal{D}}$ -maximal, then  $A$  can be uniformized. The following corollary gives a version of the converse. Its proof shows that if  $S$  is a set of ordinals such that the set  $\{(x, y) \in (\omega^\omega)^2 : y \notin L[S, x]\}$  can be uniformized, then  $S$  is not  $\leq_{\mathcal{D}}$ -maximal.

**Corollary 10.2.7** ( $\text{ZF} + \text{TD} + \text{DC}_{\mathbb{R}}$ ). *Suppose that every subset of  $\omega^\omega$  is  $\infty$ -Borel. Then Uniformization is equivalent to the assertion that  $\leq_{\mathcal{D}}$  has no greatest element.*

*Proof.* For the reverse direction, let  $B$  be any set of ordinals strictly  $\leq_{\mathcal{D}}$ -above a set of ordinals constructing  $S^\infty$ ,  $Q_S^\infty$  and  $K_S^\infty$ , with  $S \in \text{OD}_{\{B\}}$ . Since  $\text{DC}_{\mathbb{R}}$  holds, the inner model  $L(B, \mathbb{R})$  satisfies the statement that the  $\mu_S$ -ultrapower of the ordinals is wellfounded. We can then apply Theorem 10.2.6 in  $L(B, \mathbb{R})$ .

The forward direction requires only that  $\aleph_1 \not\leq 2^{\aleph_0}$  and every subset of  $\omega^\omega$  is  $\infty$ -Borel. Let  $S$  be a set of ordinals, and let  $f: \omega^\omega \rightarrow \omega^\omega$  uniformize the set  $\{(x, y) \in (\omega^\omega)^2 : y \notin L[S, x]\}$ . Let  $(T, \varphi)$  be an  $\infty$ -Borel\* code for the set  $\{(x, i, j) : (i, j) \in f(x)\}$ . Then for all  $(x, i, j)$ ,  $(i, j) \in f(x)$  if and only if  $L[T, x] \models \varphi(T, x, i, j)$ , which shows that  $f(x) \in L[T, x] \setminus L[S, x]$  for all  $x \in \omega^\omega$ .  $\square$

### 10.3 The Vopěnka algebra for $\infty$ -Borel sets

Given a set  $Y \subseteq \text{Ord}$ , the Vopěnka algebra can be modified to use  $\infty$ -Borel\* codes in  $\text{OD}_{\{Y\}}$  to identify sets of reals. One key difference between this version and the one defined in Section 10.1 is discussed in Remark 10.3.3. Theorem 10.3.6 shows, however, assuming  $\text{TD} + \text{DC}_{\mathbb{R}}$  and that  $V = L(Y, \mathbb{R})$ , that the two versions are in fact equivalent. An application of this fact is given in part (2) of Corollary 10.3.7. Part (1) of Corollary 10.3.7 shows, again assuming  $\text{TD} + \text{DC}_{\mathbb{R}}$  and that  $V = L(Y, \mathbb{R})$ , that every set of reals in  $L(Y, \mathbb{R})$  is  $\infty$ -Borel whenever  $Y$  is a set of ordinals.

As in Section 10.1, we define one-step and  $\omega$ -sequence versions of the Vopěnka algebra. As we will be using Theorem 10.2.2, we restrict to the version for

subsets of  $\omega^\omega$ . We fix a cardinal  $\kappa$  such that for each  $n \in \omega$  and each  $A \subseteq (\omega^\omega)^n$ , if there exist a set of ordinals  $S$  in  $\text{OD}_{\{Y\}}$  and a formula  $\varphi$  such that

$$A = \{\bar{x} \in (\omega^\omega)^n : L[S, \bar{x}] \models \varphi(S, \bar{x})\}$$

then there exist a set of ordinals  $S$  in  $V_\kappa \cap \text{OD}_{\{Y\}}$  and a formula  $\psi$  such that

$$A = \{\bar{x} \in (\omega^\omega)^n : L_\kappa[S, \bar{x}] \models \psi(S, \bar{x})\}.$$

Let  $P_{\infty, Y}$  be the set of pairs  $(m, S)$  such that

- $m$  is the Gödel number of a binary formula  $\varphi$ ,
- $S \in V_\kappa \cap \text{OD}_{\{Y\}}$  is a set of ordinals,
- the set  $B_{m, S} = \{x \in \omega^\omega : L_\kappa[S, x] \models \varphi(S, x)\}$  is nonempty.

Then, as in Section 10.1, we have an induced ordering  $\leq_{\infty, Y}$  on  $P_{\infty, Y}$  defined by setting

$$(m, S) \leq_{\infty, Y} (j, T)$$

if  $B_{m, S} \subseteq B_{j, T}$ . Let  $\equiv_{\infty, Y}$  be the induced equivalence relation on  $P_{\infty, Y}$ , and let  $[(m, S)]_{\infty, Y}$  denote the  $\equiv_{\infty, Y}$ -class of each  $(m, S) \in P_{\infty, Y}$ . Let  $\mathbb{V}_{\infty, Y}$  be the partial order whose domain is the set of  $\equiv_{\infty, Y}$ -classes of  $P_{\infty, Y}$ , with the order inherited from  $\leq_{\infty, Y}$ .

For the sequence version, let  $P_{\infty, Y}^\omega$  be the set of triples  $(n, m, S)$  such that

- $n \in \omega$ ,
- $m$  is the Gödel number of a binary formula  $\varphi$ ,
- $S \in V_\kappa \cap \text{OD}_{\{Y\}}$  is a set of ordinals,
- the set  $B_{n, m, S}^\omega = \{\bar{x} \in (\omega^\omega)^n : L[S, \bar{x}] \models \varphi(S, \bar{x})\}$  is nonempty.

Again we have an induced ordering  $\leq_{\infty, Y}^\omega$  on  $P_{\infty, Y}^\omega$  defined by setting

$$(n, m, S) \leq_{\infty, Y}^\omega (k, j, T)$$

if  $n \geq k$  and  $\{\bar{x} \upharpoonright k : \bar{x} \in B_{n, m, S}^\omega\} \subseteq B_{k, j, T}^\omega$ . Let  $\equiv_{\infty, Y}^\omega$  be the induced equivalence relation on  $P_{\infty, Y}^\omega$ , and let  $[(n, m, S)]_{\infty, Y}^\omega$  denote the  $\equiv_{\infty, Y}^\omega$ -class of each  $(n, m, S) \in P_{\infty, Y}^\omega$ . Let  $\mathbb{V}_{\infty, Y}^\omega$  be the partial order whose domain is the set of  $\equiv_{\infty, Y}^\omega$ -classes of  $P_{\infty, Y}^\omega$ , with the order inherited from  $\leq_{\infty, Y}^\omega$ .

**10.3.1 Remark.** By Theorem 10.2.2, for each bounded  $S \subseteq \kappa$  in  $\text{HOD}_{\{Y\}}$  and each ternary formula  $\varphi$ , there exist a bounded  $T \subseteq \kappa$  in  $\text{HOD}_{\{Y\}}$  and a binary formula  $\psi$  such that

$$\{x \in \omega^\omega : \exists y \in \omega^\omega L[S, x, y] \models \varphi(S, x, y)\} = \{x \in \omega^\omega : L[T, x] \models \psi(T, x)\}.$$

**10.3.2 Remark.** Let  $(n, m, S)$  be an element of  $P_{\infty, Y}^\omega$ , and let  $\varphi$  be the formula with Gödel number  $m$ . Let  $n'$  be an element of  $\omega \setminus n$ . Then

$$\{\bar{x} \in (\omega^\omega)^{n'} : \bar{x} \restriction n \in B_{n, m, S}\} = \{\bar{x} \in (\omega^\omega)^{n'} : L_\kappa[S, \bar{x}] \models \varphi(S, \bar{x} \restriction n)\},$$

so this set has the form  $B_{n', k, S}$  for some  $k \in \omega$ , and  $B_{n', k, S} \leq_{\infty, Y}^\omega B_{n, m, S}$ . Moreover, there is a recursive function (not depending on  $S$ ) sending each such tuple  $(n, m, n')$  to a corresponding value  $k$ .

**10.3.3 Remark.** Any set in  $\text{OD}_{\{Y\}}$  consisting of sets of ordinals can be coded by a single set of ordinals in  $\text{OD}_{\{Y\}}$ , from which it follows that for any subset  $D$  of  $P_{\infty, Y}$  in  $\text{OD}_{\{Y\}}$  the set  $\bigcup\{B_{m, S} : (m, S) \in D\}$  has the form  $B_{j, T}$  for some  $(j, T) \in P_{\infty, Y}$ . Using this fact, the first paragraph of the proof of Theorem 10.1.2 adapts to prove that for each  $x \in \omega^\omega$ , the set  $G_x$  consisting of those  $[(m, S)]_{\infty, Y} \in P_{\infty, Y}$  such that  $x \in B_{m, S}$  is a  $\text{HOD}_{\{Y\}}$ -generic filter. This fact is implicit in Lemma 10.3.5 below, using the fact that the first coordinate of a  $\mathbb{V}_{\infty, Y}^\omega$ -generic sequence is generic for  $\mathbb{V}_{\infty, Y}$ .

Given  $x \in \omega^\omega$  and  $i \in \omega$ ,  $x(i)$  is the unique  $j \in \omega$  such that  $[(\emptyset, m_{i, j})]_{\infty, Y} \in G_x$ , where  $m_{i, j}$  is the Gödel number of the formula  $\varphi(u, v)$  expressing that  $(i, j) \in v$ . It follows then that

$$\text{HOD}_{\{Y\}}[x] = \text{HOD}_{\{Y\}}[G_x],$$

since  $G_x$  is the set of  $[(m, S)]_{\infty, Y} \in \mathbb{V}_{\infty, Y}$  such that  $L_\kappa[S, x] \models \varphi(S, x)$ , where  $\varphi$  is the formula with Gödel number  $m$ , and this can be computed in  $\text{HOD}_{\{Y\}}[x]$ . Note that we did not have this equality for the version of Vopěnka algebra defined in Section 10.1 (see Remark 10.1.3).

Since every condition in  $\mathbb{V}_{\infty, Y}$  is a member of  $G_x$  for some  $x \in \omega^\omega$ , every  $\text{HOD}_{\{Y\}}$ -generic filter  $G \subseteq \mathbb{V}_{\infty, Y}$  is equal to  $G_x$ , for

$$x = \{(i, j) : [(\emptyset, m_{i, j})]_{\infty, Y} \in G\}.$$

**10.3.4 Remark.** Since the first coordinate of a  $\mathbb{V}_{\infty, Y}^\omega$ -generic sequence is generic for  $\mathbb{V}_{\infty, Y}$ , there is a  $\mathbb{V}_{\infty, Y}$ -name  $\dot{Q}$  such that  $\mathbb{V}_{\infty, Y}^\omega$  is forcing-equivalent to  $\mathbb{V}_{\infty, Y} * \dot{Q}$ , via a map which carries the generic real for  $\mathbb{V}_{\infty, Y}$  to the first coordinate of the generic sequence produced by forcing with  $\mathbb{V}_{\infty, Y}^\omega$ .

As with the partial order  $\mathbb{V}_{\bar{x}, Y}^\omega$  in Section 10.1, each injection  $f: \omega \rightarrow \omega^\omega$  defines a filter  $H_f$  on  $\mathbb{V}_{\infty, Y}^\omega$ , consisting of the  $\equiv_{\infty, Y}^\omega$ -classes of those  $(n, m, S) \in P_{\infty, Y}^\omega$  for which  $f \restriction n$  is in  $B_{n, m, S}$ . Using this notation, Lemma 10.3.5 is the version of Lemma 10.1.7 for  $\mathbb{V}_{\infty, Y}^\omega$ . Remarks 10.3.1, 10.3.2 and 10.3.3 are used in the proof of the lemma.

The remainder of this section uses the notation  $\mu_S$  and  $\mathcal{D}_S$  introduced in Definition 9.1.1, for  $S$  a set of ordinals. Recall from Remark 1.2.5 that if  $\mu_\emptyset$  is a countably complete ultrafilter on  $\mathcal{D}_\emptyset$  (i.e., the constructibility degrees), then, for each set  $S \subseteq \text{Ord}$ ,  $\mu_S$  is a countably complete ultrafilter on  $\mathcal{D}_S$ . The statement  $\aleph_1 \not\leq 2^{\aleph_0}$  implies that the relation  $\leq_S$  is locally countable for each such  $S$ .

**Lemma 10.3.5** ( $\text{ZF} + \text{DC}_{\mathbb{R}} + \aleph_1 \not\leq 2^{\aleph_0}$ ). *Suppose that  $\mu_0$  is an ultrafilter on  $\mathcal{D}_0$ , and let  $Y$  be a set of ordinals. If  $G \subseteq \text{Col}^*(\omega, \omega^\omega)$  is a  $V$ -generic filter, and  $g = \bigcup G$ , then  $H_g$  is  $\mathbb{V}_{\infty, Y}^\omega$ -generic over  $\text{HOD}_{\{Y\}}$ , and  $g$  is in  $\text{HOD}_{\{Y\}}[H_g]$ .*

*Proof.* Let  $D \subseteq \mathbb{V}_{\infty, Y}^\omega$  be a dense set in  $\text{HOD}_{\{Y\}}$ , and let  $p$  be a condition in  $\text{Col}^*(\omega, \omega^\omega)$ . Let  $n_p$  be the domain of  $p$ . By strengthening the conditions in  $D$  if necessary, we may assume, applying Remark 10.3.2, that each element of  $D$  has the form  $[(n, m, S)]_{\infty, Y}^\omega$  for some  $(n, m, S) \in P_{\infty, Y}^\omega$  with  $n \geq n_p$ . Let  $D_0$  be the set of  $(n, m, S) \in P_{\infty, Y}^\omega$  with  $[(n, m, S)]_{\infty, Y}^\omega \in D$ . To show that  $H_g$  is  $\text{HOD}_{\{Y\}}$ -generic (applying the genericity of  $G$ ), it suffices to find a condition  $p' \leq p$  and a triple  $(n, m, S)$  in  $D_0$  such that  $p'$  is in  $B_{n, m, S}$ . By Remark 10.3.1 there is an  $\text{OD}_{\{Y\}}$  function associating each  $(n, m, S) \in D_0$  to a tuple  $(n_p, k, T) \in P_{\infty, Y}^\omega$  such that  $B_{n, m, S} \restriction n_p = B_{n_p, k, T}$ . Applying the first sentence of Remark 10.3.3, there exist  $T \in V_\kappa \cap \text{OD}_{\{Y\}}$  and a binary formula  $\psi$  (with Gödel number  $k$ ) such that

$$B_{n_p, k, T} = \{\bar{x} \in (\omega^\omega)^{n_p} : \exists (n, m, S) \in D_0 \ \bar{x} \in B_{n, m, S} \restriction n_p\}.$$

Furthermore,  $B_{n_p, k, T}$  must be the set of all injections from  $n_p$  to  $\omega^\omega$ , since otherwise the complement of this set has the form  $B_{n_p, j, R}$  for some  $j \in \omega$  and  $R \in V_\kappa \cap \text{OD}_{\{Y\}}$ , and we get a contradiction by considering a  $(n, m, S) \in D_0$  with

$$\{\bar{x} \restriction n_p : \bar{x} \in B_{n, m, S}\} \subseteq B_{n_p, j, R}.$$

There exists then a triple  $(n, m, S) \in D_0$  such that  $p \in B_{n, m, S} \restriction n_p$ , and a  $p' \leq p$  in  $B_{n, m, S}$ , as desired. Since  $G$  is  $V$ -generic, this gives the genericity of  $H_g$  over  $\text{HOD}_{\{Y\}}$ .

To see that  $g$  is in  $\text{HOD}_{\{Y\}}[H_g]$ , as in the proof of Theorem 10.1.2, let  $K$  be the set of pairs

$$((i, j, k), (n, m, S)) \in (\omega \times \omega \times \omega) \times P_{\infty, Y}^\omega$$

such that  $i < n$  and  $p(i)(j) = k$  for all  $p \in B_{n, m, S}$ . Then again  $K$  is in  $\text{HOD}_{\{Y\}}$ , and  $K$  can be used to define a  $\mathbb{V}_{\infty, Y}^\omega$ -name  $\tau \in \text{HOD}_{\{Y\}}$  such that for every  $\text{Col}^*(\omega, \omega^\omega)$ -generic function  $g$  over  $V$ ,  $\tau_{H_g} = g$ .  $\square$

The following is the main theorem of this section. The theorem shows that, in models of the form  $L(Y, \mathbb{R})$  satisfying  $\text{TD} + \text{DC}_{\mathbb{R}}$ , where  $Y$  is a set of ordinals, the orders  $\mathbb{V}_{\langle Y \rangle, \omega}$  and  $\mathbb{V}_{\infty, Y}$  (defined in Sections 10.1 and 10.3 respectively) are isomorphic.

**Theorem 10.3.6** ( $\text{ZF} + \text{DC}_{\mathbb{R}} + \aleph_1 \not\leq 2^{\aleph_0}$ ). *Suppose that  $\mu_0$  is an ultrafilter on  $\mathcal{D}_0$ , and let  $Y$  be a set of ordinals such that  $V = L(Y, \mathbb{R})$ . Then for each  $\text{OD}_{\{Y\}}$  set  $A \subseteq \omega^\omega$  there exist an  $\text{OD}_{\{Y\}}$  set  $S \subseteq \text{Ord}$  and a binary formula  $\varphi$  such that  $A = \{x \in \omega^\omega : L[S, x] \models \varphi(S, x)\}$ .*

*Proof.* Let  $\bar{a}$  be a finite set of ordinals, let  $\psi$  be a ternary formula, and let  $A$  be  $\{x \in \omega^\omega : \psi(\bar{a}, x, Y)\}$ . By Theorem 10.1.5, we may fix a set  $B \subseteq \text{Ord}$  such that  $\text{HOD}_{\{Y\}} = L[B]$ .

By Remark 10.3.4,  $\mathbb{V}_{\infty,Y}^\omega$  is forcing-equivalent to an iteration of the form  $\mathbb{V}_{\infty,Y} * \dot{Q}$ , for some  $\mathbb{V}_{\infty,Y}$ -name  $\dot{Q}$ . By Remark 10.3.3, for every  $x \in \omega^\omega$  there is a  $\text{HOD}_{\{Y\}}$ -generic filter  $G_x \subseteq \mathbb{V}_{\infty,Y}$  such that  $\text{HOD}_{\{Y\}}[G_x] = \text{HOD}_{\{Y\}}[x]$ .

Let  $\tau \in \text{HOD}_{\{Y\}}$  be the  $\mathbb{V}_{\infty,Y}^\omega$ -name for the generic function  $g$  from the end of the proof of Lemma 10.3.5. Let  $\dot{R}$  be a  $\mathbb{V}_{\infty,Y}$ -name in  $\text{HOD}_{\{Y\}}$  for the range of the realization of  $\tau$ . Let  $\dot{R}_*$  be the  $\mathbb{V}_{\infty,Y} * \dot{Q}$ -name induced by  $\dot{R}$  and a map in  $\text{HOD}_{\{Y\}}$  witnessing the forcing-equivalence of  $\mathbb{V}_{\infty,Y} * \dot{Q}$  with  $\mathbb{V}_{\infty,Y}^\omega$  which carries the generic real for  $\mathbb{V}_{\infty,Y}$  to the first coordinate of the generic sequence produced by forcing with  $\mathbb{V}_{\infty,Y}^\omega$ , as in Remark 10.3.4. We want to see that for each  $x \in \omega^\omega$ , all generic  $\dot{Q}_{G_x}$ -extensions of  $\text{HOD}_{\{Y\}}[x]$  agree about whether or not  $\psi(\bar{\alpha}, x, Y)$  holds in  $L(Y, \dot{R}_*)$ . By Lemma 10.3.5,  $(\omega^\omega)^V$  is the realization of  $\dot{R}_*$  in one of these extensions. The theorem will then follow, with  $S$  a set of ordinals coding  $B, Y, \bar{\alpha}, \mathbb{V}_{\infty,Y} * \dot{Q}$  and  $\dot{R}_*$ , since then, letting  $q_0$  be the empty condition in  $\mathbb{V}_{\infty,Y} * \dot{Q}$ ,

$$A = \{x \in \omega^\omega : L[B, x] \models q_0 \Vdash "L(\check{Y}, \dot{R}_*) \models \psi(\check{\alpha}, \check{x}, \check{Y})"\}.$$

Supposing toward a contradiction that not all such extensions agreed about whether  $\psi(\bar{\alpha}, x, Y)$  holds in  $L(Y, \dot{R}_*)$ , there exist  $x \in \omega^\omega$  and  $(n, m, S), (n', m', S')$  in  $P_{\infty,Y}^\omega$  such that

- $x \in B_{n,m,S} \cap B_{n',m',S'}$ ;
- $[(n, m, S)]_{\infty,Y}^\omega \Vdash "L(\check{Y}, \dot{R}) \models \psi(\check{\alpha}, \tau(0), \check{Y})"$ ;
- $[(n', m', S')]_{\infty,Y}^\omega \Vdash "L(\check{Y}, \dot{R}) \models \neg\psi(\check{\alpha}, \tau(0), \check{Y})"$ .

Let  $p: n + n' \rightarrow \omega^\omega$  and  $\pi: \omega \rightarrow \omega$  be such that

- $p(0) = x$ ;
- $\pi(0) = 0$ ;
- $\pi \upharpoonright (n + n')$  is a permutation of  $n + n'$ ;
- $\pi(i) = i$  for all  $i \geq n + n'$ ;
- $p \upharpoonright n \in B_{n,m,S}$ ;
- the function  $p': n' \rightarrow \omega^\omega$  defined by setting  $p'(i) = p(\pi(i))$  is in  $B_{n',m',S'}$ .

Let  $g: \omega \rightarrow \omega^\omega$  be the union of a  $V$ -generic filter for  $\text{Col}^*(\omega, \omega^\omega)$  containing  $p$ , and let  $g': \omega \rightarrow \omega^\omega$  be defined by setting  $g'(i) = g(\pi(i))$  for all  $i \in \omega$ . Then  $g'$  is also the union of a  $V$ -generic filter for  $\text{Col}^*(\omega, \omega^\omega)$ ,  $\dot{R}_{H_g} = \dot{R}_{H_{g'}} = (\omega^\omega)^V$  and  $g(0) = g'(0)$ , giving a contradiction.  $\square$

Part (1) of the following corollary appears as Theorem 1.9 in [1].

**Corollary 10.3.7** ( $\text{ZF} + \text{DC}_\mathbb{R} + \aleph_1 \not\leq 2^{\aleph_0}$ ). *Suppose that  $\mu_\emptyset$  is an ultrafilter on  $\mathcal{D}_\emptyset$ . Let  $Y$  be a set of ordinals such that  $V = L(Y, \mathbb{R})$ . Then the following hold.*

1. Every subset of  $\omega^\omega$  is  $\infty$ -Borel.
2.  $\text{HOD}_{\{Y,x\}} = \text{HOD}_{\{Y\}}[x]$  for all  $x \in \omega^\omega$ .

*Proof.* For part (1), every set in  $L(Y, \mathbb{R})$  is definable from  $Y$ , a finite set of ordinals and a finite subset of  $\omega^\omega$ . It follows then that for each  $A \subseteq \omega^\omega$  in  $L(Y, \mathbb{R})$ , there exist an  $\text{OD}_{\{Y\}}$ -set  $B \subseteq \omega^\omega \times \omega^\omega$  and an  $x_0 \in \omega^\omega$  such that  $A = \{y \in \omega^\omega : (x_0, y) \in B\}$ . By Theorem 10.3.6, there exist an  $\text{OD}_{\{Y\}}$  set  $S$  of ordinals and a ternary formula  $\varphi$  such that  $B = \{(x, y) \in \omega^\omega : L[S, x, y] \models \varphi(S, x, y)\}$ . Letting  $T$  be a set of ordinals coding  $S$  and  $x_0$ , there is a binary formula  $\varphi'$  such that  $A = \{y \in \omega^\omega : L[T, y] \models \varphi'(T, y)\}$ .

For part (2), fix  $x \in \omega^\omega$ . Clearly,  $\text{HOD}_{\{Y\}}[x]$  is contained in  $\text{HOD}_{\{Y,x\}}$ . To show the reverse inclusion, let  $G_x \subseteq \mathbb{V}_{\{Y\}, \omega}$  (from Section 10.1) be the  $\text{HOD}_{\{Y\}}$ -generic filter induced by  $x$ . By Theorem 10.1.2 it suffices to show that  $G_x$  is in  $\text{HOD}_{\{Y\}}[x]$ .

Let  $\kappa_0$  be the cardinal  $\kappa$  used in the definition of  $P_{(Y), \omega}$  and, for each  $(n, \bar{\alpha}) \in P_{(Y), \omega}$  let  $B_{n, \bar{\alpha}}^0$  be the set  $B_{n, \bar{\alpha}}$  from this definition. Similarly, let  $\kappa_1$  be the cardinal  $\kappa$  used in the definition of  $P_{\infty, Y}$ , and, for each  $(m, S) \in P_{\infty, Y}$  let  $B_{m, S}^1$  be the set  $B_{m, S}$  from that definition. By Theorem 10.3.6, for each  $(n, \bar{\alpha}) \in P_{(Y), \omega}$  there is a pair  $(m, S) \in P_{\infty, Y}$  such that  $B_{n, \bar{\alpha}}^0 = B_{m, S}^1$ . Furthermore, the set of pairs  $((n, \bar{\alpha}), (m, S))$  in  $P_{(Y), \omega} \times P_{\infty, Y}$  such that  $B_{n, \bar{\alpha}}^0 = B_{m, S}^1$  is in  $\text{HOD}_{\{Y\}}$ . The set of  $(m, S) \in P_{\infty, Y}$  such that  $x \in B_{m, S}^1$  is in  $\text{HOD}_{\{Y\}}[x]$ . It follows that  $G_x$ , which is the set of  $(n, \bar{\alpha})$  such that  $x \in B_{n, \bar{\alpha}}^0$ , is in  $\text{HOD}_{\{Y\}}[x]$ .  $\square$

The following is an application of Corollary 10.3.7 to the ordinals defined in Section 8.2.

**Corollary 10.3.8** ( $\text{ZF} + \text{AD} + \text{DC}_{\mathbb{R}}$ ).  $\chi_B = \kappa_B = \lambda_B = \rho_B$

*Proof.* By Lemma 8.2.2,  $\chi_B \leq \kappa_B \leq \lambda_B$ . To show that  $\chi_B$  and  $\rho_B$  are at least  $\lambda_B$ , let  $\gamma$  be the length of some  $\infty$ -Borel prewellordering, and let  $Y \subseteq \text{Ord}$  be an  $\infty$ -Borel code for this prewellordering. By Corollary 10.3.7, every set of reals in  $L(Y, \mathbb{R})$  is  $\infty$ -Borel. The Coding Lemma and Theorem 8.2.10 then imply that  $\chi_B$  and  $\rho_B$  are at least  $\Theta^{L(Y, \mathbb{R})}$ , which is more than  $\gamma$ . The same argument, letting  $\gamma$  be the Wadge rank of some  $\infty$ -Borel subset of  $\omega^\omega$ , shows that  $\lambda_B \geq \rho_B$ .  $\square$

**10.3.9 Remark.** Suppose that  $\text{DC}_{\mathbb{R}}$  and  $\aleph_1 \not\leq 2^{\aleph_0}$  hold, and that  $\mu_\emptyset$  is an ultrafilter on  $\mathcal{D}_\emptyset$ . Let  $\kappa$  be an infinite cardinal, and suppose that  $A \subseteq \omega^\omega$  is  $\kappa$ -Borel. By Theorem 8.2.3, there is a  $\kappa$ -Borel code  $S$  for  $A$  such that  $L(S, \mathbb{R}) = L(A, \mathbb{R})$ . By Theorem 10.3.6, in  $L(S, \mathbb{R})$  every subset of  $\omega^\omega$  is  $\infty$ -Borel. By the Moschovakis Coding Lemma, not every subset of  $\omega^\omega$  in  $L(S, \mathbb{R})$  is  $\kappa$ -Borel. It follows then that there is some  $B \subseteq \omega^\omega$  in  $L(S, \mathbb{R})$  which is  $\lambda$ -Borel but not  $<\lambda$ -Borel, for some  $\lambda > \kappa$ .

# Chapter 11

## Suslin sets and strong codes

This chapter presents three applications of the material in Chapters 7-10. In Section 11.2 we show, assuming  $<\Theta$ -Determinacy +  $\text{DC}_{\mathbb{R}}$ , that every subset of  $\omega^\omega$  which has an  $\infty$ -Borel code which is not  $\leq_{\mathcal{D}}$ -maximal is Suslin (Theorem 11.2.2). In Section 11.3 we show that  $\text{AD} + \text{Uniformization}$  implies that every subset of  $\omega^\omega$  is  $\infty$ -Borel. In Section 11.4 we show that, under  $\text{AD}^+$ , the set of Suslin cardinals is closed below  $\Theta$ . The material in Section 11.2 uses the notion of generic codes, which we introduce in Section 11.1.

### 11.1 Generic codes

Given a partial order  $\mathbb{P}$ ,  $n \in \omega$  and a nice  $\mathbb{P}$ -name  $\tau$  for an element of  $\omega^\omega$ , we let  $D_{\tau,n}$  be the set of  $\mathbb{P}$ -conditions (explicitly) deciding the value of  $\tau(n)$ , i.e., the set

$$\{p \in \mathbb{P} : \exists m \in \omega (p, (n, m)) \in \tau\}.$$

When  $E$  is a set containing  $\{D_{\tau,n} : n \in \omega\}$ , we write  $A_{\mathbb{P},E,\tau}$  for the set of values  $\tau_g$ , where  $g$  ranges over the set of all  $E$ -generic filters contained in  $\mathbb{P}$  (see Section 0.2 for the definitions of nice name and  $B$ -generic filter, and our corresponding notational conventions). We write  $\text{dom}(\mathbb{P})$  for the domain of the partial order  $\mathbb{P}$ .

**11.1.1 Definition.** Given

- $A \subseteq \omega^\omega$ ,
- a partial order  $\mathbb{P}$ ,
- a function  $B$  with  $\text{dom}(B)$  a set of ordinals and  $\text{range}(B)$  a set of dense open subsets of  $\mathbb{P}$ , and
- a nice  $\mathbb{P}$ -name  $\tau$  for an element of  $\omega^\omega$ ,

we say that  $(\mathbb{P}, B, \tau)$  is a *generic code* for  $A$  if  $\{D_{\tau,n} : n \in \omega\} \subseteq \text{range}(B)$  and  $A_{\mathbb{P}, \text{range}(B), \tau} = A$ . If in addition  $\alpha$  is an ordinal containing  $\text{dom}(\mathbb{P}) \cup \text{dom}(B)$ , then we say that  $(\mathbb{P}, B, \tau)$  is a *generic  $\alpha$ -code* for  $A$ . We say that  $(\mathbb{P}, B, \tau)$  is a *generic  $\infty$ -code* if it is a generic  $\alpha$ -code for some ordinal  $\alpha$ . If we say that  $C$  is a generic code, then we mean that  $C$  is a triple of the form  $(\mathbb{P}_C, B_C, \tau_C)$  where  $\mathbb{P}_C$ ,  $B_C$  and  $\tau_C$  are as above, and write  $A_C$  for  $A_{\mathbb{P}_C, B_C, \tau_C}$ .

Theorem 10.1.2 shows that if  $S \subseteq \text{Ord}$  is an  $\infty$ -Borel code for a set  $A \subseteq \omega^\omega$ , then  $(\mathbb{V}_{\{S\}, \omega^2} \restriction p, B, \tau \restriction p)$  is a generic code for  $A$  in  $\text{HOD}_{\{S\}}$ , where, letting  $\tau$  be the  $\mathbb{V}_{\{S\}, \omega^2}$ -name for the associated generic real (given by the set  $K$  in the proof of Theorem 10.1.2),

- $p$  is the  $\mathbb{V}_{\{S\}, \omega}$ -condition corresponding to  $A_{\varphi_S}$ ,
- $\mathbb{V}_{\{S\}, \omega^2} \restriction p$  and  $\tau \restriction p$  are the corresponding restrictions of  $\mathbb{V}_{\{S\}, \omega^2}$  and  $\tau$  below  $p$  and
- $B$  the a function in  $\text{HOD}_{\{S\}}$  listing the dense open subsets of  $\mathbb{V}_{\{S\}, \omega^2} \restriction p$  in  $\text{HOD}_{\{S\}}$  in their definability order.

Using the definability order on  $\text{HOD}_{\{S\}}$ , one can convert this generic code to a generic  $\infty$ -code. In the proof of Theorem 11.2.1 we use (and prove) the fact that, by Theorem 9.1.3, for a Turing cone of  $x \in \omega^\omega$  the resulting generic  $\infty$ -code is, in  $L[S, x]$ , a generic  $\gamma$ -code for  $A \cap L[S, x]$ , for some  $\gamma$  less than the  $\omega_3$  of  $\text{HOD}_S^{L[S, x]}$ .

Definition 11.1.3 below defines a game, relative to a generic code  $C$  for a set  $A$ , in which Player *II* attempts to build a countable subcode for a subset of  $A$ . Theorem 11.1.6 shows that a winning strategy for Player *I* in this game gives a Suslin representation for  $A$ . The following definition gives the relevant notion of subcode.

**11.1.2 Definition.** Suppose that  $(\mathbb{P}, B, \tau)$  is a generic  $\infty$ -code. For any set  $X \subseteq \text{Ord}$ , we let  $\mathbb{P}_X$  be  $\mathbb{P} \restriction (\text{dom}(\mathbb{P}) \cap X)$ ,  $B_X$  be  $\langle B(\beta) \cap X : \beta \in X \cap \text{dom}(B) \rangle$  and  $\tau_X$  be  $((\text{dom}(\mathbb{P}) \cap X) \times \omega^2) \cap \tau$ . If  $C = (\mathbb{P}, B, \tau)$ , we write  $C_X$  for  $(\mathbb{P}_X, B_X, \tau_X)$ .

**11.1.3 Definition.** Suppose that  $C = (\mathbb{P}, B, \tau)$  is a generic  $\alpha$ -code for a set  $A$ , for some ordinal  $\alpha$ . We associate to  $(\mathbb{P}, B, \tau)$  a game on  $\text{dom}(\mathbb{P}) \cup \text{dom}(B)$ , called  $\mathcal{G}_{\mathbb{P}, B, \tau}$  or  $\mathcal{G}_C$ , where players *I* and *II* collaborate to build a countable subset  $\sigma \subseteq \text{dom}(\mathbb{P}) \cup \text{dom}(B)$ , and *I* wins if  $C_\sigma = (\mathbb{P}_\sigma, B_\sigma, \tau_\sigma)$  is a generic code for a subset of  $A$ . We say that  $(\mathbb{P}, B, \tau)$  is a *strong generic code* (or *strong  $\alpha$ -generic code*) for  $A_\varphi$  if *II* does not have a winning strategy in  $\mathcal{G}_{\mathbb{P}, B, \tau}$ .

Note that a strong generic  $\alpha$ -code is also a strong generic  $\beta$ -code for any ordinal  $\beta \geq \alpha$ .

**11.1.4 Remark.** For any infinite cardinal  $\kappa$  and  $\kappa$ -generic code  $(\mathbb{P}, B, \tau)$ ,  $\kappa$ -Determinacy implies the determinacy of the game  $\mathcal{G}_{\mathbb{P}, B, \tau}$ , since a run of the game continuously builds a subset of  $\omega$  which HC-codes a generic  $\omega$ -code.



**11.1.5 Remark.** Let (for this remark)  $X$  be the set of pairs  $(x, y) \in (\omega^\omega)^2$  such that  $x$  is an HC-code for a generic  $\omega$ -code  $(\mathbb{P}_x, B_x, \tau_x)$  and  $y$  is an HC-code for a  $B_x$ -generic filter  $g_y \subseteq \mathbb{P}_x$ . If  $A \subseteq \omega^\omega$  is Suslin and co-Suslin, then the set of  $x \in \omega^\omega$  for which there is a  $y \in \omega^\omega$  with  $(x, y) \in X$  and  $\tau_{x, g_y} \notin A$  is clearly Suslin. Since the pointclass of Suslin sets is  $\forall^{\omega^\omega}$ -closed (by the Second Periodicity Theorem; see Remark 6.1.6), the set of  $x \in \omega^\omega$  for which  $\tau_{x, g_y} \in A$  for every  $y \in \omega^\omega$  is also Suslin under these assumptions. It follows that if AD holds, and  $(\mathbb{P}, B, \tau)$  is an  $\alpha$ -Borel code for a set  $A$  which is Suslin and co-Suslin, for some  $\alpha < \Theta$ , then  $\mathcal{G}_{\mathbb{P}, B, \tau}$  is determined, by Theorem 7.0.1.

Unlike  $\infty$ -Borel codes, strong generic codes witness Suslinity, in the context of  $<\Theta$ -Determinacy.

**Theorem 11.1.6.** *Let  $\alpha$  be an infinite ordinal and let  $C$  be a generic  $\alpha$ -code for a set  $A \subseteq \omega^\omega$ . If player  $I$  has a winning strategy in the game  $\mathcal{G}_C$ , then  $A$  is  $\alpha$ -Suslin. Moreover, if  $\Sigma$  is a winning strategy for player  $I$  in  $\mathcal{G}_C$ , then  $A = p[T]$ , for some tree  $T \subseteq (\omega \times \alpha \times 2)^{<\omega}$  which is definable from  $C$  and  $\Sigma$ .*

*Proof.* Let  $C$  be  $(\mathbb{P}, B, \tau)$ , and fix a winning strategy  $\Sigma$  for player  $I$  in  $\mathcal{G}_C$ . Then for all  $x \in \omega^\omega$ ,  $x \in A$  if and only if there exist a countable  $\sigma \subseteq \text{dom}(\mathbb{P}) \cup \text{dom}(B)$  produced by a run of  $\mathcal{G}_C$  where  $I$  plays according to  $\Sigma$ , and a  $B_\sigma$ -generic filter  $g \subseteq \mathbb{P}_\sigma$  such that  $x = \tau_{\sigma, g}$ . To see this, note first of all that, since a  $\Sigma$  is winning strategy for player  $I$ , if there is a play against  $\Sigma$  producing  $\sigma$  with  $x$  equal to  $\tau_{\sigma, g}$  for some  $B_\sigma$ -generic  $g \subseteq \mathbb{P}_\sigma$ , then  $x \in A$ . For the other direction, fix  $x \in A$  and a  $B$ -generic filter  $G \subseteq \mathbb{P}$  with  $\tau_G = x$ . Let  $\sigma$  be a run of  $\mathcal{G}_C$  where player  $I$  plays according to  $\Sigma$  and, for each  $D \in B_\sigma$ ,  $D \cap \mathbb{P}_\sigma$  is nonempty. Then  $G \cap \mathbb{P}_\sigma$  is  $B_\sigma$ -generic, and  $\tau_{\sigma, G \cap \mathbb{P}_\sigma} = x$ .

The set of  $(x, y, z)$  such that

- $x \in \omega^\omega$ ,
- $y \in \alpha^\omega$  is a run of  $\mathcal{G}_C$  where  $I$  plays according to  $\Sigma$ ,
- $z \in 2^\omega$  and
- $y[z^{-1}[\{1\}]]$  is a filter on  $y[\omega] \cap \mathbb{P}$  for which the realization of

$$\tau_{y[\omega], y[z^{-1}[\{1\}]}$$

is  $x$

is the set of paths through a tree on  $\omega \times \alpha \times 2$  whose projection is  $A$ .  $\square$

There is a natural join operation on wellordered sequences of generic  $\infty$ -codes, which we now define. Suppose that  $I$  is a set of ordinals and  $\langle C_\alpha : \alpha \in I \rangle$  is a sequence of generic  $\infty$ -codes. For each  $\alpha \in I$ , let  $C_\alpha$  be  $(\mathbb{P}_\alpha, B_\alpha, \tau_\alpha)$  and let  $\leq_\alpha$  be the order on  $\text{dom}(\mathbb{P}_\alpha)$  given by  $\mathbb{P}_\alpha$ . Let  $X$  be the set  $\{(\alpha, \gamma) : \gamma \in \text{dom}(\mathbb{P}_\alpha)\}$  and let  $\leq_X$  the partial order on  $X$  defined by setting  $(\alpha, \gamma) \leq_X (\alpha', \gamma')$  to hold if  $\alpha = \alpha'$  and  $\gamma \leq_\alpha \gamma'$ . Let  $\mathbb{P}_X$  denote the corresponding partial order on  $X$ .

For each  $\alpha \in I$ , let  $B_\alpha$  be the sequence  $\langle b_{\alpha,\delta} : \delta \in \text{dom}(B_\alpha) \rangle$ . Let  $Y$  be the set  $\{(\alpha, \delta) : \delta \in \text{dom}(B_\alpha)\}$ . For each pair  $(\alpha, \delta)$  in  $Y$ , let  $e_{\alpha,\delta}$  be the set of  $(\beta, \gamma) \in X$  such that either  $\beta \neq \alpha$  or  $\gamma \in b_{\alpha,\delta}$ .

Let  $Z$  be the set of pairs  $((\alpha, \gamma), (n, m))$  such that  $(\alpha, \gamma) \in X$  and the pair  $(\gamma, (n, m))$  is in  $\tau_\alpha$ . Then  $Z$  is a nice  $\mathbb{P}_X$ -name for an element of  $\omega^\omega$ . Let  $E$  be

$$\{e_{\alpha,\delta} : (\alpha, \delta) \in Y\} \cup \{D_{Z,n} : n \in \omega\}.$$

Then  $(\mathbb{P}_X, E, Z)$  is almost a generic code for  $\bigcup_{\alpha \in I} A_{C_\alpha}$ , except that  $E$  is a set and not a function.

Next we map  $(\mathbb{P}_X, E, Z)$  over to a generic  $\infty$ -code. Let  $f: \eta \rightarrow X$  and  $g: \xi \rightarrow Y$  (for some ordinals  $\eta, \xi$ ) be the bijections induced by the Gödel ordering on pairs of ordinals. Let  $\mathbb{P}$  be the partial order  $f(\gamma) \leq_X f(\gamma')$  on  $\eta$ . Let  $B = \langle b_\delta : \delta < \omega + \xi \rangle$  be defined by setting each  $b_n$  ( $n \in \omega$ ) to be  $f^{-1}[D_{Z,n}]$  for each  $n \in \omega$ , and each  $b_{\omega+\nu}$  to be  $f^{-1}[b_{\alpha,\delta}]$ , where  $g(\nu) = (\alpha, \delta)$ . Let  $\tau$  be the set of pairs  $(\gamma, (n, m))$  for which  $(f(\gamma), (n, m)) \in Z$ . Let  $C$  be  $(\mathbb{P}, B, \tau)$ . Then  $C$  is a generic  $(\eta \cup \xi)$ -code for  $\bigcup_{\alpha < \kappa} A_{C_\alpha}$ . We call  $C$  the *join* of  $\langle C_\alpha : \alpha \in I \rangle$ . Note that if  $I$  is an infinite cardinal  $\kappa$ , and each  $C_\alpha$  is a generic  $\kappa$ -code, then  $(\eta \cup \xi) = \kappa$ .

**11.1.7 Remark.** A generic  $\infty$ -code  $(\mathbb{P}, B, \tau)$  induces a natural *collapse* code  $(\mathbb{P}', B', \tau')$ , induced by the unique order-preserving bijections  $h: \alpha \rightarrow \text{dom}(\mathbb{P})$  and  $k: \beta \rightarrow \text{dom}(\mathbb{B})$ , where  $\alpha$  and  $\beta$  are the respective ordertypes of  $\text{dom}(\mathbb{P})$  and  $\text{dom}(B)$ . We have  $\text{dom}(\mathbb{P}') = \alpha$ , and the order on  $\mathbb{P}'$  is given by setting  $\delta \leq_{\mathbb{P}'} \gamma$  to hold if and only if  $h(\delta) \leq_{\mathbb{P}} h(\gamma)$ . Similarly, for each  $\delta < \beta$ ,  $B'(\delta) = f^{-1}[B(h(\delta))]$ . Finally,  $\tau'$  is the set of pairs  $(\delta, (n, m))$  for which  $(f(\delta), (n, m)) \in \tau$ . Then  $(\mathbb{P}, B, \tau)$  and  $(\mathbb{P}', B', \tau')$  are generic  $\infty$ -codes for the same subset of  $\omega^\omega$ .

Theorem 11.1.8 below shows that the join operation just defined preserves strong generic codes, under the appropriate form of ordinal determinacy.

**Theorem 11.1.8** (ZF + DC $_{\mathbb{R}}$ ). *Let  $\kappa$  be an infinite cardinal below  $\Theta$  such that  $\kappa$ -Determinacy holds, and let  $\bar{C} = \langle C_\alpha : \alpha < \kappa \rangle$  be a sequence of strong  $\kappa$ -codes. Then the join of  $\bar{C}$  is a strong generic  $\kappa$ -code for  $\bigcup_{\alpha < \kappa} A_{C_\alpha}$ .*

*Proof.* Let  $C = (\mathbb{P}, B, \tau)$  be the join of  $\bar{C}$ , and let  $f$  and  $g$  be the corresponding functions from the definition of the join. To see that  $C$  is a strong generic code, fix a set  $A \subseteq \omega^\omega$  of Wadge rank greater than  $\kappa$ . It suffices to show that player  $II$  does not have a winning strategy in  $\mathcal{G}_C$  in  $L(A, \mathbb{R})$ , since  $L(A, \mathbb{R})$  contains all subsets of  $\kappa$ , by the Coding Lemma. Since DC $_{\mathbb{R}}$  holds, DC holds in  $L(A, \mathbb{R})$ . Fixing a strategy  $\Sigma$  for player  $II$ , we can find by DC a countable  $\sigma \subseteq \eta \cup \xi$  and winning strategies  $\rho_\alpha$  for player  $I$  in the games  $\mathcal{G}_{C_\alpha}$  ( $\alpha \in \sigma \cap \eta$ ) such that

- $\omega \subseteq g[\sigma]$ ,
- $\sigma$  is the result of a run of  $\mathcal{G}_C$  where  $II$  has played by  $\Sigma$  and
- for each  $\alpha \in \sigma \cap \eta$ , the set

$$\sigma_\alpha = \{\gamma \in \text{dom}(\mathbb{P}_\alpha) : (\alpha, \gamma) \in f[\sigma]\} \cup \{\delta \in \text{dom}(B_\alpha) : (\alpha, \delta) \in g[\sigma]\}$$

is the result of a run of  $\mathcal{G}_{C_\alpha}$  where player  $I$  has played by  $\rho_\alpha$ .

Then the join of  $\langle C_{\alpha, \sigma_\alpha} : \alpha \in \sigma \rangle$  is the collapse code associated to  $C_\sigma$ . It follows that

$$A_{C_\sigma} = \bigcup_{\alpha \in \sigma} A_{C_{\alpha, \sigma_\alpha}} \subseteq \bigcup_{\alpha \in \sigma} A_{C_\alpha},$$

showing that  $I$  wins this run of the game.  $\square$

Examining the proof of Theorem 7.0.1, we get an alternate version of Theorem 11.1.8 where instead of assuming  $\kappa$ -Determinacy we assume that the union is Suslin, and that there is a strong partition cardinal above  $\kappa$ .

**Theorem 11.1.9** (ZF + DC $_{\mathbb{R}}$ ). *Suppose that  $\kappa < \delta$  are infinite cardinals and that  $\delta$  is a regular cardinal such that*

$$\forall \mu < \delta (\delta \rightarrow (\delta)^\mu_\mu)$$

*holds. Let  $\bar{C} = \langle C_\alpha : \alpha < \kappa \rangle$  be a sequence of strong  $\kappa$ -codes. Let  $A$  be  $\bigcup_{\alpha < \kappa} A_{C_\alpha}$ , and suppose that  $A$  is Suslin. Then there is a strong  $\kappa$ -code for  $A$  which is definable from  $\bar{C}$ .*

*Proof.* (Sketch) Let  $C_\alpha$  be  $(\mathbb{P}_\alpha, B_\alpha, \tau_\alpha)$ , for each  $\alpha < \kappa$ . Let  $C = (\mathbb{P}, B, \tau)$  be the join of  $\bar{C}$ . Then  $A_C$  is a generic  $\kappa$ -code for  $A$ . We want to see that player  $I$  has winning strategy in  $\mathcal{G}_C$ . As in Remark 11.1.4,  $\mathcal{G}_C$  is the same as  $\mathcal{G}_{\rho, D}$  (as defined in Chapter 7) for some continuous function  $\rho: \kappa^\omega \rightarrow \omega^\omega$ , where  $D$  is the set of HC-codes for generic  $\omega$ -codes for subsets of  $A$ . As in Remark 11.1.5, since  $A_C$  is Suslin,  $D$  is Suslin. Let  $T$  be a tree on  $\omega \times \kappa$  projecting to  $D$ . To show that  $\mathcal{G}_{\rho, D}$  is determined, run the proof of Theorem 7.0.1, omitting all mentions of  $S$  there, since there is no  $S$  in our context. Since the proof of Claim 1 of Theorem 7.0.1 does not mention  $S$ , it suffices to show that player  $I$  has a winning strategy in the augmented game  $\mathcal{G}^T$  defined there. Since the game  $\mathcal{G}^T$  is closed, it suffices to see that player  $II$  does not have a winning strategy. However, a winning strategy for player  $II$  in  $\mathcal{G}^T$  gives one for player  $II$  in  $\mathcal{G}_C$ , and the second half of the proof of Theorem 11.1.9 shows that there can be no such strategy.  $\square$

Section 11.2 shows how to convert  $\infty$ -Borel codes into strong generic codes.

## 11.2 Producing strong generic codes

In this section we use the terms  $\mu_S$  and  $\mathcal{D}_S$  from Definition 9.1.1, for  $S$  a set of ordinals. Recall from Section 9.5 that  $\delta_S^\infty$  denotes  $\prod \omega_2^{L[S, x]} / \mu_S$ . Recall also from Theorem 11.1.6 that subsets of  $\omega^\omega$  with strong generic codes are Suslin. The following theorem produces strong generic codes.

**Theorem 11.2.1** (ZF + CC $_{\mathbb{R}}$ ). *Let  $S$  be a set of ordinals such that  $\leq_S$  is locally countable,  $\mu_S$  is an ultrafilter and the ultrapower  $\text{Ord}^{\mathcal{D}_S} / \mu_S$  is wellfounded. If  $S$  is an  $\infty$ -Borel code, then  $A_{\varphi_S}$  has a strong generic  $\delta_S^\infty$ -code which is definable from  $S$ .*

*Proof.* By Theorem 9.1.3 we have that for some  $x_0 \in \omega^\omega$  and all  $y \in \omega^\omega$  such that  $[y]_S \geq_S [x_0]_S$ , GCH holds in  $L[S, y]$  below  $\omega_1^V$  (which is strongly inaccessible in  $L[S, y]$ , since  $\aleph_1 \not\leq 2^{\aleph_0}$ ). For each such  $y$ , it follows that the partial order  $\mathbb{V}_{\langle S \rangle, \omega^2}$  (defined at the beginning of Section 10.1) as computed in  $L[S, y]$ , being isomorphic to the subset relation restricted to a subset of  $\mathcal{P}(\omega^\omega)$ , has cardinality at most  $\aleph_2$  in  $L[S, y]$ . Furthermore, since antichains in this partial order correspond to sequences of pairwise disjoint subsets of  $\mathcal{P}(\omega^\omega)$ ,  $\mathbb{V}_{\langle S \rangle, \omega^2}^{L[S, y]}$  is  $\aleph_2$ -c.c. in  $L[S, y]$ , and the set of all antichains of  $\mathbb{V}_{\langle S \rangle, \omega^2}^{L[S, y]}$  has cardinality  $\aleph_2$  in  $L[S, y]$ .

Since  $\omega_3^{L[S, y]}$  is the cardinal successor of  $\omega_2^{L[S, y]}$  in  $\text{HOD}_{\{S\}}^{L[S, y]}$  (see Remark 10.1.4), it follows, using Theorem 10.1.2 and the definability order in  $\text{HOD}_{\{S\}}^{L[S, y]}$  that there is an  $S$ -invariant function  $f$  on  $\omega^\omega$  such that, for an  $S$ -cone of  $y$ ,  $f(y)$  is, in  $L[S, y]$ , an  $\omega_2^{L[S, y]}$ -generic code  $C_y = (\mathbb{P}_y, B_y, \tau_y)$  for  $A_{\varphi_S}$ . Moreover,  $\mathbb{P}_y$  can be taken to be a partial order on  $\omega_2^{L[S, y]}$  which is isomorphic to  $(\mathbb{V}_{\langle S \rangle, \omega^2} \upharpoonright p)^{L[S, y]}$ , where  $p$  is the  $\mathbb{V}_{\langle S \rangle, \omega^2}^{L[S, y]}$ -condition corresponding to  $A_{\varphi_S}$ , and  $B_y$  and  $\tau_y$  are the corresponding images, respectively, of the set of all dense open subsets of  $(\mathbb{V}_{\langle S \rangle, \omega^2} \upharpoonright p)^{L[S, y]}$  in  $\text{HOD}_{\{S\}}^{L[S, y]}$  and the canonical name for the generic real added by  $(\mathbb{V}_{\langle S \rangle, \omega^2} \upharpoonright p)^{L[S, y]}$  (corresponding to the set  $K$  from the proof of Theorem 10.1.2).

One key point is for that each such  $y$ ,  $A_{C_y}$  is contained in  $A_{\varphi_S}$  as computed in  $V$ . To see this, note that if some  $\text{HOD}_{\{S\}}^{L[S, y]}$ -generic filter gave a counterexample, then, since  $S$  is in  $\text{HOD}_{\{S\}}^{L[S, y]}$ , this would be forced by some condition in  $(\mathbb{V}_{\langle S \rangle, \omega^2} \upharpoonright p)^{L[S, y]}$ . Since  $L[S, y]$  contains a  $(\mathbb{V}_{\langle S \rangle, \omega^2})^{L[S, y]}$ -generic real below each condition, such a counterexample would then exist in  $L[S, y]$ .

Now consider the wellfounded ultrapower  $V^{\mathcal{D}_S}/\mu_S$ . Let  $C = (\mathbb{P}, B, \tau)$  be the triple represented by the function  $f$ . We want to see that this is a strong generic code for  $A$ . We have that  $\text{dom}(\mathbb{P}) \cup \text{dom}(B)$  is contained in  $\delta_S^\infty$ .

We want to see then that player *II* does not have a winning strategy in the game  $\mathcal{G}_{\mathbb{P}, B, \tau}$  from Definition 11.1.3. Suppose towards a contradiction that  $\Sigma$  is such a strategy. For each  $y \in \omega^\omega$  such that  $y \geq x_0$ , let  $R_y$  be the tree of attempts to find a countable  $\sigma \subseteq \delta_S^\infty$  resulting from a play of  $\mathcal{G}_C$  according to  $\Sigma$ , and an isomorphism between  $C_y = (\mathbb{P}_y, B_y, \tau_y)$  and  $C_\sigma = (\mathbb{P}_\sigma, B_\sigma, \tau_\sigma)$ . If any of these trees is illfounded, then we have a contradiction. If not, then (since there is in each case a wellordering of  $R_y$  definable from  $\Sigma$ ,  $C$  and  $C_y$ ) we can find (in an  $S$ -invariant fashion) ranking functions  $\rho_y$  on each such tree  $R_y$ . Let  $j$  be the map sending each element of  $V$  to the element of  $V^{\mathcal{D}_S}/\mu_S$  represented by the corresponding constant function. In addition, let  $R^*$  and  $\rho^*$  be the elements of  $V^{\mathcal{D}_S}/\mu_S$  represented by the maps  $y \mapsto R_y$  and  $y \mapsto \rho_y$ . Then in the wellfounded model

$$\prod \text{HOD}_{\{S, C, \Sigma\}}^{L[S, C, \Sigma, y]} / \mu_S,$$

$\rho^*$  is a ranking function on the tree  $R^*$  of attempts to find a countable  $\sigma \subseteq j(\delta_S^\infty)$  resulting from a play of  $j(\mathcal{G}_C)$  according to  $j(\Sigma)$ , and an isomorphism between

$C$  and  $j(C)_\sigma$ .

Now,  $C$  is the collapse code corresponding to  $(j[\mathbb{P}], j[B], j[\tau])$ , which is

$$j(C)_{j[\delta_S^\infty]} = (j(\mathbb{P})_{j[\delta_S^\infty]}, j(B)_{j[\delta_S^\infty]}, j(\tau)_{j[\delta_S^\infty]}).$$

Since  $j[\delta_S^\infty]$  is closed under  $j(\Sigma)$ , there is a nonempty subset of  $R^*$  without terminal nodes, corresponding to those nodes where the isomorphism is contained in  $j$ , and the run of the game is contained in  $j[\delta_S^\infty]$ . It follows that  $R^*$  is illfounded, giving the desired contradiction.  $\square$

Theorems 11.1.6 and 11.2.1 and give Suslin representations for sets of reals having  $\infty$ -Borel codes which are not  $\leq_{\mathcal{D}}$ -maximal.

**Theorem 11.2.2** (ZF +  $\text{DC}_{\mathbb{R}}$  +  $<\Theta$ -Determinacy). *Every subset of  $\omega^\omega$  which has an  $\infty$ -Borel code which is not  $\leq_{\mathcal{D}}$ -maximal is Suslin.*

*Proof.* Suppose that  $A \subseteq \omega^\omega$  has an  $\infty$ -Borel code which is not  $\leq_{\mathcal{D}}$ -maximal. A reflection argument shows that there exist bounded subsets  $S$  and  $T$  of  $\Theta$  such that  $S <_{\mathcal{D}} T$  and  $S$  is an  $\infty$ -Borel code for  $A$ . Let  $B \subseteq \omega^\omega$  have Wadge rank greater than the supremum of  $S \cup T$ . By the Coding Lemma,  $S$  and  $T$  are in  $L(B, \mathbb{R})$ . Since AD and  $\text{DC}_{\mathbb{R}}$  hold,  $L(B, \mathbb{R})$  satisfies DC and the other hypotheses of Theorem 11.2.1 regarding  $S$ . Work in  $L(B, \mathbb{R})$  for the rest of the proof. By Theorem 9.5.1, and part (2b) of Theorem 9.3.8,  $\delta_S^\infty < \Theta$ . By Theorem 11.2.1,  $A$  has a strong generic  $\delta_S^\infty$ -code. Since  $\delta_S^\infty$ -Determinacy holds, it follows from Theorem 11.1.6 that  $A$  is Suslin.  $\square$

In addition, one gets the following equivalences under  $\text{AD}^+$ .

**Theorem 11.2.3** (ZF +  $\text{AD}^+$ ). *The following are equivalent.*

1. *There is no  $\leq_{\mathcal{D}}$ -maximal set of ordinals.*
2. *The Suslin cardinals are cofinal in  $\Theta$ .*
3. *Every subset of  $\omega^\omega$  is Suslin.*
4. **Uniformization**

*Proof.* The equivalence of (2) and (3) is Corollary 6.1.19, which requires only AD. By Theorem 11.2.2, (1) implies (3). That (3) implies (4) follows from Theorem 6.2.1. The equivalence of (4) and (1) follows from Corollary 10.2.7.  $\square$

Recall from Theorem 6.3.2 that if every subset of  $\omega^\omega$  has a Wadge rank, and Lipschitz Determinacy and Uniformization hold, then the length of the Solovay sequence is a limit ordinal. A version of the converse holds assuming  $\text{AD}^+$ .

**Theorem 11.2.4.** *If  $\text{AD}^+$  holds and the Solovay sequence has limit length, then every subset of  $\omega^\omega$  is Suslin.*

*Proof.* By Theorem 11.2.2 it suffices to show every subset of  $\omega^\omega$  has an  $\infty$ -Borel code which is not  $\leq_{\mathcal{D}}$ -maximal. Suppose that  $A \subseteq \omega^\omega$  is a counterexample. Applying Remark 8.1.12, fix an  $\infty$ -Borel code  $S$  for  $A$  which is a bounded subset of  $\Theta$ . Let  $B \subseteq \omega^\omega$  have Wadge rank greater than the supremum of  $S$ , so that, in  $L(B, \mathbb{R})$ ,  $S$  is definable from  $B$  and some  $x_0 \in \omega^\omega$ , by the Coding Lemma. Since the Solovay sequence has limit length, we may fix  $C \subseteq \omega^\omega$  such that  $C$  is not ordinal definable from  $B$  and any element of  $\omega^\omega$ . Since membership in the model  $L(C, \mathbb{R})$  is ordinal definable (using the Wadge rank of  $C$ ), the model  $L(C, \mathbb{R})$  also satisfies the statement that  $C$  is not ordinal definable from  $B$  and any element of  $\omega^\omega$ . The model  $L(C, \mathbb{R})$  satisfies  $\text{AD}^+$ , by Theorem 8.2.8, and the statement that  $S$  is  $\leq_{\mathcal{D}}$ -maximal. Work in  $L(C, \mathbb{R})$ , which satisfies DC, for the rest of the proof. The set  $S^\infty$  from Section 9.6 is definable from  $S$ , and it follows from Theorem 9.6.6 that every subset of  $\omega^\omega$  is definable from  $S^\infty$  and a real. Since  $S$  is definable from  $B$  and  $x_0$  we have a contradiction.  $\square$

**11.2.5 Remark.** Since each  $C_y$  in the proof of Theorem 11.2.1 is a generic code in  $L[S, y]$  for a subset of  $A$  containing  $A \cap L[S, y]$ , the set  $C$  produced in the proof is a generic code for  $A$ . If  $A$  happens to be Suslin and co-Suslin, and AD holds, then  $\mathcal{G}_C$  is determined, by Remark 11.1.5. In conjunction with Theorem 11.1.6, then, we have that if AD holds and  $S$  is an  $\infty$ -Borel code for a Suslin set  $A$ , then  $A$  is the projection of a tree on  $\omega \times \delta_S^\infty$  which is definable from  $S$ , a real and any prewellordering of  $\omega^\omega$  of length  $\delta_S^\infty$  (from which strategies in the corresponding game  $\mathcal{G}_C$  are definable, by the Coding Lemma).

### 11.3 $\infty$ -Borel representations from Uniformization

The main theorem of this section is Theorem 11.3.4, which says that if AD and Uniformization hold, then all sets of reals are  $\infty$ -Borel. Theorem 11.3.3 is a slightly stronger version, in which the assumption of AD is replaced with some of its consequences. Theorem 11.3.3 is Theorem 5.10 of [21], whose presentation we follow closely. Again we use the notation from Definition 9.1.1, with  $S$  as  $\emptyset$ .

Before beginning the proof of Theorem 11.3.3 we introduce some terminology for treating generic filters over countable structures in terms of descriptive set theory, using notion introduced in Section 0.2. Given a partial order  $\mathbb{P}$ , we say that a set  $Y$  consisting of filters on  $\mathbb{P}$  is *comeager* if there is a countable set  $X$  such that  $Y$  contains every  $X$ -generic filter contained in  $\mathbb{P}$ . Given  $p \in \mathbb{P}$ ,  $A \subseteq \omega^\omega$  and a nice  $\mathbb{P}$ -name  $\tau$  for an element of  $\omega^\omega$ , we write  $p \Vdash^* \tau \in A$  to mean that for a comeager set of filters  $g \subseteq \mathbb{P}$ ,  $\tau_g$  is in  $A$ . We define  $p \Vdash^* \tau \notin A$  analogously. We let  $D_{\mathbb{P}, \tau}^A$  be the set of  $p \in \mathbb{P}$  such that either  $p \Vdash^* \tau \in A$  or  $p \Vdash^* \tau \notin A$ . If  $\mathbb{P}$  is countable, and every subset of  $\omega^\omega$  has the property of Baire, then each set of the form  $D_{\mathbb{P}, \tau}^A$  is dense in  $\mathbb{P}$ .

Given  $A \subseteq \omega^\omega$ , the *term relation* for  $A$  is the class  $\text{TR}(A)$  consisting of those triples  $(\mathbb{P}, p, \tau)$  such that  $\mathbb{P}$  is a poset,  $\tau$  is a nice  $\mathbb{P}$ -name for an element of  $\omega^\omega$  and  $p \Vdash^* \tau \in A$ . Note that  $\text{TR}(A)$  is definable from  $A$ . We write  $\dot{t}_{A, \mathbb{P}}$  for the set of  $(p, \tau)$  for which  $(\mathbb{P}, p, \tau) \in \text{TR}(A)$ . Note that  $\dot{t}_{A, \mathbb{P}}$  is a  $\mathbb{P}$ -name, as is every

subset of it.

**11.3.1 Remark.** If  $\mathbb{P}$  is a countable partial order,  $T$  is a countable set of nice  $\mathbb{P}$ -names for reals and each set  $D_{\mathbb{P},\tau}^A$  ( $\tau \in T$ ) is dense in  $\mathbb{P}$ , then, for comeagerly many filters  $g \subseteq \mathbb{P}$ , for all  $\tau \in T$ ,  $\tau_g \in A$  if and only if, for some  $p \in g$ ,  $(p, \tau) \in \dot{t}_{A,\mathbb{P}}$ .

A transitive set  $M$  is said to be *weakly*  $(A, \mathbb{P})$ -closed if  $\dot{t}_{A,\mathbb{P}} \cap M \in M$ . Applying the definition to the set of names of the form  $\check{x}$  ( $x \in \omega^\omega \cap M$ ), this implies that  $A \cap M \in M$ , if  $M$  is a model of a certain weak fragment of ZF. We say that  $M$  is *strongly*  $(A, \mathbb{P})$ -closed if it is weakly  $(A, \mathbb{P})$ -closed and, for all  $M$ -generic filters  $g \subseteq \mathbb{P}$ ,  $(\dot{t}_{A,\mathbb{P}} \cap M)_g$  (that is,  $\{\tau_g : \exists p \in g (p, \tau) \in \dot{t}_{A,\mathbb{P}} \cap M\}$ ) is equal to  $A \cap M[g]$  (here we are referring to existing  $M$ -generic filters, not filters existing in a forcing extension of  $V$ ). We say that  $M$  is weakly (or strongly)  $A$ -closed if it is weakly (or strongly)  $(A, \mathbb{P})$ -closed for all partial orders  $\mathbb{P} \in M$ .

**11.3.2 Remark.** Suppose that  $M$  is an inner model of ZFC, and that  $\mathbb{P}$  is a partial order in  $M$  such that  $\mathcal{P}(\mathbb{P}) \cap M$  is countable. Let  $A \subseteq \omega^\omega$  be such that  $M$  is strongly  $(A, \mathbb{P})$ -closed. Let  $S$  be a set of ordinals in  $M$  such that  $\mathcal{P}(\mathbb{P}) \cap M$  and  $\dot{t}_{A,\mathbb{P}} \cap M$  are both definable from  $S$  in  $L[S]$ . Suppose that  $N$  is an inner model with  $S \in N$ , such that  $A \cap N \in N$  and, for every  $v \in \omega^\omega \cap N$ , there is an  $L[S]$ -generic filter  $g \subseteq \mathbb{P}$  such that  $L[S, v] = L[S][g]$ . Then, in  $N$ ,  $A \cap N$  is the set of  $v \in \omega^\omega$  such that  $L[S, v] \models v \in (\dot{t}_{A,\mathbb{P}} \cap M)_g$  for any  $L[S]$ -generic filter  $g \subseteq \mathbb{P}$  such that  $L[S, v] = L[S][g]$ . It follows that  $A \cap N$  is  $\infty$ -Borel in  $N$ .

The appendix to [27] contains a proof of the general forcing fact that whenever  $\mathbb{P}$  is a partial order and  $\tau$  is a  $\mathbb{P}$ -name for a subset of the ground model, there is a partial order  $\mathbb{P}'$  such that, whenever  $g \subseteq \mathbb{P}$  is  $V$ -generic, there is a  $V$ -generic filter  $h \subseteq \mathbb{P}'$  such that  $V[h] = V[\tau_g]$ .

**Theorem 11.3.3** (ZF + Uniformization +  $\aleph_1 \not\leq 2^{\aleph_0}$ ). *Suppose that every subset of  $\omega^\omega$  has the Baire property, and that  $\mu_\emptyset$  is an ultrafilter on  $\mathcal{D}_\emptyset$ . Then every subset of  $\omega^\omega$  is  $\infty$ -Borel.*

*Proof.* Fix  $A \subseteq \omega^\omega$ . Let  $B$  be the set of  $(x, y) \in (\omega^\omega)^2$  such that

- $x$  HC-codes a pair  $(\mathbb{P}, \tau)$  where  $\mathbb{P}$  is a countable poset and  $\tau$  is a nice  $\mathbb{P}$ -name for an element of  $\omega^\omega$ ;
- $y$  HC-codes a countable set  $\mathcal{D}$  consisting of dense open subsets of  $\mathbb{P}$ , with  $D_{\mathbb{P},\tau}^A \in \mathcal{D}$ , such that for any  $\mathcal{D}$ -generic filter  $g \subseteq \mathbb{P}$ ,
  - $\tau_g \in A$  if and only if  $\exists p \in g \ p \Vdash^* \tau \in A$ , and
  - $\tau_g \notin A$  if and only if  $\exists p \in g \ p \Vdash^* \tau \notin A$ .

The conclusion of the condition on  $y$  is equivalent to : for all  $\mathcal{D}$ -generic filters  $g \subseteq \mathbb{P}$ ,  $\tau_g \in \dot{t}_{A,\mathbb{P},g}$  if and only if  $\tau_g \in A$ . By the density of the sets  $D_{\mathbb{P},\tau}^A$  (following from the Baire property, as mentioned above), the assumption that every set of reals has the property of Baire implies that for each  $x \in \omega^\omega$  coding a pair  $(\mathbb{P}, \tau)$  as above, there is a  $y \in \omega^\omega$  such that  $(x, y) \in B$ .

Let  $f$  be a function uniformizing  $B$ , and let  $F \subseteq \omega^\omega$  code  $f$  as follows :  $x' \in F$  if and only if, letting  $x \in \omega^\omega$  be such that  $x(n) = x'(n+2)$  for all  $n \in \omega$ ,  $x \in \text{dom}(f)$  and  $f(x)(x'(0)) = x'(1)$ . For each  $w \in \omega^\omega$ , let  $N_w$  be

$$L_{\omega_1}[\text{TR}(A), \text{TR}(F), w],$$

and let  $M_w$  be

$$\text{HOD}_{\{\text{TR}(A), \text{TR}(F)\}}^{N_w}.$$

Then the models  $N_w$  and  $M_w$  are weakly  $A$ -closed and weakly  $F$ -closed, and since  $\aleph_1 \not\leq 2^{\aleph_0}$  they are models of ZFC.

**Claim 1.** *For each  $w \in \omega^\omega$ ,  $M_w$  is strongly  $A$ -closed.*

*Proof of Claim 1.* Fix  $w \in \omega^\omega$  and a partial order  $\mathbb{P} \in M_w$ . Let  $C$  be the set of nice  $\mathbb{P}$ -names in  $M_w$  for elements of  $\omega^\omega$ . Let  $\mathbb{Q}$  be  $\text{Col}(\omega, \mathbb{P})$ , and for each  $\tau \in C$ , let  $\dot{z}_\tau$  be a nice  $\mathbb{Q}$ -name in  $M_w$  for an element of  $\omega^\omega$  HC-coding the pair  $(\mathbb{P}, \tau)$ .

Since  $M_w$  is weakly  $F$ -closed,  $\dot{t}_{F, \mathbb{Q}} \cap M_w$  is in  $M_w$ . Let  $\mathcal{E}$  be a countable set of dense open subsets of  $\mathbb{Q}$ , containing each dense open subset of  $\mathbb{Q}$  from  $M_w$  and having the property that for each  $\mathcal{E}$ -generic  $h \subseteq \mathbb{Q}$  and each nice name  $\rho \in M_w$  for an element of  $\omega^\omega$ ,  $\rho_h \in F$  if and only if, for some  $p \in h$ ,  $(p, \rho) \in \dot{t}_{F, \mathbb{Q}}$  (see Remark 11.3.1). Applying this property to the collection  $J$  of nice  $\mathbb{Q}$ -names  $\rho \in M_w$  for which it is forced by  $1_{\mathbb{Q}}$  that (for some  $\tau \in C$ )  $\rho(n+2) = \dot{z}_\tau(n)$  for all  $n \in \omega$ , we get that for any  $\mathcal{E}$ -generic filter  $h \subseteq \mathbb{Q}$ , and for each  $\tau \in C$ ,  $\dot{z}_{\tau, h} \in \omega^\omega$  and  $f(\dot{z}_{\tau, h})$  is in  $M_w[h]$ , being the set of pairs  $(i, j)$  for which there exists a pair  $(p, \rho) \in \dot{t}_{F, \mathbb{Q}} \cap M_w$  with  $p \in h$ ,  $\rho \in J$  and  $(\rho_h(0), \rho_h(1)) = (i, j)$ . Fix such a  $h$ , and let  $\mathcal{D}_h$  be the collection of dense open subsets of  $\mathbb{P}$  which are HC-coded by the members of  $\{f(\dot{z}_{\tau, h}) : \tau \in C\}$ .

Let  $g \subseteq \mathbb{P}$  be an  $M_w[h]$ -generic filter. Then  $g$  is  $\mathcal{D}_h$ -generic. Since  $f(\dot{z}_{\tau, h}) \in M_w[h]$  for each  $\tau \in C$ ,  $(\dot{t}_{A, \mathbb{P}} \cap M_w)_g = A \cap M_w[g]$ . Since  $M_w[h][g] = M_w[g][h]$  (i.e.,  $(g, h)$  is generic for  $\mathbb{P} * \mathbb{Q}$ ), it follows that  $(\dot{t}_{A, \mathbb{P}} \cap M_w)_g = A \cap M_w[g]$  holds for any  $M_w$ -generic filter  $g \subseteq \mathbb{P}$ , as desired. This ends the proof of Claim 1.  $\square$

**Claim 2.** *There is a  $\emptyset$ -invariant function  $c$  on  $\omega^\omega$  such that, for each  $w \in \omega^\omega$ ,  $c(w)$  is a pair  $(S_w, \varphi_w) \in M_w$  such that, for all  $v \in \omega^\omega \cap N_w$ ,  $v \in A$  if and only if  $L[S_w, v] \models \varphi_w(S_w, v)$ .*

*Proof of Claim 2.* This follows from Remark 11.3.2, with  $M_w$  as  $M$  and  $N_w$  as  $N$ , using Theorem 10.1.2 and the fact, shown in the proof of Theorem 10.1.2, that there is a  $\mathbb{V}_{\langle \text{TR}(A), \text{TR}(F) \rangle, \omega \times \omega}$ -name  $\tau$  (using the set  $K$  defined there) such that, for all  $v \subseteq \omega \times \omega$ ,  $\tau_{G_v} = v$ .  $\square$

We complete the proof working in the inner model  $L(\text{TR}(A), \text{TR}(F), \omega^\omega)$ . Since Uniformization holds,  $\text{DC}_{\mathbb{R}}$  holds. It follows that  $L(\text{TR}(A), \text{TR}(F), \omega^\omega)$  satisfies DC, and, since  $\mu_\emptyset$  is a countably complete ultrafilter, that  $\text{Ord}^{\mathcal{D}_\emptyset} / \mu_\emptyset$  is wellfounded when computed in  $L(\text{TR}(A), \text{TR}(F), \omega^\omega)$ . The function  $w \mapsto \varphi_w$  from Claim 2 is constant on a  $\mu_\emptyset$ -cone; let  $\varphi$  be the constant value. The function



$w \mapsto S_w$  represents a set of ordinals  $S$ . The function  $w \mapsto A \cap L[w]$  represents  $A$ . For each  $v \in \omega^\omega$ , since the function  $w \mapsto S_w$  is  $\emptyset$ -invariant and each model of the form  $L[S_w, A, v]$  satisfies Choice, whenever

- $f_0, \dots, f_{n-1}$  are  $\emptyset$ -invariant functions on  $\omega^\omega$  in  $L(\text{TR}(A), \text{TR}(F), \omega^\omega)$  such that  $f_i(w) \in L[S_w, A, v]$  for all  $w \in \omega^\omega$  and  $i < n$ , and
- $\psi$  is an  $n$ -ary formula,

$L[S, A, v] \models \psi([f_0]_\emptyset, \dots, [f_{n-1}]_\emptyset)$  if and only if

$$\{w \in \omega^\omega : L[S_w, A, v] \models \psi(f_0(w), \dots, f_{n-1}(w))\} \in \mu_\emptyset.$$

For each  $v \in \omega^\omega$ ,  $v \in N_w$  for a  $\mu_\emptyset$ -cone of  $w$ . By Claim 2 then, letting  $\psi$  be the formula

$$\forall v \in \omega^\omega (v \in A \Leftrightarrow L[S, v] \models \varphi(S, v)),$$

this implies that the pair  $(S, \varphi)$  witnesses that  $A$  is  $\infty$ -Borel.  $\square$

Putting together Remark 1.1.2 (for  $\aleph_1 \not\leq 2^{\aleph_0}$  and  $\text{Baire}(\mathcal{P}(\omega^\omega))$ ) and Corollary 1.2.6 (for  $\mu_\emptyset$ ) with Theorem 11.3.3, we have the following.

**Theorem 11.3.4.** *If AD + Uniformization holds then every subset of  $\omega^\omega$  is  $\infty$ -Borel.*

## 11.4 Closure of the Suslin cardinals

Recall from Chapter 6 that a cardinal  $\kappa$  is *Suslin* if there is a set  $A \subseteq \omega^\omega$  which is  $\kappa$ -Suslin but not  $\gamma$ -Suslin for any  $\gamma < \kappa$ . In this section we prove Theorem 11.4.1 below, on the closure of the set of Suslin cardinals. Corollary 11.4.4 says that under  $\text{AD}^+$  this set is closed below  $\Theta$ . Again, we follow the presentation in [21].<sup>1</sup>

**Theorem 11.4.1** (ZF +  $\text{DC}_\mathbb{R}$ ). *Suppose that*

1.  $\kappa < \Theta$  *is a limit of Suslin cardinals,*
2.  $\kappa$ -*Determinacy holds and that*
3.  $\langle S_\alpha : \alpha < \kappa \rangle$  *is a sequence of bounded subsets of  $\kappa$  which are  $\infty$ -Borel codes for pairwise disjoint subsets of  $2^\omega$ .*

*Then  $\kappa$  is a Suslin cardinal.*

---

<sup>1</sup>Given AD,  $\text{AD}^+$  is equivalent to the Suslin cardinals being closed below  $\Theta$ .  $\Theta$ -Determinacy would follow from the existence of a strong partition cardinal above  $\Theta$ , which may not be possible. What is the relationship between the Wadge rank of a set of reals, and the least cardinal for which it is Suslin, if any?

It follows from AD alone (or, more directly part (1) of the Coding Lemma (Corollary 3.0.2) and the fact that AD implies  $\text{CC}_{\mathbb{R}}$ ; see part (1) of Remark 1.1.2) that if  $\kappa < \Theta$  is a limit of an  $\omega$ -sequence of Suslin cardinals, then  $\kappa$  is a Suslin cardinal. We will assume then for the rest of this section that  $\text{cof}(\kappa)$  is uncountable, in which case (since we are assuming  $\text{AD} + \text{DC}_{\mathbb{R}}$ ) the  $<\kappa$ -Borel sets and the  $<\kappa$ -Suslin sets coincide, by Theorem 8.1.13.

**11.4.2 Remark.** The map  $\varphi \mapsto \varphi^*$  from Section 8.2 shows that a sequence  $\langle S_\alpha : \alpha < \kappa \rangle$  as in hypothesis (3) of Theorem 11.4.1 exists if there is an  $\infty$ -Borel set which is not  $\kappa$ -Borel. By Remark 10.3.9 (assuming  $\text{AD} + \text{DC}_{\mathbb{R}}$ ) if there is an  $\infty$ -Borel set which is not  $<\kappa$ -Borel, then there is one which is not  $\kappa$ -Borel. Alternately, if  $\kappa < \Theta$ , then the existence of an  $\infty$ -Borel set which is not  $\kappa$ -Borel then follows from the assumption that every subset of  $2^\omega$  is  $\infty$ -Borel (and the Coding Lemma). In the case where  $\kappa$  is a limit of Suslin cardinals with  $\text{cof}(\kappa) > \omega$  (assuming  $\text{AD} + \text{DC}_{\mathbb{R}}$  but not necessarily that every set of reals is  $\infty$ -Borel), Theorem 8.1.13 implies that if there is a Suslin cardinal greater than or equal to  $\kappa$  then there is an  $\infty$ -Borel set which is not  $<\kappa$ -Borel, and therefore one which is not  $\kappa$ -Borel.

Let  $\mu_S$  (for  $y \in \omega^\omega$  and  $S$  a set of ordinals) be as defined in Section 9.5. Assuming AD,  $\mu_S$  is an ultrafilter on the set of  $S$ -degrees, for each such  $S$ , by Corollary 1.2.6. In this section we will use only the case  $S = \emptyset$ . While it is not necessary for the present argument, we note that the following lemma implies the corresponding version where  $\emptyset$  is replaced with an arbitrary set of ordinals.

**Lemma 11.4.3** ( $\text{ZF} + \text{AD} + \text{DC}_{\mathbb{R}}$ ). *If  $\kappa$  is a limit of Suslin cardinals of uncountable cofinality, then for all bounded  $S \subseteq \kappa$ ,  $\prod \omega_2^{L[S,x]} / \mu_\emptyset < \kappa$ .*

*Proof.* Fix a bounded  $S \subseteq \kappa$ . By Theorem 8.1.13, every  $<\kappa$ -Borel set is in  $\mathcal{S}_{<\kappa}$ . In particular the set  $\{(x, y) : y \notin L[S, x]\}$  is  $<\kappa$ -Borel by Remark 8.1.11, and therefore  $<\kappa$ -Suslin. The leftmost branch construction for uniformizing Suslin sets gives a bounded  $T \subseteq \kappa$  such that  $S <_{\mathcal{D}} T$ . It follows from part (2b) of Theorem 9.3.8 that the set of  $[x]_\emptyset$  for which  $\omega_2^{L[S,x]} < \omega_1^{L[T,x]}$  is in  $\mu_\emptyset$ . By Theorem 9.5.1,  $\prod \omega_2^{L[S,x]} / \mu_\emptyset$  is wellfounded, and isomorphic to a relation projective in the set  $\leq_\emptyset \times <_T^c$  defined there. This set is  $<\kappa$ -Borel, so also in  $\mathcal{S}_{<\kappa}$ . By Remark 6.1.5,  $\mathcal{S}_{<\kappa}$  is a projective algebra, so every relation projective in  $\leq_\emptyset \times <_T^c$  is also in  $\mathcal{S}_{<\kappa}$ . By the Kunen-Martin Theorem (Theorem 6.1.16) every prewellordering in  $\mathcal{S}_{<\kappa}$  has length less than  $\kappa$ .  $\square$

The proof of Theorem 11.4.1 is completed by appealing to Theorems 11.1.8 and 11.2.1.

*Proof of Theorem 11.4.1.* We assume that  $\kappa$  has uncountable cofinality, as the countable cofinality case was dealt with just after the statement of the theorem. For each  $\alpha < \kappa$ , let  $A_\alpha$  be the subset of  $2^\omega$  coded by  $S_\alpha$ . Using the coding at the beginning of Section 8.1, we may find  $S_{\alpha,\beta}$  ( $\alpha \leq \beta < \kappa$ ), bounded subsets of  $\kappa$ , such that each  $S_{\alpha,\beta}$  is an  $\infty$ -Borel code for the set  $A_\alpha \times A_\beta$ . Lemma

11.4.3 and Theorem 11.2.1 give a sequence  $\langle C_{\alpha,\beta} : \alpha \leq \beta < \kappa \rangle$ , where each  $C_{\alpha,\beta}$  is a strong generic  $\gamma_{\alpha,\beta}$ -code for the corresponding set  $A_\alpha \times A_\beta$ , for some  $\gamma_{\alpha,\beta} < \kappa$ . Let  $\leq$  be  $\bigcup_{\alpha \leq \beta < \kappa} A_\alpha \times A_\beta$ . Then  $\leq$  is a prewellordering of length  $\kappa$ . By the Kunen-Martin Theorem (Theorem 6.1.16),  $\leq$  cannot be  $<\kappa$ -Suslin. The set  $\bigcup_{\alpha \leq \beta < \kappa} A_\alpha \times A_\beta$  has a strong  $\kappa$ -generic code by Theorem 11.1.8, and this code induces a  $\kappa$ -Suslin representation for  $\leq$  by Theorem 11.1.6 and the assumption of  $\kappa$ -Determinacy.  $\square$

By Remark 11.4.2, Theorem 11.4.1 has the following consequence.

**Corollary 11.4.4** (ZF + AD<sup>+</sup>). *The set of Suslin cardinals is closed below  $\Theta$ .*



## Chapter 12

# Scales from Uniformization

In this chapter we will prove Theorem 12.3.1, which says that  $\text{AD} + \text{DC} + \text{Uniformization}$  implies that every subset of  $\omega^\omega$  is Suslin. The proof is a combination of arguments of Becker, Harrington, Kechris, Steel and Woodin.

### 12.1 Ordinal determinacy in the codes

The proof of Theorem 12.3.1 uses a weaker notion of strong generic code (introduced in Section 12.2) where the players play real number codes for ordinals. The determinacy of the associated game follows from  $\text{AD}$ , by Theorem 12.1.1, which is a variation of the Harrington-Kechris theorem [4] on the determinacy (under  $\text{AD}$ ) of certain real games where the payoff set depends only on the rank of the reals played in a fixed prewellordering.

In this chapter, we will say that a sequence  $\bar{P} = \langle P_i : i \in \omega \rangle$  of subsets of  $\omega^\omega$  is *good* if, for each  $i \in \omega$ , every subset of  $\omega^\omega \times \omega^\omega$  which is projective in  $P_i$  is uniformized by a function which is  $\Sigma_1^1$  in  $P_{i+1}$  (without parameters). Recall that for any set  $P \subseteq \omega^\omega$ , there is a universal  $\Sigma_1^1(P)$  set  $U \subseteq \omega \times \omega^\omega$ , in the sense that every  $\Sigma_1^1(P)$  subset of  $\omega^\omega$  equal to  $\{x : (n, x) \in U\}$  for some  $n \in \omega$ . Moreover, such a set can be defined from  $P$ , so in particular for any sequence  $\langle P_i : i \in \omega \rangle$  of subsets of  $\omega^\omega$  there is a sequence  $\langle U_i : i \in \omega \rangle$  such that each  $U_i$  is a universal  $\Sigma_1^1(P_i)$  subset of  $\omega \times \omega^\omega$ . We will use a special case of this fact below.

We say that a sequence  $\langle \leq_i : i < \omega \rangle$  of prewellorderings of  $2^\omega$  is *suitable* if there exists a good sequence  $\langle P_i : i \in \omega \rangle$  of subsets of  $\omega^\omega$  such that, for all  $i$ ,  $\leq_i$  is projective in  $P_i$ .

Now suppose that  $Q = \langle \leq_i : i \in \omega \rangle$  is a sequence of prewellorderings of  $2^\omega$ , and, for each  $i \in \omega$ , let  $\gamma_i$  be the length of  $\leq_i$  and  $\rho_i$  be the associated rank function. Given  $A \subseteq \prod_{i \in \omega} \gamma_i$ , we let  $\mathcal{G}(Q, A)$  be the game where players  $I$  and  $II$  successively pick  $u_i \in 2^\omega$  and player  $I$  wins if and only if  $\langle \rho_i(u_i) : i \in \omega \rangle \in A$ .

Theorem 12.1.1 is a version of the Harrington-Kechris theorem from [4]. The proof we give is the Harrington-Kechris proof as modified by Becker and

Woodin.

**Theorem 12.1.1** (ZF + AD). *Let  $Q = \langle \leq_i : i < \omega \rangle$  be a suitable sequence of prewellorderings of  $2^\omega$ , and for each  $i \in \omega$  let  $\gamma_i$  be the length of  $\leq_i$ . Then for each  $A \subseteq \prod_{i \in \omega} \gamma_i$  the game  $\mathcal{G}(Q, A)$  is determined.*

The rest of this section is a proof of Theorem 12.1.1, so we assume AD throughout. We fix a suitable sequence  $Q = \langle \leq_i : i < \omega \rangle$  of prewellorderings of  $2^\omega$ , as witnessed by sets  $P_i \subseteq \omega^\omega$  ( $i \in \omega$ ). For each  $i \in \omega$  we let  $\gamma_i$  be the length of  $\leq_i$  and  $\rho_i$  be the corresponding rank function. Fix  $A \subseteq \prod_{i \in \omega} \gamma_i$ .

We say that a subtree  $T$  of  $2^{<\omega}$  is *2-perfect* if each node of  $T$  has a unique shortest pair of incompatible extensions. There is then a natural bijection between  $[T]$  and  $2^\omega$ . Let  $\mathcal{T}_P$  be the set of 2-perfect subtrees of  $2^{<\omega}$ .

Using the uniform existence of universal  $\Sigma_1^1(P)$  sets (discussed above), we can fix a sequence  $\langle U_i : i \in \omega \rangle$  such that each  $U_i$  is a  $\Sigma_1^1(P_i)$ -universal subset of  $\omega \times (2^\omega)^2$ . Fix a sequence  $\langle T_i : i \in \omega \rangle$  such that each  $T_i$  is a function from  $\omega \times 2^\omega$  to  $(\mathcal{T}_P)^\omega$ , and, for all  $(i, n, x, y) \in (\omega)^2 \times (2^\omega)^2$ ,  $T_i(n, x)$  is the element of  $(\mathcal{T}_P)^\omega$  which is HC-coded by the unique  $y$  with  $(n, x, y) \in U_i$ , if there exists such a  $y$ . Then, for each  $i \in \omega$ , every  $\Sigma_1^1(P_i)$  function from  $2^\omega$  to  $(\mathcal{T}_P)^\omega$  is equal to the function  $x \mapsto T_i(n, x)$ , for some  $n \in \omega$ .

The Kuratowski-Ulam theorem (see 5A.9 of [32]) implies that if every subset of  $\omega^\omega$  has the property of Baire then every wellordered union of meager sets is meager. The proof of Lemma 12.1.2 uses the Kuratowski-Ulam theorem and a diagonalization along the lines of the Recursion Theorem (Theorem 2.4.8), which results in the double use of  $m$  in the formula  $T_{i+1}(m, y)_m$ .

**Lemma 12.1.2.** *For each  $i \in \omega$  and each  $H : \omega \times (2^\omega)^2 \rightarrow 2^\omega$  projective in  $P_i$  there exists an  $m \in \omega$  such that for each  $y \in 2^\omega$ ,  $\rho_i(H(m, y, z))$  is the same for all  $z \in [T_{i+1}(m, y)_m]$ .*

*Proof.* Fix  $i \in \omega$ . Let  $B$  be the set of  $(m, y, T)$  such that  $(m, y) \in \omega \times 2^\omega$ ,  $T \in \mathcal{T}_P$  and  $\rho_i(H(m, y, z))$  is the same for all  $z \in [T]$ . Then  $B$  is projective in  $P_i$ , and, by Kuratowski-Ulam, for each  $(m, y)$  there is a  $T$  such that  $(m, y, T) \in B$ .

Since  $\langle P_j : j \in \omega \rangle$  is good, there is a  $\Sigma_1^1(P_{i+1})$  function  $g$  on  $\omega \times 2^\omega$  such that, for all  $(m, y) \in \omega \times 2^\omega$ ,  $(m, y, g(m, y)) \in B$ . Moreover, the function sending each  $y \in 2^\omega$  to the sequence  $\langle g(n, y) : n \in \omega \rangle$  is  $\Sigma_1^1(P_{i+1})$ , so for some  $m \in \omega$  we have that  $g(n, y) = T_{i+1}(m, y)_n$  for all  $(n, y) \in \omega \times 2^\omega$ . Then for all  $y \in 2^\omega$ ,  $T_{i+1}(m, y)_m = g(m, y)$ , so  $\rho_i(H(m, y, z))$  is the same for all paths  $z \in [T_{i+1}(m, y)_m]$ .  $\square$

We next define an integer game whose determinacy (which follows from AD) implies the determinacy of  $\mathcal{G}(Q, A)$ .

Fix a recursive homeomorphism  $\chi$  between  $2^\omega$  and  $2^\omega \times \omega \times (2^\omega)^2$ . Given  $i \in \omega$ , say that  $(n, w, x) \in \omega \times 2^\omega \times \omega^\omega$  is *i-good* if  $x \in [T_i(n, w)_n]$ . For each  $i \in \omega$ , let

$$F_i : \omega \times (2^\omega)^2 \rightarrow 2^\omega \times \omega \times (2^\omega)^2$$

be the function which on input  $(n, w, x)$  returns the  $\chi$ -value of the image of  $x$  under the natural bijection between  $[T_i(n, w)_n]$  and  $2^\omega$  if  $x \in [T_i(n, w)_n]$  (i.e., if

$(n, w, x)$  is  $i$ -good), and some arbitrary fixed value otherwise. For all  $i, n \in \omega$  and  $w \in 2^\omega$ , then, the function sending  $x$  to  $F_i(n, w, x)$  is a bijection between  $[T_i(n, w)_n]$  and  $2^\omega \times \omega \times (2^\omega)^2$ . Given

$$(n, w, u, m, y, z) \in \omega \times (2^\omega)^2 \times \omega \times (2^\omega)^2,$$

we let  $F_i^*(n, w, u, m, y, z)$  be the unique  $x \in [T_i(n, w)_n]$  such that  $F_i(n, w, x) = (u, m, y, z)$ .

**12.1.3 Remark.** For each  $i \in \omega$ , the functions  $F_i$  and  $F_i^*$  are defined using  $T_i$ , which was defined from  $U_i$ , which is a universal  $\Sigma_1^1(P_i)$  set. In particular,  $F_i$  and  $F_i^*$  are projective in  $P_i$ .

We let  $\mathcal{G}'(Q, A)$  be the integer game in which player  $I$  plays  $(n_0, w_0, x_0) \in \omega \times (2^\omega)^2$  and player  $II$  plays  $(m_0, y_0, z_0) \in \omega \times (2^\omega)^2$ , both one coordinate per move, as in the following diagram.

I	$n_0$	$w_0(0)$	$x_0(0)$	$w_0(1)$	$\dots$
II	$m_0$	$y_0(0)$	$z_0(0)$	$\dots$	

The game  $\mathcal{G}'_{Q,A}$ .

To determine the winner of the game, we then let, for each  $i \in \omega$ ,

$$(u_{2i}, n_{i+1}, w_{i+1}, x_{i+1})$$

be  $F_{2i}(n_i, w_i, x_i)$  and

$$(u_{2i+1}, m_{i+1}, y_{i+1}, z_{i+1})$$

be  $F_{2i+1}(m_i, y_i, z_i)$ . Player  $I$  wins this run of  $\mathcal{G}'(Q, A)$  if and only if either

- there is an  $i \in \omega$  such that  $(m_i, y_i, z_i)$  is not  $(2i+1)$ -good, and such that, for all  $j \leq i$ ,  $(n_j, w_j, x_j)$  is  $(2j)$ -good, or
- $(n_i, w_i, x_i)$  is  $(2i)$ -good for all  $i \in \omega$ , and  $\langle \rho_i(u_i) : i < \omega \rangle \in A$ .

Since  $\mathcal{G}'(Q, A)$  is an integer game, AD implies that it is determined. We will show that if either player  $I$  has a winning strategy in  $\mathcal{G}'(Q, A)$  then the same player has one in  $\mathcal{G}(Q, A)$ . This gives the determinacy of each game  $\mathcal{G}(Q, A)$  under AD.

We first fix a winning strategy  $\sigma$  for player  $I$  in  $\mathcal{G}'(Q, A)$ , and write  $(n, w, x) = \sigma(m, y, z)$  to mean that  $(n, w, x)$  is the result of player  $I$ 's moves when he plays according to  $\sigma$  and player  $II$  plays  $(m, y, z)$ . Our desired strategy for  $\mathcal{G}(Q, A)$  will be induced by the function  $E$  given in the following lemma.

**Lemma 12.1.4.** *There is a function  $E: (2^\omega)^{<\omega} \rightarrow (\omega \times 2^\omega \times 2^\omega)^{<\omega}$  such that*

1. for all  $t, t' \in (2^\omega)^{<\omega}$ , if  $t \subseteq t'$  then  $E(t) \subseteq E(t')$ ;
2. for all  $t \in (2^\omega)^{<\omega}$ ,  $\text{length}(E(t)) = \text{length}(t) + 1$ ;
3. for all  $u_1, u_3, \dots, u_{2k-1} \in 2^\omega$ , if

$$E(u_1, u_3, \dots, u_{2k-1}) = ((m_0, y_0, v_0), (m_1, y_1, v_2), \dots, (m_k, y_k, v_{2k})),$$

then for each  $z_k \in [T_{2k+1}(m_k, y_k)_{m_k}]$ , if

- for all  $i < k$ ,  $z_i = F_{2i+1}^*(m_i, y_i, u_{2i+1}, m_{i+1}, y_{i+1}, z_{i+1})$ ;
- $(n_0, w_0, x_0) = \sigma(m_0, y_0, z_0)$ ;
- for all  $i \leq k$ ,  $(u_{2i}, n_{i+1}, w_{i+1}, x_{i+1}) = F_{2i}(n_i, w_i, x_i)$ ;

then  $\rho_{2i}(u_{2i}) = \rho_{2i}(v_{2i})$  for all  $i \leq k$ .

Granting the claim, we get a strategy for  $I$  in  $\mathcal{G}(Q, A)$  as follows: When  $II$  plays  $u_1, u_3, \dots$ ,  $I$  answers with  $v_0, v_2, \dots$ , where

$$E(u_1, u_3, \dots, u_{2k-1}) = ((m_0, y_0, v_0), (m_1, y_1, v_2), \dots, (m_k, y_k, v_{2k}))$$

for each  $k \in \omega$ .

We will show that this is indeed a winning strategy for player  $I$ . Suppose that  $II$  has played  $u_{2i+1}$  ( $i \in \omega$ ) in a run of the game and  $I$  produced  $v_{2i}$  ( $i \in \omega$ ) following this strategy. Then we have  $m_i$  and  $y_i$  ( $i \in \omega$ ) as produced by  $E$ . For each  $k \in \omega$  define  $C_0^k, C_1^k, \dots, C_k^k$  as follows :

- $C_k^k$  is the set of  $z \in [T_{2k+1}(m_k, y_k)_{m_k}]$  such that

$$F_{2k+1}(m_k, y_k, z) = (u_{2k+1}, m_{k+1}, y_{k+1}, z')$$

for some  $z' \in 2^\omega$ ;

- for each  $i < k$ ,  $C_i^k$  is the set of  $z \in [T_{2i+1}(m_i, y_i)_{m_i}]$  such that

$$F_{2i+1}(m_i, y_i, z) = (u_{2i+1}, m_{i+1}, y_{i+1}, z')$$

for some  $z' \in C_{i+1}^k$ .

Then for all  $i \leq k$  in  $\omega$ ,  $C_i^{k+1} \subseteq C_i^k$ . In particular  $\langle C_0^k : k \in \omega \rangle$  is a  $\subseteq$ -decreasing sequence of nonempty closed sets, so we may pick a  $z_0$  in their intersection. Now suppose that player  $II$  plays  $(m_0, y_0, z_0)$  in  $\mathcal{G}'(Q, A)$  against  $\sigma$ . Then (since  $z_0 \in \bigcap_{k \in \omega} C_0^k$ ) the induced values of each  $m_i$  and  $y_i$  (momentarily treating them as variables and forgetting that we have fixed values for them) according to the rules of  $\mathcal{G}'(Q, A)$  agree with the (fixed) values given by  $E$ . Moreover, the triples  $(m_i, y_i, z_i)$  are each respectively  $(2i+1)$ -good, and the corresponding values of  $u_{2i+1}$  are also as in the given run of the game.

Furthermore, if  $(n_0, w_0, x_0) = \sigma(m_0, y_0, z_0)$ , then the induced values  $(n_i, w_i, x_i)$  are each respectively  $(2i)$ -good, since  $\sigma$  is a winning strategy for  $I$ . For each



$i \in \omega$ , the triple  $(n_i, w_i, x_i)$  induces a value for  $u_{2i}$  via the function  $F_{2i}$ . For each  $k \in \omega$ , since  $z_0 \in C_0^k$ , there exist  $z_1, \dots, z_k$  satisfying the conditions in part (3) of Lemma 12.1.4, so  $\rho_{2i}(u_{2i}) = \rho_{2i}(v_{2i})$  for all  $i \leq k$ . The same is then true for all  $i \in \omega$ . Since  $I$  won the run of  $\mathcal{G}'(Q, A)$  by playing  $(n_0, w_0, x_0)$ , he wins the run of  $\mathcal{G}(Q, A)$  by playing  $v_{2i}$  ( $i \in \omega$ ).

It remains then to prove Lemma 12.1.4.

*Proof of Lemma 12.1.4.* We construct  $f(t)$  recursively on the length of  $t$ . We first consider the case where  $t = \emptyset$ , where we have to produce  $m_0, y_0$  and  $v_0$  such that, for any  $z \in [T_1(m_0, y_0)_{m_0}]$ , if  $(n_0, w_0, x_0) = \sigma(m_0, y_0, z)$ , and  $F_0(n_0, w_0, x_0) = (u, n', w', x')$ , then  $\rho_0(u) = \rho_0(v_0)$ .

For each  $y \in 2^\omega$ , let  $\sigma_y$  be the strategy for player  $I$  in  $\mathcal{G}'(Q, A)$  which is HC-coded by  $y$  if there is one, and some fixed recursive strategy otherwise. Let  $y_0 \in \text{creals}$  be such that  $\sigma = \sigma_{y_0}$ .

Let  $H$  be the function sending (a candidate)  $m \in \omega$  and  $y, z \in 2^\omega$  to the first coordinate of  $F_0(\sigma_y(m, y, z))$ . Then  $H$  is projective in  $P_0$ , so Lemma 12.1.2 gives an  $m_0 \in \omega$  such that  $\rho_0(H(m_0, y_0, z))$  is the same for all  $z \in [T_1(m_0, y_0)_{m_0}]$ . Then we can let  $v_0$  be any such value  $H(m_0, y_0, z)$ , for instance, the value produced when  $z$  is the leftmost branch of  $T_0(m_0, y_0)_{m_0}$ .

In the case where  $t \neq \emptyset$ , let  $t = (u_1, \dots, u_{2k+1})$ . We may assume that

$$E(u_1, \dots, u_{2k-1}) = ((m_0, y_0, v_0), \dots, (m_k, y_k, v_{2k}))$$

is known. We need to find  $m_{k+1}, y_{k+1}$  and  $v_{2k+2}$  such that for each  $z_{k+1} \in [T_{2k+3}(m_{k+1}, y_{k+1})_{m_{k+1}}]$ , if

- for all  $i \leq k$ ,  $z_i = F_{2i+1}^*(m_i, y_i, u_{2i+1}, m_{i+1}, y_{i+1}, z_{i+1})$ ;
- $(n_0, w_0, x_0) = \sigma(m_0, y_0, z_0)$ ;
- for all  $i \leq k+1$ ,  $(u_{2i}, n_{i+1}, w_{i+1}, x_{i+1}) = F_{2i}(n_i, w_i, x_i)$ ;

then  $\rho_{2k+2}(u_{2k+2}) = \rho_{2k+2}(v_{2k+2})$ .

Consider the function  $H$  which takes in

- $m$  (a candidate for  $m_{k+1}$ ),
- $y \in 2^\omega$  (HC-coding a tuple consisting of a strategy  $\hat{\sigma}$  for player  $I$  in  $\mathcal{G}'(Q, A)$  along with values  $(\hat{m}_0, \hat{y}_0), \dots, (\hat{m}_k, \hat{y}_k)$  in  $\omega \times 2^\omega$ ), and
- $z \in 2^\omega$

and returns the corresponding  $u_{2k+2}$  as above, with  $z$  in the role of  $z_{k+1}$ ,  $\hat{\sigma}$  in the role of  $\sigma$  and each  $\hat{m}_i$  and  $\hat{y}_i$  in the role of the corresponding  $m_i$  or  $y_i$ . By Remark 12.1.3, this function is projective in  $P_{2k+2}$ . Applying Lemma 12.1.2 (with  $y = y_{k+1}$  HC-coding the tuples consisting of  $\sigma$  and the values  $(m_0, y_0), \dots, (m_k, y_k)$ ) again we can find an  $m_{k+1}$  as desired. Again we can let  $v_{k+1}$  be the value of  $H(m_{k+1}, y_{k+1}, z)$ , where  $z$  is the leftmost branch of  $T_{2k+3}(m_{k+1}, y_{k+1})_{m_{k+1}}$ .  $\square$

We now show that if player  $II$  has a winning strategy in  $\mathcal{G}'(Q, A)$  then he also has one in  $\mathcal{G}(Q, A)$ . Fix a winning strategy  $\sigma$  for player  $II$  in  $\mathcal{G}'(Q, A)$ , and write  $(m, y, z) = \sigma(n, w, x)$  to mean that  $(m, y, z)$  is the result of player  $II$ 's moves when he plays according to  $\sigma$  and player  $I$  plays  $(n, w, x)$ . Our desired strategy for  $\mathcal{G}(Q, A)$  will be induced by the function  $E$  given in the following lemma.

**Lemma 12.1.5.** *There is a function  $E: (2^\omega)^{<\omega} \rightarrow (\omega \times 2^\omega \times 2^\omega)^{<\omega}$  such that*

1. *for all  $t, t' \in (2^\omega)^{<\omega}$ , if  $t \subseteq t'$  then  $E(t) \subseteq E(t')$ ;*
2. *for all  $t \in (2^\omega)^{<\omega}$ ,  $\text{length}(E(t)) = \text{length}(t)$ ;*
3. *for all  $u_0, u_2, \dots, u_{2k} \in 2^\omega$ , if*

$$E(u_0, u_2, \dots, u_{2k}) = ((n_1, w_1, v_1), (n_2, w_2, v_3), \dots, (n_{k+1}, w_{k+1}, v_{2k+1})),$$

*then for each  $x_{k+1} \in [T_{2k+2}(n_{k+1}, w_{k+1})_{n_{k+1}}]$ , if*

- *for all  $i \leq k$ ,  $x_i = F_{2i}^*(n_i, w_i, u_{2i}, n_{i+1}, w_{i+1}, z_{i+1})$ ;*
- *$(m_0, y_0, z_0) = \sigma(n_0, w_0, x_0)$ ;*
- *for all  $i \leq k$ ,  $(u_{2i+1}, m_{i+1}, y_{i+1}, z_{i+1}) = F_{2i+1}(m_i, y_i, z_i)$ ;*

*then  $\rho_{2i+1}(u_{2i+1}) = \rho_{2i+1}(v_{2i+1})$  for all  $i \leq k$ .*

Granting the claim, we get a strategy for  $II$  in  $\mathcal{G}(Q, A)$  as follows: When  $I$  plays  $u_0, u_2, \dots$ ,  $II$  answers with  $v_1, v_3, \dots$ , where

$$E(u_0, u_2, \dots, u_{2k}) = ((n_1, w_1, v_1), (n_2, w_2, v_3), \dots, (n_{k+1}, w_{k+1}, v_{2k+1}))$$

for each  $k \in \omega$ .

We show that this is a winning strategy for player  $II$ . Suppose that  $I$  has played  $u_{2i}$  ( $i \in \omega$ ) in a run of the game and  $II$  produced  $v_{2i+1}$  ( $i \in \omega$ ) following this strategy. Then we have  $m_i$  and  $y_i$  ( $i \in \omega$ ) as produced by  $E$ . For each  $k \in \omega$  define  $C_0^k, C_1^k, \dots, C_k^k$  as follows :

- $C_k^k$  is the set of  $x \in [T_{2k}(n_k, w_k)_{n_k}]$  such that

$$F_{2k}(n_k, w_k, x) = (u_{2k}, n_{k+1}, w_{k+1}, x')$$

for some  $x' \in 2^\omega$ ;

- for each  $i < k$ ,  $C_i^k$  is the set of  $x \in [T_{2i}(n_i, w_i)_{n_i}]$  such that

$$F_{2i}(n_i, w_i, x) = (u_{2i}, n_{i+1}, w_{i+1}, x')$$

for some  $x' \in C_{i+1}^k$ .

Then for all  $i \leq k$  in  $\omega$ ,  $C_i^{k+1} \subseteq C_i^k$ . In particular  $\langle C_0^k : k \in \omega \rangle$  is a  $\subseteq$ -decreasing sequence of nonempty closed sets, so we may pick a  $x_0$  in their intersection. Now suppose that player  $I$  plays  $(n_0, w_0, x_0)$  in  $\mathcal{G}'(Q, A)$  against  $\sigma$ . Then (since  $x_0 \in \bigcap_{k \in \omega} C_0^k$ ) the induced values of each  $n_i$  and  $w_i$  (momentarily treating them as variables and forgetting that we have fixed values for them) according to the rules of  $\mathcal{G}'(Q, A)$  agree with the (fixed) values given by  $E$ . Moreover, the triples  $(n_i, w_i, x_i)$  are each respectively  $(2i)$ -good, and the corresponding values of  $u_{2i}$  are also as in the given run of the game.

Furthermore, if  $(m_0, y_0, z_0) = \sigma(n_0, w_0, x_0)$ , then, since  $\sigma$  is a winning strategy for  $II$ , the induced values  $(m_i, y_i, z_i)$  are each respectively  $(2i+1)$ -good. For each  $i \in \omega$ , the triple  $(m_i, y_i, z_i)$  induces a value for  $u_{2i+1}$  via the function  $F_{2i+1}$ . For each  $k \in \omega$ , since  $x_0 \in C_0^k$ , there exist  $x_1, \dots, x_{k+1}$  satisfying the conditions in part (3) of Lemma 12.1.5, so  $\rho_{2i+1}(u_{2i+1}) = \rho_{2i+1}(v_{2i+1})$  for all  $i \leq k$ . The same is then true for all  $i \in \omega$ . Since  $II$  won the run of  $\mathcal{G}'(Q, A)$  by playing  $(m_0, y_0, z_0)$ , he wins the run of  $\mathcal{G}(Q, A)$  by playing  $v_{2i+1}$  ( $i \in \omega$ ).

It remains then to prove Lemma 12.1.5.

*Proof of Lemma 12.1.5.* We construct  $f(t)$  recursively on the length of  $t$ . We can let  $E(\emptyset) = \emptyset$ . Let  $t = (u_0, \dots, u_{2k})$ . We may assume that

$$E(u_1, \dots, u_{2k-2}) = ((n_1, w_1, v_0), \dots, (n_k, w_k, v_{2k-1}))$$

is known. We need to find  $n_{k+1}$ ,  $w_{k+1}$  and  $v_{2k+1}$  such that for each  $x_{k+1} \in [T_{2k+2}(n_{k+1}, w_{k+1})_{n_{k+1}}]$ , if

- for all  $i \leq k$ ,  $x_i = F_{2i}^*(n_i, w_i, u_{2i}, n_{i+1}, w_{i+1}, x_{i+1})$ ;
- $(m_0, y_0, z_0) = \sigma(n_0, w_0, x_0)$ ;
- for all  $i \leq k$ ,  $(u_{2i+1}, m_{i+1}, y_{i+1}, z_{i+1}) = F_{2i+1}(m_i, y_i, z_i)$ ;

then  $\rho_{2k+1}(u_{2k+1}) = \rho_{2k+1}(v_{2k+1})$ .

Consider the function  $H$  which takes in

- $n$  (a candidate for  $n_{k+1}$ ),
- $w \in 2^\omega$  (HC-coding a tuple consisting of a strategy  $\hat{\sigma}$  for player  $II$  in  $\mathcal{G}'(Q, A)$  along with values  $(\hat{n}_1, \hat{w}_1), \dots, (\hat{n}_k, \hat{w}_k)$  in  $\omega \times 2^\omega$ ), and
- $x \in 2^\omega$

and returns the corresponding  $u_{2k+1}$  as above, with  $x$  in the role of  $x_{k+1}$ ,  $\hat{\sigma}$  in the role of  $\sigma$  and each  $\hat{n}_i$  and  $\hat{w}_i$  in the role of the corresponding  $n_i$  or  $w_i$ . By Remark 12.1.3, this function is projective in  $P_{2k+1}$ . Applying Lemma 12.1.2 (with  $w = w_k$  HC-coding the tuple consisting of  $\sigma$  and the values  $(n_1, w_1), \dots, (n_k, w_k)$ ) again we can find an  $n_{k+1}$  as desired. We can let  $v_{2k+1}$  be the value of  $H(n_{k+1}, w_{k+1}, x)$ , where  $x$  is the leftmost branch of  $T_{2k+2}(n_{k+1}, w_{k+1})_{n_{k+1}}$ .  $\square$

## 12.2 Boundedness for $\infty$ -Borel relations

In Section 12.1 we considered a version of coded ordinal determinacy, where the players play elements of  $2^\omega$  representing ordinals via some fixed prewellorderings. The following definition gives a version of the notion of strong generic  $\infty$ -code relative to such a game.

**12.2.1 Definition.** Suppose that  $\delta$  is an infinite cardinal,  $\pi: 2^\omega \rightarrow \delta$  is a surjection and  $C$  is a generic  $\delta$ -code. Associate to  $\pi$  and  $C$  a game  $\mathcal{G}_{\pi,C}$  on  $2^\omega$ , where  $I$  and  $II$  collaborate to build a countable set  $\sigma \subseteq 2^\omega$ , and  $I$  wins if  $A_{C_{\rho[\sigma]}} \subseteq A_C$ . We say that  $\varphi$  is a  $\rho$ -strong generic  $\infty$ -code (or  $\pi$ -strong generic  $\delta$ -code) for  $A_\varphi$  if player  $I$  has a winning strategy in  $\mathcal{G}_{\pi,C}$ .

Replacing the use of  $\delta_S^\infty$ -Determinacy in the proof of Theorem 11.2.1 with AD, and using Theorem 12.1.1, we get the following. Recall that AD implies that  $\mu_S$  is an ultrafilter on  $\mathcal{D}_S$ , for each set  $S$  of ordinals (Corollary 1.2.6).

**Theorem 12.2.2 (ZF + AD).** *Let  $S$  be a set of ordinals such that the ultrapower  $\text{Ord}^{\mathcal{D}_S}/\mu_S$  is wellfounded, and  $\delta_S^\infty < \Theta$ . If  $S$  is an  $\infty$ -Borel code for a set  $A \subseteq \omega^\omega$ , then  $A$  has a  $\pi$ -strong generic  $\delta_S^\infty$ -code which is definable from  $S$ .*

We use Theorem 12.2.2 to prove Theorem 12.2.3 below, which is a boundedness result for the ranks of wellfounded relations with a suitable  $\infty$ -Borel representation. The proof uses Woodin's collapse embedding technique.

Theorem 9.5.1 and Corollary 10.2.7 together give that (under AD + DC $_{\mathbb{R}}$ ) if Uniformization holds then for each bounded subset  $S$  of  $\Theta$ ,  $\delta_S^\infty < \Theta$ . If in addition DC holds, then there are cofinally many  $\kappa < \Theta$  with the property that every subset of  $\omega^\omega$  with Wadge rank less  $\kappa$  has an  $\infty$ -Borel code  $S$  with  $\delta_S^\infty < \kappa$ .

**Theorem 12.2.3 (ZF + Lipschitz Determinacy + DC).** *Let  $\gamma < \Theta$  be an ordinal. Suppose that, for each  $i \in \omega$ ,*

- $\pi_i$  is a surjection from  $2^\omega$  onto some ordinal  $\gamma_i < \gamma$ ;
- $C_i$  is a  $\pi_i$ -strong generic  $\gamma_i$ -code for a subset of  $\omega^\omega \times \omega^\omega$ .

*Suppose also that  $R = \bigcup_{i \in \omega} A_{\varphi_{S_i}}$  is a wellfounded relation. Then the rank of  $R$  is less than  $\gamma^+$ .*

*Proof.* For each  $i \in \omega$ , let  $\Sigma_i$  be a strategy witnessing that  $C_i$  is a  $\pi_i$ -strong generic  $\gamma_i$ -code. Let  $C = (\mathbb{P}, B, \tau)$  be the join of  $\langle C_i : i \in \omega \rangle$ . Since DC holds, we may fix a suitable sequence  $\langle \leq_i : i \in \omega \rangle$  of prewellorderings of  $2^\omega$  such that the length of each  $\leq_i$  is  $\gamma$ . Let  $B \subseteq \omega^\omega$  be of Wadge rank greater than  $\gamma$ , and such that  $C$ ,  $\langle \pi_i : i \in \omega \rangle$ ,  $\langle \Sigma_i : i \in \omega \rangle$  and  $\langle \leq_i : i \in \omega \rangle$  are in  $L(B, \mathbb{R})$ . Let  $\tau$  be a  $\text{Col}(\omega, \gamma)$ -name for an element of  $\omega^\omega$  HC-coding the pair  $(\gamma, C)$ .

Let  $c: \gamma \rightarrow \text{Col}(\omega, \gamma)$  be a bijection, and for each  $i \in \omega$  let  $\rho_i$  be the rank function for  $\leq_i$ . Given a condition  $p \in \text{Col}(\omega, \gamma)$  and a set  $X \subseteq \omega^\omega$ , let  $\mathcal{G}_{\tau,p}(X)$  be the game where players  $I$  and  $II$  alternately play  $x_i \in 2^\omega$  in such a way that  $\langle c(\rho_i(x_i)) \rangle$  is a descending sequence of conditions in  $\text{Col}(\omega, \gamma)$  below  $p$ , with the

$n$ th condition deciding at least the first  $n$  digits of the realization of  $\tau$ , with player  $II$  winning if the induced realization of  $\tau$  is in  $X$  (or if player  $I$  violates the rules before player  $II$  does). By Theorem 12.1.1, each game  $\mathcal{G}_{\tau,p}(X)$  is determined.

For each condition  $p \in \text{Col}(\omega, \gamma)$ , let  $I_\tau^p$  be the set of  $X \subseteq \omega^\omega$  for which player  $I$  has a winning strategy in  $\mathcal{G}_{\tau,p}(X)$ . Then each  $I_\tau^p$  is an ideal, and  $I_\tau^p \subseteq I_\tau^q$  whenever  $p \leq q$ . Furthermore, for any  $p \in \text{Col}(\omega, \gamma)$  and  $X \subseteq \omega^\omega$ , either player  $II$  has a winning strategy in  $\mathcal{G}_{\tau,p}(X)$  or there exists a  $q \leq p$  such that player  $II$  has a winning strategy in  $\mathcal{G}_{\tau,p}(\omega^\omega \setminus X)$ . Let  $g \subset \text{Col}(\omega, \gamma)$  be  $L(B, \mathbb{R})$ -generic, and let  $U_g$  be the set of  $X \in \mathcal{P}(\omega^\omega) \cap V$  for which  $\omega^\omega \setminus X$  is in  $I_\tau^p$ , for some  $p \in g$ . Then  $U_g$  is an  $L(B, \mathbb{R})$ -ultrafilter.

Since DC holds, the ultrapower  $M_g = \text{Ult}(L(B, \mathbb{R}), U_g)$  (using functions in  $V$ ) is wellfounded, and, since Uniformization holds, the induced embedding  $j$  from  $L(B, \mathbb{R})$  to  $M_g$  is elementary. Moreover,  $j(R)$  is wellfounded. The identity function represents  $\tau_g$ , so  $\gamma$  and  $C$  are both in  $H(\aleph_1)^{M_g}$ . Since  $\gamma$  is countable in  $M_g$ ,  $\varphi$  codes a Borel relation  $R'$  in  $M_g$  which agrees with  $R$  on  $\omega^\omega \cap V$ . Since  $\text{Col}(\omega, \gamma)$  has cardinality  $\gamma$ ,  $(\gamma^+)^V \geq \omega_1^{M_g}$ . If  $R'$  is wellfounded in  $M_g$ , then we have a contradiction to  $\Sigma_1^1$ -boundedness.

To show that  $R'$  is wellfounded, we show that  $R'$  is contained in  $j(R)$ . It suffices to fix  $i \in \omega$  and show that, in  $M_g$ ,  $A_{C_i} \subseteq A_{j(C_i)}$ . Fixing such an  $i$ , consider the tree  $T$  of attempts to find a run  $\sigma$  of  $\mathcal{G}_{j(\pi_i), j(C_i)}$  according to  $j(\Sigma_i)$  where the induced  $j(C_i)_{j(\pi_i)[\sigma]}$  is isomorphic to  $C_i$ . It suffices to see that this tree has an infinite path in  $M_g$ . If there is no such path, then  $T$  has a ranking function in  $M_g$ . Since  $j(\Sigma_i)$  maps finite sequences from  $2^\omega \cap V$  to elements of  $2^\omega \cap V$ , each element of  $T \cap j[(2^\omega \cap V)^{<\omega}]$  has an extension in  $T \cap j[(2^\omega \cap V)^{<\omega}]$ . Consideration of the least rank of an element of  $T \cap j[(2^\omega \cap V)^{<\omega}]$  shows that this is impossible.  $\square$

## 12.3 Becker's argument

In this section we complete the proof of the following theorem

**Theorem 12.3.1** (ZF + AD + DC). *If Uniformization holds then every subset of  $\omega^\omega$  is Suslin.*

The rest of this section is a proof of Theorem 12.3.1. We start with an arbitrary  $A \subseteq \omega^\omega$ , which we will show to be Suslin. Since AD and Uniformization hold, we have that all subsets of  $\omega^\omega$  are  $\infty$ -Borel. Since AD, DC and Uniformization hold, there exist a suitable sequence  $\langle \leq_i : i \in \omega \rangle$  of prewellorderings of  $2^\omega$  (with lengths  $\gamma_i$  and rank functions  $\rho_i$  for  $i \in \omega$ ), as witnessed by a good sequence  $\langle P_i : i \in \omega \rangle$ , and  $\gamma < \Theta$  such that, applying Theorem 12.2.2 and the remark just before Theorem 12.2.3,

1.  $A \leq_w P_0$ ;
2.  $\text{length}(\leq_0) > \omega$ ;

3. the sequence  $\langle \text{length}(\leq_i) : i \in \omega \rangle$  is strictly increasing;
4. for all  $n \in \omega$  there is an  $a \in 2^\omega$  such that, for all  $i \in \omega$ ,  $a$  is the unique  $u \in 2^\omega$  with  $\rho_{2i}(u) = n$ ;
5.  $\gamma$  is equal to both the  $\sup_{i \in \omega} \gamma_i$  the supremum of the Wadge ranks of the sets  $P_i$ ;
6. for every subset  $A$  of  $\omega^\omega$  projective in any  $P_i$  there exist a bounded  $S \subseteq \gamma$  and a surjection  $\pi$  from  $2^\omega$  onto some ordinal below  $\gamma$  such that  $S$  is a  $\pi$ -strong  $\infty$ -Borel code for  $A$ .

Let  $\mathcal{A}$  be the set of  $a \in 2^\omega$  satisfying condition (4) for any  $n \in \omega$ .

Say that a tree on  $\gamma$  is *good* if it is a subset of  $\bigcup_{n \in \omega} \prod_{i < n} \gamma_i$ . Using a surjection  $f: \gamma \rightarrow \eta$  (and pauses) one can find a good wellfounded tree on  $\gamma$  of rank  $\eta$ , for any  $\eta < \gamma^+$ , using, for instance the tree of sequences of the form  $\langle \alpha_0, \dots, \alpha_k \rangle$  such that

- each  $\alpha_i$  is in  $\gamma_i$ ;
- if  $i < k$  and  $\alpha_i \in \omega$ , then either  $\alpha_{i+1} = \alpha_i - 1$  or  $\alpha_i = 0$  and  $\alpha_{i+1} \notin \omega$ ;
- if  $i < j \leq k$ ,  $\alpha_i = \omega + \beta$  and  $\alpha_j = \omega + \delta$ , then  $f(\beta) > f(\delta)$ .

Let  $b: 2^\omega \rightarrow (2^\omega)^\omega$  be a recursive bijection, with component functions  $b_i$  ( $i \in \omega$ ). For each  $i \in \omega$ , let  $<_i$  be the strict part of  $\leq_i$ . For each  $n \in \omega$ , let  $\prec_n^0$  be the strict prewellordering of  $(2^\omega)^n$  induced by the lexicographical order and  $<_0, \dots, <_n$ , and let  $\prec_n$  be the strict prewellordering on  $2^\omega$  defined by setting  $x \prec_n y$  to hold if and only if  $(b_0(x), \dots, b_n(x)) \prec_n^0 (b_0(y), \dots, b_n(y))$ .

Fixing a universal  $\Sigma_1^1$ -set  $U \subseteq 2^\omega \times \omega^\omega \times (2^\omega)^2$  and applying the Moschovakis Coding Lemma, we can let, for each  $n \in \omega$  and  $u \in 2^\omega$ ,  $S_n(u)$  be the subset of  $\prod_{i < n} \gamma_i$  coded by  $b_n(u)$  relative to  $\prec_n$ . That is, letting  $U(\prec_n)$  be the universal pos- $\Sigma_1^1(\prec_n)$  subset of  $(2^\omega)^3$  induced by  $U$ ,  $S_n(u)$  is the set of  $\langle \alpha_i : i < n \rangle$  such that, for some  $(v, w) \in U(\prec_n)_{b_n(u)}$ , for each  $i < n$  the  $\leq_i$ -rank of  $b_i(v)$  is  $\alpha_i$ .

Given  $m \leq n$  in  $\omega$  and  $u \in 2^\omega$ , we let  $S_n(u) \upharpoonright m$  denote the set  $\{t \upharpoonright m : t \in S_n(u)\}$ . Given  $m \leq n$  in  $\omega$  and  $u, v \in 2^\omega$ , we say that  $S_n(u)$  *extends*  $S_m(v)$  if  $S_n(u) \upharpoonright m \subseteq S_m(v)$ . We let  $\text{Tr}$  be the set of  $\langle u_n : n < m \rangle \in (2^\omega)^{<\omega}$  such that, for all positive  $n < m$ ,  $S_n(u_n)$  extends  $S_{n-1}(u_{n-1})$ . Given  $\bar{u} = \langle u_n : n \in \omega \rangle \in [\text{Tr}]$ , we write  $\text{Tr}(\bar{u})$  for  $\bigcup_{n \in \omega} S_n(u_n)$ .

Let  $\mathcal{G}_1$  be the game where players *I* and *II* alternate picking  $u_i \in 2^\omega$ , with player *II* winning if and only if either some  $u_{2i}$  is not in  $\mathcal{A}$  or

- $\langle u_{2n+1} : n \in \omega \rangle \in [\text{Tr}]$  and
- $\langle \rho_{2i}(u_{2i}) \rangle \in A$  if and only if  $\text{Tr}(\langle u_{2n+1} : n \in \omega \rangle)$  is illfounded.

If *II* has a winning strategy in this game, then  $A$  is Suslin.

Let  $\Lambda = \bigcup_{n \in \omega} \Sigma_1^1(P_n)$ , and say that a strategy  $\sigma$  for player *I* is *simple* if, for each  $n$ , the fragment of  $\sigma$  for the first  $n$  moves of the game is in  $\Lambda$ .

The results of Section 12.2 gives the following claim.

**Claim 1.** *Player I does not have a simple winning strategy in  $\mathcal{G}_1$ .*

*Proof.* Suppose that  $\Sigma$  is a simple winning strategy for player  $I$  in  $\mathcal{G}_1$ . Noting that the supremum of the ranks of the good trees on  $\gamma$  is  $\gamma^+$ , we aim to contradict Theorem 12.2.3. For each  $i \in \omega$ , let  $R_i$  be the set of triples

$$(\langle u_{2i} : i \in \omega \rangle, \langle x_{2n+1} : n < m \rangle, \langle z_j : j < m \rangle)$$

such that

- $\langle \rho_{2i}(u_{2i}) : i \in \omega \rangle \in A$ ;
- $\langle u_{2n+1} : n < m \rangle \in \text{Tr}$ ;
- $\langle u_{2i} : i \leq m+1 \rangle$  is the sequence of moves played by  $\Sigma$  against the sequence  $\langle u_{2n+1} : n < m \rangle$ ;
- for all  $n < m$ ,  $\langle \rho_{2j+1}(z_j) : j < n \rangle \in S_n(u_{2n+1})$ .

Order  $R_i$  by identity on the first coordinate, and extension in the other two. Then  $\bigcup_{i \in \omega} R_i$  is a wellfounded relation of rank  $\gamma^+$ . Item (6) above shows that the hypotheses of Theorem 12.2.3 apply to the  $R_i$ 's, giving the desired contradiction.  $\square$

We now code  $\mathcal{G}_1$  with a game  $\mathcal{G}_2$  in the same way that  $\mathcal{G}'(Q, A)$  coded  $\mathcal{G}(Q, A)$  in Section 12.1. We reuse the objects  $\langle T_i : i < \omega \rangle$ ,  $\chi$ ,  $\langle F_i : i < \omega \rangle$  and  $\langle F_i^* : i \in \omega \rangle$  defined there. Let  $Q$  be  $\langle \leq_i : i \in \omega \rangle$ .

We let  $\mathcal{G}_2$  be the integer game in which player  $I$  plays  $(n_0, w_0, x_0) \in \omega \times 2^\omega \times 2^\omega$  and player  $II$  plays  $(m_0, y_0, z_0) \in \omega \times 2^\omega \times 2^\omega$ , both one coordinate per move. To determine the winner of the game, we then let, for each  $i \in \omega$ ,

$$(u_{2i}, n_{i+1}, w_{i+1}, x_{i+1})$$

be  $F_{2i}(n_i, w_i, x_i)$  and

$$(u_{2i+1}, m_{i+1}, y_{i+1}, z_{i+1})$$

be  $F_{2i+1}(m_i, y_i, z_i)$ . Player  $I$  wins this run of  $\mathcal{G}_2(Q, A)$  if and only if either

- there is an  $i \in \omega$  such that  $(m_i, y_i, z_i)$  is not  $(2i+1)$ -good, and such that, for all  $j \leq i$ ,  $(n_j, w_j, x_j)$  is  $(2j)$ -good, or
- $(n_j, w_j, x_j)$  is  $(2j)$ -good for all  $i \in \omega$  and  $\langle u_i : i \in \omega \rangle$  is a winning run of  $\mathcal{G}_1$  for player  $I$ .

Since  $\mathcal{G}_2$  is an integer game, AD implies that it is determined.

The games  $\mathcal{G}_1$  and  $\mathcal{G}_2$  are different from the games considered in Section 12.1, in that the winning condition is not invariant under the  $\leq_{2i+1}$ -ranks of the plays made by player  $II$ . It is, however, invariant under the  $\leq_{2i}$ -ranks of the moves made by player  $I$ . Only this condition was needed in the proof of Theorem 12.1.1 to show that if player  $I$  had a winning strategy in  $\mathcal{G}'(Q, A)$  then he had one in  $\mathcal{G}(Q, A)$ . Moreover, for each  $n \in \omega$  the strategy produced

for player  $I$  in the proof of Theorem 12.1.1 used only finitely many functions of the form  $F_{2i+1}^*$  or  $F_{2i}$ . This gives the following, showing that player  $I$  does not have a winning strategy in  $\mathcal{G}_1$ , by Claim 1.

**Claim 2.** *If player  $I$  has a winning strategy in  $\mathcal{G}_2$ , then he has a simple winning strategy in  $\mathcal{G}_1$ .*

Finally, a modification of the proof in Section 12.1 gives the following claim, which completes the proof of Theorem 12.3.1.

**Claim 3.** *If player  $II$  has a winning strategy in  $\mathcal{G}_2$ , then she has one in  $\mathcal{G}_1$ .*

The rest of this section is a proof of Claim 3. For notational convenience, we will label player  $I$ 's moves as  $a_i$  ( $i \in \omega$ ), so  $a_i$  and  $u_{2i}$  refer to the same object. Suppose that  $\sigma$  is a winning strategy for player  $II$ . Given

- $\bar{a} = \langle a_0, \dots, a_k \rangle \in \omega^{<\omega}$ ,
- $\bar{s} = \langle (n_1, w_1), \dots, (n_{k+1}, w_{k+1}) \rangle \in (\omega \times 2^\omega)^{<\omega}$  and
- $x_{k+1} \in [T_{2k+2}(n_{k+1}, w_{k+1})_{n_{k+1}}]$ ,

let  $B(\bar{a}, \bar{s}, x_{k+1})$  be  $u_{2k+1}$ , where

- for all  $i \leq k$ ,  $x_i = F_{2i}^*(n_i, w_i, a_i, n_{i+1}, w_{i+1}, x_{i+1})$ ;
- $(m_0, y_0, z_0) = \sigma(n_0, w_0, x_0)$ ;
- for all  $i \leq k$ ,  $(u_{2i+1}, m_{i+1}, y_{i+1}, z_{i+1}) = F_{2i+1}(m_i, y_i, z_i)$ ;

In addition, say that  $(\bar{a}, \bar{s}, x_k)$  is coherent if the resulting sequence  $\langle u_{2i+1} : i \leq k \rangle$  is in  $\text{Tr}$ .

We define the analogous version of the function  $E$  from the proof of Theorem 12.1.1, giving us our strategy for player  $II$  in  $\mathcal{G}_1$ .

**Lemma 12.3.2.** *There is a function  $E: \mathcal{A}^{<\omega} \rightarrow (\omega \times 2^\omega)^{<\omega}$  such that*

1. *for all  $t, t' \in \mathcal{A}^{<\omega}$ , if  $t \subseteq t'$  then  $E(t) \subseteq E(t')$ ;*
2. *for all  $t \in \mathcal{A}^{<\omega}$ ,  $\text{length}(E(t)) = \text{length}(t)$ ;*
3. *for all  $\bar{a} = \langle a_0, \dots, a_k \rangle \in \mathcal{A}^{<\omega}$ , if*

$$E(a_0, \dots, a_k) = \bar{s} = \langle (n_1, w_1), \dots, (n_{k+1}, w_{k+1}) \rangle,$$

*then*

- (a) *for each  $x_{k+1} \in [T_{2k+2}(n_{k+1}, w_{k+1})_{n_{k+1}}]$  and each  $\bar{\xi} \in \prod_{i < k} \gamma_i$  there exists an  $\ell \in \omega$  such that for any*

$$x \in [T_{2k+2}(n_{k+1}, w_{k+1})_{n_{k+1}}]$$

*with  $x \restriction \ell = x'_k \restriction \ell$ ,  $\bar{\xi} \in S_k(B(\bar{a}, \bar{s}, x_{k+1}))$  and only if  $\bar{\xi} \in S_k(B(\bar{a}, \bar{s}, x))$ ;*



(b) there exist  $s_1, \dots, s_k$  such that for each

$$x_{k+1} \in [T_{2k+2}(n_{k+1}, w_{k+1})_{n_{k+1}}],$$

if for all  $i \leq k$ ,  $x_i = F_{2i}^*(n_i, w_i, a_i, n_{i+1}, w_{i+1}, x_{i+1})$ , then  $x_i \upharpoonright k = s_k$  for all  $i \leq k+1$ .

Given a function  $E$  as in Lemma 12.3.2, we define a strategy for player  $II$  in  $\mathcal{G}_1$  as follows. If at any time player  $I$  plays an element of  $2^\omega \setminus \mathcal{A}$ , then player  $II$  has won, so we can let her response be any element of  $2^\omega$ . In response to  $\bar{a} = \langle a_0, \dots, a_k \rangle \in \mathcal{A}^{<\omega}$ , letting

$$\bar{s} = E(\bar{a}) = \langle (n_1, w_1), \dots, (n_{k+1}, w_{k+1}) \rangle,$$

$II$  plays a  $v$  such that  $S_k(v)$  is the union of all sets of the form  $S_k(B(\bar{a}, E(\bar{a}), x))$  for which  $x \in [T_{2k+2}(n_{k+1}, w_{k+1})_{n_{k+1}}]$  and  $(\bar{a}, \bar{s}, x)$  is coherent. Since  $\mathcal{A}$  is countable, we can choose such a  $v$  for each such sequence from  $\mathcal{A}$ .

To see that this is a winning strategy for player  $II$ , fix  $\bar{a} = \langle a_i : i \in \omega \rangle$  and let  $\langle v_{2i+1} : i \in \omega \rangle$  be the sequence of moves played by player  $II$  in response, according to this strategy. Let

$$S_* = \bigcup_{i \in \omega} S_i(v_{2i+1}).$$

Let  $\bar{s} = \langle (n_i, w_i) : i \in \omega \setminus \{0\} \rangle$  be as given by  $\bar{a}$  and  $E$ , and for each  $k \in \omega$  let  $\bar{s} \upharpoonright k$  denote  $\langle (n_i, w_i) : i \in k \setminus \{0\} \rangle$ .

As in the proof of the Theorem 12.1.1, for each  $k \in \omega$  define  $C_0^k, C_1^k, \dots, C_k^k$  as follows :

- $C_k^k$  is the set of  $x \in [T_{2k}(n_k, w_k)_{n_k}]$  such that

$$F_{2k}(n_k, w_k, z) = \langle a_k, n_{k+1}, w_{k+1}, x' \rangle$$

for some  $z' \in 2^\omega$ ;

- for each  $i < k$ ,  $C_i^k$  is the set of  $x \in [T_{2i}(n_i, w_i)_{n_i}]$  such that

$$F_{2i}(n_i, w_i, x) = \langle a_i, n_{i+1}, w_{i+1}, x' \rangle$$

for some  $x' \in C_{i+1}^k$ .

Then  $\langle C_0^k : k \in \omega \rangle$  is a  $\subseteq$ -decreasing sequence of nonempty closed sets, so we may pick a  $x_0$  in their intersection and let player  $I$  play  $(n_0, w_0, x_0)$ . Then the induced values  $(n_i, w_i, x_i)$  according to the rules of the game agree with the values  $(n_i, w_i)$  given by  $E$ , are each respectively  $(2i)$ -good.

Let  $(m_0, y_0, z_0)$  be  $\sigma(n_0, w_0, x_0)$ , and let

$$(u_{2i+1}, m_{i+1}, y_{i+1}, z_{i+1}) = F_{2i+1}(m_i, y_i, z_i)$$

for each  $i \in \omega$ . Let  $S = \bigcup_{i \in \omega} S_i(u_{2i+1})$ . For each  $i \in \omega$ ,  $x_{i+1}$  is in

$$[T_{2i+2}(n_{i+1}, w_{i+1})_{n_{i+1}}]$$

and  $u_{2i+1}$  is equal to  $B(\bar{a} \upharpoonright (i+1), \bar{s} \upharpoonright (i+2), x_{i+1})$ . It follows that  $S \subseteq S_*$ .

Since  $\sigma$  is a winning strategy for player *II*,  $\langle \rho_{2i}(a_i) : i \in \omega \rangle$  is in  $A$  if and only if  $S$  is wellfounded. It suffices then to see that  $S$  is wellfounded if and only if  $S_*$  is. Since  $S \subseteq S_*$ , one direction of this is immediate.

For the other direction, we show that  $[S_*] \subseteq [S]$ . Suppose toward a contradiction that  $\langle \xi_i : i \in \omega \rangle$  is an infinite path through  $S_*$ , and that  $k \in \omega$  is such that  $\langle \xi_i : i < k \rangle$  is not in  $S$ . Then  $\langle \xi_i : i < k \rangle$  is not in

$$S_k(B(\bar{a} \upharpoonright (k+1), \bar{s} \upharpoonright (k+2), x_{k+1})).$$

Let  $\ell \geq k+1$  be as in conclusion (3a) of Lemma 12.3.2, with respect to  $\bar{\xi}$  and  $x_{k+1}$ . Let  $x \in [T_{2\ell}(n_\ell, w_\ell)_{n_\ell}]$  be such that  $(\bar{a} \upharpoonright (\ell+1), \bar{s} \upharpoonright (\ell+2), x)$  is coherent, and let  $x'$  be the member of  $[T_{2k+2}(n_{k+1}, w_{k+1})_{n_{k+1}}]$  induced by the functions  $F_{2\ell-2}^*, \dots, F_{2k+2}^*$ . Then  $x_{k+1} \upharpoonright \ell = x' \upharpoonright \ell$  by conclusion (3b) of Lemma 12.3.2, so  $\langle \xi_i : i < k \rangle$  is not in

$$S_k(B(\bar{a} \upharpoonright (k+1), \bar{s} \upharpoonright (k+2), x')),$$

by conclusion (3a). Since  $(\bar{a} \upharpoonright (\ell+1), \bar{s} \upharpoonright (\ell+2), x)$  is coherent  $\langle \xi_i : i < k \rangle$  is not in

$$S_\ell(B(\bar{a} \upharpoonright (\ell+1), \bar{s} \upharpoonright (\ell+2), x) \upharpoonright k).$$

It follows then that  $\bar{\xi}$  is not in  $S_\ell(v_{2\ell+1})$ . This completes the proof that  $S_*$  is wellfounded, and the proof that the strategy defined above is winning for player *II*.

It remains to prove Lemma 12.3.2.

*Proof of Lemma 12.3.2.* We can let  $E(\emptyset) = \emptyset$ . Let  $a_0, \dots, a_k \in \omega$  be given and suppose, in the case where  $k > 1$ , that  $E(a_0, \dots, a_{k-1})$  has been chosen. Fix a recursive coding  $w \mapsto e_w$  of elements of  $(\omega \times 2^\omega)^{k-1}$  by elements of  $2^\omega$ , and let  $w_k$  be such that  $E(a_0, \dots, a_{k-1}) = e_{w_k}$ . Let  $e_{w,i}$  refer to the  $i$ th pair from  $e_w$ , for each  $i < k$ .

We need to choose  $n_{k+1}$  in such a way that

$$(n_{k+1}, w_{k+1}, T_{2k+2}(n_{k+1}, w_{k+1})_{n_{k+1}}) \in D,$$

where  $D$  is the set of  $(n, w, T)$  such that  $(n, w) \in \omega \times 2^\omega$ ,  $T \subseteq \omega^{<\omega}$  is a 2-perfect tree and

- for all  $x \in [T]$  and all  $\bar{\xi} \in \prod_{i < k} \gamma_i$ , there exists an  $\ell \in \omega$  such that, for any  $x' \in [T]$  with  $x \upharpoonright \ell = x' \upharpoonright \ell$ ,

$$\bar{\xi} \in S_k(B(\langle a_0, \dots, a_k \rangle, e_w^\wedge(n, w), x))$$

if and only if

$$\bar{\xi} \in S_k(B(\langle a_0, \dots, a_k \rangle, e_w^\wedge(n, w), x'));$$

- there exist  $s_1, \dots, s_k$  such that for each  $x_k \in [T]$ , if for all  $i < k$ ,  $x_i = F_{2i}^*(e_{w,i}, a_i, e_{w,i+1}, x_{i+1})$ , then  $x_i \upharpoonright k = s_k$  for all  $i \leq k$  (letting  $e_{w,k}$  denote the pair  $(n, w)$ ).

Note that  $D$  is projective in  $P_{2k+1}$ .

For each  $(n, w)$  there is a  $T$  such that  $(n, w, T) \in D$ . To see this, note first that (since every subset of  $\omega^\omega$  has the Baire Property) for each  $\bar{\xi} \in \prod_{i < k} \gamma_i$ , the following set is comeager : the set of  $x \in \omega^\omega$  for which there is some  $\ell \in \omega$  such that, for comeagerly many  $x' \in \omega^\omega$ , if  $x \upharpoonright \ell = x' \upharpoonright \ell$ , then

$$\bar{\xi} \in S_k(B(\langle a_0, \dots, a_k \rangle, E(\langle a_0, \dots, a_{k-1} \rangle^\frown (n, w), x)))$$

if and only if

$$\bar{\xi} \in S_k(B(\langle a_0, \dots, a_k \rangle, E(\langle a_0, \dots, a_{k-1} \rangle^\frown (n, w), x'))).$$

By the Kuratowski-Ulam theorem, the intersection of these sets over all  $\bar{\xi}$  (fixing  $n$  and  $w$ ) is also comeager. Again applying the Baire Property, we can shrink this set to a somewhere comeager set of  $x$  for which the values  $x_i \upharpoonright k$  are all the same, where, for each  $i < k$ ,  $x_i = F_{2i}^*(n_i, w_i, a_i, n_{i+1}, w_{i+1}, x_{i+1})$ . Any  $T$  such that  $[T]$  is contained in this somewhere comeager set suffices.

The rest of the proof is the same as the end of the proof of Lemma 12.1.2. Since  $\langle P_i : i \in \omega \rangle$  is good, there is a  $\Sigma_1^1(P_{2k+2})$  function  $g$  on  $\omega \times 2^\omega$  such that, for all  $(n, w) \in \omega \times \omega^\omega$ ,  $(n, w, g(n, w)) \in D$ . Moreover, the function sending each  $w \in 2^\omega$  to the sequence  $\langle g(n, w) : n \in \omega \rangle$  is  $\Sigma_1^1(P_{2k+2})$ , so for some  $n_{k+1} \in \omega$  we have that  $g(n, w) = T_{2k+2}(n_{k+1}, w)_n$  for all  $(n, w) \in \omega \times 2^\omega$ . Then

$$T_{2k+2}(n_{k+1}, w_{k+1})_{n_{k+1}} = g(n_{k+1}, w_{k+1}),$$

so  $(n_{k+1}, w_{k+1}, T_{2k+2}(n_{k+1}, w_{k+1})_{n_{k+1}}) \in D$ .

□



# Chapter 13

## $\text{AD}_{\mathbb{R}}$ from scales

In this chapter we will prove the following theorem, which is due to Martin and Woodin independently.

**Theorem 13.0.1** (Martin, Woodin). *If  $\text{AD}$  holds and every subset of  $\omega^\omega$  is Suslin, then  $\text{AD}_{\mathbb{R}}$  holds.*

We will give two versions of one of the key steps in our proof of this theorem. One is a theorem of Martin (Theorem 13.0.2 below) which we will not prove. The other is a lemma which we will prove in Section 13.1, which can be used in place of Martin's theorem. Our proof of this lemma will not be entirely self-contained, either. We prove another key lemma in Section 13.2 and finish the proof of Theorem 13.0.1 in Section 13.3.

We will be using material on towers of measures, some of which we briefly review here. More thorough treatments can be found in [29, 27, 37], among other places. Given a set  $X$  and an  $i \in \omega$ , an ultrafilter  $\mu$  on  $X^{i+1}$  *projects* to the ultrafilter  $\mu'$  on  $X^i$  whose members are the set of  $A \subseteq X^i$  for which  $\{s \smallfrown \langle x \rangle : s \in A, x \in X\}$  is in  $\mu$  (we sometimes refer to ultrafilters as *measures*). Each function on  $X^i$  has a natural reinterpretation as function on  $X^{i+1}$  (ignoring the last coordinate of the input) and this reinterpretation induces a factor map  $j_{\mu', \mu}$  from a  $\mu'$ -ultrapower to a  $\mu$ -ultrapower. A *tower* of ultrafilters on  $X$  is a sequence  $\vec{\mu} = \langle \mu_i : i \in \omega \rangle$  such that each  $\mu_i$  is an ultrafilter on  $X^i$ , and each  $\mu_{i+1}$  projects to the corresponding  $\mu_i$ . There is a natural direct limit embedding associated to a tower, and the tower is said to be *wellfounded* if the corresponding image model is wellfounded. Equivalently,  $\vec{\mu}$  is wellfounded if whenever  $A_i \in \mu_i$  for each  $i \in \omega$ , there is an  $x \in X^\omega$  such that  $x \restriction i \in A_i$  for all  $i \in \omega$ .

Given a tree  $T$  on a product  $X \times Y$ , and an  $s \in X^{<\omega}$ ,  $T_s$  denotes the set  $\{t \in Y^{|s|} : (s, t) \in T\}$ . A tree  $T$  on  $\omega \times Y$  (for some set  $Y$ ) is *homogeneous* if there exists a collection of measures  $\{\mu_s : s \in \omega^{<\omega}\}$  such that,

- for each  $s \in \omega^{<\omega}$ ,  $T_s \in \mu_s$ , and

- for each  $x \in \omega^\omega$ ,  $x \in p[T]$  if and only if  $\langle \mu_{x \upharpoonright n} : n \in \omega \rangle$  is a wellfounded tower.

A tree  $T$  on  $\omega \times Y$  (for some set  $Y$ ) is *weakly homogeneous* if there exists a countable set  $\sigma$  such that, for each  $x \in \omega^\omega$ ,  $x \in p[T]$  if and only if there exists a wellfounded tower  $\langle \mu_n : n \in \omega \rangle$  such that, for each  $n \in \omega$ ,  $\mu_n \in \sigma$  and  $T_{x \upharpoonright n} \in \mu_n$ . We refer the reader to [29, 27, 37] for more on homogeneous trees and weakly homogeneous trees. A subset of  $\omega^\omega$  is *homogeneously Suslin* if it is the projection of a homogeneous tree, and *weakly homogeneously Suslin* if it is the projection of a weakly homogeneous tree (equivalently, if it is the projection, in a different sense, of a homogeneously Suslin subset of  $\omega^\omega \times \omega^\omega$ ).

The following theorem of Martin appears as Theorem 1.1 of [31].

**Theorem 13.0.2** (ZF + AD +  $\text{DC}_{\mathbb{R}}$ ; Martin). *For any  $A \subseteq \omega^\omega$ ,  $A$  is homogeneously Suslin if and only if  $A$  and its complement are Suslin.*

**13.0.3 Remark.** The construction of the Martin-Solovay tree (in [28]; see also Section 1.3 of [27] and the remarks in the proof of Lemma 13.1.4 below on adapting the arguments there to the determinacy context) implies that the complement of a weakly homogeneously Suslin set is Suslin. It is the other direction of Theorem 13.0.2 which is used in the proof of Theorem 13.0.1.

## 13.1 Weakly homogeneous trees

We start by proving a theorem of Kunen on definability of measures. Given a set  $X$ , we let  $\text{meas}(X)$  denote the set of countable complete ultrafilters on  $X$ . Recall (from part (3) of Remark 1.1.2) that AD implies that every set of reals has the property of Baire, which implies that there are no nonprincipal ultrafilters on  $\omega$ , which in turn implies that every nonprincipal ultrafilter (on any set) is countably complete. Theorem 13.1.1 shows, assuming AD +  $\text{DC}_{\mathbb{R}}$ , that every nonprincipal ultrafilter on an ordinal below  $\Theta$  is ordinal definable. Our proof is taken from [37].

In the proofs of Theorem 13.1.1 and Lemma 13.1.4 we use the ordered equivalence relation  $(\equiv_{\text{TU}}, \leq_{\text{TU}})$  given by the Turing degrees, although any ordered equivalence relation as thick as  $(\equiv_{\text{Ma}}, \leq_{\text{Ma}})$  (such as the constructibility degrees) would work just as well. By Theorem 1.2.2, AD implies TD, the statement that the  $\equiv_{\text{TU}}$ -cone measure  $\mu_{\text{TU}}$  is an ultrafilter on the set  $\mathcal{C}_{\text{TU}}$  of Turing degrees.

**Theorem 13.1.1** (ZF + AD +  $\text{DC}_{\mathbb{R}}$ ; Kunen). *For every  $\kappa < \Theta$ , every element of  $\text{meas}(\kappa)$  is ordinal definable.*

*Proof.* Fix  $\kappa < \Theta$ ,  $\mu \in \text{meas}(\kappa)$  and  $A \subseteq \omega^\omega$  of Wadge rank  $\kappa$ . By Proposition 2.5.8 and Corollary 3.0.2 (the Coding Lemma), there is a surjective function  $F: \omega^\omega \rightarrow \mathcal{P}(\kappa)$  in  $L(A, \mathbb{R})$ . Let  $\gamma < \Theta$  be greater than both  $\kappa$  and the Wadge rank of  $F^{-1}[\mu]$ . The model  $L(B, \mathbb{R})$  is the same for all  $B \subseteq \omega^\omega$  of Wadge rank  $\gamma$ , which implies that membership in this model is ordinal definable. It suffices then to fix such a  $B$  and prove that  $\mu$  is ordinal definable in  $L(B, \mathbb{R})$ . Note that  $F$  and  $\mu$  are elements of  $L(B, \mathbb{R})$ .

Since  $\text{DC}_{\mathbb{R}}$  holds,  $\text{DC}$  holds in  $L(B, \mathbb{R})$ , which means that  $(\text{Ord}^{\mathcal{C}_{\text{Tu}}} / \mu_{\text{Tu}})^{L(B, \mathbb{R})}$  is wellfounded. Let

- $F_{\mu}$  be the function on  $\omega^{\omega}$  defined by setting  $F_{\mu}(x)$  to be  $F(x)$  if  $F(x) \in \mu$ , and  $\kappa \setminus F(x)$  otherwise;
- $f_{\mu}$  be the function on  $\mathcal{C}_{\text{Tu}}$  defined by setting  $f_{\mu}(d)$  to be the least member of  $\bigcap \{F_{\mu}(x) : x \leq_{\text{Tu}} y\}$ , for any  $y \in d$  (this does not depend on the choice of  $y$ );
- $\gamma_{\mu}$  be the ordinal represented by  $f_{\mu}$  in the ultrapower  $(\text{Ord}^{\mathcal{C}_{\text{Tu}}} / \mu_{\text{Tu}})^{L(B, \mathbb{R})}$ .

Then  $\mu$  is definable from  $\gamma_{\mu}$  in  $L(B, \mathbb{R})$ , as it is the set of  $C \subseteq \kappa$  such that, for any function  $g$  representing  $\gamma_{\mu}$  in

$$(\text{Ord}^{\mathcal{C}_{\text{Tu}}} / \mu_{\text{Tu}})^{L(B, \mathbb{R})},$$

$$\{d \in \mathcal{C}_{\text{Tu}} : g(d) \in C\} \in \mu_{\text{Tu}}.$$

□

**13.1.2 Remark.** Let  $\kappa$  be an ordinal below  $\Theta$ , and let  $F: \omega^{\omega} \rightarrow \mathcal{P}(\kappa)$  be a surjection. By Theorem 13.1.1, each set  $F^{-1}[\mu]$  is ordinal definable from  $F$ . If the Wadge ranks of the members of  $\{F^{-1}[\mu] : \mu \in \text{meas}(\kappa)\}$  are not cofinal in  $\Theta$ , then there is a surjection from  $\omega^{\omega}$  onto  $\text{meas}(\kappa)$ . By Theorem 6.2.5, if the Wadge ranks of the members of  $\{F^{-1}[\mu] : \mu \in \text{meas}(\kappa)\}$  are cofinal in  $\Theta$ , then there is a counterexample to Uniformization.

Lemma 13.1.4 is our alternative to Theorem 13.0.2. By Remark 13.1.2, the lemma implies, under  $\text{AD} + \text{Uniformization}$ , that for every  $\kappa < \Theta$ , every tree on  $\omega \times \kappa$  is weakly homogeneous. The proof of the lemma, which is adapted from a MathOverflow post by Trevor Wilson,<sup>1</sup> uses the following definition.<sup>2</sup>

**13.1.3 Definition.** Given a set  $X$ , we say that an ultrafilter  $\mu$  on  $\mathcal{P}_{\aleph_1}(X)$  is *fine* if for all  $x \in X$ ,  $\{\sigma \in \mathcal{P}_{\aleph_1}(X) : x \in \sigma\}$  is in  $\mu$ .

**Lemma 13.1.4** (ZF + TD). *If  $\kappa < \Theta$  and there is a surjection from  $\omega^{\omega}$  to  $\text{meas}(\kappa^{<\omega})$ , then every tree on  $\omega \times \kappa$  is weakly homogeneous.*

*Proof.* Let  $F: \omega^{\omega} \rightarrow \text{meas}(\kappa^{<\omega})$  be a surjection, and define

$$F_*: \mathcal{C}_{\text{Tu}} \rightarrow \mathcal{P}_{\aleph_1}(\text{meas}(\kappa^{<\omega}))$$

by setting  $F_*(d)$  to be  $F[\{x \in \omega^{\omega} : \exists y \in d \ x \leq_{\text{Tu}} y\}]$ . Let  $U$  be the set of  $C \subseteq \mathcal{P}_{\aleph_1}(\text{meas}(\kappa^{<\omega}))$  such that  $F_*^{-1}[C] \in \mu_{\text{Tu}}$ . Then  $U$  is a fine, countably complete measure on  $\mathcal{P}_{\aleph_1}(\text{meas}(\kappa^{<\omega}))$ .

<sup>1</sup><https://mathoverflow.net/questions/102854/weakly-homogeneous-trees-under-ad/104632#104632>

<sup>2</sup>There is an alternate hypothesis for the following lemma, saying that  $\kappa$  is less than the sup of the Suslin cardinals. By the coding of measures theorem of Kunen, assuming  $\text{AD}$  and that there is a Suslin cardinal above  $\kappa$ , there are fewer than  $\Theta$  many measures on  $\kappa$  (or  $\kappa^{<\omega}$  for that matter.) I didn't quite say this in the previous remark.

Fix a tree  $T$  on  $\omega \times \kappa$ . As usual, given  $x \in \omega^\omega$ , we let  $T_x$  be the set  $\{s \in \kappa^{<\omega} : (x \upharpoonright |s|, s) \in T\}$ . We will show that the set of  $\sigma \in \mathcal{P}_{\aleph_1}(\text{meas}(\kappa^{<\omega}))$  witnessing the weak homogeneity of  $T$  is in  $U$ .

To do this, we define for each such infinite  $\sigma$  a game  $G_\sigma$ . This game will be closed for player  $II$ , and player  $II$  will have a winning strategy if  $\sigma$  does not witness the weak homogeneity of  $T$ . For each  $i \in \omega$ , in round  $i$  of the game  $G_\sigma$ , player  $I$  plays a measure  $\mu_i \in \sigma$  concentrating on  $\kappa^i$ , and player  $II$  plays  $k_i \in \omega$ ,  $\alpha_i \in \kappa$  and  $\beta_i < \Theta$ . Player  $I$  is required to play so that for each  $i \in \omega$ ,  $\mu_{i+1}$  projects to  $\mu_i$ . Player  $II$  is required to play so that for each  $i \in \omega$ ,  $(\langle k_0, \dots, k_i \rangle, \langle \alpha_0, \dots, \alpha_i \rangle)$  is in  $T$  and  $j_{\mu_i, \mu_{i+1}}(\beta_i) > \beta_{i+1}$ , where  $j_{\mu_i, \mu_{i+1}}$  is the factor map corresponding to the projection of  $\mu_{i+1}$  to  $\mu_i$ . The first player to break these rules loses; if there is no such player then player  $II$  wins.

I	$\mu_0$	$\mu_1$	$\mu_2$	$\dots$
II	$k_0, \alpha_0, \beta_0$	$k_1, \alpha_1, \beta_1$	$k_2, \alpha_2, \beta_2$	$\dots$

The game  $G_\sigma$ 

**Claim.** *If  $\sigma \in \mathcal{P}_{\aleph_1}(\text{meas}(\kappa^{<\omega}))$  does not witness the weak homogeneity of  $T$ , then player  $II$  has a winning strategy in  $G_\sigma$ .*

*Proof of Claim.* It follows from the definition of wellfounded tower that if  $x \in \omega^\omega \setminus p[T]$ , then there is no wellfounded tower  $\langle \mu_i : i < \omega \rangle$  consisting of measures from  $\sigma$ , with  $\{s \in \kappa^i : (x \upharpoonright i, s) \in T\} \in \mu_i$  for each  $i$ . If  $\sigma$  does not witness that  $T$  is weakly homogeneous, then, there is an  $x \in p[T]$  for which there is no wellfounded tower  $\langle \mu_i : i < \omega \rangle$  consisting of measures from  $\sigma$ , such that  $\{s \in \kappa^i : (x \upharpoonright i, s) \in T\} \in \mu_i$  for each  $i$ . We want to see that in this case there is a continuous witness to the illfoundedness of the towers of measures from  $\sigma$  concentrating on  $T_x$ . This is almost given by Lemma 1.3.8 of [27], except that the proof given there uses Choice. We will use  $\mu_{\text{Tu}}$  to avoid the use of Choice in the proof.

Let  $m: \omega \rightarrow \sigma$  be a bijection, and let  $\Sigma$  be the set of  $\langle a_0, \dots, a_\ell \rangle \in \omega^{<\omega}$  such that

- for all  $i \leq \ell$ ,  $\{s \in \kappa^i : (x \upharpoonright i, s) \in T\} \in \mu_{m(a_i)}$  and
- for all  $i < \ell$ ,  $m(a_{i+1})$  projects to  $m(a_i)$ .

Let  $K: \omega^\omega \rightarrow \mathcal{P}(\kappa^{<\omega})$  be a surjection, and define

$$K_*: \mathcal{C}_{\text{Tu}} \rightarrow \mathcal{P}_{\aleph_1}(\mathcal{P}(\kappa^{<\omega}))$$

by setting  $K_*(d)$  to be  $K[\{x \in \omega^\omega : \exists y \in d \text{ } x \leq_{\text{Tu}} y\}]$ . Let  $W$  be the set of  $D \subseteq \mathcal{P}_{\aleph_1}(\mathcal{P}(\kappa^{<\omega}))$  such that  $K_*^{-1}[D] \in \mu_{\text{Tu}}$ . Then  $W$  is a fine, countably complete measure on  $\mathcal{P}_{\aleph_1}(\mathcal{P}(\kappa^{<\omega}))$ .



Given  $\tau \in \mathcal{P}_{\aleph_1}(\mathcal{P}(\kappa^{<\omega}))$  and  $i \in \omega$ , let  $A_i^\tau = \bigcap (\tau \cap m(i))$ . Call such a  $\tau$  good if, for each infinite  $y \in [\Sigma]$ , the sequence  $\langle A_{y(i)}^\tau : i \in \omega \rangle$  witnesses that the tower  $\langle m(y(i)) : i \in \omega \rangle$  is illfounded. The leftmost branch construction shows that there is a function picking a counterexample  $y_\tau$  for each  $\tau$  which is not good. Supposing that there is no good  $\tau$ , the countable completeness of  $W$  then shows that  $y_\tau$  is the same sequence  $y_*$  for  $W$ -many  $\tau$ , which is impossible, since the tower  $\langle m(y_*(i)) : i \in \omega \rangle$  is illfounded by our assumption on  $\sigma$ .

The Coding Lemma implies that for each  $\mu \in \text{meas}(\kappa^{<\omega})$ , each function from  $\kappa^{<\omega}$  to  $\kappa^+$  represents an ordinal below  $\Theta$  in the  $\mu$ -ultrapower. Using a good  $\tau$  then, the proof of Lemma 1.3.8 of [27] gives a function  $\pi : \Sigma \rightarrow \Theta$  such that, for all  $\langle a_0, \dots, a_\ell \rangle \in \Sigma$  and all  $\ell' < \ell$ ,

$$j_{m(a_{\ell'}), m(a_\ell)}(\pi(\langle a_0, \dots, a_{\ell'} \rangle)) > \pi(\langle a_0, \dots, a_\ell \rangle).$$

This gives a winning strategy for player *II* in  $G_\sigma$ , letting  $\langle k_i : i < \omega \rangle$  be  $x$ , the sequence  $\langle \alpha_i : i < \omega \rangle$  be any fixed witness to  $x \in p[T]$ , and each value  $\beta_i$  be

$$\pi(\langle m^{-1}(\mu_0), \dots, m^{-1}(\mu_i) \rangle).$$

This ends the proof of the claim.  $\square$

It suffices now to show that for  $U$ -many  $\sigma$ , player *II* does not have a winning strategy for  $G_\sigma$ . Assume toward a contradiction that for  $U$ -almost every  $\sigma$ , player *II* does have a winning strategy. Since each game  $G_\sigma$  is closed, and since the set of possible moves for player *II* is wellorderable, we can let, for each  $\sigma$  for which *II* has a winning strategy in  $G_\sigma$ ,  $H(\sigma)$  be the winning strategy for *II* where *II* always plays the least move leading to a subgame where she still has a winning strategy.

Now let:

- $\mu_0$  be the unique measure on  $\kappa^{<\omega}$  concentrating on  $\kappa^0$ ;
- $k_0$  be the response given to  $\mu_0$  by  $H(\sigma)$  for  $U$ -many  $\sigma$ ;
- $\mu_1$  be the set of  $A \subseteq \kappa$  such that, for  $U$ -many  $\sigma$ , the ordinal  $\alpha_0^\sigma$  played by  $H(\sigma)$  in round 0 (in response to  $\mu_0$ ) is in  $A$  (then  $\mu_1$  is a countably complete measure on  $\kappa$ );
- $k_1$  be the element of  $\omega$  played by  $H(\sigma)$  in round 1 in response to  $\mu_0$  and  $\mu_1$ , for  $U$ -many  $\sigma$ ;
- $\mu_2$  be the set of  $A \subseteq \kappa^2$  such that, for  $U$ -many  $\sigma$ , letting  $\langle \alpha_0^\sigma, \alpha_1^\sigma \rangle$  be as played by  $H(\sigma)$  in rounds 0 and 1 in response to  $\mu_0$  and  $\mu_1$ ,  $\langle \alpha_0^\sigma, \alpha_1^\sigma \rangle \in A$  (then  $\mu_2$  is a countably complete measure on  $\kappa^2$  projecting to  $\mu_1$ ).

Continuing in this way, we get  $x = \langle k_i : i \in \omega \rangle \in \omega^\omega$  and a tower of measures  $\vec{\mu}$ . Each  $\mu_i$  concentrates on  $T_x$  because  $\langle \alpha_0^\sigma, \dots, \alpha_i^\sigma \rangle$  is in  $T_x$  for  $U$ -many  $\sigma$ . The tower  $\vec{\mu}$  is wellfounded since, if  $A_i \in \mu_i$  for all  $i < \omega$  then by the countable completeness of  $U$  there is a  $\sigma$  such that  $\langle \alpha_0^\sigma, \dots, \alpha_i^\sigma \rangle \in A_i$  for all  $i \in \omega$ . By the

countable completeness of  $U$  there is a  $\sigma$  containing  $\{\mu_i : i \in \omega\}$  such that  $\vec{\mu}$  is a legal play by player  $I$  against player  $II$ 's winning strategy  $H(\sigma)$ . Then player  $II$ 's moves  $\beta_i$  induced by playing  $H(\sigma)$  against  $\vec{\mu}$  witness the illfoundedness of  $\vec{\mu}$ , giving a contradiction.  $\square$

**13.1.5 Remark.** If Lipschitz Determinacy holds, and all sets of reals are weakly homogeneously Suslin, then all set of reals are homogeneously Suslin. Otherwise, the homogeneously Suslin sets would form a proper initial segment of the Wadge hierarchy, and every subset of  $\omega^\omega$  would be the continuous image of a member of this initial segment. This would induce a surjection from  $\omega^\omega$  onto  $\mathcal{P}(\omega^\omega)$ .

Remark 13.1.6 and Theorem 13.1.7 derive consequences of Lemma 13.1.4 which are not used in the remainder of this chapter.

**13.1.6 Remark.** Suppose that  $\text{AD}$  holds and that  $\kappa$  is the largest Suslin cardinal below some member  $\theta_*$  of the Solovay sequence. Let  $\Gamma$  be the set of subsets of  $\omega^\omega$  of Wadge rank less than  $\theta_*$ . Then  $\Theta^{\text{HOD}_\Gamma} = \theta_*$ , and, by the Coding Lemma,  $\mathcal{P}(\kappa)$  is contained in  $\text{HOD}_\Gamma$ . By Theorem 13.1.1, every element of  $\text{meas}(\kappa)$  is in  $\text{HOD}_\Gamma$ .

By Theorem 6.1.10 there exists in  $\text{HOD}_\Gamma$  a tree  $T$  on  $\omega \times \kappa$  projecting to a set whose complement is not Suslin in  $\text{HOD}_\Gamma$ . This tree is not weakly homogeneous in  $\text{HOD}_\Gamma$ , since otherwise the Martin-Solovay construction (see Remark 13.0.3) would produce a Suslin representation for the complement of the projection of  $T$ . There is then no surjection in  $\text{HOD}_\Gamma$  from  $\omega^\omega$  onto  $\text{meas}(\kappa^{<\omega})$  (by Lemma 13.1.4) or even onto any set of measures witnessing the weak homogeneity of  $T$  (since  $\omega^\omega \subseteq \text{HOD}_\Gamma$ , any countable subset of a surjective image of  $\omega^\omega$  in  $\text{HOD}_\Gamma$  is also in  $\text{HOD}_\Gamma$ ). If  $\theta_* < \Theta$ , then there exists by Lemma 13.1.4 a countable  $\sigma \subseteq \text{meas}(\kappa^{<\omega})$  witnessing that  $T$  is weakly homogeneously Suslin. Letting  $F: \omega^\omega \rightarrow \mathcal{P}(\kappa^{<\omega})$  be a surjection in  $\text{HOD}_\Gamma$ , the Wadge ranks of the  $F$ -preimages of the members of  $\sigma$  must be cofinal in  $\theta_*$ .

The Martin-Solovay construction (using the proof of Lemma 1.3.8 of [27]) also shows in this case that the complement of the projection of  $T$  is  $\theta_*$ -Suslin. Briefly, this follows from the fact that, for any tree  $S$  of cardinality at most  $\kappa$  (necessarily isomorphic to a tree in  $\text{HOD}_\Gamma$ ), the rank function on  $S$  maps into  $\kappa^+$ , and therefore represents an element of  $\theta_*$  in any ultrapower by an element of  $\text{meas}(\kappa^{<\omega})$ . This shows that  $\theta_*$  is the least Suslin cardinal above  $\kappa$ .

**Theorem 13.1.7.** *If  $\text{AD}^+$  holds then every nonlimit member of the Solovay sequence below  $\Theta$  has cofinality  $\omega$ .*

*Proof.* Let  $\theta_*$  be a nonlimit member of the Solovay sequence, and let  $\Gamma$  be the set of subsets of  $\omega^\omega$  of Wadge rank less than  $\theta_*$ . There exists an  $A \in \Gamma$  such that every element of  $\Gamma$  is ordinal definable from  $A$  and some element of  $\omega^\omega$ . The proof of Theorem 6.2.5 shows that Uniformization fails in  $\text{HOD}_\Gamma$ . By Theorem 8.2.8,  $\text{HOD}_\Gamma \models \text{AD}^+$ . By Corollaries 6.1.19 and 11.4.4, there is largest Suslin cardinal below  $\theta_*$ . If  $\theta_* < \Theta$ , then Remark 13.1.6 shows that  $\theta_*$  has countable cofinality.  $\square$

## 13.2 Normal measures on $\mathcal{P}_{\aleph_1}(\lambda)$

In this section we show (assuming  $\text{DC}_{\mathbb{R}}$ ) that if  $\lambda$  is a Suslin cardinal and  $\lambda$ -Determinacy holds, then there is a normal fine ultrafilter on  $\mathcal{P}_{\aleph_1}(\lambda)$ .

**13.2.1 Definition.** Given a set  $X$ , we say that an ultrafilter  $\mu$  on  $\mathcal{P}_{\aleph_1}(X)$  is *normal* if whenever  $f: \mathcal{P}_{\aleph_1}(X) \rightarrow \mathcal{P}_{\aleph_1}(X)$  is such that  $f(\sigma)$  is a nonempty subset of  $\sigma$  for each nonempty  $\sigma \in \mathcal{P}_{\aleph_1}(X)$ , there exists an  $x \in X$  such that

$$\{\sigma \in \mathcal{P}_{\aleph_1}(X) : x \in f(\sigma)\}$$

is in  $\mu$ .

**13.2.2 Remark.** Given a set  $A \subseteq \mathcal{P}_{\aleph_1}(\omega^\omega)$ , consider the game  $\mathcal{G}_A$  where players  $I$  and  $II$  alternate picking finite subsets of  $\omega^\omega$ , with player  $I$  winning if the union of all the sets chosen is in  $A$ . This game is determined under  $\text{AD}_{\mathbb{R}}$ . Solovay [36] showed that if all such games are determined, then the set of  $A$  for which player  $I$  has a winning strategy in  $\mathcal{G}_A$  is a normal fine ultrafilter on  $\mathcal{P}_{\aleph_1}(\omega^\omega)$ .

Definition 6.1.9 defines the notion of minimal tree (roughly, a tree where every node is on the leftmost branch of some member of the projection). Recall from Definition 6.1.12 that a tree  $T$  is  $\lambda$ -full if  $\{\langle(0, \alpha)\rangle : \alpha < \lambda\} \subseteq T$ , and from Lemma 6.1.14 that if  $\lambda$  is a Suslin cardinal then there exists a minimal  $\lambda$ -full tree on  $\omega \times \lambda$ .

**Theorem 13.2.3.** *If  $\lambda$  is a Suslin cardinal, and  $\lambda$ -Determinacy +  $\text{DC}_{\mathbb{R}}$  holds, then there is a normal fine measure on  $\mathcal{P}_{\aleph_1}(\lambda)$ .*

*Proof.* Fix a minimal  $\lambda$ -full tree  $T$  on  $\omega \times \lambda$ , as given by Lemma 6.1.14. For each  $\sigma \subseteq \lambda$ , let  $T| \sigma$  be the set of nodes  $(s, t)$  of  $T$  for which the range of  $t$  is contained in  $\sigma$ . We say that  $\sigma$  is *full* if, for each node  $(s, t)$  of  $T| \sigma$ , there exists an  $x \in p[T| \sigma]$  whose leftmost branch in  $T$  contains  $(s, t)$ , i.e., such that  $s \subseteq x$  and  $t \subseteq \text{lb}(T_x)$ .

Given an injection  $g: \omega \rightarrow \lambda$ , we let  $T_g$  be the tree on  $\omega \times \omega$  consisting of those  $(s, t) \in \omega^{<\omega} \times \omega^{<\omega}$  for which  $(s, g \circ t) \in T$ . We say that a function  $g: \omega \rightarrow \lambda$  is *full* if it is injective and its range is full. We call the function  $(s, t) \mapsto (s, g \circ t)$  embedding  $T_g$  into  $T$  the  *$g$ -induced map*.

**Claim.** *If  $g$  and  $g'$  are distinct full functions from  $\omega$  to  $\lambda$ , then  $T_g \neq T_{g'}$ .*

*Proof of Claim.* Since  $T$  is  $\lambda$ -full, whenever  $g$  and  $g'$  are distinct functions from  $\omega$  to  $\lambda$ , their induced maps are also distinct. It suffices then to see that, given a full function  $g: \omega \rightarrow \lambda$ , the  $g$ -induced map can be recovered from  $T_g$  without using  $g$ .

Let  $T_g^1$  be  $\{t : \exists s (s, t) \in T_g\}$ . We work recursively through  $T_g^1$  (using some wellordering of  $T_g^1$  in ordertype  $\omega$  which lists each node before its successors), building a  $\subseteq$ -increasing sequence of finite partial length-preserving embeddings  $e_k: \omega^{<\omega} \rightarrow \lambda^{<\omega}$  ( $k \in \omega$ ) such that the corresponding function  $(s, t) \mapsto (s, e_k(t))$

(which we will call  $e_k^*$ ) maps  $T_g$  (partially) into  $T$  (possibly halting if our construction breaks down, although we will eventually see that it doesn't). We let  $e_0$  be the map which sends the empty sequence to itself. Now suppose that we have built  $e_k$  (for some  $k \in \omega$ ), and that  $t$  is the least node of  $T_g^1$  in our wellordering outside the domain of  $e_k$ . Then  $t$  is of length  $n+1$ , for some  $n \in \omega$ ,  $t \restriction n$  is in the domain of  $e_k$  and, for all  $s$  such that  $(s, t \restriction n) \in T_g$ ,  $(s, e_k(t \restriction n))$  is in  $T$ . Let  $E_k$  be the set of  $\gamma < \lambda$  for which there exists a length-preserving embedding  $e: T_g^1 \rightarrow \lambda^{<\omega}$  extending  $e_k \cup \{(t, (e_k \circ (t \restriction n)) \restriction \langle \gamma \rangle)\}$  such that the map  $(s, r) \mapsto (s, e \circ r)$  embeds  $T_g$  into  $T$ . If  $E_k$  is empty, then we stop the construction. Otherwise, letting  $\gamma_*$  be the least element of  $E_k$ , we let  $e_{k+1}$  be  $e_k \cup \{(t, (e_k \circ (t \restriction n)) \restriction \langle \gamma_* \rangle)\}$ . This completes the construction.

We now verify inductively that our maps  $e_k^*$  agree with the  $g$ -induced map. For  $e_0^*$  this is clear. Supposing that this is true for a given  $k \in \omega$ , and letting  $t$  be the least node of  $T_g^1$  outside domain of  $e_k$ , we need to see that  $g(t(n))$  is the least element of  $E_k$  (where  $t$  has length  $n+1$ ). Since  $e_k^*$  agrees with the  $g$ -induced map,  $g(t(n))$  is in  $E_k$ , which means that  $e_{k+1}$  was defined. Let  $s \in \omega^{<\omega}$  be such that  $(s, t) \in T_g$ . Since  $g$  is full, there is an  $x \in p[T[\text{range}(g)]]$  whose  $T$ -leftmost branch  $(x, f)$  goes through  $(s, g \circ t)$ . This shows that  $g(t(n))$  is the least element of  $E_k$ , as desired, since the image of  $f$  under a length-preserving embedding of  $T_g^1$  into  $\lambda^{<\omega}$  witnessing the contrary would contradict the assumption that  $f$  is the leftmost branch of  $T_x$ .  $\square$

We define now a function  $F$  whose domain is the set of trees on  $\omega \times \omega$ :

- for trees of the form  $T_g$ , for  $g$  a full function from  $\omega$  to  $\lambda$ ,  $F(T_g)$  is the range of  $g$ ;
- for all other trees on  $\omega \times \omega$ ,  $F$  takes the value  $\emptyset$ .

By the claim, this function is well defined. We now apply  $\lambda$ -Determinacy.

Given a set  $A \subseteq \mathcal{P}_{\aleph_1}(\lambda)$ , let  $\mathcal{G}_A$  be the game where

- players  $I$  and  $II$  build a  $\subseteq$ -increasing sequence  $\langle p_n : n \in \omega \rangle$  of finite partial injections from  $\omega$  to  $\lambda$ , with  $n$  contained in the domain of  $p_n$  for each  $n \in \omega$ ;
- letting  $g = \bigcup_{n \in \omega} p_n$ , player  $II$  wins if  $T_g$  is  $T_h$  for some full function  $h$ , and  $F(T_g)$  is in  $A$ .

Since the tree  $T_g$  is built continuously from  $g$ ,  $\lambda$ -Determinacy implies that  $\mathcal{G}_A$  is determined. Let  $\mu$  be the set of  $A \subseteq \mathcal{P}_{\aleph_1}(\lambda)$  for which  $II$  has a winning strategy. We wish to see that  $\mu$  is a normal fine ultrafilter.

To check that  $\mu$  is fine, fix an  $\alpha < \lambda$  and a strategy  $\Sigma$  for player  $I$  in the game  $\mathcal{G}_A$ , where  $A = \{\sigma \in \mathcal{P}_{\aleph_1}(\lambda) : \alpha \in \sigma\}$ . Using  $\text{DC}_{\mathbb{R}}$  and the assumption that every node of  $T$  is part of the leftmost branch of some element of  $p[T]$ , we can find a winning play for  $II$  against  $\Sigma$  (i.e., such that the function  $g$  produced is full and has  $\alpha$  in its range).

To see that  $\mu$  is an ultrafilter, fix  $A \subseteq \mathcal{P}_{\aleph_1}(\lambda)$ . We show first that at least one of  $A$  and  $\mathcal{P}_{\aleph_1}(\lambda) \setminus A$  is in  $\mu$ . Fix strategies  $\Sigma_A$  and  $\Sigma_{\mathcal{P}_{\aleph_1}(\lambda) \setminus A}$  for player  $I$  in  $\mathcal{G}_A$

and  $\mathcal{G}_{\mathcal{P}_{\aleph_1}(\lambda) \setminus A}$ , respectively. Using  $\text{DC}_{\mathbb{R}}$  we can find a countable full set  $\sigma \subseteq \lambda$  which is closed under both  $\Sigma_A$  and  $\Sigma_{\mathcal{P}_{\aleph_1}(\lambda) \setminus A}$ , in the sense that a response by either strategy to a sequence of moves whose ranges are contained in  $\sigma$  will be a function with range contained in  $\sigma$ . Then, playing as  $II$ , we can produce runs of  $\mathcal{G}_A$  and  $\mathcal{G}_{\mathcal{P}_{\aleph_1}(\lambda) \setminus A}$  where  $I$  plays with  $\Sigma_A$  and  $\Sigma_{\mathcal{P}_{\aleph_1}(\lambda) \setminus A}$  respectively, and the range of the resulting function  $g$  is  $\sigma$  in each case. This shows that  $\Sigma_A$  and  $\Sigma_{\mathcal{P}_{\aleph_1}(\lambda) \setminus A}$  cannot both be winning strategies for  $I$ . The same argument, with  $I$  and  $II$  reversed, shows that  $A$  and  $\mathcal{P}_{\aleph_1}(\lambda) \setminus A$  cannot both be in  $\mu$ .

To see that  $\mu$  is normal, fix a function  $f$  which picks a nonempty subset of each  $\sigma \in \mathcal{P}_{\aleph_1}(\lambda)$ . For each  $\alpha \in \lambda$ , let  $A_\alpha$  be the set of  $\sigma \in \mathcal{P}_{\aleph_1}(\lambda)$  for which  $\alpha$  is in  $f(\sigma)$ . We want to see that player  $II$  has a winning strategy in some  $A_\alpha$ . Suppose toward a contradiction that the opposite holds. By the Coding Lemma, there is a surjection from  $\omega^\omega$  onto the set of all strategies for games of the form  $\mathcal{G}_A$ . Using  $\text{DC}_{\mathbb{R}}$  again we can find a countable full set  $\sigma \subseteq \lambda$  and a winning strategy  $\Sigma_\alpha$  for player  $I$  for each  $\alpha \in \sigma$ , such that  $\sigma$  is closed under  $\Sigma_\alpha$  for each  $\alpha \in \sigma$ . Then we can find runs of the games  $\mathcal{G}_{A_\alpha}$  ( $\alpha \in \sigma$ ), where in each case  $I$  plays according to  $\Sigma_\alpha$  and the range of the resulting function  $g$  is  $\sigma$ . Since  $f(\sigma)$  is a nonempty subset of  $\sigma$ , this gives a contradiction.  $\square$

We note the following theorem, whose proof is essentially the same as the proof of the claim in the proof of Theorem 13.2.3.

**Theorem 13.2.4.** *Let  $\lambda$  be an ordinal, and let  $T$  be a tree on  $\omega \times \lambda$ . Suppose that  $M$  is a transitive set such that, in  $M$ ,  $T$  is minimal. Let  $N$  be a transitive set and let  $j: M \rightarrow N$  be an elementary embedding. Then  $j[T]$  is in  $L[T, j(T)]$ .*

*Proof.* The map  $j|T$  embeds  $T$  into  $j(T)$ . We want to find this map in  $L[T, j(T)]$ . We use the fact that if

- $P$  is a wellfounded model of ZF (in particular, a forcing extension of  $L[T, j(T)]$ ),
- $S$  and  $R$  are trees in  $P$  such that, in  $P$ ,  $S$  is countable and  $T$  is wellorderable,
- $g$  is, in  $P$ , a length-preserving partial embedding of  $S$  into  $R$ , and
- there is a length-preserving embedding of  $S$  into  $R$  extending  $g$  (in  $V$ ),

then there is, in  $P$ , a length-preserving embedding of  $S$  into  $R$  extending  $g$ .

Let  $T_1$  be  $\{t \in \exists s(s, t) \in T\}$ . We work recursively through  $T_1$ , using some wellordering  $\langle t_\alpha : \alpha < \delta \rangle$  of  $T_1$  which lists each node before its successors, building a continuous  $\subseteq$ -increasing sequence of partial length-preserving functions  $e_\beta: T_1 \rightarrow j(\lambda)^{<\omega}$  ( $\beta < \delta$ ) such that

- the domain of each  $e_\beta$  is  $\{t_\alpha : \alpha < \beta\}$ ;
- for all  $\alpha < \beta$  and all  $s \in \omega^{<\omega}$ , if  $(s, t_\alpha) \in T$  then  $(s, e_\beta(t_\alpha)) \in j(T)$ .

Each  $e_\alpha$  will have  $\{(s_\beta, t_\beta) : \beta < \alpha\}$  as its domain. Then  $e_0$  is the empty function. We let  $e_1$  be the map which sends the empty sequence to itself. Now suppose that we have built  $e_\beta$ , for some  $\alpha \in [1, \delta)$ . Then  $t_\beta$  is of length  $n+1$ , for some  $n \in \omega$ , and  $t_\beta \upharpoonright n$  is in the domain of  $e_\beta$ . Let  $E_\beta$  be the set of  $\gamma < \lambda$  for which, in any generic extension of  $L[T, j(T)]$  in which  $T$  is countable, there exists a length-preserving function  $e: T_1 \rightarrow j(\lambda)^{<\omega}$  extending  $e_\beta$  such that,

- $e(t_\beta) = e_\beta(t_\beta \upharpoonright n) \smallfrown \langle \gamma \rangle$ ;
- for all  $(s, t) \in T$ ,  $(s, e(t)) \in j(T)$ .

If  $E_\beta$  is empty, then we stop the construction. Otherwise, letting  $\gamma_*$  be the least element of  $E_\beta$ , we let  $e_{\beta+1}$  be  $e_\beta \cup \{(t_\beta, e_\beta(t_\beta \upharpoonright n) \smallfrown \langle \gamma_* \rangle)\}$ . This completes the construction.

We now verify inductively that our maps  $e_\beta$  agree with  $j \upharpoonright T_1$ . For  $e_0$  and  $e_1$  this is clear. The limit case of the induction also follows immediately. Supposing that it is true for a given  $\beta < \delta$ , we need to see that  $j(t_\beta(n))$  is the least element of  $E_\beta$  (where  $t_\beta$  has length  $n+1$ ). Our induction hypothesis implies that  $j(t_\beta(n))$  is in  $E_\beta$ , which means that  $e_{\beta+1}$  was defined. Fix an  $x \in \omega^\omega \cap M$  such that  $t_\beta$  is an initial segment of  $\text{lb}(T_x)$ . Then  $j(t_\beta)$  is an initial segment of  $\text{lb}(j(T)_x)$  (as computed in any wellfounded model of  $\text{ZF}$ ). This shows that  $j(t_\alpha(n))$  is the least element of  $E_\alpha$ , as desired, since a length-preserving embedding of  $T$  into  $j(T)$  witnessing the contrary would produce a path through  $j(T)_x$  contradicting the fact that  $j(t_\beta)$  is an initial segment of  $\text{lb}(j(T)_x)$ .  $\square$

### 13.3 $\text{AD}$ implies $\text{AD}_{\mathbb{R}}$ if all sets of reals are Suslin

In this section we complete our proofs of Theorem 13.0.1. Our proofs will show that quasi- $\text{AD}_{\mathbb{R}}$  holds. Then we will be done by Remark 6.2.4. Recall that if every subset of  $\omega^\omega$  is Suslin, then Uniformization holds, and that Uniformization implies  $\text{DC}_{\mathbb{R}}$ .

We fix a bijection  $\pi: \omega \times \omega \rightarrow \omega$  such that  $\pi(i, j) < \pi(m, n)$  whenever  $i+j < m+n$ . It follows that for all odd  $n \in \omega$ ,  $\pi^{-1}[(n+1)/2] \subseteq (n+1) \times \omega$ . Let  $\pi_*$  be the induced bijection from  $(\omega^\omega)^\omega \rightarrow \omega^\omega$  (so  $\pi_*(\langle x_n : n \in \omega \rangle)(\pi(i, j)) = x_i(j)$  for all  $i, j \in \omega$ ). We fix a real game  $\mathcal{G}$ , and let  $A$  be the  $\pi_*$  image of its payoff set. By either Theorem 13.0.2 or Lemma 13.1.4 and Remark 13.1.5, we may fix  $\kappa$ -homogeneous trees  $S$  and  $T$  (on  $\omega \times \kappa$ , for some cardinal  $\kappa$ ) such that  $p[S] = A$  and  $p[T] = \omega^\omega \setminus A$ .

The key remaining step is given in Lemma 13.3.1 below. Note that in the statement of the lemma the trees  $S$  and  $T$  are homogeneous in  $V$ , but not necessarily in  $M$ .

**Lemma 13.3.1** ( $\text{ZF} + \text{AD}$ ). *Let  $\kappa$  be an ordinal. Let  $\mathcal{G}$  be a real game with payoff set  $A \subseteq (\omega^\omega)^\omega$ , and let  $S$  and  $T$  be homogeneous trees on  $\omega \times \kappa$  such that  $p[S] = \pi_*[A]$  and  $p[T] = \omega^\omega \setminus \pi_*[A]$ . Let  $M$  be an inner model of  $\text{ZF}$  containing  $S$  and  $T$ , such that  $\omega^\omega \cap M$  is countable. Then, in  $M$ , the real game with payoff set  $\pi_*^{-1}[p[S]]$  is quasi-determined.*

*Proof.* Let  $\mathcal{G}_M$  be the real game with payoff set  $\pi_*^{-1}[\text{p}[S]]$ , as computed in  $M$ . We define games  $\mathcal{G}_I^*$  and  $\mathcal{G}_{II}^*$ , both in  $M$ . In  $\mathcal{G}_I^*$ , players  $I$  and  $II$  alternate playing  $x_i \in \omega^\omega \cap M$ . In each turn,  $I$  also plays  $\alpha_{i/2} \in \kappa$ . Player  $I$  wins if and only if  $(\pi_*(\langle x_i : i < \omega \rangle), \langle \alpha_i : i < \omega \rangle)$  is in  $[S]$ .

I	$x_0, \alpha_0$	$x_2, \alpha_1$	$x_4, \alpha_2$	...
II		$x_1$	$x_3$	...

The game  $\mathcal{G}_I^*$ .

In the game  $\mathcal{G}_{II}^*$ , players  $I$  and  $II$  alternate playing  $x_i \in \omega^\omega \cap M$ . In each turn,  $II$  also plays  $\alpha_{(i-1)/2} \in \kappa$ . Player  $II$  wins if and only if

$$(\pi_*(\langle x_i : i < \omega \rangle), \langle \alpha_i : i < \omega \rangle)$$

is in  $[T]$ .

I	$x_0$	$x_2, \alpha_1$	$x_4$	...
II		$x_1, \alpha_0$	$x_3, \alpha_1$	...

The game  $\mathcal{G}_{II}^*$ .

The first of these games is closed, and the second is open, so they are both quasi-determined. Moreover, for each game there is a quasi-strategy in  $M$  which is a winning quasi-strategy in  $V$ , since  $M$  and  $V$  compute the same ranking function when applying the proof of open determinacy to these games. Since  $\omega^\omega \cap M$  is countable, such a winning quasi-strategy can be converted in  $V$  into a winning strategy.

We want to see that, in  $M$ , either  $I$  has a winning quasi-strategy in  $\mathcal{G}_I^*$  or  $II$  has a winning quasi-strategy in  $\mathcal{G}_{II}^*$ , since the corresponding player could use his strategy to win the game  $\mathcal{G}_M$ . It suffices then to derive a contradiction from the assumption that, in  $V$ , player  $II$  has a winning strategy  $\Sigma_{II}$  in  $\mathcal{G}_I^*$  and player  $I$  has a winning strategy  $\Sigma_I$  in  $\mathcal{G}_{II}^*$ . This contradiction follows from the fact that, in  $V$ , if player  $II$  has a winning strategy in  $\mathcal{G}_I^*$ , then player  $II$  has a winning strategy in  $\mathcal{G}_M$  and, similarly that if player  $I$  has a winning strategy in  $\mathcal{G}_{II}^*$ , then player  $I$  has a winning strategy in  $\mathcal{G}_M$ . These facts are essentially the same, and essentially the same as Martin's theorem that homogeneously Suslin sets are determined (see Exercise 32.2 of [10], for instance). We give a proof for the case where player  $I$  has a winning strategy in  $\mathcal{G}_{II}^*$ .

Let  $\Sigma_I$  be such a strategy. Let  $\{\nu_s : s \in \omega^{<\omega}\}$  be a set of measures witnessing that  $T$  is homogeneously Suslin. Then for each  $s \in \omega^{<\omega}$ ,  $\nu_s$  is a countably

complete ultrafilter on  $\kappa^{|s|}$ , and  $\{t \in \kappa^{|s|} : (s, t) \in T\}$  is in  $\nu_s$ . Define a strategy  $\Sigma_*$  for player  $I$  in  $\mathcal{G}_M$  as follows. Let  $\Sigma_*(\langle \rangle) = \Sigma_I(\langle \rangle)$ . Now fix a sequence  $\bar{x} = \langle x_0, \dots, x_n \rangle \in (\omega^\omega \cap M)^{<\omega}$  with  $n$  odd, and let  $s_{\bar{x}} \in \omega^{(n+1)/2}$  be the common initial segment of length  $(n+1)/2$  of the  $\pi_*$ -values of the elements of  $(\omega^\omega)^\omega$  extending  $\bar{x}$ . Let  $\Sigma_*(\bar{x})$  be the unique  $y \in \omega^\omega \cap M$  such that the set

$$R_{\bar{x}}^y = \{t \in \kappa^{(n+1)/2} : \Sigma_I(\langle x_0, (x_1, t(0)), \dots, (x_n, t((n-1)/2)) \rangle) = y\}$$

is in  $\nu_{s_{\bar{x}}}$ . Now suppose that  $\langle x_i : i < \omega \rangle$  is a run of  $G_M$  where  $I$  has played according to  $\Sigma_*$  and lost. Since the measures  $\nu_s$  witness the homogeneity of  $T$ , there is an  $f \in \kappa^\omega$  such that  $f \restriction ((n+1)/2)$  is in  $R_{\langle x_i : i < n \rangle}^{x_{n+1}}$  for each odd  $n \in \omega$ . Then  $\langle x_0, (x_1, f(0)), x_2, (x_3, f(1)), \dots \rangle$  is a run of  $G_{II}^*$  where  $I$  plays according to  $\Sigma_I$  and loses, giving a contradiction.  $\square$

*Proof of Theorem 13.0.1.* Fix  $A \subseteq (\omega^\omega)^\omega$  and let  $S$  and  $T$  be homogeneous trees projecting to  $\pi_*[A]$  and its complement respectively, on  $\omega \times \kappa$ , for some  $\kappa < \Theta$ . Let  $\lambda$  be a Suslin cardinal greater than  $\Theta^{\text{HOD}_{\{S, T\}}}$  (which exists by Remark 6.1.7 if all subsets of  $\omega^\omega$  are Suslin), and let  $B \subseteq \omega^\omega$  have Wadge rank greater than  $\lambda$ . We work in  $L(B, \mathbb{R})$ , which satisfies DC and  $\lambda$ -Determinacy (by Corollary 7.0.3 and Remark 7.0.4).

By the proof of Lemma 10.1.7, we may fix a partial order  $P$  on  $\Theta^{\text{HOD}_{\{S, T\}}}$  and a set  $K \subseteq \Theta^{\text{HOD}_{\{S, T\}}}$ , both in  $\text{HOD}_{\{S, T\}}$ , such that, in any forcing extension of  $V$  by  $\text{Col}^*(\omega, \mathbb{R})$ , there is an  $L[P, K]$ -generic filter  $G \subseteq P$  such that  $\mathbb{R}^V$  is contained in  $L[P, K][G]$ .

Let  $\mu_\lambda$  be a normal fine measure on  $\mathcal{P}_{\aleph_1}(\lambda)$ , as given by Theorem 13.2.3. For each countable  $\sigma \subseteq \lambda$ , let  $P_\sigma$  and  $K_\sigma$  be the restrictions of  $P$  to  $\sigma \cap \Theta^{\text{HOD}_{\{S, T\}}}$ . By Lemma 13.3.1, for any such  $\sigma$ , in any inner model of any forcing extension of  $L[S, T, P_\sigma, K_\sigma]$  by  $P_\sigma$ , the real game with payoff set  $\pi_*^{-1}[p[S]]$  is determined.

Since  $\mu_\lambda$  is a normal fine measure, the ultraproduct

$$\prod_{\sigma \in \mathcal{P}_{\aleph_1}(\lambda)} L[S, T, P_\sigma, K_\sigma] / \mu_\lambda$$

is a wellfounded model  $L[S', T', P, K]$ , and by elementarity, in any inner model of any forcing extension of  $L[S', T', P, K]$  by  $P$ , the real game with payoff set  $\pi_*^{-1}[p[S']]$  is determined. However,  $p[S'] = p[S] = \pi_*[A]$ , and  $L[S', T', P, K](\mathbb{R})$  is such an inner model, which shows that the real game with payoff set  $\pi_*^{-1}[p[S]]$  is determined.  $\square$

Combining Theorem 11.2.3 and 13.0.1 with Remark 6.2.2 (i.e.,  $\text{AD}_{\mathbb{R}}$  implies Uniformization) gives the following.

**Theorem 13.3.2** ( $\text{ZF} + \text{AD}^+$ ). *The following are equivalent.*

1. *There is no  $\leq_{\mathcal{D}}$ -maximal set of ordinals.*
2. *The Suslin cardinals are cofinal in  $\Theta$ .*
3. *Every subset of  $\omega^\omega$  is Suslin.*



4.  $\text{AD}_{\mathbb{R}}$ 

## 5. Uniformization

By Theorems 11.4.1 and 13.3.2, the context of the following theorem is exactly the case where  $\text{AD}^+$  holds and  $\text{AD}_{\mathbb{R}}$  fails.

**Theorem 13.3.3** ( $\text{ZF} + \text{AD}^+$ ). *If there is a largest Suslin cardinal then there is a set of ordinals  $S$  such that  $\mathcal{P}(\omega^\omega)$  is contained in  $L(S, \mathbb{R})$ .*

*Proof.* By the Moschovakis Coding Lemma, if there is a largest Suslin cardinal, then some subset of  $\omega^\omega$  is not Suslin. It follows by Theorem 11.2.3 that there is a  $\leq_{\mathcal{D}}$ -maximal set of ordinals. From this and Theorem 9.6.6 it follows that there is a set  $S$  of ordinals such that  $\mathcal{P}(\omega^\omega)$  is contained in  $L(S, \mathbb{R})$ .  $\square$

Our proof of Theorem 13.0.1 (using Theorem 13.0.2) gives the following local version.

**Theorem 13.3.4** ( $\text{ZF} + \text{AD} + \text{DC}_{\mathbb{R}}$ ). *Suppose that*

- $A \subseteq (\omega^\omega)^\omega$  has the property that  $\pi_*^{-1}[A]$  and  $\omega^\omega \setminus \pi_*^{-1}[A]$  are both  $\kappa$ -homogeneously Suslin as witnessed by trees  $S$  and  $T$  on  $\omega \times \kappa$ , and
- $\lambda \geq \Theta^{\text{HOD}_{\{S, T\}}}$  is a Suslin cardinal such that  $\lambda$ -Determinacy holds.

*Then the real game with payoff set  $A$  is determined.*

3

---

<sup>3</sup>The rest of this section is from an email from Howard.  $\text{ZF} + \text{AD}$  does not imply  $\text{AD}_{\mathbb{R}}$  for Suslin-co-Suslin sets. In fact,  $\text{ZF} + \text{AD}$  does not imply that  $\Pi_1^1$  games on reals are determined. The real theorem is this.

**Theorem 13.3.5** ( $\text{ZF} + \text{DC}$ ). *All  $\Pi_1^1$  games on reals are determined if and only if the sharp of  $L(\mathbb{R})$  exists.*

Therefore, in  $L(\mathbb{R})$ , there is a non-determined  $\Pi_1^1$  game. ( $\text{AD}$  is irrelevant.) This theorem was proved sometime in the 20th century by somebody (but not Howard). The theorem may not be published, and the proof may not even be available anywhere. What follows is a weaker theorem which is enough to answer Question 12.

**Theorem 13.3.6** ( $\text{ZF} + \text{DC} + V = L(\mathbb{R})$ ). *There exists a  $\Pi_1^1$  subset of  $(\omega^\omega)^\omega$  such that the corresponding real game is not determined.*

*Proof.* Suppose the theorem is false. Let  $M$  be the minimal model of  $\text{ZF}^- + \text{DC} + V = L(\mathbb{R})$  + “all  $\Pi_1^1$  games on reals are determined” containing all reals.

Work in  $V (= L(\mathbb{R}))$ . A Lowenheim-Skolem argument, using  $\text{DC}$ , shows that given any strategy  $\sigma$  for either player, for a game on reals:

(\*) There exists a countable elementary substructure  $M'$  of  $M$  and there exists a countable set of reals,  $S$ , such that  $\mathbb{R} \cap M' = S$  and  $S$  is closed under  $\sigma$ .

Let  $\mathcal{G}$  be the following game. On each move both players play a real and Player II also plays two elements of 2. After  $\omega$  moves, the two players together have played a countable set of reals,  $S$ , and Player II has played  $w, x$  in  $2^\omega$ . Player II wins the run of  $\mathcal{G}$  if

- $w$  is a code for a countable ordinal  $\alpha$ ,
- $(\mathbb{R} \cap L_\alpha(S)) = S$ ,

- 
- $L_\alpha(S)$  is a model of “ $V$  is the minimal model of  $\text{ZF}^- + \text{DC} + V = L(\mathbb{R}) + \text{all } \Pi_1^1 \text{ games on reals are determined containing all reals}$ ” and
  - $x$  is the set of true sentences of  $N$ .

Now (\*) implies that Player I cannot have a winning strategy. And (\*) implies that if  $\sigma$  is ANY winning strategy for Player II, then the  $x$  played by  $\sigma$  is constant, i.e., independent of Player I's moves, and is the set of true sentences of  $M$ .

A winning strategy in  $M$  is a winning strategy in  $V$ . Since there exists a winning strategy for  $\mathcal{G}$  in  $M$ , truth in  $M$  is definable in  $M$ . Contradiction.  $\square$

# Chapter 14

## Questions

1. Does TD follow from  $\aleph_1 \not\leq 2^{\aleph_0}$  plus the assumption that the cone measure on the constructibility degrees is an ultrafilter?
2. Does AD imply  $\text{DC}_{\mathbb{R}}$ ?
3. Does AD imply that the Wadge hierarchy is wellfounded? (yes if  $\text{DC}_{\mathbb{R}}$  is also assumed)
4. Does  $\text{AD} + \text{DC}_{\mathbb{R}}$  imply that all subsets of  $\omega^\omega$  are  $\infty$ -Borel?
5. Does  $\text{AD} + \text{DC}_{\mathbb{R}}$  imply that the collection of  $\infty$ -Borel subsets of  $\omega^\omega$  is closed under wellordered unions?
6. Does  $\text{AD} + \text{DC}_{\mathbb{R}}$  imply  $<\Theta$ -Determinacy?
7. Does  $\text{AD} + \text{DC}_{\mathbb{R}}$  imply  $\omega_2$ -Determinacy?
8. Does AD imply  $\text{WF}_{\text{Tu}}$ ?
9. Does AD imply that  $\chi_{\text{B}} = \kappa_{\text{B}}$ ? (yes if  $\text{DC}_{\mathbb{R}}$  is also assumed)
10. Does AD imply that  $\kappa_{\text{B}} = \lambda_{\text{B}}$ ? (yes if  $\text{DC}_{\mathbb{R}}$  is also assumed)
11. Does  $\text{TD} + \text{CC}_{\mathbb{R}}$  imply AD?
12. Does TD imply  $\text{DC}_{\mathbb{R}}$ ?
13. Does  $\text{AD}_{\mathbb{R}}$  imply  $<\Theta$ -Determinacy?
14. Does  $\text{AD} + \text{DC}_{\mathbb{R}}$  imply that there are no strong partition cardinals greater than (or equal to)  $\Theta$ ?
15. Does  $\text{AD}^+$  imply that for each set of ordinals  $S$  there is a bounded  $T \subseteq \Theta$  such that  $S \equiv_{\mathcal{D}} T$ ?
16. Does AD imply that each pointclass  $\Sigma_1^2(A)$  is  $\forall^{\omega^\omega}$ -closed? ( $\text{AD}^+$  does.)

17. Must Uniformization hold if AD holds and the Solovay sequence has limit length?

# Bibliography

- [1] Andrés Eduardo Caicedo and Richard Ketchersid. A trichotomy theorem in natural models of  $\text{AD}^+$ . In *Set theory and its applications*, volume 533 of *Contemp. Math.*, pages 227–258. Amer. Math. Soc., Providence, RI, 2011.
- [2] Sean Cody. Note on  $\text{TD} + \text{DC} + V = L(\mathbb{R})$  implies  $\text{AD}$ .
- [3] David Gale and F. M. Stewart. Infinite games with perfect information. In *Contributions to the theory of games, vol. 2*, Annals of Mathematics Studies, no. 28, pages 245–266. Princeton University Press, Princeton, N.J., 1953.
- [4] Leo A. Harrington and Alexander S. Kechris. On the determinacy of games on ordinals. *Ann. Math. Logic*, 20(2):109–154, 1981.
- [5] Steve Jackson. Suslin cardinals, partition properties, homogeneity. Introduction to Part II. In *Games, scales, and Suslin cardinals. The Cabal Seminar. Vol. I*, volume 31 of *Lect. Notes Log.*, pages 273–313. Assoc. Symbol. Logic, Chicago, IL, 2008.
- [6] Steve Jackson. Structural consequences of  $\text{AD}$ . In *Handbook of set theory. Vols. 1, 2, 3*, pages 1753–1876. Springer, Dordrecht, 2010.
- [7] Steve Jackson and Donald A. Martin. Pointclasses and well-ordered unions. In *Cabal seminar 79–81*, volume 1019 of *Lecture Notes in Math.*, pages 56–66. Springer, Berlin, 1983.
- [8] Thomas Jech. *Set theory*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2003. The third millennium edition, revised and expanded.
- [9] R. B. Jensen and R. M. Solovay. Some applications of almost disjoint sets. In *Mathematical Logic and Foundations of Set Theory (Proc. Internat. Colloq., Jerusalem, 1968)*, pages 84–104. North-Holland, Amsterdam, 1970.
- [10] Akihiro Kanamori. *The higher infinite*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, second edition, 2003. Large cardinals in set theory from their beginnings.

- [11] Alexander S. Kechris. Souslin cardinals,  $\kappa$ -Souslin sets and the scale property in the hyperprojective hierarchy. In *Cabal Seminar 77–79 (Proc. Caltech-UCLA Logic Sem., 1977–79)*, volume 839 of *Lecture Notes in Math.*, pages 127–146. Springer, Berlin, 1981.
- [12] Alexander S. Kechris. The axiom of determinacy implies dependent choices in  $L(\mathbf{R})$ . *J. Symbolic Logic*, 49(1):161–173, 1984.
- [13] Alexander S. Kechris. “AD + UNIFORMIZATION” is equivalent to “half  $\text{AD}_{\mathbf{R}}$ ”. In *Cabal Seminar 81–85*, volume 1333 of *Lecture Notes in Math.*, pages 98–102. Springer, Berlin, 1988.
- [14] Alexander S. Kechris, Eugene M. Kleinberg, Yiannis N. Moschovakis, and W. Hugh Woodin. The axiom of determinacy, strong partition properties and nonsingular measures. In *Cabal Seminar 77–79 (Proc. Caltech-UCLA Logic Sem., 1977–79)*, volume 839 of *Lecture Notes in Math.*, pages 75–99. Springer, Berlin, 1981.
- [15] Alexander S. Kechris, Benedikt Löwe, and John R. Steel, editors. *Games, scales, and Suslin cardinals. The Cabal Seminar. Vol. I*, volume 31 of *Lecture Notes in Logic*. Association for Symbolic Logic, Chicago, IL; Cambridge University Press, Cambridge, 2008.
- [16] Alexander S. Kechris, Benedikt Löwe, and John R. Steel, editors. *Wadge degrees and projective ordinals. The Cabal Seminar. Volume II*, volume 37 of *Lecture Notes in Logic*. Association for Symbolic Logic, La Jolla, CA; Cambridge University Press, Cambridge, 2012.
- [17] Alexander S. Kechris, Benedikt Löwe, and John R. Steel, editors. *Ordinal definability and recursion theory. The Cabal Seminar. Vol. III*, volume 43 of *Lecture Notes in Logic*. Association for Symbolic Logic, Ithaca, NY; Cambridge University Press, Cambridge, 2016. Including reprinted papers from the Caltech-UCLA Cabal seminars held in Los Angeles, CA.
- [18] Alexander S. Kechris, Robert M. Solovay, and John R. Steel. The axiom of determinacy and the prewellordering property. In *Cabal Seminar 77–79 (Proc. Caltech-UCLA Logic Sem., 1977–79)*, volume 839 of *Lecture Notes in Math.*, pages 101–125. Springer, Berlin-New York, 1981.
- [19] A.S. Kechris. *Classical descriptive set theory*, volume 156 of *Graduate texts in mathematics*. Springer, 1995.
- [20] A.S. Kechris. *Classical descriptive set theory*, volume 156 of *Graduate texts in mathematics*. Springer, 1995.
- [21] Richard Ketchersid. More structural consequences of AD. In *Set theory and its applications*, volume 533 of *Contemp. Math.*, pages 71–105. Amer. Math. Soc., Providence, RI, 2011.

- [22] Peter Koellner and W. Hugh Woodin. Large cardinals from determinacy. In *Handbook of set theory. Vols. 1, 2, 3*, pages 1951–2119. Springer, Dordrecht, 2010.
- [23] K. Kunen. *An Introduction to Independence Proofs*. North-Holland, 1980.
- [24] Kenneth Kunen. *Set theory*, volume 34 of *Studies in Logic (London)*. College Publications, London, 2011.
- [25] Paul B. Larson. Wadge determinacy implies countable choice for reals.
- [26] Paul B. Larson. A brief history of determinacy. In *Sets and extensions in the twentieth century*, volume 6 of *Handb. Hist. Log.*, pages 457–507. Elsevier/North-Holland, Amsterdam, 2012.
- [27] P.B. Larson. *The stationary tower*, volume 32 of *University Lecture Series*. American Mathematical Society, Providence, RI, 2004. Notes on a course by W. Hugh Woodin.
- [28] D. A. Martin and R. M. Solovay. A basis theorem for  $\sum_3^1$  sets of reals. *Ann. of Math. (2)*, 89:138–159, 1969.
- [29] D.A. Martin and J. Steel. A proof of projective determinacy. *Journal of the American Mathematical Society*, 2:71–125, 1989.
- [30] Donald A. Martin and John R. Steel. The extent of scales in  $L(\mathbf{R})$ . In *Cabal seminar 79–81*, volume 1019 of *Lecture Notes in Math.*, pages 86–96. Springer, Berlin, 1983.
- [31] Donald A. Martin and John R. Steel. The tree of a Moschovakis scale is homogeneous. In *Games, scales, and Suslin cardinals. The Cabal Seminar. Vol. I*, volume 31 of *Lect. Notes Log.*, pages 404–420. Assoc. Symbol. Logic, Chicago, IL, 2008.
- [32] Yiannis N. Moschovakis. *Descriptive set theory*, volume 155 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, second edition, 2009.
- [33] Yinhe Peng and Liang Yu. TD implies  $\text{CC}_{\mathbb{R}}$ . *Adv. Math.*, 384:Paper No. 107755, 7, 2021.
- [34] Gerald E. Sacks. Forcing with perfect closed sets. In *Axiomatic Set Theory (Proc. Sympos. Pure Math., Vol. XIII, Part I, Univ. California, Los Angeles, Calif., 1967)*, pages 331–355. Amer. Math. Soc., Providence, R.I., 1971.
- [35] Ralf Schindler. *Set theory*. Universitext. Springer, Cham, 2014. Exploring independence and truth.

- [36] Robert M. Solovay. The independence of DC from AD. In *Cabal Seminar 76–77 (Proc. Caltech-UCLA Logic Sem., 1976–77)*, volume 689 of *Lecture Notes in Math.*, pages 171–183. Springer, Berlin, 1978.
- [37] J. R. Steel. The derived model theorem. In *Logic Colloquium 2006*, volume 32 of *Lect. Notes Log.*, pages 280–327. Assoc. Symbol. Logic, Chicago, IL, 2009.
- [38] John R. Steel. Determinateness and the separation property. *J. Symbolic Logic*, 46(1):41–44, 1981.
- [39] John R. Steel. A classification of jump operators. *J. Symbolic Logic*, 47(2):347–358, 1982.
- [40] Robert Van Wesep. Wadge degrees and descriptive set theory. In *Cabal Seminar 76–77 (Proc. Caltech-UCLA Logic Sem., 1976–77)*, volume 689 of *Lecture Notes in Math.*, pages 151–170. Springer, Berlin, 1978.



# Index

- $(\mu, \kappa)$ -club directedness, 66
- 2-perfect tree, 150
- $<_{\text{LS}}$ , 83
- $<_{\mathcal{D}}$ , 107
- $A \oplus B$ , 35
- $A^{(i)}$ , 24
- $A_{(i)}$ , 24
- $A_\varphi$ , 89
- $A_x$ , 31
- $B$ -generic filter, 9
- $E$ -invariant function, 101
- $H(\kappa)$ , 9
- $S$ -cone, 103
- $S$ -pointed tree, 112
- $S$ -positive set, 112
- $T_s$ , 69
- $U_n(A_1, \dots, A_m)$ , 36
- $[T]$ , 69
- $[X]^\alpha$ , 8, 61
- $[f]_{\mu_E}$ , 101
- AD, 15
- $\text{AD}_X$ , 15
- Axiom of Determinacy, 15
- Baire( $\Gamma$ ), 17
- CC, 10
- $\text{CC}_X$ , 10
- Countable Choice, 10
- DC, 10
- $\text{DC}_X$ , 10
- $\Delta_\omega(A)$ , 35
- Dependent Choice, 10
- $\Gamma$ -complete set, 29
- $\Gamma$ -norm, 50
- $\Gamma$ -separability, 45
- $\Gamma$ -universal set, 31
- HC, 9
- HC-coding, 9
- HF, 9
- $\text{HOD}_X$ , 119
- $\text{LR}(A)$ , 24
- Lipschitz Determinacy, 22
- $\text{OD}_X$ , 76, 119
- Ord, 8
- $\text{Red}(\Gamma)$ , 47
- $\text{Sep}(\Gamma)$ , 45
- $\text{TC}(X)$ , 9
- TD, 19
- $\Theta$ , 36
- $\text{WF}_E$ , 101
- $\text{WR}(A)$ , 24
- $\aleph_1 \not\preceq 2^{\aleph_0}$ , 17
- $\mathbb{E}_0$ , 90
- $\mathbb{P}_S$ , 114
- $\bigcup_\delta \Gamma$ , 30
- $\mathcal{D}$ , 107
- $\mathcal{D}_E(x)$ , 18
- $\mathcal{F}^{rmc}$ , 36
- $\mathcal{L}_{\gamma, \delta}^0$ , 89
- $\mathcal{L}_\gamma^0$ , 89
- $\mathcal{P}_A$ , 96
- $\mathcal{S}_\gamma$ , 70
- $\mathcal{S}_{<\gamma}$ , 70
- $\mathcal{U}_E(x)$ , 18
- $\check{A}$ , 29
- $\check{\Gamma}$ , 29
- $\delta(\Gamma)$ , 50
- $\delta_S^\infty$ , 112, 139
- $\delta_A$ , 96
- $\delta_\omega(A)$ , 102
- $\equiv_{\mathcal{D}}$ , 107
- $\equiv_{\text{Ma}}$ , 18
- $\equiv_{\text{Tu}}$ , 9
- $\eta$ -full tree, 72

- $\exists^{\omega^\omega}$ -closed pointclass, 30
- $\exists^{\omega^\omega} A$ , 30
- $\forall^{\omega^\omega}$ -closed pointclass, 30
- $\forall^{\omega^\omega} A$ , 30
- $\gamma$ -Borel set, 90
- $\gamma$ -Borel\* code, 94
- $\infty$ -Borel code, 89
- $\infty$ -Borel set, 90
- $\infty$ -Borel subset of  $\omega^\delta$ , 93
- $\infty$ -Borel\* code, 94
- $\lambda$ -Determinacy, 87
- $\lambda$ -reasonable ordinal, 63
- $\text{lb}_\leq(T)$ , 69
- $\leq_L$ , 21
- $\leq_{\text{Wa}}$ , 21
- $\leq_{\text{Gö}}$ , 9
- $\leq_{\text{Tu}}$ , 9
- $<\gamma$ -Determinacy, 87
- $<\gamma$ -Borel set, 90
- $\mathcal{X}$ , 28
- $\prec\alpha, \beta\succ$ , 9
- $\text{meas}(X)$ , 166
- $\text{pos-}\widetilde{\Sigma}_1^1(A)$ , 35
- $\prod V/\mu S$ , 103
- $\prod \text{Ord}/\mu S$ , 103
- $\prod f(x)/\mu S$ , 103
- $\text{rank}_T$ , 78
- $\text{rank}_\leq$ , 8
- $\text{Ma}$ , 18
- $\text{p}[T]$ , 69
- $\sigma * x$ , 16
- $\sigma \circ x$ , 39
- $\Delta_n^1(A)$ , 35
- $\Delta_1^2(A)$ , 52
- $\Pi_n^1(A)$ , 35
- $\Pi_1^2(A)$ , 52
- $\Sigma_n^1(A)$ , 35
- $\Sigma_1^2(A)$ , 52
- $\delta_1^2(A)$ , 52
- $f_x^c$ , 36
- $o(\Gamma)$ , 57
- $o(\varphi)$ , 50
- $s^\frown x$ , 24
- $x * \sigma$ , 16
- $x \circ \sigma$ , 39
- almost-disjoint coding, 10
- base of a cone, 18
- basic open interval of a space in  $\mathcal{X}$ , 29
- boldface pointclass, 28
- canonical rank function for a wellfounded tree, 78
- Coding Lemma, 39
- cofinality of a pointclass, 28
- computing a set of reals on a cone, 115
- constructibility degree, 19
- determined set, 15
- diffuse function, 109
- downward cone, 18
- fine measure, 167
- generic  $\alpha$ -code, 136
- generic  $\infty$ -code, 136
- generic code, 136
- hereditarily countable set, 9
- hereditarily finite set, 9
- homogeneous tree, 165
- homogeneously Suslin set, 166
- illfounded tree, 69
- induced map, 171
- join of a sequence of generic codes, 138
- leftmost branch, 69
- length of a (strict) prewellordering, 50
- length of a norm, 50
- lightface pointclass, 29
- Lipschitz class, 21
- Lipschitz function, 21
- Lipschitz game, 22
- Lipschitz rank, 24
- Lipschitz reducibility, 21
- locally countable relation, 102
- Lusin-Sierpiński order, 83
- measure, 165

- minimal tree, 71
- minimization of a tree, 71
- nice name, 8
- nonselfdual class, 21
- nonselfdual pointclass, 29
- norm, 50
- normal measure, 171
- ordered equivalence relation, 18
- payoff set, 15
- pointclass, 28
- pointset, 28
- preorder, 8
- prewellordering, 50
- prewellordering property, 51
- projection of a tree, 69
- projection of an ultrafilter, 165
- projective algebra, 57
- projective in  $A$ , 35
- projectively closed pointclass, 35
- quasi- $\text{AD}_X$ , 76
- quasi-determined set, 76
- quasi-strategy, 76
- rank function, 8
- rank of a norm, 50
- real, 8
- recursion property, 32
- Recursion Theorem, 34
- recursive function, 29
- recursive subset of HF, 9
- reduction property, 47
- regular norm, 50
- s-m-n property, 32
- selfdual class, 21
- selfdual pointclass, 29
- semi-recursive subset of HF, 9
- separation property, 45
- strategy, 16
- strict prewellordering, 50
- strict supremum, 8
- strong partition cardinal, 61
- strongly Lipschitz function, 45
- strongly maximal set of ordinals, 117
- Suslin cardinal, 70
- Suslin set, 69
- tower of ultrafilters, 165
- Turing cone, 18
- Turing Determinacy, 19
- Turing equivalence, 9
- Turing measure, 18
- Turing reducibility, 9
- Uniform Coding Lemma, 41
- uniformizing, 75
- upward cone, 18
- Vopěnka algebra, 120
- Wadge class, 21
- Wadge rank, 24
- Wadge reducibility, 21
- Wadge's Theorem, 22
- weakly  $\infty$ -Borel set, 74
- weakly  $\kappa$ -Borel set, 74
- weakly  $<\kappa$ -Borel set, 74
- weakly homogeneous tree, 166
- weakly homogeneously Suslin set, 166
- wellfounded preorder, 8
- wellfounded tower of ultrafilters, 165
- wellfounded tree, 69
- winning quasi strategy, 76
- winning strategy, 16
- witness for a  $\Sigma_1^2$  statement, 52