



HENDRIX

COLLEGE

Homework 7: Section 15

Algebra

Author

Paul Beggs

BeggsPA@Hendrix.edu

Instructor

Dr. Christopher Camfield, Ph.D.

Due

NOVEMBER 25, 2025



Section 15

In Exercises 1, 3, 4, and 7, classify the given group according to the fundamental theorem of finitely generated abelian groups.

1. $(\mathbb{Z}_2 \times \mathbb{Z}_4)/\langle(0, 1)\rangle$

Solution. $\langle(0, 1)\rangle$ has order 4, so $(\mathbb{Z}_2 \times \mathbb{Z}_4)/\langle(0, 1)\rangle$ has order 2. Therefore, this leaves only one choice: $(\mathbb{Z}_2 \times \mathbb{Z}_4)/\langle(0, 1)\rangle \simeq \mathbb{Z}_2$.

3. $(\mathbb{Z}_2 \times \mathbb{Z}_4)/\langle(1, 2)\rangle$

Solution. $\langle(1, 2)\rangle$ has order 2, so $(\mathbb{Z}_2 \times \mathbb{Z}_4)/\langle(1, 2)\rangle$ has order 4. This leaves us with two choices: \mathbb{Z}_4 or $\mathbb{Z}_2 \times \mathbb{Z}_2$. Since $(1, 1) + \langle(1, 2)\rangle$ has order 4 in the factor group, we must have $(\mathbb{Z}_2 \times \mathbb{Z}_4)/\langle(1, 2)\rangle \simeq \mathbb{Z}_4$.

4. $(\mathbb{Z}_4 \times \mathbb{Z}_8)/\langle(1, 2)\rangle$

Solution. $|\langle(1, 2)\rangle| = 4 \implies |(\mathbb{Z}_4 \times \mathbb{Z}_8)/\langle(1, 2)\rangle| = 8$. Therefore, either \mathbb{Z}_8 or $\mathbb{Z}_4 \times \mathbb{Z}_2$. Since $|(0, 1) + \langle(1, 2)\rangle| = 8$, $(\mathbb{Z}_4 \times \mathbb{Z}_8)/\langle(1, 2)\rangle \simeq \mathbb{Z}_8$.

7. $(\mathbb{Z} \times \mathbb{Z})/\langle(1, 2)\rangle$

Solution. In the factor group, everything in the subgroup becomes the identity, so $(1, 2) = (0, 0)$. This implies $x + 2y = 0$, or $x = -2y$. Therefore, every element in the factor group can be written as $(-2y, y) = y(-2, 1)$. Since x depends on y , the factor group is isomorphic to \mathbb{Z} .

In Exercises 20 and 21, let F be the additive group of all functions mapping \mathbb{R} into \mathbb{R} , and let F^* be the multiplicative group of all elements of F that do not assume the value 0 at any point of \mathbb{R} .

20. Let K be the subgroup of F consisting of the constant functions. Find a subgroup of F to which F/K is isomorphic.

Solution. Define a homomorphism $\varphi : F \rightarrow F$ by $\varphi(f) = f(x) - f(0)$. The kernel of this map consists of all functions where $f(x) - f(0) = 0$, which implies $f(x)$ is constant, so $\ker(\varphi) = K$. The image of the map, $\varphi[F]$, is the set of all functions that evaluate to 0 at $x = 0$. By the fundamental theorem of homomorphisms, F/K is isomorphic to the subgroup $\{f \in F \mid f(0) = 0\}$.



21. Let K^* be the subgroup of F^* consisting of the nonzero constant functions. Find a subgroup of F^* to which F^*/K^* is isomorphic.

Solution. This problem is almost exactly like the last one: Define a homomorphism $\varphi : F^* \rightarrow F^*$ by $\varphi(f) = f(x)/f(0)$. The kernel of this map consists of all functions where $f(x)/f(0) = 1$, which implies $f(x)$ is constant, so $\ker(\varphi) = K^*$. The image of the map, $\varphi[F^*]$, is the set of all functions that evaluate to 1 at $x = 0$. By the fundamental theorem of homomorphisms, F^*/K^* is isomorphic to the subgroup $\{f \in F^* \mid f(0) = 1\}$.

28. Give an example of a group G having no elements of finite order > 1 but having a factor group G/H , all of whose elements are of finite order.

Proof. The group $G = \mathbb{Z}$ contains only the identity element, 0, with finite order. Every other integer has infinite order. Thus, G has no elements of finite order > 1 . Now, consider G/H where $H = 2\mathbb{Z}$. Since this factor group has order 2, it is isomorphic to \mathbb{Z}_2 . Then, since \mathbb{Z}_2 is itself finite, all of its elements must also have finite order. \square

29. Let H and K be normal subgroups of a group G . Give an example showing that we may have $H \simeq K$ while G/H is not isomorphic to G/K .

Proof. Let $G = \mathbb{Z}_4 \times \mathbb{Z}_2$. Consider the subgroups $H = \langle (2, 0) \rangle$ and $K = \langle (0, 1) \rangle$. Both H and K are subgroups of order 2. Since there is only one group of order 2 up to isomorphism, $H \simeq K$. Notice, however, that $G/H \simeq \mathbb{Z}_2 \times \mathbb{Z}_2$, while $G/K \simeq \mathbb{Z}_4$. Therefore, $G/H \not\simeq G/K$. \square