

Paul Gustafson
 Texas A&M University - Math 607
 Instructor: Thomas Schlumprecht

HW 7

- 1 Assume that $(f_n) \subset L_1(\mu)$ and $f_n \rightarrow f$ uniformly.
 a) If $\mu(X) < \infty$, then $f \in L_1$ and $\lim_{n \rightarrow \infty} \int f_n d\mu = \int f d\mu$.
 b) If $\mu(X) = \infty$, then the conclusion of (a) may fail.

Proof. For (a), we have

$$\begin{aligned} \left| \int f d\mu - \int f_n d\mu \right| &\leq \int |f - f_n| d\mu \\ &\leq \mu(X) \sup_{x \in X} |f(x) - f_n(x)| \\ &\rightarrow 0. \end{aligned}$$

For (b), let $(f_n) = 1/n\chi_{[0,n]}$. Then $f_n \rightarrow 0$ uniformly on \mathbb{R} , but $\int f_n dx = 1$ for all n . \square

- 2 Let $f_n, g_n, g \in L_1$, $n \in \mathbb{N}$, and assume that $f_n \rightarrow f$, f measurable, and $g_n \rightarrow g$ μ -a.e., and that $|f_n| \leq g_n$ and $\int g_n d\mu \rightarrow \int g d\mu$.

Then $\int f_n d\mu \rightarrow \int f d\mu$.

Proof. Following the proof of the DCT, since $f_n \leq g_n$, we have $f \leq g$, so $f \in L_1$. We also have $g_n + f_n \geq 0$ a.e. and $g_n - f_n \geq 0$ a.e. Hence by Fatou's lemma and linearity of the integral on L_1 ,

$$\int g + \int f \leq \liminf \int (g_n + f_n) = \liminf \int g_n + \int f_n = \int g + \liminf \int f_n$$

The last inequality follows from the fact that if $(a_n) \rightarrow a$ and $(b_n) \subset \mathbb{R}$, then $\liminf a_n + b_n = a + \liminf b_n$. To see this, pick $\epsilon > 0$ and N such that $|a - a_n| < \epsilon$ for all $n \geq N$. Hence $\liminf a_n + b_n = \liminf (a_n - a) + a + b_n \leq \liminf \epsilon + a + b_n = \epsilon + a + \liminf b_n$, and similarly $\liminf a_n + b_n \geq -\epsilon + a + \liminf b_n$. Hence $\liminf a_n + b_n = a + \liminf b_n$.

Similarly,

$$\int g - \int f \leq \liminf \int (g_n - f_n) = \liminf \int g_n - \int f_n = \int g - \limsup \int f_n$$

Hence $\limsup \int f_n \leq \int f \leq \liminf \int f_n$, so $\int f = \lim \int f_n$. \square

- 3 Suppose that for $n \in \mathbb{N}$, $f_n = \chi_{E_n}$ for some $E_n \subset \mathbb{R}$, and assume that $f(x) = \lim_{n \rightarrow \infty} f_n(x)$ exists a.e.

a) Show that $f = \chi_E$ a.e. for some measurable set $E \subset \mathbb{R}$.

b) Show that for any $g \in L_1$:

$$\int_E g \, dx = \lim_{n \rightarrow \infty} \int_{E_n} g \, dx.$$

c) Establish a necessary and sufficient condition for $f_n \rightarrow f$ in L_1 .

Proof. For (a), we have $\chi_{E_n} \rightarrow f$ on N^c for some null set N . Let $x \in N^c$. Since $(\chi_{E_n}(x))_n$ is a convergent discrete-valued sequence, it must be eventually constant. Thus, $f(x) \in \{0, 1\}$. Let $E = f^{-1}(1) \cap N^c$. Hence $f = \chi_E$ on N^c , so $f = \chi_E$ a.e. on \mathbb{R} . By a previous homework problem, f is measurable since it is the limit of measurable functions. Hence E is measurable.

For (b), we have $\chi_{E_n} g \rightarrow \chi_E g$ pointwise a.e. by part (a). Moreover, $\chi_{E_n} g \leq |g| \in L_1$. Hence, by the DCT, we have the desired conclusion.

For (c), one such condition is that $m(E_n) \rightarrow m(E)$ with $m(E_n), m(E) < \infty$. Clearly, the latter condition is necessary for f_n, f to be in L_1 . For the necessity of the former condition, suppose $f_n \rightarrow f$ in L_1 . Then $|m(E_n) - m(E)| = |\int f_n - \int f| \leq \int |f_n - f| \rightarrow 0$.

For sufficiency, suppose $m(E_n) \rightarrow m(E)$ with $m(E_n), m(E) < \infty$. Then $|f_n - f| \leq |f_n| + |f| = \chi_{E_n} + \chi_E$ a.e. Moreover, $\chi_{E_n} + \chi_E \rightarrow 2\chi_E$ and $\int(\chi_{E_n} + \chi_E) = m(E_n) + m(E) \rightarrow 2m(E) = \int(2\chi_E)$. Hence, by the Generalized DCT (Exercise 2), we have $\int |f_n - f| \rightarrow \int \lim_n |f_n - f| = 0$. \square

4 Let $L_0([0, 1])$ be the space of all measurable functions $f : [0, 1] \rightarrow \mathbb{R}$.

a) for $f, g \in L_0([0, 1])$ put

$$d(f, g) = \int_0^1 \min\{1, |f - g|\} \, dx.$$

Show that $(L_0([0, 1]), d)$ is a metric space and that for $f, f_n \in L_0([0, 1])$:

$$f_n \rightarrow f \text{ in } (L_0([0, 1]), d) \iff f_n \rightarrow f \text{ in measure.}$$

b) Is there a metric d' on $L_0([0, 1])$ for which

$$f_n \rightarrow f \text{ in } (L_0([0, 1]), d') \iff f_n \rightarrow f \text{ a.e.}$$

Proof. For (a), to see that d is a metric, we need to show that d is positive definite, symmetric, and satisfies the triangle inequality. The function d is clearly nonnegative and $0 = d(f, g) = \int_0^1 \min\{1, |f - g|\} \, dx$ implies that $f = g$ a.e. The function d is obviously symmetric. For the triangle inequality, I first claim that for $x, y, z \in \mathbb{R}$ we have $\min\{1, |x - y|\} \leq \min\{1, |x - z|\} + \min\{1, |y - z|\}$.

We have four cases from the RHS of the inequality.

Case $|x - z| \leq 1$ and $|y - z| \leq 1$. We have $\min\{1, |x - y|\} \leq |x - y| \leq |x - z| + |y - z| = \min\{1, |x - z|\} + \min\{1, |y - z|\}$.

Case $|x - z| \leq 1$ and $|y - z| > 1$. We have $\min\{1, |x - y|\} \leq \min\{1, |x - z| + |y - z|\} \leq \min\{1, 1 + |y - z|\} = 1 + |y - z| = \min\{1, |x - z|\} + \min\{1, |y - z|\}$.

Case $|x - z| > 1$ and $|y - z| \leq 1$. Analogous to previous case.

Case $|x - z| > 1$ and $|y - z| > 1$. We have $\min\{1, |x - y|\} \leq \min\{1, |x - z| + |z - y|\} = 1 \leq \min\{1, |x - z|\} + \min\{1, |z - y|\}$.

Hence $\min\{1, |x - y|\} \leq \min\{1, |x - z|\} + \min\{1, |y - z|\}$ for all $x, y, z \in \mathbb{R}$. Thus, if $f, g, h \in L_0([0, 1])$ then $d(f, g) = \int_0^1 \min\{1, |f - g|\} dx \leq \int_0^1 \min\{1, |f - h| + \min\{1, |h - g|\}\} dx \leq \int_0^1 \min\{1, |f - h|\} dx + \int_0^1 \min\{1, |h - g|\} dx = d(f, h) + d(g, h)$. Thus, d is a metric.

Suppose $f_n \rightarrow f$ in $(L_0([0, 1]), d)$. Let $0 < \epsilon < 1$. Pick N such that $d(f, f_n) < \epsilon^2$ for all $n \geq N$. Then for all $n \geq N$, we have

$$\begin{aligned} m(\{|f - f_n| \geq \epsilon\}) &= \int_{\{|f - f_n| \geq \epsilon\}} dx \\ &= \int_{\{\epsilon \leq |f - f_n| < 1\}} dx + \int_{\{|f - f_n| \geq 1\}} dx \\ &\leq \int_{\{\epsilon \leq |f - f_n| < 1\}} \frac{|f - f_n|}{\epsilon} dx + \int_{\{|f - f_n| \geq 1\}} \min\{1, |f - f_n|\} dx \\ &\leq \int_{\{\epsilon \leq |f - f_n| < 1\}} \epsilon^{-1} \min\{1, |f - f_n|\} dx + \int_{\{|f - f_n| \geq 1\}} \min\{1, |f - f_n|\} dx \\ &\leq \epsilon^{-1} \int \min\{1, |f - f_n|\} dx \\ &< \epsilon \end{aligned}$$

Conversely, suppose $f_n \rightarrow f$ in measure. Let $0 < \epsilon < 1$. Pick N such that $m(\{|f - f_n| \geq \epsilon\}) \leq \epsilon$ for all $n \geq N$. Then for all $n \geq N$ we have

$$\begin{aligned} \int \min\{1, |f_n - f|\} dx &= \int_{\{0 \leq |f_n - f| < \epsilon\}} |f_n - f| dx + \int_{\{\epsilon \leq |f_n - f| < 1\}} |f_n - f| dx + \int_{\{|f_n - f| \geq 1\}} dx \\ &\leq \int_{\{0 \leq |f_n - f| < \epsilon\}} \epsilon dx + \int_{\{\epsilon \leq |f_n - f| < 1\}} dx + \int_{\{|f_n - f| \geq 1\}} dx \\ &\leq \epsilon + m(\{|f - f_n| \geq \epsilon\}) \\ &< 2\epsilon. \end{aligned}$$

For (b), I use a fact about convergence in metric spaces. Let (M, d) be a metric space, $(x_n)_{n \in \mathbb{N}} \subset M$, and $x \in M$. I claim that if every subsequence of (x_n) has a further subsequence converging to x , then $x_n \rightarrow x$. Suppose $x_n \not\rightarrow x$. Then there exists $\epsilon > 0$ and a subsequence $(x_n)_{n \in N_1}$ such that $d(x, x_n) \geq \epsilon$ for all $n \in N_1$. This subsequence cannot have a further subsequence converging to x , contradicting the hypothesis.

Thus it suffices to find a sequence that does not converge pointwise a.e., but each subsequence has a subsequence that converges to the 0 function. For $n \in \mathbb{N}$, write n as $n = 2^j + k$ for $j \geq 0$ and $0 \leq k < 2^j$. Let $E_n = [k2^{-j}, (k+1)2^{-j}]$ and $f_n = \chi_{E_n}$. Every element of $[0, 1]$ is contained in infinitely many E_n and infinitely many E_n^c , so f_n do not converge pointwise a.e.

On the other hand, suppose $(f_n)_{n \in N_1}$ is a subsequence of $(f_n)_{n \in \mathbb{N}}$. For each n , pick $x_n \in E_n$. Then by the sequential compactness of $[0, 1]$, there exists an infinite set $N_2 \subset N_1$ and $x_0 \in [0, 1]$ such that $x_n \rightarrow x_0$ as $n \rightarrow \infty, n \in N_2$. Since $\text{diam}(E_n) \rightarrow 0$, we have $f_n(x) \rightarrow 0$ as $n \rightarrow \infty, n \in N_2$ for $x \neq x_0$. Hence, $f_n \rightarrow 0$ as $n \rightarrow \infty, n \in N_2$ pointwise a.e. \square

5

(a)

$$\lim_{n \rightarrow \infty} \int_0^\infty \left(1 + \frac{x}{n}\right)^{-n} \sin\left(\frac{x}{n}\right) dx = 0$$

(b)

$$\lim_{n \rightarrow \infty} \int_0^1 \frac{1 + nx^2}{(1 + x^2)^2} = 0$$

(c)

$$\lim_{n \rightarrow \infty} \int_0^\infty n \sin\left(\frac{x}{n}\right) [x(1 + x^2)]^{-1} dx = \frac{\pi}{2}$$

(d)

$$\lim_{n \rightarrow \infty} \int_a^\infty \frac{n}{1 + n^2 x^2} dx = \begin{cases} 0 & \text{if } a > 0 \\ \pi/2 & \text{if } a = 0 \\ \pi & \text{if } a < 0 \end{cases}$$

Proof. For (a), we have $(1 + \frac{x}{n})^n = \sum_{j=0}^n \binom{n}{j} \left(\frac{x}{n}\right)^j$

\square