Name:	
Student ID:	
Section:	
Instructor: Paul Gustafson	

## Math 131 (Principles of Calculus) Exam 3A

RED

## **Instructions:**

- For questions which require a written answer, show all your work. Full credit will be given only if the necessary work is shown justifying your answer.
- Simplify your answers.
- Calculators are allowed.
- Should you have need for more space than is allocated to answer a question, use the back of the exam.
- Please do not talk about the test with other students until exams are handed back.
- Honor Code:

An Aggie does not lie, cheat, or steal or tolerate those who do.			
Signature			

## For Instructor use only.

#	Possible	Earned
MC	40	
9	10	
10	12	
Sub	62	

#	Possible	Earned
11	20	
12	10	
Sub	30	
Total	92	

Part I: Multiple Choice (5 points each) Mark the correct answer on the bubble sheet.

- 1. Find the absolute maximum and minimum values for the function  $f(x) = 3x^2 6x + 3$  on the interval [-1,3]
  - a) maximum value = 0, minimum value = -2
  - b) maximum value = 9, minimum value = -2
  - c) maximum value = 12, minimum value = 0
  - d) maximum value = 9, minimum value = 0
  - e) maximum value = 12, minimum value = -2
- 2. If  $f'(x) = \frac{1}{\sqrt{x}} + 3x^2$  and f(4) = 38

a) 
$$f(x) = \sqrt{x} + 3x^3 - 30$$

b) 
$$f(x) = 2\sqrt{x} + x^3 - 30$$

c) 
$$f(x) = 2\sqrt{x} + x^3 + 30$$

d) 
$$f(x) = \frac{2}{3}x^{3/2} + x^3 + 38$$

e) 
$$f(x) = \frac{2}{3}x^{3/2} + x^3 - 30$$

3. A particle moves along a wire with velocity  $v(t) = \sin(t) + 3$ . Find the net change in position between times t = 0 and  $t = \pi$ 

a) 
$$1 + 3\pi$$

b) 
$$-2 + 3\pi$$

c) 
$$2 + 3\pi$$

e) 
$$3\pi$$

4. Calculate the indefinite integral  $\int \frac{4}{x} + \sec^2(3x) dx$ 

a) 
$$\frac{2}{x^2} + \frac{1}{3}\tan(3x) + C$$

$$b) \quad 4 + 3\tan(3x) + C$$

c) 
$$4 \ln |x| + \tan(3x) + C$$

$$d) \quad 4 + \frac{1}{3}\tan(3x) + C$$

e) 
$$4 \ln |x| + \frac{1}{3} \tan(3x) + C$$

- 5. Use the fundamental theorem of calculus to find the derivative of  $f(x) = \int_1^x \frac{t^3 e^t}{\cos^2(t)} dt$ 
  - a)  $\frac{2x^2 e^x}{2\cos(x)\sin(x)}$

b)  $\frac{(2x^2 - e^x)\cos^2(x) - 2\cos(x)\sin(x)(x^3 - e^x)}{\cos^4(x)}$ 

c)  $\frac{t^4 - e^t}{\cos^2(t)}$ 

 $d) \quad \frac{x^3 - e^x}{\cos^2(x)}$ 

- $e) \quad \frac{3t^2 e^t}{\cos^4(t)}$
- 6. Use the geometric shape of the graph to find the integral  $\int_{-3}^{3} f(x)$  where

$$f(x) = \begin{cases} 3 - x, & x \le 0\\ \sqrt{9 - x^2}, & x > 0 \end{cases}$$

a)  $\frac{9}{2} + \frac{9}{4}\pi$ 

b)  $\frac{27}{2} + \frac{3}{4}\pi$ 

c)  $\frac{9}{2} + 3\pi$ 

d)  $\frac{27}{2} + \frac{9}{4}\pi$ 

- e)  $\frac{27}{2} + 9\pi$
- 7. The acceleration of a particle is given by a(t) = 6t 2. The position of the particle at times t = 0 and t = 1 are s(0) = 2 and s(1) = 5, respectively. The position function for the particle is
  - a)  $s(t) = 3t^2 2t + 2$

b)  $s(t) = 3t^2 - 2t + 4$ 

c)  $s(t) = t^3 - t^2 + 5t + 2$ 

d)  $s(t) = t^3 - t^2 + 3t + 2$ 

- e)  $s(t) = t^3 2t + 4$
- 8. Calculate  $\int_{1}^{e^2} \frac{\ln(x)}{x} dx$ .
  - a)  $e^{-4} 1$

b)  $2e^{-4} - 1$ 

c)  $2e^{-4}$ 

d)  $e^{-2}$ 

e) 2

## Part II: Free Response Show all work

9. (10 points) Use four approximating rectangles with left endpoints to estimate the definite integral

$$\int_2^{10} \frac{1}{\sqrt{x}+1} \, dx$$

10. (12 points) A glassblower wants to make a cylindrical vase with one end covered and one end open. He has enough molten glass to cover a surface area of 30 square centimeters. Determine the dimensions of the vase that will maximize its volume.

- 11. (20 points) Let  $f(x) = \frac{1}{3}x^3 \frac{1}{2}x^2 6x + 1$ .
  - a.) (5 points) Find the intervals on which f(x) is **increasing** and the intervals where it is **decreasing**.

b.) (5 points) Find the x-coordinates where f(x) has a **local max or min**. Make sure to specify which are maxes and which are mins.

c.) (5 points) Find the x-coordinates of the **inflection points** of f(x).

d.) (5 points) Find the intervals where f(x) is **concave up** and where it is **concave down**.

12. (10 points) Find the exact value of the definite integral. Show all your work.

$$\int_0^1 3x \sin(x^2 - 1) \, dx$$