

FINITENESS FOR MAPPING CLASS GROUP REPRESENTATIONS FROM TWISTED DIJKGRAAF-WITTEN THEORY

PAUL GUSTAFSON

ABSTRACT. Any twisted Dijkgraaf-Witten representation of a mapping class group of a closed surface has finite image.

1. INTRODUCTION

Given a spherical category \mathcal{A} over a field k and an oriented compact surface M , possibly with boundary, the Turaev-Viro-Barrett-Westbury (TVBW) construction gives a projective representation of the mapping class group $\mathrm{MCG}(M)$ [1, 2]. A natural question is to determine the image of these representations. In particular, when does such a representation have finite image?

It is conjectured that these representations have finite image if and only if \mathcal{A} is weakly integral. This conjecture is a modification of the Property F conjecture [3, 4], which states that braid group representations coming from a braided monoidal category \mathcal{C} should have finite image if and only if \mathcal{C} is weakly integral. Instead of only considering braid group representations, one can consider mapping class groups of arbitrary orientable surfaces. In this case, the input categories to construct the representations must be more specialized than just braided monoidal. One can either apply the Reshetikhin-Turaev construction to a modular tensor category, or apply the TVBW construction to a spherical category. The former is more general than the latter since the Reshetikhin-Turaev construction for the Drinfeld center $Z(\mathcal{A})$ of a spherical category \mathcal{A} yields the same representation as the TVBW construction for \mathcal{A} . However, for the case considered in this paper, the simpler TVBW construction suffices.

In this paper, our input category is $\mathcal{A} = \mathrm{Vec}_G^\omega$, the spherical category of G -graded vector spaces with associativity modified by a cocycle $\omega \in Z^3(G, k^\times)$. In this case, the TVBW construction corresponds to the twisted Dijkgraaf-Witten theory of [5]. The category Vec_G^ω is integral, so one expects that its associated mapping class group representations have finite image. Our main contribution is to verify this in the case when the surface M is closed and (G, ω) are arbitrary.

Acknowledgments. This paper would not have been written without the guidance of my advisor, Eric Rowell. I am also grateful to Zhengang Wang and my father Robert Gustafson for their advice.

2. RELATED WORK

The closest related work is a result of Fjelstad and Fuchs [6] showing that, given a surface with at most one boundary component, the mapping class group representations corresponding to the untwisted (i.e. $\omega = 1$) Dijkgraaf-Witten theory have finite image. Their paper uses an algebraic method of Lyubashenko [7] that gives a projective mapping class group representation to any factorizable ribbon Hopf algebra, in their case, the double $D(G)$. In our case, we instead consider the mapping class group action on a vector space of Vec_G^ω -colored embedded graphs defined by Kirillov [8], yielding a simpler, more geometric proof.

In [9], Bantay defined representations of mapping class groups on the Hilbert space of an orbifold model associated to $D^\omega(G)$. These representations appear to coincide with the twisted Dijkgraaf-Witten representations. However, the precise details of the connection are not clear to me.

More is known when we fix a particular surface M . In the case where M is a torus, Ng and Schauenburg showed that any Reshetikhin-Turaev representation of the mapping class group of the torus is always finite [10]. In the case where M is an n -punctured disk, the mapping class group of M relative to the boundary of the disk is the braid group B_n . In this case, Etingof, Rowell, and Witherspoon proved that the representations associated to $\mathrm{Mod}(D^\omega(G))$ are finite [4].

3. DEFINITIONS

Let G be a finite group, and let Vec_G^ω denote the skeletal spherical category of G -graded vector spaces with associativity defined by the 3-cocycle $\omega \in Z^3(G, k^\times)$. We will follow [11] in the choice of structural morphisms. The associator $\alpha_{g,h,k} : (g \otimes h) \otimes k \rightarrow g \otimes (h \otimes k)$ is defined to be

$$\alpha_{g,h,k} = \omega(g, h, k) \text{id}_{ghk}.$$

The evaluator $\text{ev}_g : g^* \otimes g \rightarrow 1$ is

$$\text{ev}_g = \omega(g^{-1}, g, g^{-1}) \text{id}_1.$$

The coevaluator $\text{coev}_g : 1 \rightarrow g \otimes g^*$ is

$$\text{coev}_g = \text{id}_1.$$

The pivotal structure $j_g : g^{**} \rightarrow g$ is

$$j_g = \omega(g^{-1}, g, g^{-1}) \text{id}_g.$$

The following definitions and theorem are from Kirillov's paper [8]. For any spherical category \mathcal{A} and surface M , he gives the following presentation of the Levin-Wen model as a vector space of colored graphs modulo local relations. He also proves that this space is canonically isomorphic to the TVBW vector space associated to M . It is straightforward to check that this isomorphism, which amounts to replacing a triangulation with its dual graph, commutes with the mapping class group action.

We will consider finite graphs embedded in the surface M ; for such a graph Γ , let $E(\Gamma)$ be the set of edges. Note that edges are not oriented. Let E^{or} be the set of oriented edges, i.e. pairs $\mathbf{e} = (e, \text{orientation of } e)$; for such an oriented edge \mathbf{e} , we denote by $\bar{\mathbf{e}}$ the edge with opposite orientation.

Definition 3.1. Let $\Gamma \subset M$ be an embedded graph as defined above. A coloring of Γ is the following data:

- Choice of an object $V(\mathbf{e}) \in \text{Obj } \mathcal{A}$ for every oriented edge $\mathbf{e} \in E^{or}(\Gamma)$ so that $V(\bar{\mathbf{e}}) = V(\mathbf{e})^*$.
- Choice of a vector $\varphi(v) \in \text{Hom}(1, V(\mathbf{e}_1) \otimes \cdots \otimes V(\mathbf{e}_n))$ for every vertex v , where $\mathbf{e}_1, \dots, \mathbf{e}_n$ are edges incident to v , taken in counterclockwise order and with outward orientation.

The following theorem is a variation of result of Reshetikhin and Turaev.

Theorem 3.2. There is a unique way to assign to every colored planar graph Γ in a disk $D \subset \mathbb{R}^2$ a vector

$$(1) \quad \langle \Gamma \rangle_D \in \text{Hom}(1, V(\mathbf{e}_1) \otimes \cdots \otimes V(\mathbf{e}_n))$$

where $\mathbf{e}_1, \dots, \mathbf{e}_n$ are the edges of Γ meeting the boundary of D (legs), taken in counterclockwise order and with outgoing orientation, so that that following conditions are satisfied:

- (1) $\langle \Gamma \rangle$ only depends on the isotopy class of Γ .
- (2) If Γ is a single vertex colored by $\varphi \in \text{Hom}(1, V(\mathbf{e}_1) \otimes \cdots \otimes V(\mathbf{e}_n))$, then $\langle \Gamma \rangle = \varphi$.
- (3) Local relations shown in Figure 1 hold.




FIGURE 1. Local relations for colored graphs. Here $\varphi \circ \psi = (\varphi \otimes \psi) \circ \text{ev}_X$.

Local relations should be understood as follows: for any pair Γ, Γ' of colored graphs which are identical outside a subdisk $D' \subset D$, and in this disk are homeomorphic to the graphs shown in Figure 1, we must have $\langle \Gamma \rangle = \langle \Gamma' \rangle$.

Moreover, so defined $\langle \Gamma \rangle$ satisfies the following properties:

- (1) $\langle \Gamma \rangle$ is linear in color of each vertex v (for fixed colors of edges and other vertices).
- (2) $\langle \Gamma \rangle$ is additive in colors of edges as shown in Figure 2.




FIGURE 2. Linearity of $\langle \Gamma \rangle$. Here φ_1, φ_2 are compositions of φ with projector $X_1 \oplus X_2 \rightarrow X_1$ (respectively, $X_1 \oplus X_2 \rightarrow X_2$), and similarly for ψ_1, ψ_2 .

- (3) *Composition property:* if $D' \subset D$ is a subdisk such that $\partial D'$ does not contain vertices of Γ and meets edges of Γ transversally, then $\langle \Gamma \rangle_D$ will not change if we replace subgraph $\Gamma \cap D'$ by a single vertex colored by $\langle \Gamma \cap D' \rangle_{D'}$.

The vector $\langle \Gamma \rangle$ is called the evaluation of Γ .

To define local relations between graphs, Kirillov defines the space of null graphs as follows. Let $D \subset M$ be an embedded disk, and let $\Gamma = c_1 \Gamma_1 + \dots + c_n \Gamma_n$ be a formal linear combination of colored graphs in M such that

- (1) Γ is transversal to ∂D (i.e., no vertices of Γ_i are on the boundary of D and edges of each Γ_i meet ∂D transversally).
- (2) All Γ_i coincide outside of D .
- (3) $\langle \Gamma \rangle_D = \sum c_i \langle \Gamma_i \cap D \rangle_D = 0$.

In this case Γ is called a null graph.

Definition 3.3. The representation space associated to a closed, oriented surface M by a spherical category \mathcal{A} is the vector space of formal linear combinations of \mathcal{A} -colored graph embeddings modulo the subspace spanned by the null graphs.

4. RESULT

Theorem 4.1. The image of the twisted Dijkgraaf-Witten representation of a mapping class group of any closed surface M is finite.

Proof. Let Γ be a $\text{Vec}_G^{\mathcal{O}}$ -colored graph embedding, and let g be the genus of M . Thinking of M as a quotient of its fundamental $4g$ -gon, by isotopy we may assume vertices of Γ lie in the interior of the polygon and that all the edges of Γ do not intersect corners and meet the sides transversally. Evaluating on the interior of the polygon shows that Γ is equivalent to a graph with a single vertex whose edges are simple closed curves, each of which intersect the boundary of the polygon precisely once. By using the local relations, we can replace all the edges intersecting a side with a single edge labeled by the tensor product of their labels. If there are no edges intersecting a side, we can insert a single edge labeled by the group identity into Γ that intersects only that side. Thus, Γ is equivalent to a colored graph with one vertex v , outgoing edges e_1, \dots, e_{2g} , and $2g$ incoming edges with the same labels as in Figure 3.

By the definition of the evaluation of a string-net and the definition of the quotient map identifying the sides of the fundamental polygon, the vertex v is colored by an element $\phi(v) \in \text{Hom}(1, \bigotimes_{i=1}^n V(e_{2i-1}) \otimes V(e_{2i}) \otimes V(e_{2i-1}^*) \otimes V(e_{2i}^*))$, where $V(e_i) \in \text{Obj}(\text{Vec}_G^{\mathcal{O}})$ is the coloring of the edge e_i . Since string-net evaluation is additive in the direct sum and linear in the vertex color, it follows that the representation space H is spanned by the set of colored graphs

$$S := \{\Gamma \in H : V(e_i) \in \text{Irr}(\text{Vec}_G^{\mathcal{O}}), \phi(v) = 1\},$$

where $\text{Irr}(\text{Vec}_G^{\mathcal{O}})$ is the set of simple objects of $\text{Vec}_G^{\mathcal{O}}$, which correspond to the elements of G .

The mapping class group of M is generated by the Lickorish generating set consisting of Dehn twists around $3g - 1$ simple closed curves. These can be divided into two types of twists: the ones around a single hole (the blue and green curves in Figure 4), and the ones connecting two holes (the red curves).



FIGURE 3. Element of the spanning set S for a genus 2 surface




FIGURE 4. Lickorish generating set (Image source: https://en.wikipedia.org/wiki/Dehn_twist)

Using the local moves as in Tables 1 and 2, one sees that the result of each of these Dehn twists lies in $\text{Im}(\omega)S$ since the structural morphisms of Vec_G^ω have coefficients in $\text{Im}(\omega)$. It is a basic result in group cohomology that, by replacing ω with a cohomologous cocycle if necessary, the image of ω lies in $\mu_{|G|}$, the set of $|G|$ -th roots of unity. Since cohomologous cocycles give rise to equivalent spherical categories Vec_G^ω , this replacement does not incur any loss in generality.

Thus, the image of any such mapping class group representation is a quotient of the group of permutations of the finite set $\mu_{|G|}S$, hence finite. \square

5. BOUNDARY CASE

6. EXAMPLE CALCULATION

This section contains a calculation of the coefficient for the first Dehn twist, shown in Table 1. WLOG we may assume that ω is normalized, i.e. that $\omega(1, x, y) = \omega(x, 1, y) = \omega(x, y, 1) = 1$ for all x, y . We also assume that the main vertex is initially labeled with an element of $\text{Hom}(1, h \otimes g \otimes h^{-1} \otimes hg^{-1}h^{-1})$. In the following, we will abbreviate this by saying that the vertex is in state $h \otimes g \otimes h^{-1} \otimes hg^{-1}h^{-1}$.

We add the upper left vertex, which is labeled by coev_g . We then connect the vertices with an unlabelled edge, which is shorthand for labelling by the object 1 . At this point, the vertices are in states $h \otimes g \otimes 1 \otimes h^{-1} \otimes hg^{-1}h^{-1}$ and $g \otimes g^{-1} \otimes 1$. To compose the two vertices, we use the spherical structure on the former vertex and reassociate until it is in the state $h^{-1} \otimes hg^{-1}h^{-1} \otimes h \otimes g \otimes 1$. In doing so, we pick up a factor of

$$\omega(h, g, h^{-1})\omega(h, gh^{-1}, hg^{-1}h^{-1})\omega(g, h^{-1}, hg^{-1}h^{-1})\omega(g, g^{-1}h^{-1}, h)$$

After performing the composition, we are in state $h^{-1} \otimes hg^{-1}h^{-1} \otimes h \otimes g \otimes (g \otimes g^{-1})$. To get rid of the last pair of parentheses, we get a factor of $\omega^{-1}(h^{-1}hg^{-1}h^{-1}hg, g, g^{-1}) = \omega^{-1}(1, g, g^{-1}) = 1$.

To tensor the parallel g and h edges together, we add coev_g and coev_h vertices in the middle of those edges and connect them with a 1 . Composing along the 1 , we get a vertex in state $g \otimes g^{-1} \otimes h^{-1} \otimes h$. To put this vertex in state $g^{-1}h^{-1} \otimes hg$ we pick up a factor of $\omega(g, g^{-1}, h^{-1})\omega(g, g^{-1}h^{-1}, h)\omega(g^{-1}h^{-1}, h, g)$. We also put the original vertex in the state $g^{-1}h^{-1} \otimes hg^{-1}h^{-1} \otimes hg \otimes g$ with a factor of

$$\omega^{-1}(g^{-1}, g^{-1}, g)\omega^{-1}(g^{-1}, g^{-1}h^{-1}, h)\omega^{-1}(g^{-1}, h^{-1}, hg^{-1}h^{-1})\omega(g^{-2}h^{-1}, h, g).$$




TABLE 1. First type of Dehn twist. The figures are ordered from left to right, then top to bottom.

To compose the two vertices, we rotate the original vertex to the state $g \otimes g^{-1}h^{-1} \otimes hg^{-1}h^{-1} \otimes hg$ which yields a factor of

$$\omega^{-1}(g, h^{-1}, hg)\omega^{-1}(g, g^{-1}h^{-1}, hg^{-1}h^{-1})\omega^{-1}(g, g^{-1}, h^{-1}).$$

We are then in a position to compose the two vertices, giving a factor $\omega(g^{-1}h^{-1}, hg, g^{-1}h^{-1}) \text{ev}_{g^{-1}h^{-1}} = 1$ and a vertex in state $g \otimes g^{-1}h^{-1} \otimes hg^{-1}h^{-1} \otimes hg$. Rotating the vertex into its initial configuration $hg \otimes g \otimes g^{-1}h^{-1} \otimes hg^{-1}h^{-1}$ gives a factor of

$$\omega^{-1}(hg, h^{-1}, hg^{-1}h^{-1})\omega^{-1}(hg, g, g^{-1}h^{-1}).$$

Thus, we have an overall factor of

$$\begin{aligned} & \frac{\omega(h, g, h^{-1})\omega(h, gh^{-1}, hg^{-1}h^{-1})\omega(g, h^{-1}, hg^{-1}h^{-1})\omega(g, g^{-1}h^{-1}, h)}{\omega(g^{-1}, g^{-1}, g)\omega(g^{-1}, g^{-1}h^{-1}, h)\omega(g^{-1}, h^{-1}, hg^{-1}h^{-1})\omega(g, g^{-2}h^{-1}, hg)} \\ & \frac{\omega(g, g^{-1}, h^{-1})\omega(g, g^{-1}h^{-1}, h)\omega(g^{-1}h^{-1}, h, g)\omega(g^{-2}h^{-1}, h, g)}{\omega(g, g^{-1}h^{-1}, hg^{-1}h^{-1})\omega(hg, h^{-1}, hg^{-1}h^{-1})\omega(hg, g, g^{-1}h^{-1})} \\ & = \frac{\omega(h, g, g^{-1}h^{-1})\omega(g, g^{-1}h^{-1}, h)}{\omega(g^{-1}, g^{-1}, g)\omega(g^{-1}, g^{-1}h^{-1}, h)\omega(g^{-1}, h^{-1}, hg^{-1}h^{-1})\omega(g, g^{-2}h^{-1}, hg)} \\ & \frac{\omega(g, g^{-1}, h^{-1})\omega(g, g^{-1}h^{-1}, h)\omega(g^{-1}h^{-1}, h, g)\omega(g^{-2}h^{-1}, h, g)}{\omega(g, g^{-1}h^{-1}, hg^{-1}h^{-1})\omega(hg, g, g^{-1}h^{-1})} \\ & = \frac{\omega(h, g, g^{-1}h^{-1})\omega(g, g^{-1}h^{-1}, h)}{\omega(g^{-1}, g^{-1}, g)\omega(g^{-1}, g^{-1}h^{-1}, h)\omega(g, g^{-2}h^{-1}, hg)} \\ & \frac{\omega^2(g, g^{-1}, h^{-1})\omega(g, g^{-1}h^{-1}, h)\omega(g^{-1}h^{-1}, h, g)\omega(g^{-2}h^{-1}, h, g)}{\omega(g, g^{-1}, g^{-1}h^{-1})\omega(hg, g, g^{-1}h^{-1})} \\ & = \frac{\omega(h, g, g^{-1}h^{-1})\omega^2(g, g^{-1}h^{-1}, h)\omega^2(g, g^{-1}, h^{-1})\omega^2(g^{-1}h^{-1}, h, g)}{\omega(g, g^{-2}h^{-1}, hg)\omega(g, g^{-1}, g^{-1}h^{-1})\omega(hg, g, g^{-1}h^{-1})\omega(g^{-1}, g^{-1}h^{-1}, hg)} \\ & = \frac{\omega(h, g, g^{-1}h^{-1})\omega^2(g, g^{-1}h^{-1}, h)\omega^2(g, g^{-1}, h^{-1})\omega^2(g^{-1}h^{-1}, h, g)}{\omega(hg, g, g^{-1}h^{-1})} \end{aligned}$$




TABLE 2. Second type of Dehn twist. The figures are ordered from left to right, then top to bottom.

$$= \frac{\omega(h, g, g^{-1}h^{-1})\omega^2(g^{-1}h^{-1}, h, g)}{\omega^2(g^{-1}, h^{-1}, h)\omega(hg, g, g^{-1}h^{-1})}$$

7. FURTHER DIRECTIONS

Given the fact that the braid group representations associated to $D^\omega(G)$ are also finite [4], it should be straightforward to prove that mapping class group representations associated to Vec_G^ω for any orientable surface, including boundary, are finite. The basic idea is that one labels the boundary components with elements of the double category, in this case $\text{Mod}(D^\omega(G))$, and the Vec_G^ω -colored embedded graphs must be compatible with the boundary labels, as detailed in [8].

One could also consider more complicated spherical categories than Vec_ω^G , such as the Tambara-Yamagami categories. The main additional complication here is the appearance of multifusion channels, i.e. the tensor product of two simple objects can be a direct sum of multiple simple objects.

REFERENCES

- [1] V.G. Turaev and O.Y. Viro. State sum invariants of 3-manifolds and quantum 6j-symbols. *Topology*, 31(4):865 – 902, 1992.
- [2] John W. Barrett and Bruce W. Westbury. Invariants of piecewise-linear 3-manifolds. 1993.
- [3] Deepak Naidu and Eric C. Rowell. A finiteness property for braided fusion categories. 2009.

- [4] Pavel Etingof, Eric C. Rowell, and Sarah Witherspoon. Braid group representations from twisted quantum doubles of finite groups. *Pacific J. Math.* 234 no. 1 (2008) 33-42, 2007.
- [5] Robbert Dijkgraaf and Edward Witten. Topological gauge theories and group cohomology. *Comm. Math. Phys.*, 129(2):393–429, 1990.
- [6] Jens Fjelstad and Jürgen Fuchs. Mapping class group representations from Drinfeld doubles of finite groups, 2015.
- [7] V. Lyubashenko. Ribbon abelian categories as modular categories. *J. Knot Theory Ramifications*, 05(03):311–403, Jun 1996.
- [8] Alexander Kirillov. String-net model of Turaev-Viro invariants. 2011.
- [9] Peter Bantay. Algebraic aspects of orbifold models. 1993.
- [10] Siu-Hung Ng and Peter Schauenburg. Congruence subgroups and generalized Frobenius-Schur indicators. 2008.
- [11] Siu-Hung Ng and Peter Schauenburg. Frobenius-Schur indicators and exponents of spherical categories. *Adv. Math.* 211 (2007) no. 1, 34–71, 2005.

E-mail address: pgustafs@math.tamu.edu

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TX U.S.A.