

Paul Gustafson  
Texas A&M University - Math 467  
Instructor: Stephen Fulling

## 1 $\mathbb{F}_2$ and the Fano plane

### 1.1 Introduction

The purpose of this paper is to answer Exercise 2.5 (p. 96) of Greenberg [1]:

Let  $\mathbb{F}_2$  be the field of two elements  $\{0, 1\}$ , whose multiplication and addition have the usual tables except that  $1 + 1 = 0$ . Show that  $\mathbb{F}_2^2$  is isomorphic to the smallest affine plane. Show that  $P^2(\mathbb{F}_2)$  is isomorphic to the Fano plane.

We will need a few preliminary definitions from Greenberg.

**Definition 1.** An *incidence geometry*  $(\mathcal{P}, \mathcal{L}, \mathcal{I})$  consists of a set of points  $\mathcal{P}$ , a set of lines  $\mathcal{L}$ , and an incidence relation  $\mathcal{I} \subset \mathcal{P} \times \mathcal{L}$  such that:

1. Every pair of distinct points is incident to a unique line.
2. Every line is incident to at least two distinct points.
3. There exist three distinct noncollinear points.

**Definition 2.** Two lines are *parallel* if there is no point incident to both lines.

**Definition 3.** A *projective plane* is an incidence geometry in which:

1. No two lines are parallel.
2. Every line is incident to at least three distinct points.

**Definition 4.** An *affine plane* is an incidence geometry in which, for every line  $l$  and point  $P$  not incident to  $l$ , there exists a unique line  $m$  incident to  $P$  and parallel to  $l$ .

### 1.2 The affine plane $\mathbb{F}_2^2$

As in  $\mathbb{R}^2$ , the points in  $\mathbb{F}_2^2$  are simply the elements of the vector space  $\mathbb{F}_2^2$ , i.e. ordered pairs of elements of  $\mathbb{F}_2$ .

Also analogous to  $\mathbb{R}^2$ , the lines in  $\mathbb{F}_2^2$  are cosets of 1-dimensional subspaces of  $\mathbb{F}_2^2$ . That is, every line in  $\mathbb{F}_2^2$  can be written as  $V + h$  for some 1-dimensional subspace  $V \subset \mathbb{F}_2^2$  and  $h \in \mathbb{F}_2^2$ .

Incidence in  $\mathbb{F}_2^2$  corresponds to inclusion. For example, the point  $(1, 1) \in \mathbb{F}_2^2$  is incident to the line  $\{(1, 0)t + (0, 1) : t \in \mathbb{F}_2\}$ , since  $(1, 1) = (1, 0)(1) + (0, 1)$ .

As Greenberg notes, the smallest affine plane, call it  $\mathcal{A}$ , consists of a set of four points  $\{A, B, C, D\}$  and a set of four lines  $\{\{A, B\}, \{B, C\}, \{C, D\}, \{D, A\}\}$ ,

where incidence corresponds to inclusion. For example, the point  $B$  is incident to the line  $\{A, B\}$ .

To see that  $\mathcal{A}$  and  $\mathbb{F}_2^2$  are isomorphic, first note that each 1-dimensional subspace over  $\mathbb{F}_2$  has exactly 2 elements, so each line in  $\mathbb{F}_2^2$  has 2 elements. Conversely, given two elements  $a, b \in \mathbb{F}_2^2$ , the line  $L((b-a)t, a)$  passes through  $a$  and  $b$ . Thus, the lines in  $\mathbb{F}_2^2$  are precisely the two-element subsets of  $\mathbb{F}_2^2$ .

Therefore, an arbitrary bijection  $f$  from the points of  $\mathbb{F}_2^2$  to the points of  $\mathcal{A}$  induces a bijection of lines (two-element subsets), and since inclusion is preserved under  $f$ , incidence is also preserved.

### 1.3 $P^2(\mathbb{F}_2)$ as the Fano plane

For an arbitrary field  $K$ , the points of the projective space  $P^2(K)$  are the 1-dimensional subspaces of  $K^3$ . The lines are the 2-dimensional subspaces of  $K^3$ . Incidence corresponds to containment.

Projective points in  $P^2(K)$  are usually denoted  $(a:b:c)$  for some generator  $(a, b, c) \in K^3 \setminus \{0\}$ . Then  $(a:b:c) = (d:e:f)$  iff  $(a, b, c)$  is a nonzero multiple of  $(d, e, f)$ . For example, the projective line  $\{x + y + z = 0 : (x:y:z) \in P^2(\mathbb{R})\}$  is incident to the point  $(1:0:-1) \in P^2(\mathbb{R})$  since  $t + 0 - t = 0$  for all  $t \in \mathbb{R}$ .

Note that a strange thing occurs in  $P^2(\mathbb{F}_2)$ : there is a correspondence between each point in  $P^2(\mathbb{F}_2)$  as a 1-dimensional subspace of  $\mathbb{F}_2^3$  and its unique nonzero element in  $\mathbb{F}_2^3$ . Since, on the other hand, each non-zero element in  $\mathbb{F}_2^3$  generates a 1-dimensional subspace of  $\mathbb{F}_2^3$ , this correspondence defines a bijection from  $P^2(\mathbb{F}_2)$  to  $\mathbb{F}_2^3 \setminus \{0\}$ . Hence, there are  $2^3 - 1 = 7$  points in  $P^2(\mathbb{F}_2)$ .

Since every 2-dimensional subspace of  $\mathbb{F}_2^3$  contains 0, a 1-dimensional subspace  $V \subset \mathbb{F}_2^3$  lies within a 2-dimensional subspace  $W \subset \mathbb{F}_2^3$  iff the unique nonzero element in  $V$  lies within  $W$ . Hence, since each 2-dimensional subspace of  $\mathbb{F}_2^3$  contains exactly  $2^2 - 1 = 3$  nonzero  $\mathbb{F}_2^3$ -elements, each line in  $P^2(\mathbb{F}_2)$  is incident to precisely 3 projective points.

Since we have seen that no two nonzero elements of an  $\mathbb{F}_2$ -vector space are linearly dependent, each pair of distinct nonzero elements in  $\mathbb{F}_2^3$  determines a 2-dimensional subspace. Hence, since each 2-dimensional subspace of  $\mathbb{F}_2^3$  contains exactly  $\binom{3}{2} = 3$  pairs of distinct points, there are  $(1/3)\binom{7}{2} = 7$  2-dimensional subspaces of  $\mathbb{F}_2^3$ .

## References

- [1] Marvin J Greenberg. *Euclidean and non-Euclidean geometries: Development and history*. WH Freeman, 2007.




Figure 1: An isomorphism between  $P^2(\mathbb{F}_2)$  and the Fano plane