

# Computing Quantum Mapping Class Group Representations with Haskell

Paul Gustafson

Why should we care about quantum mapping class group representations?

- Topological quantum computation

Why should we care about quantum mapping class group representations?

- Topological quantum computation
- Understand mapping class groups

Why should we care about quantum mapping class group representations?

- Topological quantum computation
- Understand mapping class groups
- Intrinsic beauty

# Mapping class group definition

- Let  $\Sigma = \Sigma_{g,b}^m$  be the oriented compact surface of genus  $g$  with  $b$  boundary components and  $m$  marked points in its interior.

# Mapping class group definition

- Let  $\Sigma = \Sigma_{g,b}^m$  be the oriented compact surface of genus  $g$  with  $b$  boundary components and  $m$  marked points in its interior.
- The mapping class group of  $\Sigma$ ,  $MCG(\Sigma)$ , is the group of isotopy classes of orientation-preserving homeomorphisms of  $\Sigma$  that preserve the boundary *pointwise* and preserve the marked points *setwise*.

# Mapping class group definition

- Let  $\Sigma = \Sigma_{g,b}^m$  be the oriented compact surface of genus  $g$  with  $b$  boundary components and  $m$  marked points in its interior.
- The mapping class group of  $\Sigma$ ,  $MCG(\Sigma)$ , is the group of isotopy classes of orientation-preserving homeomorphisms of  $\Sigma$  that preserve the boundary *pointwise* and preserve the marked points *setwise*.
- Examples
  - $MCG(\Sigma_{0,1}^m) = B_m$
  - $MCG(\Sigma_{1,0}^0) = SL(2, \mathbb{Z})$

# Mapping class group definition

- Let  $\Sigma = \Sigma_{g,b}^m$  be the oriented compact surface of genus  $g$  with  $b$  boundary components and  $m$  marked points in its interior.
- The mapping class group of  $\Sigma$ ,  $MCG(\Sigma)$ , is the group of isotopy classes of orientation-preserving homeomorphisms of  $\Sigma$  that preserve the boundary *pointwise* and preserve the marked points *setwise*.
- Examples
  - $MCG(\Sigma_{0,1}^m) = B_m$
  - $MCG(\Sigma_{1,0}^0) = SL(2, \mathbb{Z})$
- Birman (1969) found “nice” finite generating set for the mapping class group of any compact surface.



# $(2+1)$ -Topological Quantum Field Theories

- Quantum mapping class group representations come from  $(2+1)$ -TQFTs

# $(2+1)$ -Topological Quantum Field Theories

- Quantum mapping class group representations come from  $(2+1)$ -TQFTs
- A  $(2+1)$ -TQFT assigns a vector space to oriented 2-manifolds
- It also assigns a vector to oriented 3-manifolds. This vector inhabits the vector space corresponding to the 3-manifold's boundary.
- To get a mapping class group representation, apply the TQFT to mapping cylinders.

# $(2+1)$ -Topological Quantum Field Theories

- Quantum mapping class group representations come from  $(2+1)$ -TQFTs
- A  $(2+1)$ -TQFT assigns a vector space to oriented 2-manifolds
- It also assigns a vector to oriented 3-manifolds. This vector inhabits the vector space corresponding to the 3-manifold's boundary.
- To get a mapping class group representation, apply the TQFT to mapping cylinders.
- History: Witten and Atiyah (1980s)

# $(2+1)$ -Topological Quantum Field Theories

- Quantum mapping class group representations come from  $(2+1)$ -TQFTs
- A  $(2+1)$ -TQFT assigns a vector space to oriented 2-manifolds
- It also assigns a vector to oriented 3-manifolds. This vector inhabits the vector space corresponding to the 3-manifold's boundary.
- To get a mapping class group representation, apply the TQFT to mapping cylinders.
- History: Witten and Atiyah (1980s)
- Examples of mathematical  $(2+1)$ -TQFTs:
  - Reshetikhin-Turaev TQFT (input: modular category)
  - Turaev-Viro-Barret-Westbury TQFT (input: spherical fusion category)

# Monoidal categories

A **monoidal category** is a category  $\mathcal{C}$  equipped with

- a tensor product – a bifunctor  $\otimes : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$
- an associativity isomorphism – a natural isomorphism  $\alpha : (\cdot \otimes \cdot) \otimes \cdot \rightarrow \cdot \otimes (\cdot \otimes \cdot)$
- a unit object  $1 \in \mathcal{C}$
- a left unitor – a natural isomorphism  $\lambda_X : 1 \otimes X \rightarrow X$
- a right unitor – a natural isomorphism  $\rho_X : X \otimes 1 \rightarrow X$ ,

satisfying certain coherence conditions (the triangle and pentagon axioms).

# Rigid monoidal categories

Let  $X$  be an object of a monoidal category  $\mathcal{C}$ . A **left dual** to  $X$  is an object  $X^*$  equipped with

- an evaluation morphism,  $\text{ev}_X : X^* \otimes X \rightarrow 1$
- a coevaluation morphism,  $\text{coev}_X : 1 \rightarrow X \otimes X^*$ ,

satisfying certain coherence conditions (the snake equations).

# Rigid monoidal categories

Let  $X$  be an object of a monoidal category  $\mathcal{C}$ . A **left dual** to  $X$  is an object  $X^*$  equipped with

- an evaluation morphism,  $\text{ev}_X : X^* \otimes X \rightarrow 1$
- a coevaluation morphism,  $\text{coev}_X : 1 \rightarrow X \otimes X^*$ ,

satisfying certain coherence conditions (the snake equations).

Right duals are defined similarly. An object  $X$  is **rigid** if it has both a left and right dual. A monoidal category is rigid if all of its objects are rigid.

A **fusion category** is a rigid semisimple linear monoidal category (tensor category), with only finitely many isomorphism classes of simple objects, such that the endomorphisms of the unit object form just the ground field  $k$ .



- A **pivotal category** is a rigid monoidal category equipped with a pivotal structure, i.e. a tensor natural automorphism  $j_X : X \rightarrow (X^*)^*$ .

# Spherical categories

- A **pivotal category** is a rigid monoidal category equipped with a pivotal structure, i.e. a tensor natural automorphism  $j_X : X \rightarrow (X^*)^*$ .
- A pivotal structure defines categorical left and right traces  $\text{End}(X) \rightarrow \text{End}(1)$  for every object  $X$ . A **spherical category** is a monoidal category such that all left and right traces coincide.

## Example: $\text{Vect}_G^\omega$

- Let  $G$  be a finite group, and  $\omega : G \times G \times G \rightarrow \mathbb{C}$  be a 3-cocycle. The spherical fusion category  $\text{Vect}_G^\omega$  is the skeletal category of  $G$ -graded finite-dimensional vector spaces with the following modified structural morphisms, where  $V_g$  is the simple object:

- The associator  $a_{g,h,k} : (V_g \otimes V_h) \otimes V_k \rightarrow V_g \otimes (V_h \otimes V_k)$

$$a_{g,h,k} = \omega(g, h, k)$$

- The evaluator  $ev_g : V_g^* \otimes V_g \rightarrow 1$

$$ev_g = \omega(g^{-1}, g, g^{-1})$$

- The pivotal structure  $j_g : V_g^{**} \rightarrow V_g$

$$j_g = \omega(g^{-1}, g, g^{-1})$$

# The TVBW space associated to a 2-manifold

- Let  $\mathcal{A}$  be a spherical fusion category, and  $\Sigma$  an oriented compact surface with boundary.
- Using Kirillov's definitions, the representation space we consider is

$$H := \frac{\mathcal{A}\text{-colored graphs in } \Sigma}{\text{local relations}}$$

# The TVBW space associated to a 2-manifold

- Let  $\mathcal{A}$  be a spherical fusion category, and  $\Sigma$  an oriented compact surface with boundary.
- Using Kirillov's definitions, the representation space we consider is

$$H := \frac{\mathcal{A}\text{-colored graphs in } \Sigma}{\text{local relations}}$$

- The vector space  $H$  is canonically isomorphic to the Turaev-Viro state sum vector space associated to  $\Sigma$ . This isomorphism commutes with the mapping class group action.

# Colored graphs in $\Sigma$

- Let  $\Gamma \subset \Sigma$  be an undirected finite graph embedded in  $\Sigma$ .

# Colored graphs in $\Sigma$

- Let  $\Gamma \subset \Sigma$  be an undirected finite graph embedded in  $\Sigma$ .
- Define  $E^{or}$  to be the set of orientation edges of  $\Gamma$ , i.e. pairs  $\mathbf{e} = (e, \text{orientation of } e)$ ; for such an oriented edge  $\mathbf{e}$ , we denote by  $\bar{\mathbf{e}}$  the edge with opposite orientation.

# Colored graphs in $\Sigma$

- Let  $\Gamma \subset \Sigma$  be an undirected finite graph embedded in  $\Sigma$ .
- Define  $E^{or}$  to be the set of orientation edges of  $\Gamma$ , i.e. pairs  $\mathbf{e} = (e, \text{orientation of } e)$ ; for such an oriented edge  $\mathbf{e}$ , we denote by  $\bar{\mathbf{e}}$  the edge with opposite orientation.
- A *coloring* of  $\Gamma$  is the following data:
  - Choice of an object  $V(\mathbf{e}) \in \text{Obj } \mathcal{A}$  for every oriented edge  $\mathbf{e} \in E^{or}$  so that  $V(\bar{\mathbf{e}}) = V(\mathbf{e})^*$ .



# Colored graphs in $\Sigma$

- Let  $\Gamma \subset \Sigma$  be an undirected finite graph embedded in  $\Sigma$ .
- Define  $E^{or}$  to be the set of orientation edges of  $\Gamma$ , i.e. pairs  $\mathbf{e} = (e, \text{orientation of } e)$ ; for such an oriented edge  $\mathbf{e}$ , we denote by  $\bar{\mathbf{e}}$  the edge with opposite orientation.
- A *coloring* of  $\Gamma$  is the following data:
  - Choice of an object  $V(\mathbf{e}) \in \text{Obj } \mathcal{A}$  for every oriented edge  $\mathbf{e} \in E^{or}$  so that  $V(\bar{\mathbf{e}}) = V(\mathbf{e})^*$ .
  - Choice of a vector  $\varphi(v) \in \text{Hom}_{\mathcal{A}}(1, V_1 \otimes \cdots \otimes V_n)$  for every interior vertex  $v$ , where  $\mathbf{e}_1, \dots, \mathbf{e}_n$  are edges incident to  $v$ , taken in counterclockwise order and with outward orientation.

# Local relations

- Isotopy of the graph embedding

# Local relations

- Isotopy of the graph embedding
- Linearity in the vertex colorings

# Local relations

- Isotopy of the graph embedding
- Linearity in the vertex colorings





Figure : The remaining local relations.


# Consequences of the local relations



**Figure :** Additivity in edge colorings. Here  $\varphi_1, \varphi_2$  are compositions of  $\varphi$  with projector  $X_1 \oplus X_2 \rightarrow X_1$  (respectively,  $X_1 \oplus X_2 \rightarrow X_2$ ), and similarly for  $\psi_1, \psi_2$ .

- Additivity in edge colorings

# Consequences of the local relations



**Figure :** Additivity in edge colorings. Here  $\varphi_1, \varphi_2$  are compositions of  $\varphi$  with projector  $X_1 \oplus X_2 \rightarrow X_1$  (respectively,  $X_1 \oplus X_2 \rightarrow X_2$ ), and similarly for  $\psi_1, \psi_2$ .

- Additivity in edge colorings
- A colored graph may be evaluated on any disk  $D \subset S$ , giving an equivalent colored graph  $\Gamma'$  such that  $\Gamma'$  is identical to  $\Gamma$  outside of  $D$ , has the same colored edges crossing  $\partial D$ , and contains at most one colored vertex within  $D$ .

# Overall Strategy

- Find a basis of colored graphs for the representation space for a surface
- “Calculate” the representation of each mapping class group generator with respect to this basis
- Analyze the image of the representation (Is it finite? Can we do universal quantum computation with it (possibly adding extra measurements)?)

# Modified Property F conjecture

## Conjecture (Rowell)

*A TVBW mapping class group representation associated to a spherical fusion category  $\mathcal{A}$  has finite image iff  $\mathcal{A}$  is weakly integral, i.e. the squared dimension of every simple object is an integer.*



## Theorem (Ng–Schauenberg)

*Every modular representation associated to a modular category has finite image.*

## Theorem (Ng–Schauenberg)

*Every modular representation associated to a modular category has finite image.*

## Theorem (Etingof–Rowell–Witherspoon)

*The braid group representation associated to the modular category  $\text{Mod}(D^\omega(G))$  has finite image.*

## Theorem (Ng–Schauenberg)

*Every modular representation associated to a modular category has finite image.*

## Theorem (Etingof–Rowell–Witherspoon)

*The braid group representation associated to the modular category  $\text{Mod}(D^\omega(G))$  has finite image.*

ERW also asked if, more generally, all mapping class group representations associated to  $\text{Mod}(D^\omega(G))$  have finite image.

## Theorem (Ng–Schauenberg)

*Every modular representation associated to a modular category has finite image.*

## Theorem (Etingof–Rowell–Witherspoon)

*The braid group representation associated to the modular category  $\text{Mod}(D^\omega(G))$  has finite image.*

ERW also asked if, more generally, all mapping class group representations associated to  $\text{Mod}(D^\omega(G))$  have finite image.

## Theorem (Fjelstad–Fuchs)

*Every mapping class group representation of a closed surface with at most one marked point associated to  $\text{Mod}(D(G))$  has finite image.*

# Answering ERW's question




TABLE 1. First type of Dehn twist. Unlabeled interior edges are colored by the group identity element.

# Answering ERW's question




TABLE 2. Second type of Dehn twist.

# Answering ERW's question




TABLE 3. A braid generator. Unlabeled interior edges are colored by the group identity element.

# Answering ERW's question




TABLE 4. Pulling a boundary component along a generator for the fundamental group of the corresponding closed surface. Unlabeled interior edges are colored by the group identity element.



## Theorem

*The image of any  $\text{Vect}_G^\omega$  TVBW representation  $\rho$  of a mapping class group of an orientable, compact surface  $\Sigma$  with boundary is finite.*

*Sketch of proof.*

- For any  $k$ , let  $\mu_{|G|}$  denote the set of  $|G|$ -th roots of unity. Then  $\omega$  is cohomologous to a cocycle taking values in  $\mu_{|G|}$ . Since cohomologous cocycles give rise to equivalent spherical categories  $\text{Vect}_G^\omega$ , WLOG  $\omega$  takes values in  $\mu_{|G|}$ .

# First result

## Theorem

*The image of any  $\text{Vect}_G^\omega$  TVBW representation  $\rho$  of a mapping class group of an orientable, compact surface  $\Sigma$  with boundary is finite.*

*Sketch of proof.*

- For any  $k$ , let  $\mu_{|G|}$  denote the set of  $|G|$ -th roots of unity. Then  $\omega$  is cohomologous to a cocycle taking values in  $\mu_{|G|}$ . Since cohomologous cocycles give rise to equivalent spherical categories  $\text{Vect}_G^\omega$ , WLOG  $\omega$  takes values in  $\mu_{|G|}$ .
- Let  $L \subset \text{MCG}(\Sigma)$  be the Birman generating set. From the figures and the definitions of structural morphisms of  $\text{Vect}_G^\omega$ ,  $\rho(L)$  acts by permutations on  $\mu_{|G|}S$  for a spanning set  $S$ .

# First result

## Theorem

*The image of any  $\text{Vect}_G^\omega$  TVBW representation  $\rho$  of a mapping class group of an orientable, compact surface  $\Sigma$  with boundary is finite.*

*Sketch of proof.*

- For any  $k$ , let  $\mu_{|G|}$  denote the set of  $|G|$ -th roots of unity. Then  $\omega$  is cohomologous to a cocycle taking values in  $\mu_{|G|}$ . Since cohomologous cocycles give rise to equivalent spherical categories  $\text{Vect}_G^\omega$ , WLOG  $\omega$  takes values in  $\mu_{|G|}$ .
- Let  $L \subset \text{MCG}(\Sigma)$  be the Birman generating set. From the figures and the definitions of structural morphisms of  $\text{Vect}_G^\omega$ ,  $\rho(L)$  acts by permutations on  $\mu_{|G|}S$  for a spanning set  $S$ .
- Let  $B \subset S$  be a basis for  $H$ . Then  $\rho(\text{MCG}(\Sigma))B \subset \rho(\text{MCG}(\Sigma))S \subset \mu_{|G|}S$ .

# First result

## Theorem

*The image of any  $\text{Vect}_G^\omega$  TVBW representation  $\rho$  of a mapping class group of an orientable, compact surface  $\Sigma$  with boundary is finite.*

*Sketch of proof.*

- For any  $k$ , let  $\mu_{|G|}$  denote the set of  $|G|$ -th roots of unity. Then  $\omega$  is cohomologous to a cocycle taking values in  $\mu_{|G|}$ . Since cohomologous cocycles give rise to equivalent spherical categories  $\text{Vect}_G^\omega$ , WLOG  $\omega$  takes values in  $\mu_{|G|}$ .
- Let  $L \subset \text{MCG}(\Sigma)$  be the Birman generating set. From the figures and the definitions of structural morphisms of  $\text{Vect}_G^\omega$ ,  $\rho(L)$  acts by permutations on  $\mu_{|G|}S$  for a spanning set  $S$ .
- Let  $B \subset S$  be a basis for  $H$ . Then  $\rho(\text{MCG}(\Sigma))B \subset \rho(\text{MCG}(\Sigma))S \subset \mu_{|G|}S$ .
- Thus,  $|\text{Im}(\rho)| < \infty$ .

# Next steps

- Tambara-Yamagami categories
- Calculate actual matrices

# Calculations = Hard

Easiest one:

$$\begin{aligned} & \frac{\omega(h, g, h^{-1})\omega(h, gh^{-1}, hg^{-1}h^{-1})\omega(g, h^{-1}, hg^{-1}h^{-1})\omega(g, g^{-1}h^{-1}, h)}{\omega(g^{-1}, g^{-1}, g)\omega(g^{-1}, g^{-1}h^{-1}, h)\omega(g^{-1}, h^{-1}, hg^{-1}h^{-1})\omega(g, g^{-2}h^{-1}, hg)} \\ & \quad \frac{\omega(g, g^{-1}, h^{-1})\omega(g, g^{-1}h^{-1}, h)\omega(g^{-1}h^{-1}, h, g)\omega(g^{-2}h^{-1}, h, g)}{\omega(g, g^{-1}h^{-1}, hg^{-1}h^{-1})\omega(hg, h^{-1}, hg^{-1}h^{-1})\omega(hg, g, g^{-1}h^{-1})} \\ & = \frac{\omega(h, g, g^{-1}h^{-1})\omega(g, g^{-1}h^{-1}, h)}{\omega(g^{-1}, g^{-1}, g)\omega(g^{-1}, g^{-1}h^{-1}, h)\omega(g^{-1}, h^{-1}, hg^{-1}h^{-1})\omega(g, g^{-2}h^{-1}, hg)} \\ & \quad \frac{\omega(g, g^{-1}, h^{-1})\omega(g, g^{-1}h^{-1}, h)\omega(g^{-1}h^{-1}, h, g)\omega(g^{-2}h^{-1}, h, g)}{\omega(g, g^{-1}h^{-1}, hg^{-1}h^{-1})\omega(hg, g, g^{-1}h^{-1})} \\ & = \frac{\omega(h, g, g^{-1}h^{-1})\omega(g, g^{-1}h^{-1}, h)}{\omega(g^{-1}, g^{-1}, g)\omega(g^{-1}, g^{-1}h^{-1}, h)\omega(g, g^{-2}h^{-1}, hg)} \\ & \quad \frac{\omega^2(g, g^{-1}, h^{-1})\omega(g, g^{-1}h^{-1}, h)\omega(g^{-1}h^{-1}, h, g)\omega(g^{-2}h^{-1}, h, g)}{\omega(g, g^{-1}, g^{-1}h^{-1})\omega(hg, g, g^{-1}h^{-1})} \\ & = \frac{\omega(h, g, g^{-1}h^{-1})\omega^2(g, g^{-1}h^{-1}, h)\omega^2(g, g^{-1}, h^{-1})\omega^2(g^{-1}h^{-1}, h, g)}{\omega(g, g^{-2}h^{-1}, hg)\omega(g, g^{-1}, g^{-1}h^{-1})\omega(hg, g, g^{-1}h^{-1})\omega(g^{-1}, g^{-1}h^{-1}, hg)} \\ & = \frac{\omega(h, g, g^{-1}h^{-1})\omega^2(g, g^{-1}h^{-1}, h)\omega^2(g, g^{-1}, h^{-1})\omega^2(g^{-1}h^{-1}, h, g)}{\omega(hg, g, g^{-1}h^{-1})} \\ & = \frac{\omega(h, g, g^{-1}h^{-1})\omega^2(g^{-1}h^{-1}, h, g)}{\omega^2(g^{-1}, h^{-1}, h)\omega(hg, g, g^{-1}h^{-1})} \end{aligned}$$