

Coordinate-free Mathematics

Paul Gustafson

May 2025

The first section develops algebraic constructions independent of a basis; the second develops geometric constructions independent of coordinates; the third develops analytic constructions from operators and positivity.

1 Algebra

Linear algebra studies linear maps between vector spaces. A **vector space** V over a field k is a set equipped with vector addition and scalar multiplication operations satisfying straightforward compatibility conditions. A **linear map** $f : V \rightarrow U$ between k -vector spaces is a map of underlying sets such that $f(\alpha v + \beta w) = \alpha f(v) + \beta f(w)$ for all $\alpha, \beta \in k$ and $v, w \in V$. Examples include differential operators, group representations, coefficient matrices of linear systems of equations, and local approximations of globally nonlinear functions [1, 2].

The **free k -vector space on a set** X , denoted kX , is the set of finitely supported set-theoretical functions $X \rightarrow k$ with pointwise addition and scalar multiplication. The **direct sum** vector space $V \oplus W$ is the set $V \times W$ with pointwise addition and scalar multiplication defined by $\alpha(v \oplus w) = \alpha v \oplus \alpha w$. The **quotient** of V by a subspace W , denoted V/W , is the set of equivalence classes of V modulo the relation $x \sim y$ iff $x - y \in W$. The **tensor product** $V \otimes W$ is the vector space $k(V \times W)/N$ where N is the subspace generated by the obvious linearity relations on V and W .

Exercise 1 An **isomorphism** of vector spaces $V \cong W$ consists of a pair of linear maps $f : V \rightarrow W$ and $g : W \rightarrow V$ such that $f \circ g = 1_W$ and $g \circ f = 1_V$. Prove the following:

- $k(X \sqcup Y) \cong kX \oplus kY$
- $k(X \times Y) \cong kX \otimes kY$
- The first isomorphism theorem for vector spaces (and second and third if you want).
- Find universal properties determining up to canonical isomorphism the free k -vector space, direct sum, quotient, tensor product, and dual space (defined below), respectively.

The **dual space** V^* is the space of linear maps from a vector space V to its ground field k . The evaluation map $\text{ev}_V : V^* \otimes V \rightarrow k$ is the linear extension of the evaluation morphism $f \otimes x \mapsto f(x)$. We say that V is **finite dimensional** if there is a co-evaluation morphism $\text{coev}_V : k \rightarrow V \otimes V^*$ satisfying the snake equations $(1_V \otimes \text{ev}_V) \circ (\text{coev}_V \otimes 1_V) = 1_V$ and $(\text{ev}_V \otimes 1_{V^*}) \circ (1_{V^*} \otimes \text{coev}_V) = 1_{V^*}$. We define the **dimension** $\dim(V) = \text{tr}(1_V)$, where $\text{tr}(f) : \text{End}(V) \rightarrow k$ is given by $f \mapsto (\text{ev}_V \circ s_{V,V^*} \circ (f \otimes 1_{V^*}) \circ \text{coev}_V)(1)$ where $s_{X,Y} : X \otimes Y \rightarrow Y \otimes X$ is the symmetric braiding $x \otimes y \mapsto y \otimes x$ [5].

Exercise 2 Prove the following for finite-dimensional vector spaces V, W and linear maps $f, g : V \rightarrow V$.

- $\text{tr}(fg) = \text{tr}(gf)$
- $\text{tr}(f \oplus g) = \text{tr}(f) + \text{tr}(g)$
- $\text{tr}(f \otimes g) = \text{tr}(f) \text{tr}(g)$
- $\dim(V) \in \mathbf{N}$
- $V \cong W$ iff $\dim(V) = \dim(W)$

The simplest linear maps are scalar multiples of the identity. Although most linear maps are not of this form, it is useful to decompose them into sums of scalar maps when possible. One condition under which we can decompose a linear map into a direct sum of scalars is if the map is self-adjoint. Suppose V is finite dimensional over $\mathbb{C}$. A Hermitian inner product $\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{C}$ is a positive-definite sesquilinear form (linear in the first variable, anti-linear in the second variable, and conjugate symmetric). Write $|v| = \sqrt{\langle v, v \rangle}$. For a linear map $f : V \rightarrow W$ between Hermitian spaces, the adjoint $f^* : W \rightarrow V$ is characterized by $\langle fv, w \rangle = \langle v, f^*w \rangle$. An endomorphism f is **self-adjoint** if $f = f^*$. An isomorphism f is **unitary** if it preserves inner products, equivalently if $f^* = f^{-1}$.

For maps $f : V \rightarrow V'$ and $g : W \rightarrow W'$, define $(f \otimes g)(v \otimes w) = f(v) \otimes g(w)$. The tensor product of Hermitian spaces has inner product

$$\langle v_1 \otimes w_1, v_2 \otimes w_2 \rangle = \langle v_1, v_2 \rangle \langle w_1, w_2 \rangle.$$

The Cauchy–Schwarz inequality is $|\langle v, w \rangle| \leq |v||w|$, with equality iff v and w are linearly dependent.

Exercise 3 (No-cloning theorem) Let V be a finite-dimensional complex inner-product space, let $b \in V$ be a unit vector, and suppose that $C : V \otimes V \rightarrow V \otimes V$ is unitary. Prove that if [18]

$$C(v \otimes b) = v \otimes v, \quad C(w \otimes b) = w \otimes w$$

for unit vectors $v, w \in V$, then

$$\langle v, w \rangle = \langle v, w \rangle^2.$$

Deduce that either $\langle v, w \rangle = 0$ or $v = w$. Conclude that no unitary map can satisfy $C(v \otimes b) = v \otimes v$ for every unit vector $v \in V$.

A scalar $\lambda \in k$ is an **eigenvalue** for a linear map $f : V \rightarrow V$ if there exists a nonzero vector $v \in V$ for which $fv = \lambda v$. The **eigenspace** of λ is

$$\ker(f - \lambda 1_V),$$

whose nonzero elements are eigenvectors; its dimension is called the **geometric multiplicity** of λ .

Exercise 4 *Prove that any self-adjoint linear map f on a finite-dimensional complex vector space V is diagonalizable:*

$$f = \sum \lambda_i p_i,$$

where $\sum p_i = 1_V$ and $p_i p_j = \delta_{ij} p_i$. Use this fact to prove the existence of the singular value decomposition for an arbitrary linear map between complex finite-dimensional vector spaces.

For $\psi \in V \otimes W$, define

$$T_\psi : V^* \rightarrow W, \quad T_\psi(\phi) = (\phi \otimes 1_W)(\psi).$$

The map $j_V : V \rightarrow V^*$ given by $j_V(v) = \langle \cdot, v \rangle$ is conjugate-linear. Equip V^* with the Hermitian inner product determined by $\langle j_V(v), j_V(w) \rangle = \langle w, v \rangle$. With $T_\psi^* : W \rightarrow V^*$ denoting the adjoint, define

$$\rho_V = j_V^{-1} T_\psi^* T_\psi j_V, \quad \rho_W = T_\psi T_\psi^*.$$

Write $U(V)$ for the group of unitary endomorphisms of V , acting with $U(W)$ on $V \otimes W$ by tensor products. A nonzero tensor is **decomposable** if it has the form $v \otimes w$. The rank of an endomorphism is the dimension of its image.

Exercise 5 (Schmidt decomposition and reduced endomorphisms)

Let V and W be finite-dimensional Hermitian vector spaces and let $0 \neq \psi \in V \otimes W$. Prove that there exist orthonormal families (v_i) in V and (w_i) in W , and positive scalars s_i , such that [17]

$$\psi = \sum_{i=1}^r s_i v_i \otimes w_i.$$

Prove the following:

- The nonzero eigenvalues of ρ_V and ρ_W are the numbers s_i^2 , with the same multiplicities.
- ψ is decomposable iff exactly one s_i is nonzero.
- ρ_V has rank one iff ρ_W has rank one iff ψ is decomposable.
- The multiset (s_i) is invariant under the action of $U(V) \times U(W)$.

For a unit vector $\psi \in V$ and a self-adjoint $A \in \text{End}(V)$, define

$$m_\psi(A) = \langle \psi, A\psi \rangle, \quad \Delta_\psi(A) = |(A - m_\psi(A)1_V)\psi|.$$

Exercise 6 (Uncertainty principle) Let $A, B \in \text{End}(V)$ be self-adjoint and let $\psi \in V$ be a unit vector. Prove that [19]

$$\Delta_\psi(A)\Delta_\psi(B) \geq \frac{1}{2} |\langle \psi, [A, B]\psi \rangle|, \quad [A, B] = AB - BA.$$

Determine the condition for equality from the equality condition in the Cauchy-Schwarz inequality.

For $K \in \text{End}(V)$, define $e^K = \sum_{n \geq 0} K^n/n!$. If H is self-adjoint, write $U_t = e^{-itH}$.

Exercise 7 (Conservation from commutators)

Let $A, H \in \text{End}(V)$ be self-adjoint, let $U_t = e^{-itH}$, and let $\psi \in V$. Prove that [17]

$$\frac{d}{dt} \langle U_t\psi, A U_t\psi \rangle = -i \langle U_t\psi, [H, A] U_t\psi \rangle.$$

Deduce that $[H, A] = 0$ implies that $\langle U_t\psi, A U_t\psi \rangle$ is independent of t for every $\psi \in V$. Prove the converse.

We can solve for the eigenvalues of a linear operator on a finite-dimensional vector space via the determinant. Define the tensor algebra and **exterior algebra** of V by [4]

$$T(V) = \bigoplus_{k \geq 0} V^{\otimes k}, \quad \Lambda V = T(V)/\langle v \otimes v : v \in V \rangle.$$

The space $\Lambda^k V$ is the degree- k component of ΛV .

Exercise 8 Show that the highest graded component of ΛV is one-dimensional.

The **determinant** $\det(f)$ is the coefficient of the linear map on the highest-graded component of ΛV given by $v_1 \wedge \cdots \wedge v_{\dim(V)} \mapsto f(v_1) \wedge \cdots \wedge f(v_{\dim(V)})$.

For an n -dimensional real vector space V , its **determinant line** is $\det(V^*) = \Lambda^n V^*$. An **orientation** is a choice of one of the two connected components of $\det(V^*) \setminus \{0\}$; the elements of the chosen component are positive.

Exercise 9 Prove the following:

- $\det(f \oplus g) = \det(f) \det(g)$
- If $V = \mathbb{R}^n$, then $\det(f)$ is the signed volume of $f(I^n)$ where $I = [0, 1]$.
- Cramer's rule

For $f \in \text{End}(V)$, the **characteristic polynomial** is the polynomial $\lambda \mapsto \det(f - \lambda 1_V)$.

Exercise 10 Assume that the characteristic polynomial of f splits over k . Prove the following for a linear operator f on a finite-dimensional vector space V :

- Every eigenvalue of f is a root of its characteristic polynomial.
- The geometric multiplicity of each eigenvalue is at most equal to its **algebraic multiplicity** (i.e. multiplicity as a root of the characteristic polynomial)
- $\text{tr}(f) = \sum_i \lambda_i$ and $\det(f) = \prod_i \lambda_i$, where each eigenvalue λ_i is repeated according to its algebraic multiplicity
- The operator f satisfies its own characteristic polynomial (replacing λ with f)
- The existence of the Jordan normal form for a linear operator on a finite-dimensional complex vector space

Let $\mathcal{C}$ be a $\mathbb{C}$ -linear monoidal category with tensor product $\otimes$, tensor unit 1 , and finite direct sums. Write $\text{Hom}_{\mathcal{C}}(x, y)$ for its complex vector space of morphisms from x to y . An object x is **simple** if $\text{End}(x) \cong \mathbb{C}$, and $\mathcal{C}$ is **semisimple** if every object is a finite direct sum of simple objects. Assume that 1 is simple, that there are finitely many isomorphism classes of simple objects, and that $\mathcal{C}$ is **rigid**: every x has a dual x^* with evaluation and coevaluation maps satisfying the snake identities [26].

The definitions of duality, trace, and dimension above extend verbatim to $\mathcal{C}$. Write $d_x = \text{tr}(1_x)$. Then

$$d_{x \oplus y} = d_x + d_y, \quad d_{x \otimes y} = d_x d_y.$$

Assume that the categorical dimensions of nonzero objects are positive real numbers.

For $x, y, z \in \mathcal{C}$, write

$$F^{xyz} : (x \otimes y) \otimes z \longrightarrow x \otimes (y \otimes z)$$

for the associator. A **braiding** is a family of isomorphisms

$$c_{x,y} : x \otimes y \longrightarrow y \otimes x.$$

For $f : x \rightarrow x'$ and $g : y \rightarrow y'$, it satisfies $(g \otimes f)c_{x,y} = c_{x',y'}(f \otimes g)$. The associator satisfies the pentagon identity, and the associator and braiding satisfy the hexagon identities. Suppressing associators, the latter are

$$\begin{aligned} c_{x,y \otimes z} &= (1_y \otimes c_{x,z})(c_{x,y} \otimes 1_z), \\ c_{x \otimes y,z} &= (c_{x,z} \otimes 1_y)(1_x \otimes c_{y,z}). \end{aligned}$$

The **monodromy** of x and y is the double braiding

$$M_{x,y} = c_{y,x}c_{x,y} : x \otimes y \longrightarrow x \otimes y.$$

Thus $c_{x,y}$ is one crossing, while $M_{x,y}$ is a full braid. If $x \otimes y$ is simple, identify $M_{x,y}$ with its scalar in $\mathbb{C}^\times = \mathbb{C} \setminus \{0\}$. Decompositions of tensor products of simple objects are **fusion rules**, and the spaces $\text{Hom}_{\mathcal{C}}(z, x \otimes y)$ are **fusion spaces**.

An object x is **invertible** if $x \otimes x^* \cong 1$. Assume first that every simple object is invertible and that their isomorphism classes form the cyclic group

$$\{1, a, a^2, \dots, a^{m-1}\}, \quad a^r \otimes a^s \cong a^{r+s \pmod{m}}.$$

Write $U(1) = \{z \in \mathbb{C} : |z| = 1\}$ and $M_{r,s} = M_{a^r, a^s} \in U(1)$.

Exercise 11 (Abelian braiding phase)

Use the hexagon identities to prove that

$$M_{r+s,t} = M_{r,t}M_{s,t}, \quad M_{r,s+t} = M_{r,s}M_{r,t}.$$

Deduce that there is an integer k modulo m such that

$$M_{r,s} = \exp\left(\frac{2\pi i k r s}{m}\right).$$

For $k = 1$, prove that the monodromy of a about N copies of a is multiplication by

$$\exp\left(\frac{2\pi i N}{m}\right).$$

In particular, for $m = 3$, increasing N by one changes the phase by $2\pi/3$.

The phase $2\pi/3$ is the value observed by interferometry in the $\nu = 1/3$ fractional quantum Hall state [28].

Now suppose that $\mathcal{C}$ has two simple objects $1, \tau$ satisfying the fusion rule

$$\tau \otimes \tau \cong 1 \oplus \tau.$$

Exercise 12 (Fibonacci categorical dimension)

Let $d_\tau = \text{tr}(1_\tau)$. Prove that [26]

$$d_\tau^2 = 1 + d_\tau$$

and, using positivity, that

$$d_\tau = \frac{1 + \sqrt{5}}{2}.$$

Let

$$V_n = \text{Hom}_{\mathcal{C}}(1, \tau^{\otimes n}).$$

Show that the dimensions of these fusion spaces satisfy the Fibonacci recurrence and determine their asymptotic growth rate.

Let

$$V = \text{Hom}_{\mathcal{C}}(\tau, \tau^{\otimes 3}).$$

This fusion space is two-dimensional, with basis indexed by the intermediate objects $1, \tau$ in $(\tau \otimes \tau) \otimes \tau$. Set $\varphi = (1 + \sqrt{5})/2$ and choose [27]

$$F = \begin{pmatrix} \varphi^{-1} & \varphi^{-1/2} \\ \varphi^{-1/2} & -\varphi^{-1} \end{pmatrix}, \quad R = \begin{pmatrix} e^{-4\pi i/5} & 0 \\ 0 & e^{3\pi i/5} \end{pmatrix}.$$

Define $\sigma_1 = R$ and $\sigma_2 = F^{-1}RF$. Let

$$B_3 = \langle b_1, b_2 \mid b_1 b_2 b_1 = b_2 b_1 b_2 \rangle.$$

Exercise 13 (Non-Abelian braid representation)

Working exactly, verify that F and R are unitary and prove that

$$\sigma_1 \sigma_2 \sigma_1 = \sigma_2 \sigma_1 \sigma_2.$$

Prove also that $\sigma_1 \sigma_2 \neq \sigma_2 \sigma_1$. Deduce that $b_i \mapsto \sigma_i$ defines a non-Abelian representation

$$B_3 \longrightarrow U(V).$$

Such Fibonacci fusion and braiding data have been implemented on superconducting quantum processors [30].¹

2 Geometry

A **smooth manifold** is a topological space that is locally diffeomorphic to Euclidean space $\mathbb{R}^n$. A **smooth vector bundle** $p : E \rightarrow M$ consists of a smooth surjection p between smooth manifolds E (the total space) and M (the base space) such that all fibers are isomorphic to a single vector space V , and smooth local trivializations exist (locally, the fiber bundle looks like a product space). For a smooth map $f : N \rightarrow M$, the **pullback bundle** is

$$f^*E = \{(x, e) \in N \times E : f(x) = p(e)\} \longrightarrow N.$$

A prototypical example is the **tangent bundle** TM which consists of all tangent spaces of points of a given manifold. The **cotangent bundle** T^*M is the dual bundle of TM . A **vector field** $X \in \Gamma(TM)$ is a smooth section of the tangent bundle (i.e. a smooth map $s : M \rightarrow TM$ such that $p \circ s = 1_M$) [3]. An orientation of M is a smoothly varying choice of orientation of each tangent space, using the determinant lines defined in the Algebra section.

Exercise 14 *Write down precise definitions for the following:*

¹This is an engineered quantum simulation of Fibonacci topological order, not evidence that naturally occurring Fibonacci quasiparticles have been conclusively observed. A related trapped-ion realization of non-Abelian topological order is reported in [29].

- smooth manifold, smooth map, vector bundle, tangent bundle.
- bundle map, pullback, direct sum, tensor product, and dual bundle
- derivative of a smooth function between manifolds

For a compact Hausdorff space X , let $\text{Vect}(X)$ be the category of finite-rank complex vector bundles with continuous local trivializations over X . It has direct sums, tensor products, duals, and pullbacks, and is an additive symmetric monoidal category with duals [7]. The **Grothendieck group** $K^0(X)$ is generated by isomorphism classes $[E]$ subject to

$$[E \oplus F] = [E] + [F].$$

Tensor product defines $[E][F] = [E \otimes F]$, making $K^0(X)$ a commutative ring. For a continuous map $f : X \rightarrow Y$ between compact Hausdorff spaces, pullback defines

$$f^* : K^0(Y) \longrightarrow K^0(X), \quad [E] \longmapsto [f^*E].$$

Exercise 15 (Grothendieck group of vector bundles)

Let X and Y be compact Hausdorff spaces and let $f : X \rightarrow Y$ be continuous. Prove that direct sum and tensor product induce a commutative ring structure on $K^0(X)$ and that f^* is a ring homomorphism. Prove that

$$K^0(\text{pt}) \cong \mathbb{Z}$$

via the rank map.

Fix a basepoint $x_0 \in X$, regarded as a map $x_0 : \text{pt} \rightarrow X$. The **reduced K-theory** of the pointed space (X, x_0) is

$$\tilde{K}^0(X) = \ker \left(K^0(X) \xrightarrow{x_0^*} K^0(\text{pt}) \right).$$

For a bundle E of rank r , its **reduced class** is

$$[E] - [\underline{\mathbb{C}}^r],$$

where $\underline{\mathbb{C}}^r = X \times \mathbb{C}^r$ is the trivial rank- r bundle. Topological K-theory extends K^0 to generalized cohomology groups $K^n(X)$. Bott periodicity gives

$$K^{n+2}(X) \cong K^n(X),$$

so complex K-theory is naturally $\mathbb{Z}/2$ -graded.

Exercise 16 (Stable equivalence)

Let E and F be complex vector bundles over a compact Hausdorff space X . Prove that $[E] = [F]$ in $K^0(X)$ if and only if there is a vector bundle G such that

$$E \oplus G \cong F \oplus G.$$

Deduce that the reduced class of E is unchanged after replacing E by $E \oplus \underline{\mathbb{C}}^n$.

For the gapped families below, assume that X is connected, so vector-bundle ranks are constant. Let V be a finite-dimensional Hermitian vector space. A continuous family is a map

$$H : X \longrightarrow \text{End}(V)$$

and we write $H_x = H(x)$. It is **gapped** if every H_x is self-adjoint and invertible. Compactness gives numbers $\delta, M > 0$ such that every eigenvalue of every H_x lies in $[-M, -\delta] \cup [\delta, M]$. Choose a positively oriented contour Γ enclosing $[-M, -\delta]$ and excluding $[\delta, M]$, and define

$$P_x^- = \frac{1}{2\pi i} \int_{\Gamma} (z - H_x)^{-1} dz, \quad E_x^- = \text{im}(P_x^-).$$

Exercise 17 (Negative spectral bundle)

Let H be a gapped family. Prove that

$$(P_x^-)^2 = P_x^-, \quad (P_x^-)^* = P_x^-,$$

that $x \mapsto P_x^-$ is continuous, and that

$$E_H^- = \coprod_{x \in X} E_x^-$$

is a complex vector bundle over X . Prove that its rank is locally constant.

For a gapped family $H : X \times [0, 1] \rightarrow \text{End}(V)$, let E_j^- be the negative spectral bundle of $x \mapsto H(x, j)$, and let $i_j : X \rightarrow X \times [0, 1]$ be given by $i_j(x) = (x, j)$.

Exercise 18 (Gap-preserving homotopy)

Prove that the negative spectral subspaces of H form a vector bundle $E^- \rightarrow X \times [0, 1]$. Prove that the endpoint pullbacks of any vector bundle over $X \times [0, 1]$ are isomorphic, and apply this to $i_0^* E^-$ and $i_1^* E^-$. Deduce that

$$[E_0^-] = [E_1^-] \in K^0(X).$$

Let $Q \in \text{End}(W)$ be a fixed invertible self-adjoint endomorphism of a finite-dimensional Hermitian space W , let W^- be its negative eigenspace, and write $\underline{W}^- = X \times W^-$.

Exercise 19 (Stabilization)

For a gapped family H , prove that

$$E_{H \oplus Q}^- \cong E_H^- \oplus \underline{W}^-.$$

Deduce that

$$[E_H^-] - [\mathbb{C}^{\text{rank } E_H^-}] \in \tilde{K}^0(X)$$

is unchanged by adjoining a constant gapped summand.

Let $E \subseteq X \times V$ be a subbundle and let P_x be the orthogonal projection onto E_x .

Exercise 20 (Flattened family)

Define $H_x = 1_V - 2P_x$. Prove that $x \mapsto H_x$ is a gapped family, that

$$H_x^2 = 1_V,$$

and that its negative spectral bundle is E . Deduce that every subbundle of a trivial bundle arises as the negative spectral bundle of a gapped family.

Thus a finite-dimensional gapped family determines a class in $\tilde{K}^0(X)$ that is invariant under gap-preserving homotopy and stabilization. Conversely, every subbundle of a trivial bundle occurs as a negative spectral bundle.

In a finite-dimensional gapped family, the negative spectral subspaces form a vector bundle. Gap-preserving deformation preserves its stable isomorphism class, so the associated reduced K-theory class is a topological invariant. This construction underlies the classification of noninteracting topological insulators and superconductors under specified symmetry assumptions [8].

The exterior powers $\Lambda^k T_p^* M$ are those defined in the Algebra section. To define integration in a coordinate-free way, we will integrate differential forms over chains. The space of **differential n -forms** is $\Omega^n(M) = \Gamma(\Lambda^n T^* M)$. On an oriented n -manifold, a **volume form** is a nowhere-vanishing element of $\Omega^n(M)$ that is positive with respect to the orientation. The exterior derivative $d : \Omega^n(M) \rightarrow \Omega^{n+1}(M)$ extends the differential of smooth functions and satisfies [6]

$$d(\alpha \wedge \beta) = d\alpha \wedge \beta + (-1)^p \alpha \wedge d\beta$$

for $\alpha \in \Omega^p(M)$. A **singular n -simplex** is a smooth map $\sigma : \Delta^n \rightarrow M$ where Δ^n is the standard n -simplex. The collection of singular n -simplices in M freely generates the abelian group $C_n(M)$ of **singular n -chains** in M . The **boundary map** $\partial : C_n(M) \rightarrow C_{n-1}(M)$ is defined on each simplex $\sigma : \Delta^n \rightarrow M$ by

$$\partial\sigma = \sum_{j=0}^n (-1)^j \sigma \circ f_j,$$

where $f_j : \Delta^{n-1} \rightarrow \Delta^n$ is the inclusion of the face opposite the j -th vertex of Δ^n .

A **chain complex** $(A_n, d_n)_n$ is a sequence of abelian groups A_n and homomorphisms $d_n : A_n \rightarrow A_{n-1}$ such that $d_{n-1} \circ d_n = 0$ for all n . The n -th **homology group** H_n of the chain complex is

$$H_n = \ker(d_n) / \operatorname{im}(d_{n+1}).$$

The prototypical example is singular homology which corresponds to the chain complex of singular chains with the boundary map defined above.

Cohomology and cochains are defined similarly, using superscripts instead of subscripts and boundary maps $d^n : A^n \rightarrow A^{n+1}$ increase degree instead of

decrease it. Singular cohomology corresponds to the cochain complex given by dualizing the singular chain complex above.

The **de Rham cohomology** of M is

$$H_{\text{dR}}^n(M) = \ker(d: \Omega^n(M) \rightarrow \Omega^{n+1}(M)) / \text{im}(d: \Omega^{n-1}(M) \rightarrow \Omega^n(M)).$$

Exercise 21 *Prove the following:*

- Show that $d^2 = 0$, so differential forms on a manifold M form a cochain complex.
- Define the integral of an n -form on an n -chain. Prove that $\int_c d\omega = \int_{\partial c} \omega$ for any n -chain c and differential $n-1$ -form ω .
- Singular cohomology with real coefficients and de Rham cohomology are naturally isomorphic.
- Bounded chain complexes of finite-dimensional vector spaces form a symmetric monoidal abelian category with duals, the trace of an endomorphism is its Lefschetz number, and the dimension of a chain complex is its Euler characteristic.

For a vector field X , contraction is the map $\iota_X: \Omega^p(M) \rightarrow \Omega^{p-1}(M)$ defined by

$$(\iota_X \alpha)(X_2, \dots, X_p) = \alpha(X, X_2, \dots, X_p).$$

The Lie derivative of forms is determined by Cartan's formula $\mathcal{L}_X = d\iota_X + \iota_X d$.

A **symplectic form** is a two-form $\omega \in \Omega^2(M)$ such that $d\omega = 0$ and $X \mapsto \iota_X \omega$ is a bundle isomorphism $TM \rightarrow T^*M$. Write $C^\infty(M)$ for the smooth real-valued functions on M . For $H \in C^\infty(M)$, define X_H by

$$\iota_{X_H} \omega = dH.$$

Exercise 22 (Liouville's theorem)

Let (M, ω) be $2n$ -dimensional with ω symplectic. Prove that $\omega^n/n!$ is a volume form. For $H \in C^\infty(M)$, prove using Cartan's formula that [20]

$$\mathcal{L}_{X_H} \omega = 0.$$

Deduce that $\mathcal{L}_{X_H} \omega^n = 0$ and hence that the local flow of X_H preserves the volume form

$$\frac{\omega^n}{n!}.$$

For $f, g \in C^\infty(M)$, define

$$\{f, g\} = \omega(X_f, X_g).$$

The following is a Hamiltonian symmetry-conservation statement, not the full variational theorem of [21].

Exercise 23 (Hamiltonian Noether theorem)

Let (M, ω) be symplectic. Prove that [20]

$$X_g(f) = \{f, g\}.$$

Let Φ_t be the local flow of X_f . Prove that the following are equivalent:

- $H \circ \Phi_t = H$ whenever both sides are defined.
- $X_f(H) = 0$.
- $\{H, f\} = 0$.
- f is constant along the integral curves of X_H .

A **pseudo-Riemannian metric** is a smooth nondegenerate symmetric section $g \in \Gamma(T^*M \otimes T^*M)$. On an oriented pseudo-Riemannian n -manifold, the metric induces a bilinear form $\langle \cdot, \cdot \rangle_g$ on differential forms. The **metric volume form** vol_g is the unique positive volume form satisfying

$$|\langle \text{vol}_g, \text{vol}_g \rangle_g| = 1.$$

The Hodge star $\star$ is determined by

$$\alpha \wedge \star \beta = \langle \alpha, \beta \rangle_g \text{vol}_g.$$

Exercise 24 (Charge conservation from Maxwell's equations)

Let M be an oriented four-dimensional pseudo-Riemannian manifold. Suppose that $F \in \Omega^2(M)$ and $J \in \Omega^3(M)$ satisfy [22]

$$dF = 0, \quad d \star F = J.$$

Prove that $dJ = 0$. If C is an oriented compact four-dimensional chain, deduce that

$$\int_{\partial C} J = 0.$$

If the three-dimensional pieces of its boundary satisfy

$$\partial C = \Sigma_1 - \Sigma_0 + B,$$

prove that

$$\int_{\Sigma_1} J - \int_{\Sigma_0} J = - \int_B J.$$

For a closed form A , write $[A]$ for its de Rham cohomology class. The next exercise is a modern de Rham formulation; the original paper does not use this language [25, 22].

Exercise 25 (Aharonov–Bohm cohomological obstruction) Let M be a compact smooth manifold and let $A \in \Omega^1(M)$ satisfy $dA = 0$. Prove the following:

- For every contractible open set $U \subseteq M$, there exists $\lambda \in C^\infty(U)$ such that $A|_U = d\lambda$.
- For every smooth closed curve γ , the number $\int_\gamma A$ depends only on the homology class of γ .
- There exists a global $\lambda \in C^\infty(M)$ with $A = d\lambda$ iff $[A] = 0 \in H_{\text{dR}}^1(M)$.
- If $[A] \neq 0$, then there exists a smooth closed curve γ such that $\int_\gamma A \neq 0$.

An **affine connection** is a map $\nabla : \Gamma(TM) \times \Gamma(TM) \rightarrow \Gamma(TM)$ such that

$$\nabla_{fX}Y = f\nabla_XY, \quad \nabla_X(fY) = X(f)Y + f\nabla_XY.$$

For a covector field α , set

$$(\nabla_X\alpha)(Y) = X(\alpha(Y)) - \alpha(\nabla_XY).$$

The connection extends uniquely to all tensor fields by the product rule and by commuting with contractions; this defines the **covariant derivative** of a tensor field.

The **Levi-Civita connection** of a pseudo-Riemannian metric is the unique affine connection satisfying

$$\nabla_XY - \nabla_YX = [X, Y], \quad \nabla g = 0.$$

An affinely parametrized curve $\gamma : I \rightarrow M$ is a **geodesic** if $\nabla_{\dot{\gamma}}\dot{\gamma} = 0$. The Lie derivative of the metric is

$$(\mathcal{L}_Xg)(Y, Z) = X(g(Y, Z)) - g([X, Y], Z) - g(Y, [X, Z]),$$

and X is a **Killing vector field** if $\mathcal{L}_Xg = 0$.

Let (M, g) be a pseudo-Riemannian manifold with Levi-Civita connection ∇ . Curvature governs the relative acceleration of geodesics, while symmetries of g determine quantities preserved along them [23, 24].

Exercise 26 (Killing fields and conserved quantities)

Let X be a Killing vector field and let γ be an affinely parametrized geodesic. Prove that

$$\frac{d}{dt}g(\dot{\gamma}, \dot{\gamma}) = 0, \quad \frac{d}{dt}g(X, \dot{\gamma}) = 0.$$

Deduce that $g(X, \dot{\gamma})$ is constant along γ .

The **curvature** of ∇ is the map

$$R(X, Y)Z = \nabla_X\nabla_YZ - \nabla_Y\nabla_XZ - \nabla_{[X, Y]}Z.$$

Exercise 27 (Curvature as a tensor)

For $f \in C^\infty(M)$ and vector fields X, Y, Z , prove that

$$R(fX, Y)Z = fR(X, Y)Z, \quad R(X, fY)Z = fR(X, Y)Z,$$

$$R(X, Y)(fZ) = fR(X, Y)Z.$$

Verify explicitly that every derivative of f cancels. Prove that $R(X, Y) = -R(Y, X)$ and deduce that

$$R \in \Gamma(\Lambda^2 T^*M \otimes T^*M \otimes TM).$$

Let $\Gamma : (-\varepsilon, \varepsilon) \times I \rightarrow M$ be a smooth variation through affinely parametrized geodesics, and define the vector fields along Γ by

$$T = \frac{\partial \Gamma}{\partial t}, \quad J = \frac{\partial \Gamma}{\partial s}.$$

Then $[T, J] = 0$.

Exercise 28 (Geodesic deviation) For a variation Γ as above, prove that

$$\nabla_T \nabla_T J = R(T, J)T.$$

Thus curvature determines the relative acceleration of nearby geodesics.

The trace used below is the ordinary trace defined in the Algebra section. Define the **Ricci tensor**, **scalar curvature**, and **Einstein tensor** by

$$\text{Ric}(X, Y) = \text{tr}(Z \mapsto R(Z, X)Y), \quad S = \text{tr}_g(\text{Ric}), \quad G = \text{Ric} - \frac{1}{2}Sg.$$

For a symmetric covariant two-tensor A , its **divergence** is the covector field

$$(\text{div } A)(Y) = \text{tr}_g((X, Z) \mapsto (\nabla_X A)(Z, Y)).$$

Exercise 29 (Contracted Bianchi identity and conservation)

Assume the second Bianchi identity

$$(\nabla_X R)(Y, Z) + (\nabla_Y R)(Z, X) + (\nabla_Z R)(X, Y) = 0.$$

Contract this identity to prove that

$$\text{div } G = 0.$$

Deduce that if T is a symmetric covariant two-tensor satisfying

$$G + \Lambda g = \kappa T$$

for constants Λ, κ with $\kappa \neq 0$, then $\text{div } T = 0$.

3 Analysis

A **unital C^* -algebra** is a complex unital Banach algebra A with an involution $a \mapsto a^*$ satisfying

$$\|a^*a\| = \|a\|^2.$$

Its positive cone is $A_+ = \{a^*a : a \in A\}$. A **state** is a linear map $\varphi : A \rightarrow \mathbb{C}$ such that [9]

$$\varphi(a) \geq 0 \quad (a \in A_+), \quad \varphi(1) = 1.$$

Exercise 30 Let φ be a positive linear functional on a unital C^* -algebra A . Prove that

$$|\varphi(b^*a)|^2 \leq \varphi(a^*a)\varphi(b^*b).$$

Deduce that $\|\varphi\| = \varphi(1)$.

For a state φ , define

$$\langle a, b \rangle_\varphi = \varphi(b^*a), \quad N_\varphi = \{a \in A : \varphi(a^*a) = 0\},$$

and let H_φ be the Hilbert-space completion of A/N_φ . For a Hilbert space H , write $\mathcal{B}(H)$ for the C^* -algebra of bounded operators on H . A vector Ω is **cyclic** for a representation π if the span of $\pi(A)\Omega$ is dense.

Exercise 31 (GNS construction)

Let φ be a state on a unital C^* -algebra A . Prove that left multiplication induces a unital $*$ -representation

$$\pi_\varphi : A \longrightarrow \mathcal{B}(H_\varphi)$$

with a cyclic vector Ω_φ satisfying

$$\varphi(a) = \langle \pi_\varphi(a)\Omega_\varphi, \Omega_\varphi \rangle_\varphi.$$

Prove that $(H_\varphi, \pi_\varphi, \Omega_\varphi)$ is unique up to unitary equivalence.

For a linear map $\Phi : A \rightarrow B$ between unital C^* -algebras, its n -th **amplification** is

$$\Phi_n : M_n(A) \longrightarrow M_n(B), \quad [a_{ij}] \longmapsto [\Phi(a_{ij})].$$

The map Φ is **completely positive** if every Φ_n is positive. Its **completely bounded norm** is

$$\|\Phi\|_{\text{cb}} = \sup_{n \geq 1} \|\Phi_n\|.$$

Exercise 32 For $n \geq 2$, let

$$T : M_n(\mathbb{C}) \longrightarrow M_n(\mathbb{C}), \quad T(a) = a^\top.$$

Prove that T is positive but not completely positive, and prove that

$$\|T\|_{\text{cb}} = n.$$

For the standard matrix units $E_{ij} \in M_n(\mathbb{C})$ and a linear map $\Phi : M_n(\mathbb{C}) \rightarrow M_m(\mathbb{C})$, define the **Choi matrix**

$$C_\Phi = \sum_{i,j=1}^n E_{ij} \otimes \Phi(E_{ij}).$$

Exercise 33 (Choi theorem)

Let $\Phi : M_n(\mathbb{C}) \rightarrow M_m(\mathbb{C})$ be linear. Prove that the following are equivalent [10]:

- Φ is completely positive.
- $C_\Phi \geq 0$.
- There are linear maps $V_r : \mathbb{C}^m \rightarrow \mathbb{C}^n$ such that

$$\Phi(a) = \sum_r V_r^* a V_r.$$

Exercise 34 (Stinespring dilation)

Let $\Phi : A \rightarrow \mathcal{B}(H)$ be completely positive. Prove that there are a Hilbert space K , a $*$ -representation $\pi : A \rightarrow \mathcal{B}(K)$, and a bounded operator $V : H \rightarrow K$ such that [11]

$$\Phi(a) = V^* \pi(a) V.$$

Prove that V is an isometry when Φ is unital. Call the dilation minimal if the span of $\pi(A) V H$ is dense in K , and prove that a minimal dilation is unique up to unitary equivalence.

For a unital completely positive map $\Phi : A \rightarrow B$, define its **multiplicative domain** by

$$\text{MD}(\Phi) = \left\{ a \in A : \begin{array}{l} \Phi(a^*a) = \Phi(a)^* \Phi(a), \\ \Phi(aa^*) = \Phi(a) \Phi(a)^* \end{array} \right\}.$$

Exercise 35 (Multiplicative domain)

Let $\Phi : A \rightarrow B$ be unital and completely positive. Prove the Kadison–Schwarz inequality

$$\Phi(a)^* \Phi(a) \leq \Phi(a^*a).$$

Prove that $\text{MD}(\Phi)$ is a C^* -subalgebra of A and that, for $a \in \text{MD}(\Phi)$ and $b \in A$,

$$\Phi(ab) = \Phi(a) \Phi(b), \quad \Phi(ba) = \Phi(b) \Phi(a).$$

For $\mathcal{S} \subseteq \mathcal{B}(H)$, its **commutant** is

$$\mathcal{S}' = \{T \in \mathcal{B}(H) : Tx = xT \text{ for every } x \in \mathcal{S}\}.$$

A **von Neumann algebra** is a unital $*$ -subalgebra $M \subseteq \mathcal{B}(H)$ satisfying $M = M''$ [12]. A net (T_λ) converges to T in the **strong operator topology** if $T_\lambda \xi \rightarrow T\xi$ for every $\xi \in H$, and in the **weak operator topology** if

$$\langle T_\lambda \xi, \eta \rangle \longrightarrow \langle T\xi, \eta \rangle$$

for every $\xi, \eta \in H$.

Exercise 36 (Bicommutant theorem)

Let $A \subseteq \mathcal{B}(H)$ be a unital $*$ -subalgebra. Prove that

$$A'' = \overline{A}^{\text{SOT}} = \overline{A}^{\text{WOT}}.$$

The **center** of a von Neumann algebra is $Z(M) = M \cap M'$, and M is a **factor** if $Z(M) = \mathbb{C}1$. A positive functional or positive map is **normal** if it preserves suprema of bounded increasing nets of positive elements, and a positive functional ψ is **faithful** if $\psi(a^*a) = 0$ implies $a = 0$. A **type II₁ factor** is an infinite-dimensional factor with a faithful normal tracial state

$$\tau : M \longrightarrow \mathbb{C}, \quad \tau(ab) = \tau(ba), \quad \tau(1) = 1.$$

The normalized trace is an analytic analogue of the categorical trace in Algebra, not an identification with it.

For projections $p, q \in M$, write $p \sim q$ if there is $v \in M$ such that

$$v^*v = p, \quad vv^* = q.$$

Write $p \precsim q$ if p is equivalent to a projection r satisfying $rq = r$.

Exercise 37 (Projection dimension)

Let M be a type II₁ factor with normalized trace τ . Prove that

$$p \sim q \implies \tau(p) = \tau(q),$$

and prove the comparison theorem

$$p \precsim q \iff \tau(p) \leq \tau(q).$$

Deduce that $\tau(p) = \tau(q)$ if and only if $p \sim q$.

A von Neumann algebra is **finite** if $v^*v = 1$ implies $vv^* = 1$. Let $N \subseteq M$ be finite von Neumann algebras with a common normalized trace τ . A **trace-preserving conditional expectation** is a normal unital completely positive map $E_N : M \rightarrow N$ such that

$$\tau \circ E_N = \tau, \quad E_N(n_1 x n_2) = n_1 E_N(x) n_2$$

for $x \in M$ and $n_1, n_2 \in N$. Let $L^2(M, \tau)$ be the completion of M for

$$\langle x, y \rangle_\tau = \tau(y^*x).$$

Let M act on $L^2(M, \tau)$ by left multiplication.

Exercise 38 *Prove that E_N extends to the orthogonal projection*

$$e_N : L^2(M, \tau) \longrightarrow L^2(N, \tau).$$

Prove that

$$e_N = e_N^* = e_N^2, \quad e_N x e_N = E_N(x) e_N$$

for every $x \in M$.

The **basic construction** is

$$M_1 = \langle M, e_N \rangle'' \subseteq \mathcal{B}(L^2(M, \tau)),$$

the double commutant of the algebra generated by M and e_N . For a projection $p = [p_{ij}] \in M_k(N)$ acting by left multiplication, define

$$\dim_N(pL^2(N, \tau)^k) = \sum_{i=1}^k \tau(p_{ii}).$$

An inclusion $N \subseteq M$ has **finite Jones index** if $L^2(M, \tau)$ is isomorphic as a right N -module to such a module; then

$$[M : N] = \dim_N L^2(M, \tau).$$

Exercise 39 (Basic construction)

Let $N \subseteq M$ be an inclusion of type II_1 factors of finite Jones index. Prove that M_1 is a finite factor and that its normalized trace satisfies [13]

$$\tau_1(xe_N y) = \frac{1}{[M : N]} \tau(xy)$$

for $x, y \in M$.

Iterating the basic construction gives the **Jones tower**

$$N = M_{-1} \subset M = M_0 \subset M_1 \subset M_2 \subset \cdots.$$

Let e_i be the Jones projection used to construct M_i from M_{i-1} , and set

$$\delta = \sqrt{[M : N]}.$$

The **Temperley–Lieb algebra** with loop parameter δ is generated by U_i subject to

$$U_i^2 = \delta U_i, \quad U_i U_j = U_j U_i \quad (|i - j| \geq 2), \quad U_i U_{i \pm 1} U_i = U_i.$$

Exercise 40 (Temperley–Lieb relations)

Prove that the Jones projections satisfy [16, 14]

$$e_i^2 = e_i = e_i^*, \quad e_i e_j = e_j e_i \quad (|i - j| \geq 2),$$

$$e_i e_{i \pm 1} e_i = \delta^{-2} e_i.$$

Deduce that $U_i \mapsto \delta e_i$ defines a representation of the Temperley–Lieb algebra.

Let $q \in \mathbb{C}^\times$ satisfy $\delta = q + q^{-1}$ and define

$$\sigma_i = q 1 - U_i.$$

These operators reconnect the Temperley–Lieb construction with the braid groups introduced in Algebra.

Exercise 41 (Braid representation)

Prove that σ_i is invertible with inverse $q^{-1}1 - U_i$, and prove that

$$\sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1},$$

$$\sigma_i \sigma_j = \sigma_j \sigma_i \quad (|i - j| \geq 2).$$

Deduce that the σ_i define a representation of the braid group [15].

The compatible normalized traces on the Jones tower restrict to a positive **Markov trace** on the Temperley–Lieb algebras satisfying

$$\mathrm{tr}(xU_i) = \delta^{-1} \mathrm{tr}(x)$$

whenever x belongs to the algebra generated by the preceding generators. Define

$$P_0(\delta) = 1, \quad P_1(\delta) = \delta, \quad P_{n+1}(\delta) = \delta P_n(\delta) - P_{n-1}(\delta).$$

Exercise 42 Prove that, when $\sin \theta \neq 0$,

$$P_n(2 \cos \theta) = \frac{\sin((n+1)\theta)}{\sin \theta}.$$

Use positivity of the Markov trace to prove that if $0 < \delta < 2$, then

$$\delta = 2 \cos \left(\frac{\pi}{m} \right)$$

for some integer $m \geq 3$.

Exercise 43 (Jones index theorem)

Let $N \subseteq M$ be an inclusion of type II_1 factors of finite Jones index. Prove that [14, 13]

$$[M : N] \in \left\{ 4 \cos^2 \left(\frac{\pi}{n} \right) : n \geq 3 \right\} \cup [4, \infty).$$

References

- [1] Paul R. Halmos, *Finite-Dimensional Vector Spaces*, 2nd ed., Springer, 1974.
- [2] Kenneth Hoffman and Ray Kunze, *Linear Algebra*, 2nd ed., Prentice Hall, 1971.
- [3] Michael Spivak, *Calculus on Manifolds: A Modern Approach to Classical Theorems of Advanced Calculus*, Addison-Wesley, 1965.
- [4] Serge Lang, *Algebra*, 3rd ed., Springer, 2002.
- [5] Kate Ponto and Michael Shulman, *Traces in symmetric monoidal categories*, *Expositiones Mathematicae*, 32(3):248–273, 2014.

- [6] Raoul Bott and Loring W. Tu, *Differential Forms in Algebraic Topology*, Graduate Texts in Mathematics, Vol. 82, Springer, 1982.
- [7] Allen Hatcher, *Vector Bundles and K-Theory*, Version 2.2, November 2017.
- [8] Alexei Kitaev, *Periodic table for topological insulators and superconductors*, AIP Conference Proceedings, 1134:22–30, 2009. doi:10.1063/1.3149495.
- [9] Vern Paulsen, *Completely Bounded Maps and Operator Algebras*, Cambridge Studies in Advanced Mathematics, Vol. 78, Cambridge University Press, 2002. doi:10.1017/CBO9780511546631.
- [10] Man-Duen Choi, *Completely positive linear maps on complex matrices*, Linear Algebra and its Applications, 10(3):285–290, 1975. doi:10.1016/0024-3795(75)90075-0.
- [11] W. Forrest Stinespring, *Positive functions on C^* -algebras*, Proceedings of the American Mathematical Society, 6:211–216, 1955. doi:10.1090/S0002-9939-1955-0069403-4.
- [12] Masamichi Takesaki, *Theory of Operator Algebras I*, Encyclopaedia of Mathematical Sciences, Vol. 124, Springer, 2002.
- [13] Vaughan F. R. Jones and V. S. Sunder, *Introduction to Subfactors*, London Mathematical Society Lecture Note Series, Vol. 234, Cambridge University Press, 1997. doi:10.1017/CBO9780511566219.
- [14] Vaughan F. R. Jones, *Index for subfactors*, Inventiones Mathematicae, 72(1):1–25, 1983. doi:10.1007/BF01389127.
- [15] Vaughan F. R. Jones, *A polynomial invariant for knots via von Neumann algebras*, Bulletin of the American Mathematical Society, 12(1):103–111, 1985. doi:10.1090/S0273-0979-1985-15304-2.
- [16] H. N. V. Temperley and E. H. Lieb, *Relations between the “percolation” and “colouring” problem and other graph-theoretical problems associated with regular planar lattices*, Proceedings of the Royal Society A, 322(1549):251–280, 1971. doi:10.1098/rspa.1971.0067.
- [17] Michael A. Nielsen and Isaac L. Chuang, *Quantum Computation and Quantum Information*, 10th Anniversary ed., Cambridge University Press, 2010.
- [18] W. K. Wootters and W. H. Zurek, *A Single Quantum Cannot Be Cloned*, Nature, 299:802–803, 1982. doi:10.1038/299802a0.
- [19] H. P. Robertson, *The Uncertainty Principle*, Physical Review, 34:163–164, 1929. doi:10.1103/PhysRev.34.163.
- [20] V. I. Arnold, *Mathematical Methods of Classical Mechanics*, 2nd ed., Springer, 1989.

- [21] Emmy Noether, *Invariant Variation Problems*, translated by M. A. Tavel, Transport Theory and Statistical Physics, 1(3):186–207, 1971; originally published 1918. doi:10.1080/00411457108231446.
- [22] Theodore Frankel, *The Geometry of Physics: An Introduction*, 3rd ed., Cambridge University Press, 2011. doi:10.1017/CBO9781139061377.
- [23] Barrett O’Neill, *Semi-Riemannian Geometry with Applications to Relativity*, Pure and Applied Mathematics, Vol. 103, Academic Press, 1983.
- [24] Robert M. Wald, *General Relativity*, University of Chicago Press, 1984.
- [25] Yakir Aharonov and David Bohm, *Significance of Electromagnetic Potentials in the Quantum Theory*, Physical Review, 115(3):485–491, 1959. doi:10.1103/PhysRev.115.485.
- [26] Pavel Etingof, Shlomo Gelaki, Dmitri Nikshych, and Victor Ostrik, *Tensor Categories*, Mathematical Surveys and Monographs, Vol. 205, American Mathematical Society, 2015. doi:10.1090/surv/205.
- [27] Parsa H. Bonderson, *Non-Abelian Anyons and Interferometry*, Ph.D. thesis, California Institute of Technology, 2007. doi:10.7907/5NDZ-W890.
- [28] J. Nakamura, S. Liang, G. C. Gardner, and M. J. Manfra, *Direct observation of anyonic braiding statistics*, Nature Physics, 16:931–936, 2020. doi:10.1038/s41567-020-1019-1.
- [29] M. Iqbal et al., *Non-Abelian topological order and anyons on a trapped-ion processor*, Nature, 626:505–511, 2024. doi:10.1038/s41586-023-06934-4.
- [30] S. Xu et al., *Non-Abelian braiding of Fibonacci anyons with a superconducting processor*, Nature Physics, 20:1469–1475, 2024. doi:10.1038/s41567-024-02529-6.