

Emergent Mathematics

Paul Gustafson

Contents

I	Symmetry	4
1	Linear Spaces, Tensor Products, and Duality	4
2	Hilbert Spaces and Trace Ideals	5
3	Measure, Integration, and L^p Spaces	6
4	C^*-Algebras and Functional Calculus	6
5	Spectral Decomposition	7
6	Abelian Harmonic Analysis	8
6.1	Fourier series and Fourier transform	8
7	Lattices, Poisson Summation, and Diffraction	9
8	Groups, Representations, and Abelian Duality	9
9	Compact Harmonic Analysis: Peter–Weyl, Characters, and Spin	11
10	Semisimple Harmonic Analysis: Harish–Chandra Theory	12
11	Unitary Coefficient Spaces and the Fourier Algebra	14
11.1	Unitary coefficient spaces	14
11.2	The Fourier algebra	15
12	Dual Algebras and Noncommutative Harmonic Analysis	15
13	Fourier–Wigner Analysis	16
14	The Heisenberg Group and Weyl Quantization	17
14.1	Central extension	17
14.2	Stone–von Neumann theorem	17
14.3	Weyl quantization	17
14.4	Heisenberg Plancherel theory	18
15	Theta Functions and Modular Forms	19
15.1	Lattice theta series and Poisson duality	19
15.2	Modular forms, theta functions, and eta products	19
15.3	Eisenstein series and arithmetic applications	20

16 Automorphic Harmonic Analysis and Arithmetic Langlands	21
16.1 Places, local fields, and adèles	21
16.2 Automorphic representations and the trace formula	22
16.3 Modular eigenforms as automorphic representations	23
16.4 Satake parameters and the dual group	23
16.5 Galois representations and matching L -functions	24
16.6 The Langlands correspondence and its status	24
 II Deformation	 24
17 Differential Geometry	25
17.1 Bundles, sections, and geometric structures	25
17.2 Differential forms, cohomology, and Hodge theory	25
17.3 Connections, curvature, and gauge fields	26
18 Symplectic Geometry	27
18.1 Projective representations and multipliers	27
18.2 Alternating commutator form	27
18.3 Linear symplectic geometry	28
18.4 Symplectic manifolds and Hamiltonian reduction	28
19 Classical and Quantum Dynamics	29
20 Geometric Quantization	30
20.1 Prequantization and polarization	30
20.2 Equivariant quantization and reduction	30
21 Deformation Quantization	31
22 Integrable Systems and the Yang–Baxter Equation	31
23 Rational Hypergeometric Functions	32
24 Trigonometric/Basic Hypergeometric Deformation	33
25 Poisson–Lie Groups and Quantum Groups	34
25.1 Poisson–Lie groups and Lie bialgebras	34
25.2 Drinfeld–Jimbo quantization	35
26 Compact Quantum Groups	35
27 Hecke Algebras and Quantum Schur–Weyl Duality	38
28 DAHA and Macdonald Theory	40
28.1 Double affine Hecke algebras	40
28.2 Macdonald theory	41
29 Quantum Spherical Functions and qKZ Transport	42
30 Elliptic Hypergeometric Functions and Ruijsenaars Theory	43
31 Elliptic Star–Triangle and Yang–Baxter Identities	45
32 Root-of-Unity Degeneration and Tensor Categories	46

III	Counting	47
33	Partitions and Generating Functions	47
33.1	Partition generating functions and finite boxes	47
33.2	Rogers–Ramanujan identities and exclusion rules	48
34	Characters and Exact State Counting	48
34.1	Virasoro representations and character formulas	48
34.2	Modular invariance and exact state counting	49
34.3	Integrable lattice models	50
35	Probability Spaces and Convergence	50
36	Random Matrices and Statistical Universality	51
36.1	Correlation functions and point processes	53
36.2	Determinantal/Pfaffian processes and Fredholm determinants	54
36.3	Orthogonal-polynomial ensembles	54
36.4	Christoffel–Darboux and integrable kernels	55
36.5	Bulk, soft-edge, and hard-edge universality	55
37	Integrable Probability and KPZ Universality	56
37.1	RSK and last-passage percolation	56
37.2	Schur and Macdonald processes	56
37.3	Dyson Brownian motion and KPZ-type dynamics	57
38	Free Probability and Asymptotic Freeness	59
38.1	Free independence and additive convolution	59
38.2	Random matrices and asymptotic freeness	60

Part I

Symmetry

1 Linear Spaces, Tensor Products, and Duality

Definition 1.1 (Linear spaces and maps). Let k be $\mathbb{R}$ or $\mathbb{C}$. A *linear space* over k is an abelian group V with scalar multiplication satisfying

$$a(v + w) = av + aw, \quad (a + b)v = av + bv, \quad (ab)v = a(bv), \quad 1v = v.$$

A map $T : V \rightarrow W$ is *linear* when $T(av + bw) = aT(v) + bT(w)$. Its kernel and image are

$$\ker T = \{v : T(v) = 0\}, \quad \operatorname{im} T = \{T(v) : v \in V\}.$$

For $T : V \rightarrow V$, a scalar λ is an *eigenvalue* when $\ker(T - \lambda I) \neq 0$; a nonzero vector in this kernel is an *eigenvector*.

Definition 1.2 (Dual spaces and tensor products). The *dual space* of V is $V^\vee = \operatorname{Hom}_k(V, k)$. Its canonical evaluation pairing is

$$\operatorname{ev}_V : V^\vee \times V \longrightarrow k, \quad (\varphi, v) \longmapsto \varphi(v).$$

The algebraic transpose of $T : V \rightarrow W$ is $T^\vee : W^\vee \rightarrow V^\vee$, defined by $T^\vee(\varphi) = \varphi \circ T$.

The *tensor product* $V \otimes W$ is characterized by a bilinear map $(v, w) \mapsto v \otimes w$ with the following universal property: every bilinear map $b : V \times W \rightarrow X$ factors uniquely as

$$V \times W \longrightarrow V \otimes W \xrightarrow{\tilde{b}} X.$$

The evaluation pairing induces a linear map $\operatorname{ev}_V : V^\vee \otimes V \rightarrow k$.

Definition 1.3 (Dualizability and finite dimensionality). A linear space V is *dualizable* when there is a coevaluation map

$$\operatorname{coev}_V : k \longrightarrow V \otimes V^\vee$$

satisfying the two snake identities

$$(\operatorname{id}_V \otimes \operatorname{ev}_V) \circ (\operatorname{coev}_V \otimes \operatorname{id}_V) = \operatorname{id}_V,$$

$$(\operatorname{ev}_V \otimes \operatorname{id}_{V^\vee}) \circ (\operatorname{id}_{V^\vee} \otimes \operatorname{coev}_V) = \operatorname{id}_{V^\vee}.$$

A linear space is *finite-dimensional* if it is dualizable.

Definition 1.4 (Categorical trace and dimension). Let V be dualizable, let $s_{V, V^\vee} : V \otimes V^\vee \rightarrow V^\vee \otimes V$ exchange the factors, and let $T \in \operatorname{End}(V)$. The *categorical trace* is the scalar

$$\operatorname{tr}(T) = \operatorname{ev}_V \circ s_{V, V^\vee} \circ (T \otimes \operatorname{id}_{V^\vee}) \circ \operatorname{coev}_V, \quad \dim V = \operatorname{tr}(\operatorname{id}_V).$$

Exercise 1.1 (Dualizability recovers ordinary finite dimension). Construct $V \otimes W$ as a quotient of the free linear space on $V \times W$ and prove its universal property. If V has basis $e_1, \dots, e_n$ with dual basis $e^1, \dots, e^n$, show that

$$\operatorname{coev}_V(1) = \sum_{j=1}^n e_j \otimes e^j$$

is independent of the chosen basis and satisfies the snake identities. Conversely, write any proposed coevaluation as $\text{coev}_V(1) = \sum_{j=1}^r v_j \otimes \varphi_j$ and use a snake identity to prove that the v_j span V . Conclude that dualizability is equivalent to ordinary finite dimensionality. Finally prove

$$\text{tr}(T) = \sum_{j=1}^n e^j(Te_j), \quad \text{tr}(ST) = \text{tr}(TS), \quad \dim V = n.$$

2 Hilbert Spaces and Trace Ideals

Definition 2.1 (Hilbert spaces and orthogonality). An *inner product* on a complex linear space is conjugate-linear in its first argument, linear in its second, positive definite, and satisfies $\langle x, y \rangle = \overline{\langle y, x \rangle}$. Its norm is $\|x\| = \sqrt{\langle x, x \rangle}$. A *Hilbert space* is an inner-product space complete for this norm.

Vectors are *orthogonal* when their inner product is zero. For a subspace $M \subseteq H$, define

$$M^\perp = \{x : \langle m, x \rangle = 0 \text{ for every } m \in M\}.$$

Definition 2.2 (Bounded operators and adjoints). A linear map $T : H \rightarrow K$ between Hilbert spaces is *bounded* when

$$\|T\| = \sup_{\|x\| \leq 1} \|Tx\| < \infty.$$

An *adjoint* of T is an operator $T^* : K \rightarrow H$ satisfying

$$\langle x, Ty \rangle = \langle T^*x, y \rangle.$$

An operator is self-adjoint when $T = T^*$, unitary when $T^*T = TT^* = I$, and normal when $T^*T = TT^*$. An *orthogonal projection* is an operator P satisfying $P = P^* = P^2$.

Definition 2.3 (Trace and Hilbert–Schmidt classes). For $x, y \in H$, let

$$\theta_{x,y}(\xi) = \langle y, \xi \rangle x.$$

The algebra $\mathcal{K}(H)$ of *compact operators* is the norm closure of the span of the operators $\theta_{x,y}$. The *trace class* $S_1(H)$ consists of operators admitting a representation

$$A = \sum_{j=1}^{\infty} \theta_{x_j, y_j}, \quad \sum_{j=1}^{\infty} \|x_j\| \|y_j\| < \infty,$$

where the series converges in operator norm, with norm the infimum of the displayed sums. Define

$$\text{Tr}(A) = \sum_{j=1}^{\infty} \langle y_j, x_j \rangle.$$

The *Hilbert–Schmidt class* $S_2(H)$ consists of operators satisfying

$$\|A\|_2^2 = \sum_j \|Ae_j\|^2 < \infty$$

for an orthonormal basis (e_j) of H .

Exercise 2.1 (Hilbert-space foundations). Prove the Cauchy–Schwarz inequality, the projection theorem, and the Riesz representation theorem. Deduce the existence of adjoints and prove

$$\|T\|^2 = \|T^*T\|, \quad (\text{im } T)^\perp = \ker T^*.$$

For every closed subspace $M \subseteq H$, prove $H = M \oplus M^\perp$ and construct the orthogonal projection $P_M : H \rightarrow M$. Show that an orthonormal family (e_j) is complete exactly when $\|x\|^2 = \sum_j |\langle e_j, x \rangle|^2$ for every $x \in H$.

Exercise 2.2 (Trace ideals). Prove that the definitions of $S_1(H)$, Tr , and $S_2(H)$ are independent of the displayed decompositions and orthonormal basis. Prove that $\langle A, B \rangle_{S_2} = \text{Tr}(A^*B)$ makes $S_2(H)$ a Hilbert space and that the pairing $(A, T) \mapsto \text{Tr}(AT)$ identifies $S_1(H)^*$ isometrically with $B(H)$.

3 Measure, Integration, and L^p Spaces

Definition 3.1 (Measure spaces and Haar measure). A σ -algebra $\mathcal{F}$ on a set X contains $\emptyset$ and is closed under complements and countable unions. A *measure* is a countably additive map $\mu : \mathcal{F} \rightarrow [0, \infty]$ with $\mu(\emptyset) = 0$, and $(X, \mathcal{F}, \mu)$ is a *measure space*. A map is *measurable* when inverse images of measurable sets are measurable. For $1 \leq p < \infty$, define

$$L^p(X, \mu) = \left\{ f : \int_X |f|^p d\mu < \infty \right\} / \text{equality a.e.}$$

with its usual norm, and let $L^\infty(X, \mu)$ consist of essentially bounded measurable functions.

A *topological group* is a group with a Hausdorff topology for which multiplication and inversion are continuous. It is *locally compact* when every point has a compact neighborhood. A *Radon measure* is a Borel measure finite on compact sets, outer regular on Borel sets, and inner regular on open sets. A *left Haar measure* on a locally compact group is a nonzero left-invariant Radon measure, so $\mu(gE) = \mu(E)$. Its *modular function* is determined by

$$\mu(Eg) = \Delta_G(g)^{-1} \mu(E).$$

Write $C_b(G)$, $C_0(G)$, and $C_c(G)$ for the continuous functions that are bounded, vanish at infinity, and have compact support, respectively.

4 C^* -Algebras and Functional Calculus

Definition 4.1 (C^* -algebras, spectrum, and positivity). A C^* -algebra is a complete normed complex algebra A with a conjugate-linear involution satisfying

$$(ab)^* = b^*a^*, \quad \|ab\| \leq \|a\|\|b\|, \quad \|a^*a\| = \|a\|^2.$$

For a unital C^* -algebra, define

$$\text{spec}_A(a) = \{\lambda \in \mathbb{C} : a - \lambda 1 \text{ is not invertible}\}.$$

An element is *positive*, written $a \geq 0$, when $a = b^*b$ for some $b \in A$. It is *self-adjoint* when $a = a^*$ and *normal* when $a^*a = aa^*$.

Definition 4.2 (Gelfand spectrum). For a commutative C^* -algebra A , let $\widehat{A}$ be the space of nonzero homomorphisms $\chi : A \rightarrow \mathbb{C}$ with its weak- $*$ topology. The *Gelfand transform* is

$$\Gamma : A \longrightarrow C_0(\widehat{A}), \quad \Gamma(a)(\chi) = \chi(a).$$

Exercise 4.1 (Spectrum, Gelfand duality, and functional calculus). Prove that $\text{spec}_A(a)$ is nonempty and compact and that

$$r(a) = \lim_{n \rightarrow \infty} \|a^n\|^{1/n}.$$

For normal a , prove $r(a) = \|a\|$. Prove that positive elements have unique positive square roots and that $a \geq 0$ if and only if $a = a^*$ and $\text{spec}_A(a) \subseteq [0, \infty)$. Prove that the Gelfand transform is an isometric $*$ -isomorphism. For every normal element a of a unital C^* -algebra, prove that there is a unique unital isometric $*$ -homomorphism

$$C(\text{spec}(a)) \longrightarrow C^*(1, a), \quad f \longmapsto f(a),$$

sending the coordinate function to a , and prove the continuous spectral mapping theorem.

5 Spectral Decomposition

Definition 5.1 (Projection-valued measures). A *projection-valued measure* on a measurable space X is a map E from measurable subsets of X to orthogonal projections on H satisfying

$$E(\emptyset) = 0, \quad E(X) = I, \quad E\left(\bigsqcup_{j=1}^{\infty} B_j\right) = \sum_{j=1}^{\infty} E(B_j)$$

for pairwise disjoint measurable sets, where the sum converges in the strong operator topology. The spectral integral $\int_X f dE$ for bounded measurable f is defined by uniform approximation from simple functions. For a topological space X , the measure is *regular* when every scalar measure $B \mapsto \langle x, E(B)y \rangle$ is a regular Borel measure.

Exercise 5.1 (Representations of $C_0(X)$). Let X be locally compact Hausdorff. A representation $\pi : C_0(X) \rightarrow B(H)$ is *nondegenerate* when $\overline{\pi(C_0(X))H} = H$. Prove that every nondegenerate $*$ -representation has a unique regular projection-valued measure E_π satisfying

$$\pi(f) = \int_X f(x) dE_\pi(x), \quad f \in C_0(X).$$

Exercise 5.2 (Bounded normal spectral theorem). For a normal operator $T \in B(H)$, apply the preceding representation theorem to the continuous functional calculus $C(\text{spec}(T)) \rightarrow B(H)$ and prove that there is a unique projection-valued measure E_T on $\text{spec}(T)$ satisfying

$$T = \int_{\text{spec}(T)} z dE_T(z), \quad f(T) = \int_{\text{spec}(T)} f(z) dE_T(z).$$

Deduce that a normal operator on a finite-dimensional Hilbert space is unitarily diagonalizable.

Exercise 5.3 (Compact spectral theorem). Prove that every trace-class operator is compact. For a compact normal operator K , prove that every nonzero spectral value is an eigenvalue of finite multiplicity, that the nonzero eigenvalues can accumulate only at zero, and that

$$K = \sum_j \lambda_j P_j$$

in operator norm for pairwise orthogonal finite-rank spectral projections P_j . For $K \geq 0$ in $S_1(H)$, prove

$$K = \sum_j \lambda_j \theta_{e_j, e_j}, \quad \lambda_j \geq 0, \quad \sum_j \lambda_j = \text{Tr } K.$$

Definition 5.2 (Unbounded self-adjoint operators). An unbounded operator is a linear map $A : \mathcal{D}(A) \rightarrow H$ on a dense domain. Its adjoint has domain

$$\mathcal{D}(A^*) = \{y \in H : x \mapsto \langle y, Ax \rangle \text{ is bounded on } \mathcal{D}(A)\}.$$

For $y \in \mathcal{D}(A^*)$, the vector A^*y is determined by $\langle A^*y, x \rangle = \langle y, Ax \rangle$ for $x \in \mathcal{D}(A)$. The operator is *self-adjoint* when $A = A^*$, including equality of domains, and *closed* when its graph is closed in $H \oplus H$.

Exercise 5.4 (Unbounded spectral theorem and Stone's theorem). For a self-adjoint operator A , prove that $A \pm iI$ are bijective from $\mathcal{D}(A)$ to H with bounded inverses and that the Cayley transform

$$C_A = (A - iI)(A + iI)^{-1}$$

is unitary with $\ker(I - C_A) = 0$. Apply the bounded normal spectral theorem to C_A and the inverse Cayley map to prove that there is a unique projection-valued measure E_A on $\mathbb{R}$ satisfying

$$A = \int_{\mathbb{R}} \lambda dE_A(\lambda), \quad \mathcal{D}(A) = \left\{ x : \int_{\mathbb{R}} \lambda^2 d\langle x, E_A(\lambda)x \rangle < \infty \right\}.$$

Use E_A to prove that e^{-itA} is a strongly continuous unitary group and that every strongly continuous one-parameter unitary group has a unique self-adjoint generator.

Exercise 5.5 (Spectral lines and time evolution). If $A\psi_n = \lambda_n\psi_n$ and an observable B has matrix elements $B_{mn} = \langle \psi_m, B\psi_n \rangle$, prove

$$\langle \psi_m, e^{itA/\hbar} B e^{-itA/\hbar} \psi_n \rangle = e^{it(\lambda_m - \lambda_n)/\hbar} B_{mn}.$$

Extend the calculation from pure point spectrum to the projection-valued spectral measure.

6 Abelian Harmonic Analysis

6.1 Fourier series and Fourier transform

Definition 6.1 (Fourier series). With normalized Haar measure on $\mathbb{T} = \mathbb{R}/2\pi\mathbb{Z}$, define

$$\hat{f}(k) = \int_{\mathbb{T}} f(x) e^{-ikx} dx, \quad k \in \mathbb{Z}.$$

The *Fourier series* of f is $\sum_{k \in \mathbb{Z}} \hat{f}(k) e^{ikx}$.

Definition 6.2 (Fourier transform). For $f \in \mathcal{S}(\mathbb{R}^n)$, the semiclassical Fourier transform is

$$(\mathcal{F}_\hbar f)(p) = (2\pi\hbar)^{-n/2} \int_{\mathbb{R}^n} e^{-ip \cdot x/\hbar} f(x) dx.$$

Exercise 6.1 (Fourier inversion and Plancherel). Prove that $(e^{ikx})_{k \in \mathbb{Z}}$ is an orthonormal basis of $L^2(\mathbb{T})$ and derive

$$f = \sum_{k \in \mathbb{Z}} \hat{f}(k) e^{ikx}, \quad \|f\|_2^2 = \sum_{k \in \mathbb{Z}} |\hat{f}(k)|^2.$$

Prove Fourier inversion on $\mathcal{S}(\mathbb{R}^n)$ and prove that $\mathcal{F}_\hbar$ extends uniquely to a unitary operator on $L^2(\mathbb{R}^n)$. Show that convolution becomes multiplication and differentiation becomes multiplication by frequency.

7 Lattices, Poisson Summation, and Diffraction

Definition 7.1 (Lattices and reciprocal lattices). A *lattice* in $\mathbb{R}^n$ is a subgroup $\Gamma = A\mathbb{Z}^n$ for some invertible linear map $A : \mathbb{R}^n \rightarrow \mathbb{R}^n$. Its covolume is $\text{vol}(\Gamma) = |\det A|$, and a fundamental cell is $F = A[0, 1)^n$. Its *reciprocal lattice* is the annihilator

$$\Gamma^* = \{\xi \in \mathbb{R}^n : \xi \cdot \gamma \in 2\pi\mathbb{Z} \text{ for every } \gamma \in \Gamma\}.$$

Definition 7.2 (Periodic summation and structure factors). For $f \in \mathcal{S}(\mathbb{R}^n)$, its Γ -*periodization* is

$$P_\Gamma f(x) = \sum_{\gamma \in \Gamma} f(x + \gamma).$$

For a finite motif $B \subset \mathbb{R}^n$ with scattering weights c_b , its *structure factor* is

$$S_B(q) = \sum_{b \in B} c_b e^{-iq \cdot b}.$$

Exercise 7.1 (Poisson summation). Prove that $P_\Gamma f$ converges with all derivatives and compute its Fourier coefficients. With the Fourier-transform normalization above, prove

$$P_\Gamma f(x) = \frac{(2\pi)^{n/2}}{\text{vol}(\Gamma)} \sum_{\xi \in \Gamma^*} \mathcal{F}f(\xi) e^{i\xi \cdot x}.$$

Set $x = 0$ to obtain the Poisson summation formula. Derive the Jacobi theta transformation by applying it to a Gaussian.

Exercise 7.2 (Crystal diffraction and Bragg peaks). Let Γ_N be a finite rectangular portion of Γ . In the single-scattering approximation, prove that the amplitude at momentum transfer q factors as

$$\mathcal{A}_N(q) = S_B(q) \sum_{\gamma \in \Gamma_N} e^{-iq \cdot \gamma}.$$

Use geometric sums and Poisson summation to show that the normalized intensity concentrates at $q \in \Gamma^*$ as the crystal grows. For elastic scattering with incident and outgoing wavevectors of magnitude $2\pi/\lambda$, derive the Laue condition and, for planes spaced by d , the Bragg relation

$$2d \sin \theta = m\lambda.$$

Show how zeros of S_B produce missing reflections. Compare the peak condition with W. H. Bragg and W. L. Bragg, *Proceedings of the Royal Society A* **88** (1913).

8 Groups, Representations, and Abelian Duality

Definition 8.1 (Groups, homomorphisms, and quotients). A *group* is a set G with an associative multiplication, an identity, and inverses. A *homomorphism* $\phi : G \rightarrow H$ preserves multiplication. A subgroup $N \leq G$ is *normal* if $gNg^{-1} = N$ for every $g \in G$; the *quotient group* G/N consists of cosets with induced multiplication.

Definition 8.2 (Actions, orbits, and stabilizers). A *left action* of G on X is a homomorphism $G \rightarrow \text{Bij}(X)$. The *orbit* and *stabilizer* of $x \in X$ are $Gx = \{gx : g \in G\}$ and $G_x = \{g : gx = x\}$.

Definition 8.3 (Representations and intertwiners). A *representation* of G on V is a homomorphism $\rho : G \rightarrow \text{GL}(V)$. An *intertwiner* $T : (V, \rho) \rightarrow (W, \sigma)$ satisfies $T\rho(g) = \sigma(g)T$. A subspace is *invariant* if it is preserved by every $\rho(g)$; a nonzero representation is *irreducible* if it has no proper nonzero invariant subspace. Direct sums and tensor products carry the actions $\rho \oplus \sigma$ and $g \mapsto \rho(g) \otimes \sigma(g)$.

Exercise 8.1 (Maschke's theorem). Let G be finite and let the characteristic of k not divide $|G|$. Average an arbitrary projection over G to construct an invariant complement to every invariant subspace. Deduce that every finite-dimensional representation is a direct sum of irreducible representations.

Exercise 8.2 (Schur's lemma). Prove that a nonzero intertwiner between irreducible complex finite-dimensional representations is an isomorphism and that every endomorphism of an irreducible representation is scalar. Deduce uniqueness of irreducible multiplicities in a decomposition.

Definition 8.4 (Characters). The *character* of a finite-dimensional representation V is the class function $\chi_V(g) = \text{Tr}(\rho_V(g))$. For class functions on a finite group, define

$$\langle f, h \rangle_G = \frac{1}{|G|} \sum_{g \in G} \overline{f(g)} h(g).$$

Exercise 8.3 (Character orthogonality and decomposition). Use averaging of linear maps and Schur's lemma to prove orthonormality of the irreducible characters. Prove that they form a complete orthonormal family in the class functions and that $\langle \chi_U, \chi_V \rangle_G = \dim \text{Hom}_G(U, V)$. Deduce the multiplicity formula and the character table constraints.

Definition 8.5 (Topological spaces). A *topology* on X is a family of subsets containing $\emptyset, X$ and closed under arbitrary unions and finite intersections. A map is *continuous* when inverse images of open sets are open. A space is *Hausdorff* when distinct points have disjoint neighborhoods, *second countable* when its topology is generated by a countable family of open sets, *compact* when every open cover has a finite subcover, and *locally compact* when every point has a neighborhood with compact closure. The *Borel sets* form the σ -algebra generated by the open sets. A Borel measure is *regular* when it is inner regular on open sets and outer regular on Borel sets.

Definition 8.6 (Topological and smooth manifolds). A *topological manifold* of dimension n is a Hausdorff, second-countable space locally homeomorphic to $\mathbb{R}^n$. A *smooth manifold* is a topological manifold with a maximal atlas whose transition maps are smooth. A map of smooth manifolds is *smooth* when its chart representatives are smooth; it is a *diffeomorphism* when it has a smooth inverse.

Definition 8.7 (Tangent and cotangent spaces). The *tangent space* $T_p M$ is the vector space of derivations $v : C^\infty(M) \rightarrow \mathbb{R}$ at p , satisfying $v(fg) = f(p)v(g) + g(p)v(f)$. The *cotangent space* is $T_p^* M = (T_p M)^*$. The differential of $F : M \rightarrow N$ is $dF_p(v)(f) = v(f \circ F)$. The disjoint union $TM = \coprod_{p \in M} T_p M$ has a canonical smooth vector-bundle structure: if $x = (x^1, \dots, x^n)$ is a coordinate chart on U , its bundle chart is

$$TM|_U \longrightarrow x(U) \times \mathbb{R}^n, \quad v \in T_p M \longmapsto (x(p), v(x^1), \dots, v(x^n)).$$

The cotangent bundle T^*M has the dual bundle charts.

Definition 8.8 (Lie groups). A *Lie group* is a topological group with a smooth manifold structure for which multiplication and inversion are smooth. Its *Lie algebra* is $\mathfrak{g} = T_e G$ with bracket induced by left-invariant vector fields. A *one-parameter subgroup* is a smooth homomorphism $\mathbb{R} \rightarrow G$. The *exponential map* $\exp : \mathfrak{g} \rightarrow G$ sends X to the value at 1 of the one-parameter subgroup with initial velocity X .

Exercise 8.4 (One-parameter subgroups and the exponential map). Prove existence and uniqueness of the one-parameter subgroup generated by $X \in \mathfrak{g}$. For a closed matrix group, prove that it is $t \mapsto e^{tX}$, identify the commutator bracket, and compute the first nontrivial terms of the Baker–Campbell–Hausdorff formula.

Definition 8.9 (Infinitesimal representations). For a finite-dimensional smooth representation $\rho : G \rightarrow \mathrm{GL}(V)$, its *infinitesimal representation* is

$$d\rho : \mathfrak{g} \rightarrow \mathrm{End}(V), \quad d\rho(X) = \left. \frac{d}{dt} \right|_0 \rho(\exp tX).$$

Definition 8.10 (Convolution). Let G be a locally compact group with left Haar measure μ . For $f, g \in C_c(G)$, define

$$(f * g)(x) = \int_G f(y)g(y^{-1}x) d\mu(y).$$

Exercise 8.5 (Haar measure and the regular representation). Identify Haar measure on $\mathbb{R}^n$, $\mathbb{Z}^n$, and $\mathbb{T}^n$. Construct normalized Haar measure on a compact Lie group from a left-invariant metric. Prove associativity of convolution, the L^1 convolution inequality, and $\|f * g\|_2 \leq \|f\|_1 \|g\|_2$. Prove that $g \mapsto \lambda_g$ is a strongly continuous unitary representation. For a finite-dimensional representation of a compact group, average an inner product over G to obtain an invariant one.

Definition 8.11 (Pontryagin dual). For a locally compact abelian group G , its *Pontryagin dual* $\widehat{G}$ is the group of continuous characters $\chi : G \rightarrow U(1)$, with pointwise multiplication and the compact-open topology. For $f \in L^1(G)$, define

$$\widehat{f}(\chi) = \int_G f(x) \overline{\chi(x)} d\mu(x).$$

The *double-dual map* sends $x \in G$ to the character $\chi \mapsto \chi(x)$ of $\widehat{G}$.

Exercise 8.6 (Abelian Fourier duality). Identify

$$\widehat{\mathbb{R}^n} \cong \mathbb{R}^n, \quad \widehat{\mathbb{Z}^n} \cong \mathbb{T}^n, \quad \widehat{\mathbb{T}^n} \cong \mathbb{Z}^n,$$

and verify the double-dual map in each case. Show that the group Fourier transform recovers Fourier transforms on $\mathbb{R}^n$ and Fourier series on $\mathbb{T}^n$. For a closed subgroup $H \leq G$, define its annihilator $H^\perp \subseteq \widehat{G}$ and prove $\widehat{G/H} \cong H^\perp$.

9 Compact Harmonic Analysis: Peter–Weyl, Characters, and Spin

Exercise 9.1 (Peter–Weyl theorem). Let G be compact with normalized Haar measure. Prove that matrix coefficients of irreducible unitary representations span a dense subspace of $C(G)$ and $L^2(G)$ and that

$$L^2(G) \cong \widehat{\bigoplus_{\pi \in \widehat{G}} H_\pi \otimes H_\pi^*}.$$

For $f \in L^2(G) \subseteq L^1(G)$, define $\widehat{f}(\pi) = \int_G f(g) \pi(g)^* dg$ and prove

$$f = \sum_{\pi \in \widehat{G}} (\dim H_\pi) \mathrm{Tr}(\widehat{f}(\pi) \pi(\cdot)), \quad \|f\|_2^2 = \sum_{\pi \in \widehat{G}} (\dim H_\pi) \|\widehat{f}(\pi)\|_{\mathrm{HS}}^2$$

, where the first sum converges in $L^2(G)$. For $f \in C(G)$, construct finite matrix-coefficient approximate identities and prove uniform convergence of the corresponding summability means to f .

Definition 9.1 (Rotations and spin). Define

$$\begin{aligned} SO(3) &= \{R \in \text{End}(\mathbb{R}^3) : R^*R = \text{id}, \det R = 1\}, \\ SU(2) &= \{U \in \text{End}(\mathbb{C}^2) : U^*U = \text{id}, \det U = 1\}. \end{aligned}$$

An *angular-momentum representation* is a unitary representation of $SU(2)$; with Planck constant $\hbar$, its infinitesimal generators J_1, J_2, J_3 are normalized by $[J_i, J_j] = i\hbar \varepsilon_{ijk} J_k$.

Exercise 9.2 ($SU(2)$ double-covers $SO(3)$). Let $SU(2)$ act by conjugation on the real space of traceless Hermitian 2×2 operators. Prove that the resulting homomorphism $SU(2) \rightarrow SO(3)$ is surjective with kernel $\{\pm \text{id}\}$. Deduce the effect of a 2π rotation in integer- and half-integer-spin representations and the 4π restoration observed by neutron interferometry.

Exercise 9.3 (Irreducible representations of $SU(2)$). Using raising and lowering operators, prove that every finite-dimensional irreducible smooth complex representation is indexed by $j \in \{0, \frac{1}{2}, 1, \dots\}$, has dimension $2j + 1$, and has J_3 -eigenvalues $m\hbar$ for $m = -j, -j + 1, \dots, j$. Compute the spectrum of $J^2 = J_1^2 + J_2^2 + J_3^2$. For $j = \frac{1}{2}$, relate the two J_3 outcomes to the two Stern–Gerlach beams.

Exercise 9.4 (Clebsch–Gordan decomposition). Prove

$$V_{j_1} \otimes V_{j_2} \cong \bigoplus_{j=|j_1-j_2|}^{j_1+j_2} V_j.$$

Construct coupled angular-momentum eigenstates from highest-weight vectors and compute the singlet and triplet decomposition of $V_{1/2} \otimes V_{1/2}$.

Exercise 9.5 (Spherical harmonics and atomic orbitals). Identify S^2 with $SO(3)/SO(2)$ and $L^2(S^2)$ with the right $SO(2)$ -invariant subspace of $L^2(SO(3))$. Use Peter–Weyl to obtain

$$L^2(S^2) \cong \widehat{\bigoplus_{\ell=0}^{\infty} V_{\ell}}$$

and identify a basis of V_{ℓ} with $Y_{\ell m} \propto D_{m0}^{\ell}$. For the unit round sphere, set

$$\Delta_{S^2} = \frac{1}{\sin \theta} \partial_{\theta} (\sin \theta \partial_{\theta}) + \frac{1}{\sin^2 \theta} \partial_{\phi}^2.$$

Derive $\Delta_{S^2} Y_{\ell m} = -\ell(\ell + 1) Y_{\ell m}$ and completeness of the spherical harmonics. Separate the Coulomb Hamiltonian in spherical coordinates, identify its angular factor, and recover $\psi_{n\ell m}(r, \theta, \phi) = R_{n\ell}(r) Y_{\ell m}(\theta, \phi)$. Determine the angular nodes of the s , p , and d orbitals and relate them to symmetry-enforced zeros in photoelectron angular distributions in the plane-wave final-state approximation. Compare with Matsui et al., *Physical Review B* **72** (2005).

10 Semisimple Harmonic Analysis: Harish–Chandra Theory

Definition 10.1 (Cartan and Iwasawa decompositions). Let G be a connected real semisimple Lie group with finite center and let $\mathfrak{g}$ be its Lie algebra. A *Cartan involution* gives a vector space decomposition

$$\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$$

into its $+1$ and -1 eigenspaces; the subgroup K with Lie algebra $\mathfrak{k}$ is maximal compact, and G/K is a noncompact Riemannian symmetric space. Choose a maximal abelian subspace $\mathfrak{a} \subseteq \mathfrak{p}$, put $A = \exp \mathfrak{a}$, choose positive restricted roots, and let N be the subgroup generated by their root spaces. The *Iwasawa decomposition* is the diffeomorphism

$$K \times A \times N \longrightarrow G, \quad (k, a, n) \longmapsto kan.$$

If $\mathfrak{a}^+$ is a positive Weyl chamber, the *Cartan decomposition* is $G = K\overline{A^+}K$. Write $M = Z_K(A)$ and $W = N_K(A)/M$, and let ρ be half the sum of the positive restricted roots, counted with multiplicity.

Definition 10.2 (Spherical principal series and spherical transform). For $\lambda \in \mathfrak{a}^*$, the normalized *spherical principal series* is

$$\pi_\lambda = \text{Ind}_{MAN}^G(1_M \otimes e^{i\lambda} \otimes 1_N),$$

where normalized induction incorporates the modular factor determined by ρ . It contains a unit K -fixed vector v_λ , and its *zonal spherical function* is

$$\varphi_\lambda(g) = \langle v_\lambda, \pi_\lambda(g)v_\lambda \rangle.$$

It is K -bi-invariant, satisfies $\varphi_\lambda(e) = 1$, and is a joint eigenfunction of the G -invariant differential operators on G/K .

For $f \in C_c^\infty(K \backslash G/K)$, its *spherical transform* is

$$\tilde{f}(\lambda) = \int_G f(g) \varphi_{-\lambda}(g) dg.$$

Harish-Chandra's c -function is the coefficient governing the leading asymptotics of $\varphi_\lambda(\exp H)$ as $H \rightarrow \infty$ in a positive chamber.

Exercise 10.1 (Harish-Chandra transform on $\mathbb{H}^2$). For the sign convention in which the Laplace–Beltrami operator is

$$\Delta_{\mathbb{H}^2} = y^2(\partial_x^2 + \partial_y^2),$$

derive its radial part

$$\Delta_{\text{rad}} = \frac{d^2}{dr^2} + \coth r \frac{d}{dr}.$$

Show that the normalized radial eigenfunction satisfying

$$\Delta \varphi_\lambda = -(\lambda^2 + \tfrac{1}{4})\varphi_\lambda, \quad \varphi_\lambda(0) = 1,$$

is the conical Legendre function

$$\varphi_\lambda(r) = P_{-1/2+i\lambda}(\cosh r).$$

For a smooth compactly supported radial function, define

$$\tilde{f}(\lambda) = 2\pi \int_0^\infty f(r) \varphi_\lambda(r) \sinh r dr.$$

Prove directly from the differential equation that distinct real values of λ^2 give orthogonal generalized eigenfunctions after truncation to $[0, R]$ with a common self-adjoint boundary condition. Compare the large- r equation with a constant-coefficient wave equation and show that its oscillation parameter is continuous. Contrast this with the discrete Dirichlet spectrum in the finite hyperbolic-drum exercise below.

Exercise 10.2 (A finite hyperbolic drum). Discretize the Laplace–Beltrami operator by a real symmetric weighted graph Laplacian on a finite regular hyperbolic tiling with Dirichlet boundary. Prove that the finite operator has a discrete orthogonal eigenbasis, and explain why those eigenvalues approximate a bounded domain with a boundary rather than the continuous spectrum of all $\mathbb{H}^2$. Compare the predicted mode ordering and propagation along discrete geodesics with the circuit measurements in Lenggenhager et al., *Nature Communications* **13** (2022). State which observations support the effective hyperbolic metric and which do not by themselves establish the infinite-space Plancherel theorem.

Definition 10.3 (Harish-Chandra characters and discrete series). A *distribution* on a Lie group G is a continuous linear functional on $C_c^\infty(G)$ with its usual test-function topology. Let π be an irreducible admissible unitary representation of a connected real reductive group. Although the infinite-dimensional operator $\pi(g)$ usually has no trace, for $f \in C_c^\infty(G)$ the integrated operator

$$\pi(f) = \int_G f(g) \pi(g) dg$$

is trace class. The *Harish-Chandra character* is the conjugation-invariant distribution

$$\Theta_\pi(f) = \text{Tr}(\pi(f)).$$

An irreducible unitary representation belongs to the *discrete series* if it has a nonzero matrix coefficient square-integrable modulo the center.

Exercise 10.3 (From compact characters to the semisimple Plancherel formula). For a compact group, prove

$$\Theta_\pi(f) = \int_G f(g) \text{Tr}(\pi(g)) dg$$

and recover the ordinary character used in Peter–Weyl theory. Show that a unitary operator on an infinite-dimensional Hilbert space is not trace class, which explains the distributional definition above. Prove directly that Θ_π is invariant under conjugation.

For $SL_2(\mathbb{R})$, identify the K -types of the spherical principal series and show that they contain a K -fixed vector. Contrast them with the holomorphic and antiholomorphic discrete series, whose nonzero one-sided K -types imply that they have no K -fixed vector. Deduce that spherical analysis on G/K sees the class-one principal series but not the discrete series. Compare the two families tabulated in Harish-Chandra, *Proceedings of the National Academy of Sciences* **43** (1957) for spherical functions and Harish-Chandra, *Transactions of the American Mathematical Society* **119** (1965) for invariant eigendistributions.

11 Unitary Coefficient Spaces and the Fourier Algebra

11.1 Unitary coefficient spaces

Definition 11.1 (Unitary representations and coefficient spaces). Let G be a locally compact group and let $\pi : G \rightarrow U(H_\pi)$ be a strongly continuous unitary representation. Strong continuity means that $s \mapsto \pi(s)x$ is continuous for every $x \in H_\pi$. Define

$$\Phi_\pi : S_1(H_\pi) \longrightarrow C_b(G), \quad \Phi_\pi(A)(s) = \text{Tr}(A\pi(s)).$$

The *coefficient space* of π is

$$A_\pi(G) = \Phi_\pi(S_1(H_\pi))$$

with the quotient norm from $S_1(H_\pi)/\ker \Phi_\pi$.

Exercise 11.1 (Coefficient-space factorization). Prove that $A_\pi(G)$ consists of the functions

$$u(s) = \sum_{j=1}^{\infty} \langle y_j, \pi(s)x_j \rangle, \quad \sum_{j=1}^{\infty} \|x_j\| \|y_j\| < \infty,$$

and that its norm is the infimum of the displayed sums. For unitary representations π and ρ , prove that $uv \in A_{\pi \otimes \rho}(G)$ for $u \in A_\pi(G)$ and $v \in A_\rho(G)$, with $\|uv\| \leq \|u\| \|v\|$.

11.2 The Fourier algebra

Definition 11.2 (Left regular representation and Fourier algebra). Let G be a locally compact group with left Haar measure. Its *left regular representation* on $L^2(G)$ is

$$(\lambda_s \xi)(t) = \xi(s^{-1}t).$$

The *Fourier algebra* is

$$A(G) = A_\lambda(G).$$

Definition 11.3 (Pontryagin dual). For a locally compact abelian group G , its *Pontryagin dual* is

$$\widehat{G} = \{\chi : G \rightarrow \mathbb{T} : \chi \text{ is a continuous homomorphism}\}$$

with pointwise multiplication and the compact-open topology.

Exercise 11.2 (Fell absorption and the Fourier algebra). Prove Fell's absorption principle

$$\lambda \otimes \pi \cong \lambda \otimes I_{H_\pi}$$

for every unitary representation π of G . Deduce that $A(G)$ is a contractive Banach algebra under pointwise multiplication and that $A(G) \subseteq C_0(G)$. For abelian G , prove the isometric algebra identification

$$A(G) \cong L^1(\widehat{G})$$

under the inverse Fourier transform.

12 Dual Algebras and Noncommutative Harmonic Analysis

Definition 12.1 (Operator topologies and commutants). The *weak operator topology* on $B(H)$ is generated by $T \mapsto |\langle x, Ty \rangle|$, and the *strong operator topology* is generated by $T \mapsto \|Tx\|$. For $S \subseteq B(H)$, define

$$S' = \{T \in B(H) : Ts = sT \text{ for every } s \in S\}, \quad S'' = (S')'.$$

A *von Neumann algebra* is a unital $*$ -subalgebra $M \subseteq B(H)$ satisfying $M = M''$.

Definition 12.2 (Coefficient von Neumann algebras). For a unitary representation $\pi : G \rightarrow U(H_\pi)$, define

$$VN_\pi(G) = \{\pi(s) : s \in G\}''.$$

The *group von Neumann algebra* is

$$VN(G) = VN_\lambda(G).$$

For concrete von Neumann algebras $M \subseteq B(H)$ and $N \subseteq B(K)$, define

$$M \overline{\otimes} N = (M \odot N)'' \subseteq B(H \otimes K).$$

Definition 12.3 (Normal functionals and maps). A positive linear functional $\varphi : M \rightarrow \mathbb{C}$ on a von Neumann algebra is *normal* when

$$a_i \uparrow a \implies \varphi(a_i) \uparrow \varphi(a)$$

for every bounded increasing net in M_+ . The linear span of the normal positive functionals is the *predual* M_* . The *ultraweak topology* is $\sigma(M, M_*)$. A bounded linear map between von Neumann algebras is *normal* when it is ultraweakly continuous.

Exercise 12.1 (Coefficient-space duality). Prove the bicommutant theorem. Prove the isometric identifications

$$A_\pi(G)^* \cong VN_\pi(G), \quad A(G)^* \cong VN(G),$$

with pairing induced by $(A, T) \mapsto \text{Tr}(AT)$. Prove that the induced weak-* topology on $VN_\pi(G)$ is its ultraweak operator topology and that its predual is $A_\pi(G)$. Prove $M \cong (M_*)^*$ isometrically for every von Neumann algebra M .

13 Fourier–Wigner Analysis

Definition 13.1 (Unitary systems and coefficient transforms). Let (X, μ) be a measure space and let $U : X \rightarrow U(H)$ be a weakly measurable map. The *coefficient transform* of the unitary system U is

$$\mathcal{V}_U : S_1(H) \longrightarrow L^\infty(X), \quad \mathcal{V}_U(A)(x) = \text{Tr}(AU_x).$$

Definition 13.2 (Weyl system and Fourier–Wigner transform). Let V be an n -dimensional real linear space with Lebesgue measure dq , and equip V^* with the dual measure dp for which $(2\pi)^{-n/2} \int_V e^{-ip(q)} f(q) dq$ is unitary on L^2 . For $q \in V$ and $p \in V^*$, define

$$(T_q \xi)(x) = \xi(x - q), \quad (M_p \xi)(x) = e^{ip(x)} \xi(x).$$

For $z = (q, p) \in V \oplus V^*$, define

$$W(z) = e^{-ip(q)/2} M_p T_q.$$

The *Fourier–Wigner transform* is

$$\mathcal{W} : S_1(L^2(V)) \longrightarrow C_b(V \oplus V^*), \quad \mathcal{W}(A)(z) = (2\pi)^{-n/2} \text{Tr}(AW(z)).$$

Exercise 13.1 (Weyl relations and the Moyal identity). Prove

$$M_p T_q = e^{ip(q)} T_q M_p, \quad W(z)^* = W(-z),$$

and

$$W(q, p)W(q', p') = \exp\left(\frac{i}{2}(p(q') - p'(q))\right) W(q + q', p + p').$$

For $A, B \in S_1(L^2(V)) \cap S_2(L^2(V))$, prove the Moyal identity

$$\int_{V \oplus V^*} \overline{\mathcal{W}(A)(z)} \mathcal{W}(B)(z) dz = \text{Tr}(A^* B).$$

Deduce that $\mathcal{W}$ extends uniquely to a unitary map

$$S_2(L^2(V)) \longrightarrow L^2(V \oplus V^*)$$

and, whenever $\mathcal{W}(A) \in L^1(V \oplus V^*)$, prove

$$A = (2\pi)^{-n/2} \int_{V \oplus V^*} \mathcal{W}(A)(z) W(z)^* dz$$

in the weak operator topology.

14 The Heisenberg Group and Weyl Quantization

14.1 Central extension

Definition 14.1 (Heisenberg group). For a symplectic space (E, ω) , its *Heisenberg group* is

$$\mathbb{H}(E, \omega) = E \times \mathbb{R}$$

with multiplication

$$(z, s)(z', s') = (z + z', s + s' + \frac{1}{2}\omega(z, z')).$$

Write $\mathbb{H}_n = \mathbb{H}(\mathbb{R}^n \oplus (\mathbb{R}^n)^*, \omega)$ for the canonical symplectic form on $\mathbb{R}^n \oplus (\mathbb{R}^n)^*$. For $E = V \oplus V^*$ and $\lambda \neq 0$, the *Schrödinger representation* on $L^2(V)$ is

$$(\pi_\lambda(q, p, s)\xi)(x) = e^{i\lambda(s+p(x-q/2))}\xi(x - q).$$

Exercise 14.1 (Central extensions and projective representations). Prove that $\mathbb{H}(E, \omega)$ is a locally compact group with center $\{0\} \times \mathbb{R}$ and that

$$\pi_\lambda(0, s) = e^{i\lambda s} I.$$

Prove that projective representations of E with multiplier $e^{i\lambda\omega/2}$ correspond to representations of $\mathbb{H}(E, \omega)$ with central character $e^{i\lambda s}$.

14.2 Stone–von Neumann theorem

Exercise 14.2 (Stone–von Neumann theorem). Prove that π_λ is irreducible. Prove that every strongly continuous irreducible unitary representation of $\mathbb{H}(V \oplus V^*, \omega)$ with central character $e^{i\lambda s}$, $\lambda \neq 0$, is unitarily equivalent to π_λ .

14.3 Weyl quantization

Definition 14.2 (Weyl quantization). For $a \in \mathcal{S}(\mathbb{R}^{2n})$, define $\text{Op}_h^W(a) : \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}(\mathbb{R}^n)$ by

$$(\text{Op}_h^W(a)\psi)(x) = \frac{1}{(2\pi\hbar)^n} \int_{\mathbb{R}^n \times \mathbb{R}^n} e^{i(x-y)\cdot p/\hbar} a\left(\frac{x+y}{2}, p\right) \psi(y) dy dp.$$

For polynomial symbols, use the same formula as an oscillatory integral on $\mathcal{S}(\mathbb{R}^n)$. For the coordinate functions q_j and p_j , set

$$Q_j = \text{Op}_h^W(q_j), \quad P_j = \text{Op}_h^W(p_j).$$

The canonical Poisson bracket on $\mathbb{R}^{2n}$ is

$$\{a, b\} = \sum_{j=1}^n \left(\frac{\partial a}{\partial q_j} \frac{\partial b}{\partial p_j} - \frac{\partial a}{\partial p_j} \frac{\partial b}{\partial q_j} \right).$$

Definition 14.3 (Wigner distributions). For $\psi, \varphi \in \mathcal{S}(\mathbb{R}^n)$, define

$$W_h(\psi, \varphi)(q, p) = \frac{1}{(2\pi\hbar)^n} \int_{\mathbb{R}^n} e^{-ip\cdot y/\hbar} \overline{\psi(q - y/2)} \varphi(q + y/2) dy.$$

Write $W_h[\psi] = W_h(\psi, \psi)$.

Exercise 14.3 (Weyl calculus and the Moyal product). Prove on $\mathcal{S}(\mathbb{R}^n)$ that

$$Q_j \psi(x) = x_j \psi(x), \quad P_j \psi(x) = -i\hbar \partial_j \psi(x), \quad [Q_j, P_k] = i\hbar \delta_{jk} I.$$

Derive the Weyl relations and recover the Schrödinger representation of the Heisenberg group. Prove

$$\text{Op}_\hbar^W(a)^* = \text{Op}_\hbar^W(\bar{a}), \quad \|\text{Op}_\hbar^W(a)\|_{\text{HS}} = (2\pi\hbar)^{-n/2} \|a\|_2,$$

and

$$\langle \psi, \text{Op}_\hbar^W(a) \varphi \rangle = \int_{\mathbb{R}^{2n}} a(q, p) W_\hbar(\psi, \varphi)(q, p) dq dp.$$

Prove that for $a, b \in \mathcal{S}(\mathbb{R}^{2n})$ there is a unique $a \star_\hbar b \in \mathcal{S}(\mathbb{R}^{2n})$ satisfying

$$\text{Op}_\hbar^W(a \star_\hbar b) = \text{Op}_\hbar^W(a) \text{Op}_\hbar^W(b).$$

Derive the oscillatory-integral formula for $\star_\hbar$ and prove

$$a \star_\hbar b = ab + \frac{i\hbar}{2} \{a, b\} + O(\hbar^2), \quad \frac{a \star_\hbar b - b \star_\hbar a}{i\hbar} = \{a, b\} + O(\hbar^2)$$

in the Schwartz-symbol topology.

Exercise 14.4 (Groenewold–van Hove obstruction). Prove that Weyl quantization carries Poisson brackets to $(i\hbar)^{-1}$ times commutators exactly for affine and quadratic polynomials. Quantize $q^2 p^2$ using

$$q^2 p^2 = \frac{1}{9} \{q^3, p^3\} = \frac{1}{3} \{q^2 p, q p^2\}$$

and prove that there is no linear quantization of the polynomial Poisson algebra satisfying

$$\mathcal{Q}(1) = I, \quad \mathcal{Q}(q) = Q, \quad \mathcal{Q}(p) = P, \quad \mathcal{Q}(q^m) = Q^m, \quad \mathcal{Q}(p^m) = P^m,$$

together with

$$\mathcal{Q}(\{f, g\}) = (i\hbar)^{-1} [\mathcal{Q}(f), \mathcal{Q}(g)]$$

on a common invariant domain.

14.4 Heisenberg Plancherel theory

Exercise 14.5 (Heisenberg Plancherel and inversion). For $V = \mathbb{R}^n$ and $f \in L^1(\mathbb{H}_n) \cap L^2(\mathbb{H}_n)$, use Haar measure $dq dp ds$ and the scalar Fourier transform $\widehat{h}(\lambda) = \int_{\mathbb{R}} h(s) e^{-i\lambda s} ds$, and define

$$\widehat{f}(\lambda) = \int_{\mathbb{H}_n} f(g) \pi_\lambda(g)^* dg.$$

Derive the Plancherel measure $(2\pi)^{-(n+1)} |\lambda|^n d\lambda$ and prove

$$\|f\|_2^2 = (2\pi)^{-(n+1)} \int_{\mathbb{R} \setminus \{0\}} \|\widehat{f}(\lambda)\|_{\text{HS}}^2 |\lambda|^n d\lambda.$$

Prove the inversion formula and the distributional orthogonality of Weyl operators. Prove that representations with trivial central character have zero Plancherel measure.

15 Theta Functions and Modular Forms

15.1 Lattice theta series and Poisson duality

Definition 15.1 (Integral lattices and theta series). A *Euclidean lattice* $L \subset \mathbb{R}^d$ is the integer span of a basis of $\mathbb{R}^d$. Its dual is

$$L^* = \{y \in \mathbb{R}^d : \langle x, y \rangle \in \mathbb{Z} \text{ for every } x \in L\}.$$

It is *integral* if $L \subseteq L^*$, *unimodular* if $L = L^*$, and *even* if $\|x\|^2 \in 2\mathbb{Z}$ for every $x \in L$. For τ in the upper half-plane and $q = e^{2\pi i\tau}$, its theta series is

$$\Theta_L(\tau) = \sum_{x \in L} q^{\|x\|^2/2}.$$

Its coefficient of q^n counts lattice vectors of squared length $2n$.

Exercise 15.1 (Poisson duality for lattice theta series). Apply Exercise 7.1 to a Gaussian and prove

$$\Theta_L(-1/\tau) = \frac{(-i\tau)^{d/2}}{\text{vol}(L)} \Theta_{L^*}(\tau),$$

using the branch positive on the positive imaginary axis. Show that evenness gives $\Theta_L(\tau+1) = \Theta_L(\tau)$. Interpret the first identity as momentum–winding duality for a compact free boson and as the reciprocal-lattice relation behind diffraction. Explain why a general lattice theta series transforms with its dual rather than being a scalar modular form by itself.

15.2 Modular forms, theta functions, and eta products

Definition 15.2 (The modular group and modular forms). The modular group $SL_2(\mathbb{Z})$ acts on the upper half-plane by

$$\gamma\tau = \frac{a\tau + b}{c\tau + d}, \quad \gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix},$$

and is generated by $S : \tau \mapsto -1/\tau$ and $T : \tau \mapsto \tau + 1$. A *modular form of integral weight k* for $SL_2(\mathbb{Z})$ is a holomorphic function f on the upper half-plane satisfying

$$f(\gamma\tau) = (c\tau + d)^k f(\tau)$$

and holomorphic at the cusp, meaning that its Fourier expansion $f(\tau) = \sum_{n \geq 0} a_n q^n$ has no negative powers. It is a *cusp form* if $a_0 = 0$.

Definition 15.3 (Jacobi theta functions and the Dedekind eta function). Set

$$\begin{aligned} \vartheta_2(\tau) &= \sum_{n \in \mathbb{Z}} q^{(n+1/2)^2/2}, \\ \vartheta_3(\tau) &= \sum_{n \in \mathbb{Z}} q^{n^2/2}, & \eta(\tau) &= q^{1/24}(q; q)_\infty. \\ \vartheta_4(\tau) &= \sum_{n \in \mathbb{Z}} (-1)^n q^{n^2/2}, \end{aligned}$$

These functions have weight 1/2 transformation laws with phases or permutations, rather than being integral-weight scalar modular forms for the full modular group. In particular,

$$\eta(\tau + 1) = e^{\pi i/12} \eta(\tau), \quad \eta(-1/\tau) = (-i\tau)^{1/2} \eta(\tau).$$

Exercise 15.2 (Theta transformations and the Jacobi triple product). Use Poisson summation on shifted and phase-twisted Gaussians to derive

$$\vartheta_2(-1/\tau) = (-i\tau)^{1/2}\vartheta_4(\tau), \quad \vartheta_3(-1/\tau) = (-i\tau)^{1/2}\vartheta_3(\tau), \quad \vartheta_4(-1/\tau) = (-i\tau)^{1/2}\vartheta_2(\tau).$$

Prove the T -transformation laws directly from the series. Establish the Jacobi triple product, first as an identity of formal Laurent series,

$$\sum_{n \in \mathbb{Z}} z^n q^{n^2/2} = \prod_{m \geq 1} (1 - q^m)(1 + zq^{m-1/2})(1 + z^{-1}q^{m-1/2}),$$

and use specializations to recover product forms for the theta functions. Explain how the two sides encode charge sectors and independent oscillator modes, respectively.

15.3 Eisenstein series and arithmetic applications

Definition 15.4 (Eisenstein series and the discriminant). Let $\sigma_r(n) = \sum_{d|n} d^r$. The normalized Eisenstein series of weights four and six are

$$E_4(\tau) = 1 + 240 \sum_{n \geq 1} \sigma_3(n)q^n, \quad E_6(\tau) = 1 - 504 \sum_{n \geq 1} \sigma_5(n)q^n.$$

The modular discriminant is the weight-twelve cusp form

$$\Delta(\tau) = \eta(\tau)^{24} = q \prod_{n \geq 1} (1 - q^n)^{24} = \frac{E_4(\tau)^3 - E_6(\tau)^2}{1728}.$$

Exercise 15.3 (Eisenstein series and the E_8 shell). Construct Eisenstein series by summing $(m\tau + n)^{-k}$ over nonzero lattice points for even $k > 2$, prove their modular transformation law, and obtain the displayed Fourier expansions for E_4 and E_6 . For a positive-definite even unimodular lattice, first use consistency of the S and T laws to prove that its dimension d is divisible by eight, and then show that its theta series is a modular form of weight $d/2$. Construct the E_8 lattice as

$$\{x \in \mathbb{Z}^8 : \sum_i x_i \equiv 0 \pmod{2}\} \cup \{x \in (\mathbb{Z} + \tfrac{1}{2})^8 : \sum_i x_i \equiv 0 \pmod{2}\}.$$

Enumerate its vectors of squared length two and obtain 240, matching the coefficient of q in E_4 .

Exercise 15.4 (Partition asymptotics from modular inversion). Put $q = e^{-t}$ and use the modular transformation of η to prove

$$P(e^{-t}) \sim \sqrt{\frac{t}{2\pi}} \exp\left(\frac{\pi^2}{6t}\right) \quad (t \downarrow 0).$$

Apply a saddle-point argument, or an appropriate Tauberian theorem with its hypotheses stated, to derive the Hardy–Ramanujan formula

$$p(n) \sim \frac{1}{4n\sqrt{3}} \exp\left(\pi\sqrt{\frac{2n}{3}}\right).$$

Interpret $\log p(n)$ as the microcanonical entropy of the bosonic modes in the first exercise of this part. Distinguish this asymptotic result from an exact coefficient formula; see Hardy–Ramanujan, *Proceedings of the London Mathematical Society* **17** (1918).

16 Automorphic Harmonic Analysis and Arithmetic Langlands

16.1 Places, local fields, and adeles

Definition 16.1 (p -adic completions). For a prime p and nonzero $x \in \mathbb{Q}$, write $x = p^{v_p(x)}a/b$ with $p \nmid ab$ and define

$$|x|_p = p^{-v_p(x)}, \quad |0|_p = 0.$$

This is a non-Archimedean absolute value:

$$|x + y|_p \leq \max\{|x|_p, |y|_p\}.$$

The completion is the locally compact field $\mathbb{Q}_p$, and its compact open subring of integers is

$$\mathbb{Z}_p = \{x \in \mathbb{Q}_p : |x|_p \leq 1\}.$$

The ordinary absolute value gives the Archimedean completion $\mathbb{R}$. A *number field* is a finite extension $F/\mathbb{Q}$. It has completions F_v indexed by its Archimedean and non-Archimedean places v ; at a finite place, $\mathcal{O}_v \subset F_v$ is the ring of integers and the residue field has some cardinality q_v .

Definition 16.2 (Adeles and ideles). The *adele ring* of $\mathbb{Q}$ is the restricted product

$$\mathbb{A}_{\mathbb{Q}} = \mathbb{R} \times \prod'_p \mathbb{Q}_p = \{(x_{\infty}, x_2, x_3, \dots) : x_p \in \mathbb{Z}_p \text{ for all but finitely many } p\}.$$

The restriction makes $\mathbb{A}_{\mathbb{Q}}$ a locally compact ring. Its group of units is the *idele group*

$$\mathbb{A}_{\mathbb{Q}}^{\times} = \mathbb{R}^{\times} \times \prod'_p \mathbb{Q}_p^{\times},$$

where the finite components lie in $\mathbb{Z}_p^{\times}$ for almost every p . For a number field F , define $\mathbb{A}_F$ and $\mathbb{A}_F^{\times}$ in the same way. The diagonal embedding of $F^{\times}$ allows one to define the *idele class group*

$$C_F = F^{\times} \backslash \mathbb{A}_F^{\times}.$$

Exercise 16.1 (The product formula). Compute the p -adic absolute values of $x = 12/25$. For arbitrary $x \in \mathbb{Q}^{\times}$, use unique factorization to prove

$$|x|_{\infty} \prod_p |x|_p = 1,$$

noting that only finitely many factors differ from one. Show that the diagonal image of x is an idele, and prove that the idelic norm

$$|(x_v)|_{\mathbb{A}} = |x_{\infty}|_{\infty} \prod_p |x_p|_p$$

is trivial on $\mathbb{Q}^{\times}$. Explain why it therefore descends to the idele class group.

Definition 16.3 (Hecke characters and their L -functions). A *Hecke character* is a continuous character

$$\chi : F^{\times} \backslash \mathbb{A}_F^{\times} \longrightarrow \mathbb{C}^{\times}$$

with the usual unitary or quasicharacter growth condition. It factors into local characters $\chi_v : F_v^\times \rightarrow \mathbb{C}^\times$. At an unramified finite place its local Euler factor is

$$L_v(s, \chi_v) = \frac{1}{1 - \chi_v(\varpi_v)q_v^{-s}},$$

where ϖ_v is a uniformizer, characterized up to a unit by $|\varpi_v|_v = q_v^{-1}$. The global L -function is the product of these factors, supplemented at ramified and Archimedean places by the appropriate local factors. For the trivial character of $\mathbb{Q}$, its finite part is the Riemann zeta function.

Definition 16.4 (Schwartz–Bruhat functions). A *Schwartz–Bruhat function* on $\mathbb{A}_F$ is a finite linear combination of restricted tensor products

$$\Phi = \bigotimes_v \Phi_v,$$

where Φ_v is an ordinary Schwartz function at each Archimedean place, is locally constant with compact support at each finite place, and equals the indicator of $\mathcal{O}_v$ for almost every finite v .

16.2 Automorphic representations and the trace formula

Definition 16.5 (Automorphic forms and representations). Let G be a reductive group over a number field F . An *automorphic form* is, roughly, a smooth function on

$$G(F) \backslash G(\mathbb{A}_F)$$

that is finite under a maximal compact subgroup and under the center of the Archimedean universal enveloping algebra, has moderate growth, and satisfies an appropriate central-character condition. It is *cuspidal* when its constant term

$$\int_{N_P(F) \backslash N_P(\mathbb{A}_F)} \varphi(ng) \, dn$$

vanishes for every proper parabolic subgroup P with unipotent radical N_P .

Right translation by $G(\mathbb{A}_F)$ acts on automorphic forms. An *automorphic representation* is an irreducible constituent π of this representation.

Exercise 16.2 (Poisson summation as an abelian trace formula). For $f \in \mathcal{S}(\mathbb{R})$, use the Fourier normalization

$$\hat{f}(m) = \int_{\mathbb{R}} f(x) e^{-2\pi i m x} \, dx$$

and define on $L^2(\mathbb{R}/\mathbb{Z})$ the integral operator with kernel

$$K_f(x, y) = \sum_{n \in \mathbb{Z}} f(x - y + n).$$

Diagonalize it in the Fourier basis $e^{2\pi i m x}$. Compute its trace once as the sum of its eigenvalues and once as $\int_0^1 K_f(x, x) \, dx$, obtaining

$$\sum_{m \in \mathbb{Z}} \hat{f}(m) = \sum_{n \in \mathbb{Z}} f(n).$$

Identify the two sides as the spectral and geometric expansions. Explain which features of this calculation depend on commutativity and which survive for an operator defined by convolution on a nonabelian quotient.

16.3 Modular eigenforms as automorphic representations

Definition 16.6 (Classical Hecke operators). Let $f(\tau) = \sum_{n \geq 1} a_n q^n$ be a cusp form of weight k for $SL_2(\mathbb{Z})$. For a prime p , define

$$(T_p f)(\tau) = p^{k-1} f(p\tau) + \frac{1}{p} \sum_{a=0}^{p-1} f\left(\frac{\tau+a}{p}\right).$$

Then

$$T_p f = \sum_{n \geq 1} (a_{pn} + p^{k-1} a_{n/p}) q^n,$$

where $a_{n/p} = 0$ if $p \nmid n$. For $n \geq 1$, define T_n on Fourier coefficients by

$$T_n f = \sum_{m \geq 1} \left(\sum_{d|(m,n)} d^{k-1} a_{mn/d^2} \right) q^m.$$

Definition 16.7 (Normalized Hecke eigenform). A *normalized Hecke eigenform* has $a_1 = 1$ and $T_n f = a_n f$ for every n .

Exercise 16.3 (Hecke recurrences and the Euler product). Derive the displayed Fourier-coefficient formula for T_p . If f is a normalized eigenform, compare coefficients to prove

$$a_{p^{r+1}} = a_p a_{p^r} - p^{k-1} a_{p^{r-1}} \quad (r \geq 1).$$

Derive $T_m T_n = \sum_{d|(m,n)} d^{k-1} T_{mn/d^2}$ from the coefficient formula, and use it to prove $a_{mn} = a_m a_n$ for $(m, n) = 1$, and then show

$$L(f, s) = \sum_{n \geq 1} \frac{a_n}{n^s} = \prod_p \frac{1}{1 - a_p p^{-s} + p^{k-1-2s}}$$

in a right half-plane. Factor the denominator as $(1 - \alpha_p p^{-s})(1 - \beta_p p^{-s})$ and derive $\alpha_p + \beta_p = a_p$ and $\alpha_p \beta_p = p^{k-1}$.

Exercise 16.4 (The GL_2 Hecke correspondence at p). Let $K_p = GL_2(\mathbb{Z}_p)$. Show that index- p sublattices of $\mathbb{Z}_p^2$ are parametrized by $\mathbb{P}^1(\mathbb{F}_p)$ and use this to obtain a decomposition of

$$K_p \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix} K_p$$

into $p+1$ one-sided cosets. Relate these representatives to the $p+1$ terms in the formula for T_p . Explain why simultaneous Hecke diagonalization resembles a complete family of commuting observables, and why this analogy does not make Hecke symmetry generic in quantum mechanics.

16.4 Satake parameters and the dual group

Definition 16.8 (Spherical Hecke algebra). Let F_v be a non-Archimedean local field, let G be split reductive over F_v , and choose a hyperspecial compact subgroup $K_v = G(\mathcal{O}_v)$. The *spherical Hecke algebra* is

$$\mathcal{H}_v(G, K_v) = C_c(K_v \backslash G(F_v) / K_v)$$

with convolution. An irreducible representation π_v with a nonzero K_v -fixed vector is *unramified*. A one-dimensional simultaneous eigenspace for $\mathcal{H}_v(G, K_v)$ defines a *spherical Hecke character*.

16.5 Galois representations and matching L -functions

Definition 16.9 (ℓ -adic representations and Frobenius). Let

$$G_F = \text{Gal}(\overline{F}/F)$$

be the absolute Galois group. An ℓ -adic representation is a continuous homomorphism

$$\rho_\ell : G_F \longrightarrow GL_n(\overline{\mathbb{Q}}_\ell).$$

At a finite place $v \nmid \ell$ where inertia acts trivially, the representation is *unramified*; a Frobenius element Frob_v is then defined up to conjugacy. Fix the Frobenius convention by declaring the local polynomial to be

$$P_v(X) = \det(1 - X\rho_\ell(\text{Frob}_v)).$$

The Galois local factor is $P_v(q_v^{-s})^{-1}$.

Exercise 16.5 (Local zeta function of an elliptic curve). Let α_p, β_p be roots of $X^2 - a_pX + p$ and define a sequence N_r by

$$N_r = p^r + 1 - \alpha_p^r - \beta_p^r,$$

use $\sum_{r \geq 1} z^r/r = -\log(1 - z)$ to prove

$$\exp\left(\sum_{r \geq 1} N_r \frac{T^r}{r}\right) = \frac{1 - a_p(E)T + pT^2}{(1 - T)(1 - pT)}.$$

For the curve $E : y^2 = x^3 - x$ at $p = 5$, count $E(\mathbb{F}_5)$, determine $a_5 = 5 + 1 - \#E(\mathbb{F}_5)$, and compute the first three N_r . Compare the resulting degree-two factor with any weight-two Hecke eigenform having the same coefficient a_5 .

16.6 The Langlands correspondence and its status

Definition 16.10 (Langlands matching and functoriality). An automorphic representation π of $GL_n(\mathbb{A}_F)$ and an n -dimensional ℓ -adic representation ρ are *matched* at an unramified place v when their local polynomials agree:

$$L_v(s, \pi_v) = \det(1 - q_v^{-s}\rho(\text{Frob}_v))^{-1}.$$

They are *globally matched* when this holds at all but finitely many places. For reductive groups, an L -homomorphism

$${}^L H \longrightarrow {}^L G$$

defines *functorial transfer data* when it maps almost all unramified conjugacy classes compatibly and preserves their associated local L -factors.

Part II

Deformation

17 Differential Geometry

17.1 Bundles, sections, and geometric structures

Definition 17.1 (Smooth bundles and sections). A *smooth fiber bundle* with fiber F is a smooth map $\pi : E \rightarrow M$ locally isomorphic over M to $U \times F \rightarrow U$. A *smooth vector bundle* has vector-space fibers and linear transition maps. A *smooth principal G -bundle* has a free right G -action, transitive on each fiber, and local equivariant trivializations. A *section* is a smooth map $s : M \rightarrow E$ with $\pi s = \text{id}_M$. For a representation $G \rightarrow \text{GL}(V)$, the *associated vector bundle* is $P \times_G V = (P \times V)/((pg, v) \sim (p, gv))$.

Definition 17.2 (Frame bundles and G -structures). The *frame bundle* of an n -manifold is the principal $\text{GL}(n, \mathbb{R})$ -bundle

$$\text{Fr}(M)_p = \text{Iso}(\mathbb{R}^n, T_p M).$$

For a closed subgroup $G \leq \text{GL}(n, \mathbb{R})$, a G -*structure* is a principal G -subbundle of $\text{Fr}(M)$. It is a choice of the frames related by the transformations in G . Riemannian, Lorentzian, and almost symplectic structures correspond respectively to reductions to $O(n)$, $O(1, n-1)$, and $\text{Sp}(2n, \mathbb{R})$; an oriented, time-oriented spacetime has structure group $SO^+(1, 3)$.

Definition 17.3 (Tensor fields and metrics). A (r, s) -*tensor field* is a smooth section of $TM^{\otimes r} \otimes T^*M^{\otimes s}$. Equivalently, an $O(p, q)$ -structure is a smooth nondegenerate symmetric tensor g of signature (p, q) . The cases $(n, 0)$ and $(1, n-1)$ give Riemannian and Lorentzian metrics. The length of a Riemannian curve is $\int \sqrt{g(\dot{\gamma}, \dot{\gamma})} dt$; the proper time of a timelike curve is $c^{-1} \int \sqrt{-g(\dot{\gamma}, \dot{\gamma})} dt$. A metric and an orientation define a volume form vol_g .

17.2 Differential forms, cohomology, and Hodge theory

Definition 17.4 (Integration of forms). An *orientation* of an n -manifold is a reduction of its frame bundle to $\text{GL}^+(n, \mathbb{R})$, equivalently a continuous choice of component of nonzero elements in $\Lambda^n T_p^* M$. The integral $\int_M \omega$ of a compactly supported top form is defined using orientation-preserving charts and a partition of unity.

Exercise 17.1 (Stokes' theorem). Prove that the integral is independent of charts and partition of unity. For an oriented manifold with boundary, prove

$$\int_M d\omega = \int_{\partial M} \omega.$$

Deduce the fundamental theorem of calculus, Green's theorem, the divergence theorem, and the classical curl theorem. For time-dependent magnetic two-form B and electric one-form E , prove the equivalence of

$$\int_{\partial S} E = -\frac{d}{dt} \int_S B \quad \text{and} \quad dE = -\partial_t B,$$

recovering the observed flux–emf law.

Definition 17.5 (de Rham cohomology). A differential form ω is *closed* if $d\omega = 0$ and *exact* if $\omega = d\eta$ for some form η . Since $d^2 = 0$, the *de Rham cohomology* of M is

$$H_{\text{dR}}^k(M) = \frac{\ker(d : \Omega^k(M) \rightarrow \Omega^{k+1}(M))}{\text{im}(d : \Omega^{k-1}(M) \rightarrow \Omega^k(M))}.$$

Pullback induces maps on de Rham cohomology, and the wedge product makes $H_{\text{dR}}^*(M)$ a graded-commutative algebra.

Exercise 17.2 (de Rham classes and periods). On a star-shaped open subset of $\mathbb{R}^n$, construct a homotopy operator and prove the Poincaré lemma: every closed form of positive degree is exact. Use Stokes' theorem to prove that an exact k -form has zero integral over every closed oriented k -manifold mapped smoothly into M . Prove $H_{\text{dR}}^1(S^1) \cong \mathbb{R}$, with the isomorphism given by integration.

On $\mathbb{R}^2 \setminus \{0\}$, show that

$$\alpha = \frac{-y dx + x dy}{x^2 + y^2}$$

is closed and has integral 2π around the unit circle, hence is not exact. Compute $H_{\text{dR}}^1(\mathbb{R}^2 \setminus \{0\}) = \mathbb{R} \cdot [\alpha]$. Deduce that a closed electromagnetic field strength has local potentials, while it has a global potential $F = dA$ exactly when $[F] = 0$ in de Rham cohomology.

Definition 17.6 (Hodge star). On an oriented pseudo-Riemannian n -manifold, the metric induces a nondegenerate pairing $\langle \cdot, \cdot \rangle_g$ on differential forms. The *Hodge star* is the bundle map

$$* : \Omega^p(M) \longrightarrow \Omega^{n-p}(M)$$

characterized by

$$\alpha \wedge * \beta = \langle \alpha, \beta \rangle_g \text{ vol}_g$$

for p -forms α, β .

Exercise 17.3 (Metric volume and Hodge star). Construct vol_g and the Hodge star intrinsically and derive their coordinate expressions. For a metric with q negative directions, prove on p -forms that

$$*^2 = (-1)^{p(n-p)+q} \text{id}.$$

Compute $*$ on the standard coordinate forms of Euclidean three-space and Minkowski four-space, stating the orientation and signature conventions.

17.3 Connections, curvature, and gauge fields

Definition 17.7 (Connections and parallel transport). A *connection* on a vector bundle $E \rightarrow M$ is an $\mathbb{R}$ -bilinear map $\nabla : \Gamma(TM) \times \Gamma(E) \rightarrow \Gamma(E)$, linear over smooth functions in the first variable and satisfying $\nabla_X(fs) = X(f)s + f\nabla_X s$. Along a curve γ it induces a covariant derivative D/dt ; its solutions $Ds/dt = 0$ define *parallel transport* between endpoint fibers. On TM , the torsion is $T(X, Y) = \nabla_X Y - \nabla_Y X - [X, Y]$, and a *geodesic* satisfies $D\dot{\gamma}/dt = 0$.

Definition 17.8 (Curvature and holonomy). The *curvature* of ∇ is the $\text{End}(E)$ -valued two-form

$$R^\nabla(X, Y)s = \nabla_X \nabla_Y s - \nabla_Y \nabla_X s - \nabla_{[X, Y]} s.$$

The *holonomy group* at p consists of the automorphisms of E_p obtained by parallel transport around loops based at p .

Exercise 17.4 (Parallel transport and curvature). Prove existence, uniqueness, composition, and inverse properties of parallel transport. Prove that curvature is tensorial and gives the leading failure of transport around an infinitesimal parallelogram to return a vector to itself. Deduce that a flat connection has path-independent transport on every simply connected coordinate domain.

Definition 17.9 (Principal connections and gauge fields). Let $P \rightarrow M$ be a principal G -bundle. A *principal connection* is an equivariant $\mathfrak{g}$ -valued one-form A on P reproducing fundamental vertical fields. It induces connections on every associated bundle, and its *curvature* is $F_A = dA + \frac{1}{2}[A \wedge A]$. A *gauge transformation* is a G -equivariant bundle automorphism covering id_M . A reduction of the frame bundle describes spacetime geometry; a separate principal bundle describes an internal gauge symmetry.

Exercise 17.5 (Gauge covariance and Bianchi identity). Choose local sections s_i and derive transition maps $g_{ij} : U_i \cap U_j \rightarrow G$ and the transformation laws for the local potentials $A_i = s_i^* A$ and curvatures $F_i = s_i^* F_A$. Prove $d_A F_A = 0$ and gauge invariance of holonomy up to conjugacy. For $G = U(1)$, recover $A \mapsto A + d\chi$ and $F = dA$. Show that a flat potential is locally pure gauge, and prove that removing it globally can fail on a nonsimply connected base by computing holonomy around a generator of the fundamental group.

18 Symplectic Geometry

18.1 Projective representations and multipliers

Definition 18.1 (Multipliers and projective representations). A *normalized multiplier* on a locally compact group G is a continuous map $\sigma : G \times G \rightarrow \mathbb{T}$ satisfying

$$\sigma(x, e) = \sigma(e, x) = 1, \quad \sigma(x, y)\sigma(xy, z) = \sigma(y, z)\sigma(x, yz).$$

A σ -*projective unitary representation* is a strongly continuous map $U : G \rightarrow U(H)$ satisfying

$$U_x U_y = \sigma(x, y) U_{xy}, \quad U_e = I.$$

Multipliers σ and σ' are *cohomologous* when there is a continuous map $b : G \rightarrow \mathbb{T}$, $b(e) = 1$, such that

$$\sigma'(x, y) = b(x)b(y)\overline{b(xy)}\sigma(x, y).$$

Exercise 18.1 (The Weyl multiplier). Prove that the phase in the Weyl product is a normalized multiplier on $V \oplus V^*$.

18.2 Alternating commutator form

Definition 18.2 (Commutator bicharacter). For a multiplier σ on an abelian group A , define

$$c_\sigma(x, y) = \sigma(x, y)\overline{\sigma(y, x)}.$$

Exercise 18.2 (Commutator bicharacter). Prove that c_σ is an alternating bicharacter and depends only on the cohomology class of σ . For the Weyl multiplier on $V \oplus V^*$, prove

$$c_\sigma((q, p), (q', p')) = e^{i(p(q') - p'(q))}.$$

18.3 Linear symplectic geometry

Definition 18.3 (Symplectic phase space). A *symplectic form* on a finite-dimensional real linear space E is a nondegenerate alternating bilinear form $\omega : E \times E \rightarrow \mathbb{R}$. On $E = V \oplus V^*$, define

$$\omega((q, p), (q', p')) = p(q') - p'(q).$$

For $\lambda \in \mathbb{R}$, define

$$\sigma_{\lambda\omega}(z, z') = e^{i\lambda\omega(z, z')/2}.$$

The Weyl multiplier is σ_ω .

Exercise 18.3 (Linear Darboux theorem). Prove that every symplectic space has a basis in which its form has matrix

$$\begin{pmatrix} 0 & -I \\ I & 0 \end{pmatrix}.$$

Prove that $\sigma_{\lambda\omega}$ is a multiplier and that $c_{\sigma_{\lambda\omega}}(z, z') = e^{i\lambda\omega(z, z')/2}$.

18.4 Symplectic manifolds and Hamiltonian reduction

Definition 18.4 (Differential forms and exterior derivative). A *differential k -form* on M is a smooth section of $\Lambda^k T^*M$; write $\Omega^k(M) = \Gamma(\Lambda^k T^*M)$. For a smooth map $F : M \rightarrow N$, define

$$(F^*\alpha)_p(v_1, \dots, v_k) = \alpha_{F(p)}(dF_p v_1, \dots, dF_p v_k).$$

The *exterior derivative* is the family $d : \Omega^k(M) \rightarrow \Omega^{k+1}(M)$ satisfying $df(X) = Xf$, the graded Leibniz rule, $d^2 = 0$, and $F^*d = dF^*$.

Exercise 18.4 (Cartan calculus). Construct d locally and prove its uniqueness. For contraction ι_X , prove Cartan's identity

$$\mathcal{L}_X = d\iota_X + \iota_X d.$$

Definition 18.5 (Symplectic manifolds and canonical transformations). A *symplectic form* is a closed nondegenerate two-form ω . For $f \in C^\infty(M)$, define

$$\iota_{X_f}\omega = -df, \quad \{f, g\} = -\omega(X_f, X_g).$$

A diffeomorphism $F : (M, \omega_M) \rightarrow (N, \omega_N)$ is a *canonical transformation* when $F^*\omega_N = \omega_M$. For a manifold Q , the tautological one-form on T^*Q is

$$\theta_\alpha(v) = \alpha(d\pi(v)),$$

and the canonical symplectic form is $\omega = d\theta$.

Definition 18.6 (Classical states and observables). A *pure classical state* is a point of M , a *classical observable* is a function $f \in C^\infty(M, \mathbb{R})$, and a *statistical state* is a Borel probability measure μ with expectation

$$\mathbb{E}_\mu[f] = \int_M f d\mu.$$

For Hamiltonian flow Φ_t , define

$$\mu_t = (\Phi_t)_*\mu_0, \quad f_t = f \circ \Phi_t.$$

Exercise 18.5 (Darboux theorem and Hamiltonian mechanics). Prove the Darboux theorem and identify the canonical form on T^*Q in local coordinates. Derive Hamilton's equations and prove that the Poisson bracket is antisymmetric, satisfies the Leibniz rule and Jacobi identity, and is preserved by canonical transformations. Prove

$$\frac{d}{dt}f(\Phi_t(x)) = \{f, H\}(\Phi_t(x)).$$

Definition 18.7 (Hamiltonian actions and reduction). A symplectic action of a Lie group G on (M, ω) is *Hamiltonian* when it has an equivariant map $J : M \rightarrow \mathfrak{g}^*$ satisfying

$$d\langle J, \xi \rangle = -\iota_{\xi_M} \omega \quad (\xi \in \mathfrak{g}).$$

Let $\eta \in \mathfrak{g}^*$ be a regular value and let $G_\eta = \{g : \text{Ad}_g^* \eta = \eta\}$. If G_η acts freely and properly on $J^{-1}(\eta)$, define

$$M//_\eta G = J^{-1}(\eta)/G_\eta, \quad q^* \omega_\eta = i^* \omega,$$

where $i : J^{-1}(\eta) \hookrightarrow M$ and $q : J^{-1}(\eta) \rightarrow M//_\eta G$.

Exercise 18.6 (Noether and Marsden–Weinstein theorems). Prove that the components of J Poisson-commute with every invariant Hamiltonian. Prove that ω_η exists uniquely and is symplectic and that invariant Hamiltonian flows descend to $M//_\eta G$. Compute the momentum maps for translations and rotations on $T^*\mathbb{R}^3$.

Exercise 18.7 (Liouville's theorem). On a $2n$ -dimensional symplectic manifold, prove that Hamiltonian flow preserves ω , the volume form $\omega^n/n!$, and every density depending only on the Hamiltonian.

19 Classical and Quantum Dynamics

Definition 19.1 (Classical and quantum evolution). Let $H \in C^\infty(\mathbb{R}^{2n}, \mathbb{R})$ have bounded derivatives of order at least two, let Φ_t be its Hamiltonian flow, and let $\text{Op}_h^W(H)$ denote its self-adjoint realization. Define

$$U_h(t) = e^{-it \text{Op}_h^W(H)/\hbar}.$$

For $a \in \mathcal{S}(\mathbb{R}^{2n})$, define

$$a_t = a \circ \Phi_t, \quad A_h(t) = U_h(t)^* \text{Op}_h^W(a) U_h(t).$$

Exercise 19.1 (Heisenberg equation and Egorov's theorem). Prove

$$\frac{d}{dt}a_t = \{a_t, H\}, \quad \frac{d}{dt}A_h(t) = \frac{i}{\hbar}[\text{Op}_h^W(H), A_h(t)].$$

For every $T > 0$, prove Egorov's theorem

$$\sup_{|t| \leq T} \|A_h(t) - \text{Op}_h^W(a \circ \Phi_t)\| = O_T(\hbar).$$

Prove exact equality for quadratic H . Derive Ehrenfest's equations for the expectations of Q_j and P_j .

20 Geometric Quantization

20.1 Prequantization and polarization

Definition 20.1 (Prequantum line bundles). A *prequantum line bundle* over (M, ω) is a Hermitian line bundle $L \rightarrow M$ with a unitary connection ∇ satisfying

$$F_{\nabla} = -\frac{i}{\hbar}\omega.$$

For $f \in C^\infty(M)$, its *Kostant–Souriau operator* is

$$\mathcal{Q}_{\text{pre}}(f) = -i\hbar\nabla_{X_f} + f.$$

Definition 20.2 (Polarizations and metaplectic correction). A *complex polarization* is an involutive Lagrangian subbundle $P \subset T_{\mathbb{C}}M$. Its canonical line is $K_P = \Lambda^{\text{top}}P^*$, and a *metaplectic correction* is a line bundle δ_P with $\delta_P^{\otimes 2} \cong K_P$. The corrected quantum space is the Hilbert completion of sections of $L \otimes \delta_P$ covariantly constant along P .

Exercise 20.1 (Prequantization and polarization). Prove that a prequantum line bundle exists if and only if

$$\left[\frac{\omega}{2\pi\hbar} \right] \in H^2(M; \mathbb{Z}).$$

Prove

$$[\mathcal{Q}_{\text{pre}}(f), \mathcal{Q}_{\text{pre}}(g)] = i\hbar\mathcal{Q}_{\text{pre}}(\{f, g\}).$$

For $T^*\mathbb{R}^n$ with the vertical polarization, recover $L^2(\mathbb{R}^n)$, $Q_j\psi = x_j\psi$, and $P_j\psi = -i\hbar\partial_j\psi$. For a compact Kähler manifold with a positive prequantum line bundle, identify the corrected quantum space with $H^0(M, L \otimes K_M^{1/2})$.

20.2 Equivariant quantization and reduction

Definition 20.3 (Equivariant quantization). Let a Lie group G act on a prequantized Hamiltonian manifold $(M, \omega, J, L, \nabla)$ and preserve a polarization P and metaplectic correction δ_P . An *equivariant quantization* is a lift of the G -action to $L \otimes \delta_P$ preserving its Hermitian structure, connection, and polarization. Its quantum representation is the induced unitary representation on the polarized quantum space.

Exercise 20.2 (Quantization commutes with reduction). For a compact connected group acting holomorphically and Hamiltonianly on a compact prequantized Kähler manifold, assume that 0 is a regular value of J , that G acts freely on $J^{-1}(0)$, and that the prequantum and half-form data descend to the quotient. Prove that the derived quantum representation is generated by the quantized components of J and prove

$$\mathcal{Q}(M//_0G) \cong \mathcal{Q}(M)^G$$

with the metaplectic correction. Derive the multiplicity formula obtained by reduction at integral coadjoint orbits.

21 Deformation Quantization

Definition 21.1 (Poisson manifolds). A *Poisson manifold* is a smooth manifold M with a bracket on $C^\infty(M)$ that is a Lie bracket and a derivation in each argument. Equivalently, there is a bivector $\Pi \in \Gamma(\Lambda^2 TM)$ satisfying

$$\{f, g\} = \Pi(df, dg), \quad [\Pi, \Pi]_{\text{Sch}} = 0.$$

Definition 21.2 (Star products). A *differential star product* on a Poisson manifold is an associative $\mathbb{C}[[\hbar]]$ -bilinear product on $C^\infty(M)[[\hbar]]$ of the form

$$f \star g = fg + \sum_{r \geq 1} \hbar^r C_r(f, g),$$

where the C_r are bidifferential operators, 1 is a unit, and

$$C_1(f, g) - C_1(g, f) = i\{f, g\}.$$

Two star products are *equivalent* when they are intertwined by an operator $T = \text{id} + \sum_{r \geq 1} \hbar^r T_r$ with differential T_r .

Exercise 21.1 (Fedosov and Kontsevich quantization). Prove that the Moyal product is a differential star product on $\mathbb{R}^{2n}$. Prove that every symplectic manifold admits a Fedosov star product and that its equivalence class is determined by its formal characteristic class. Prove the Kontsevich formality theorem and deduce that every finite-dimensional Poisson manifold admits a deformation quantization.

22 Integrable Systems and the Yang–Baxter Equation

Exercise 22.1 (A direct optical Yang–Baxter test). Consider the unitary polarization transformations

$$A(\theta) = \begin{pmatrix} e^{-i\theta} & 0 \\ 0 & e^{i\theta} \end{pmatrix}, \quad B(\theta) = \begin{pmatrix} \cos \theta & -i \sin \theta \\ -i \sin \theta & \cos \theta \end{pmatrix}.$$

Multiply the matrices and prove that, away from singular boundary cases,

$$A(\theta_1)B(\theta_2)A(\theta_3) = B(\theta_3)A(\theta_2)B(\theta_1)$$

exactly when

$$\tan \theta_2 = \frac{\sin(\theta_1 + \theta_3)}{\cos(\theta_1 - \theta_3)}.$$

For $\theta_1 = 56^\circ$ and $\theta_3 = 23^\circ$, obtain $\theta_2 \approx 49.49^\circ$. For an arbitrary input qubit $|\psi\rangle$, show that the output-state overlap has absolute value one when the relation holds and is generally smaller away from it. Compare these predictions with the wave-plate experiment of Zheng et al., *Journal of the Optical Society of America B* **30** (2013). State clearly that this experiment tests a two-dimensional Temperley–Lieb reduction of the Yang–Baxter equation, not the full representation theory of a quantum group or DAHA.

23 Rational Hypergeometric Functions

Definition 23.1 (Ordinary hypergeometric series). For $m \in \mathbb{Z}_{\geq 0}$, define

$$(a)_0 = 1, \quad (a)_m = a(a+1) \cdots (a+m-1) \quad (m \geq 1).$$

Where the quotient is defined, $(a)_m = \Gamma(a+m)/\Gamma(a)$; the product gives its continuation at nonpositive integral a . If no denominator parameter b_j is a nonpositive integer, the *generalized hypergeometric series* is

$${}_rF_s \left(\begin{matrix} a_1, \dots, a_r \\ b_1, \dots, b_s \end{matrix}; z \right) = \sum_{m=0}^{\infty} \frac{(a_1)_m \cdots (a_r)_m}{(b_1)_m \cdots (b_s)_m} \frac{z^m}{m!}.$$

It is interpreted as a finite sum when a numerator parameter is a nonpositive integer. Otherwise it converges for every z when $r \leq s$, for $|z| < 1$ when $r = s+1$, and only at $z = 0$ when $r > s+1$. The Gauss hypergeometric operator is

$$\mathcal{L}_{a,b,c} = z(1-z) \frac{d^2}{dz^2} + (c - (a+b+1)z) \frac{d}{dz} - ab.$$

Exercise 23.1 (Gauss hypergeometric theorem). Prove

$${}_1F_0 \left(\begin{matrix} a \\ - \end{matrix}; z \right) = (1-z)^{-a}$$

and prove that ${}_2F_1(a, b; c; z)$ is the solution of $\mathcal{L}_{a,b,c}f = 0$ analytic at $z = 0$ with $f(0) = 1$. Derive Euler's integral and transformation

$${}_2F_1(a, b; c; z) = \frac{\Gamma(c)}{\Gamma(b)\Gamma(c-b)} \int_0^1 u^{b-1} (1-u)^{c-b-1} (1-zu)^{-a} du,$$

$${}_2F_1(a, b; c; z) = (1-z)^{c-a-b} {}_2F_1(c-a, c-b; c; z),$$

and Gauss's summation

$${}_2F_1(a, b; c; 1) = \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)}.$$

State the parameter conditions for each identity and extend them by analytic continuation.

Definition 23.2 (Jacobi and Wilson polynomials). For $\alpha, \beta > -1$, the *Jacobi polynomial* is

$$P_n^{(\alpha, \beta)}(x) = \frac{(\alpha+1)_n}{n!} {}_2F_1 \left(\begin{matrix} -n, n+\alpha+\beta+1 \\ \alpha+1 \end{matrix}; \frac{1-x}{2} \right).$$

For positive a, b, c, d , the *Wilson polynomial* is

$$W_n(x^2; a, b, c, d) = (a+b)_n(a+c)_n(a+d)_n \times {}_4F_3 \left(\begin{matrix} -n, n+a+b+c+d-1, a+ix, a-ix \\ a+b, a+c, a+d \end{matrix}; 1 \right).$$

Exercise 23.2 (Rational rank-one spectral theory). Prove

$$\left((1-x^2) \frac{d^2}{dx^2} + (\beta - \alpha - (\alpha + \beta + 2)x) \frac{d}{dx} \right) P_n^{(\alpha, \beta)} = -n(n + \alpha + \beta + 1) P_n^{(\alpha, \beta)}.$$

Derive Jacobi orthogonality on $[-1, 1]$ with weight $(1-x)^\alpha(1+x)^\beta$. Prove the Wilson difference equation and orthogonality on $[0, \infty)$ with weight

$$w_W(x) = \left| \frac{\Gamma(a+ix)\Gamma(b+ix)\Gamma(c+ix)\Gamma(d+ix)}{\Gamma(2ix)} \right|^2.$$

Derive the three-term recurrences of both families.

24 Trigonometric/Basic Hypergeometric Deformation

Definition 24.1 (Basic hypergeometric series). For $m \in \mathbb{N}$, define

$$(a; q)_m = \prod_{j=0}^{m-1} (1 - aq^j), \quad (a; q)_\infty = \prod_{j=0}^{\infty} (1 - aq^j),$$

and

$$(a_1, \dots, a_r; q)_m = \prod_{i=1}^r (a_i; q)_m, \quad (a_1, \dots, a_r; q)_\infty = \prod_{i=1}^r (a_i; q)_\infty.$$

The *basic hypergeometric series* is

$${}_r\phi_s \left(\begin{matrix} a_1, \dots, a_r \\ b_1, \dots, b_s \end{matrix}; q, z \right) = \sum_{m=0}^{\infty} \frac{(a_1, \dots, a_r; q)_m}{(q, b_1, \dots, b_s; q)_m} \left((-1)^m q^{m(m-1)/2} \right)^{1+s-r} z^m.$$

It is *terminating* when one numerator parameter is q^{-n} .

Exercise 24.1 (Basic hypergeometric summations). Prove the q -binomial theorem

$${}_1\phi_0 \left(\begin{matrix} a \\ - \end{matrix}; q, z \right) = \frac{(az; q)_\infty}{(z; q)_\infty}, \quad |z| < 1,$$

the q -Gauss summation

$${}_2\phi_1 \left(\begin{matrix} a, b \\ c \end{matrix}; q, \frac{c}{ab} \right) = \frac{(c/a, c/b; q)_\infty}{(c, c/(ab); q)_\infty}, \quad \left| \frac{c}{ab} \right| < 1,$$

and the terminating q -Pfaff–Saalschütz summation

$${}_3\phi_2 \left(\begin{matrix} a, b, q^{-n} \\ c, abq^{1-n}/c \end{matrix}; q, q \right) = \frac{(c/a, c/b; q)_n}{(c, c/(ab); q)_n}.$$

Definition 24.2 (Askey–Wilson operator and polynomials). For $0 < q < 1$ and nonzero a, b, c, d , define

$$A(z) = \frac{(1-az)(1-bz)(1-cz)(1-dz)}{(1-z^2)(1-qz^2)}$$

and, on $f(z) = f(z^{-1})$,

$$(\mathcal{D}_{AW}f)(z) = A(z)(f(qz) - f(z)) + A(z^{-1})(f(q^{-1}z) - f(z)).$$

For $x = (z + z^{-1})/2$, the *Askey–Wilson polynomial* is

$$p_n(x; a, b, c, d \mid q) = a^{-n}(ab, ac, ad; q)_n \times {}_4\phi_3 \left(\begin{matrix} q^{-n}, abcdq^{n-1}, az, az^{-1} \\ ab, ac, ad \end{matrix}; q, q \right).$$

Its weight on the unit circle is

$$w_{AW}(z) = \frac{(z^2, z^{-2}; q)_\infty}{(az, a/z, bz, b/z, cz, c/z, dz, d/z; q)_\infty}.$$

Exercise 24.2 (Rank-one DAHA and the Askey–Wilson theorem). Construct the basic representation of $\widehat{H}_q(C_1^\vee, C_1)$ from the involutions

$$(s_1f)(z) = f(z^{-1}), \quad (s_0f)(z) = f(qz^{-1})$$

and first-order reflection- q -difference operators. Verify the four Hecke relations and the product relation. Prove that the spherical subalgebra is generated in its polynomial representation by multiplication by $z + z^{-1}$ and $\mathcal{D}_{AW}$. Prove

$$\mathcal{D}_{AW}p_n = (q^{-n} - 1)(1 - abcdq^{n-1})p_n.$$

For $a, b, c, d \in (-1, 1)$, prove that $\mathcal{D}_{AW}$ is symmetric for

$$\langle f, g \rangle_{AW} = \int_{|z|=1} f(z) \overline{g(z)} w_{AW}(z) \frac{dz}{2\pi iz},$$

and deduce the orthogonality and three-term recurrence of the Askey–Wilson polynomials.

25 Poisson–Lie Groups and Quantum Groups

25.1 Poisson–Lie groups and Lie bialgebras

Definition 25.1 (Poisson–Lie groups and Lie bialgebras). A *Poisson–Lie group* is a Lie group G with a Poisson structure for which multiplication $G \times G \rightarrow G$ is Poisson. A *Lie bialgebra* is a Lie algebra $\mathfrak{g}$ with a linear map

$$\delta : \mathfrak{g} \rightarrow \mathfrak{g} \wedge \mathfrak{g}$$

that is a 1-cocycle for the adjoint action and whose transpose defines a Lie bracket on $\mathfrak{g}^*$. It is *coboundary* when

$$\delta(x) = [x \otimes 1 + 1 \otimes x, r]$$

for some $r \in \mathfrak{g} \otimes \mathfrak{g}$.

Exercise 25.1 (Infinitesimal Poisson–Lie duality). Differentiate a Poisson–Lie structure at the identity and prove that it defines a Lie bialgebra. Prove the converse for connected simply connected groups. For a coboundary Lie bialgebra, define

$$\text{CYB}(r) = [r_{12}, r_{13}] + [r_{12}, r_{23}] + [r_{13}, r_{23}].$$

Prove that the co-Jacobi identity is equivalent to $\text{CYB}(r)$ being invariant under the diagonal adjoint action. Under the additional triangular hypothesis $\text{CYB}(r) = 0$, recover the classical Yang–Baxter equation; state the modified classical Yang–Baxter equation as $\text{CYB}(r) = \Omega$ for an invariant tensor Ω and distinguish the case $\Omega \neq 0$.

25.2 Drinfeld–Jimbo quantization

Definition 25.2 (Quantized enveloping algebras). A *quantized enveloping algebra* of $(\mathfrak{g}, \delta)$ is a topologically free Hopf algebra $U_\hbar(\mathfrak{g})$ over $\mathbb{C}[[\hbar]]$ satisfying

$$U_\hbar(\mathfrak{g})/\hbar U_\hbar(\mathfrak{g}) \cong U(\mathfrak{g}), \quad \delta(x) = \frac{\Delta_\hbar(x) - \Delta_\hbar^{\text{op}}(x)}{\hbar} \bmod \hbar.$$

For generic q , $U_q(\mathfrak{sl}_2)$ is generated by E, F, K, K^{-1} with

$$KEK^{-1} = q^2 E, \quad KFK^{-1} = q^{-2} F, \quad [E, F] = \frac{K - K^{-1}}{q - q^{-1}},$$

and

$$\Delta(K) = K \otimes K, \quad \Delta(E) = E \otimes 1 + K \otimes E, \quad \Delta(F) = F \otimes K^{-1} + 1 \otimes F.$$

Exercise 25.2 (Drinfeld–Jimbo theorem). Construct $U_q(\mathfrak{g})$ from the Cartan matrix of a complex semisimple Lie algebra and prove that it is a Hopf deformation of $U(\mathfrak{g})$. For $\mathfrak{sl}_2$, verify the displayed relations and coproduct and compute the semiclassical Lie bialgebra.

26 Compact Quantum Groups

Definition 26.1 (Compact quantum groups and Haar states). A *compact quantum group* $\mathbb{G}$ is a unital C^* -algebra $C(\mathbb{G})$ with a unital $*$ -homomorphism

$$\Delta : C(\mathbb{G}) \longrightarrow C(\mathbb{G}) \otimes_{\min} C(\mathbb{G})$$

where $\otimes_{\min}$ is the closure of the algebraic tensor product in $B(H \otimes K)$ for faithful representations on H and K . The coproduct satisfies

$$(\Delta \otimes \text{id})\Delta = (\text{id} \otimes \Delta)\Delta$$

and the linear spans of $\Delta(C(\mathbb{G}))(1 \otimes C(\mathbb{G}))$ and $\Delta(C(\mathbb{G}))(C(\mathbb{G}) \otimes 1)$ are dense in $C(\mathbb{G}) \otimes_{\min} C(\mathbb{G})$. A *Haar state* is a state h satisfying

$$(h \otimes \text{id})\Delta(a) = h(a)1 = (\text{id} \otimes h)\Delta(a).$$

Definition 26.2 (Corepresentations and representative functions). A finite-dimensional *unitary corepresentation* on H_U is a unitary $U \in B(H_U) \otimes C(\mathbb{G})$ satisfying

$$(\text{id} \otimes \Delta)(U) = U_{12}U_{13}.$$

If $U = (u_{ij})$, then $\Delta(u_{ij}) = \sum_k u_{ik} \otimes u_{kj}$. Denote the linear span of the matrix coefficients of all finite-dimensional unitary corepresentations by $\text{Pol}(\mathbb{G})$.

Definition 26.3 (Drinfeld–Jimbo and Woronowicz duality). Let $0 < q < 1$, let G be a compact connected simply connected semisimple Lie group, and let $\mathfrak{g}_{\mathbb{C}}$ be its complexified Lie algebra. The Hopf $*$ -algebra $\text{Pol}(G_q)$ is spanned by the matrix coefficients $c_{\varphi, v}$ of finite-dimensional type-1 unitary $U_q(\mathfrak{g}_{\mathbb{C}})$ -modules for the compact real-form involution, with pairing

$$\langle x, c_{\varphi, v} \rangle = \varphi(xv).$$

Its universal and reduced C^* -completions are denoted $C_u(G_q)$ and $C_r(G_q)$.

Exercise 26.1 (Woronowicz Peter–Weyl theorem). Prove existence and uniqueness of the Haar state. Choose one representative from each equivalence class of irreducible unitary corepresentations and prove that all of their matrix coefficients are linearly independent and that $\text{Pol}(\mathbb{G})$ is a dense Hopf $*$ -subalgebra of $C(\mathbb{G})$. For each irreducible U^α , construct the normalized positive matrix Q_α satisfying

$$(\text{id} \otimes S^2)(U^\alpha) = (Q_\alpha \otimes 1)U^\alpha(Q_\alpha^{-1} \otimes 1), \quad \text{Tr}(Q_\alpha) = \text{Tr}(Q_\alpha^{-1}),$$

and derive the orthogonality relations in the Haar GNS space and the resulting Hilbert direct-sum decomposition. Prove that the displayed pairing between $U_q(\mathfrak{g}_\mathbb{C})$ and $\text{Pol}(G_q)$ is a nondegenerate Hopf pairing.

Exercise 26.2 ($SU_q(2)$). Let $C(SU_q(2))$ be generated by α, γ so that

$$u = \begin{pmatrix} \alpha & -q\gamma^* \\ \gamma & \alpha^* \end{pmatrix}$$

is unitary, and define $\Delta(u_{ij}) = \sum_k u_{ik} \otimes u_{kj}$. Derive the relations

$$\begin{aligned} \alpha^* \alpha + \gamma^* \gamma &= 1, & \alpha \alpha^* + q^2 \gamma \gamma^* &= 1, & \gamma^* \gamma &= \gamma \gamma^*, \\ \alpha \gamma &= q \gamma \alpha, & \alpha \gamma^* &= q \gamma^* \alpha, \end{aligned}$$

and prove that Δ is a coassociative $*$ -homomorphism satisfying the cancellation conditions.

Definition 26.4 (Representation-category duality). Let $\text{Rep}_{\text{type } 1}^{\text{fd}}(U_q(\mathfrak{g}_\mathbb{C}))$ be the category of finite-dimensional type-1 unitary modules and intertwiners, and let $\text{Corep}^{\text{fd}}(\text{Pol}(G_q))$ be the category of finite-dimensional unitary corepresentations and intertwiners. For a module (V, π_V) , define $U_V \in \text{End}(V) \otimes \text{Pol}(G_q)$ by

$$(\text{id} \otimes \langle x, \cdot \rangle)(U_V) = \pi_V(x) \quad (x \in U_q(\mathfrak{g}_\mathbb{C})).$$

Define

$$\mathcal{F} : \text{Rep}_{\text{type } 1}^{\text{fd}}(U_q(\mathfrak{g}_\mathbb{C})) \longrightarrow \text{Corep}^{\text{fd}}(\text{Pol}(G_q)), \quad \mathcal{F}(V) = U_V, \quad \mathcal{F}(T) = T,$$

with identity tensor constraints on the underlying spaces $V \otimes W$.

Exercise 26.3 (Matrix coefficients and tensor products). For $v \in V$, $w \in W$, $\varphi \in V^*$, and $\psi \in W^*$, prove

$$c_{\varphi, v} c_{\psi, w} = c_{\varphi \otimes \psi, v \otimes w}.$$

Use the Hopf-pairing identities to prove

$$(\text{id} \otimes \Delta)(U_V) = U_{V,12} U_{V,13}, \quad U_{V \otimes W} = U_{V,13} U_{W,23}.$$

Prove that $\mathcal{F}$ is fully faithful and essentially surjective and that its tensor constraints are intertwiners. Conclude

$$\text{Rep}_{\text{type } 1}^{\text{fd}}(U_q(\mathfrak{g}_\mathbb{C})) \simeq \text{Corep}^{\text{fd}}(\text{Pol}(G_q))$$

as monoidal $*$ -categories.

Definition 26.5 (Conjugates and quantum dimension). For a finite-dimensional H -module V , its *left dual* $V^\vee$ has action

$$(x\varphi)(v) = \varphi(S(x)v),$$

evaluation $\text{ev}_V(\varphi \otimes v) = \varphi(v)$, and coevaluation $\text{coev}_V(1) = \sum_i e_i \otimes e^i$ for a basis (e_i) and its dual basis (e^i) . In a rigid C^* -tensor category, a *conjugate* of V is an object $\bar{V}$ with morphisms

$$R_V : \mathbf{1} \longrightarrow \bar{V} \otimes V, \quad \bar{R}_V : \mathbf{1} \longrightarrow V \otimes \bar{V}$$

satisfying

$$(\bar{R}_V^* \otimes \text{id}_V)(\text{id}_V \otimes R_V) = \text{id}_V, \quad (R_V^* \otimes \text{id}_{\bar{V}})(\text{id}_{\bar{V}} \otimes \bar{R}_V) = \text{id}_{\bar{V}}.$$

The adjoints R_V^* and $\bar{R}_V^*$ are evaluation maps, and R_V and $\bar{R}_V$ are coevaluation maps. A solution is *standard* when

$$R_V^*(\text{id}_{\bar{V}} \otimes T)R_V = \bar{R}_V^*(T \otimes \text{id}_{\bar{V}})\bar{R}_V \quad (T \in \text{End}(V)).$$

For a standard solution, define

$$\text{tr}_q(T) = R_V^*(\text{id}_{\bar{V}} \otimes T)R_V, \quad \dim_q(V) = \text{tr}_q(\text{id}_V).$$

Exercise 26.4 ($SU_q(2)$ fusion and dimensions). Let V_n be the irreducible type-1 $U_q(\mathfrak{sl}_2)$ -module of highest weight n . Construct its conjugate equations and prove

$$V_m \otimes V_n \cong \bigoplus_{k=0}^{\min(m,n)} V_{m+n-2k}.$$

Prove that the normalized matrices from the Woronowicz orthogonality relations implement the standard solutions and compute

$$\dim_q(V_n) = [n+1]_q = \frac{q^{n+1} - q^{-(n+1)}}{q - q^{-1}}.$$

Prove

$$\dim V_n = n+1, \quad \lim_{q \rightarrow 1} \dim_q(V_n) = n+1, \quad \dim_q(V_n) > n+1$$

for $0 < q < 1$ and $n \geq 1$.

Definition 26.6 (Tannaka–Krein reconstruction). Let $\mathcal{C}$ be an essentially small rigid C^* -tensor category and let $F : \mathcal{C} \rightarrow \text{Hilb}_{\text{fd}}$ be a faithful unitary tensor functor. Define

$$\mathcal{A}_F = \left(\bigoplus_{V \in \text{Ob } \mathcal{C}} F(V)^* \otimes F(V) \right) / \langle (\psi \circ F(T)) \otimes v - \psi \otimes F(T)v \rangle,$$

where $T : V \rightarrow W$, $v \in F(V)$, and $\psi \in F(W)^*$. Its multiplication is induced by tensor products, and for a basis (e_i) of $F(V)$ define

$$\Delta[\varphi \otimes v]_V = \sum_i [\varphi \otimes e_i]_V \otimes [e_i^* \otimes v]_V, \quad \varepsilon[\varphi \otimes v]_V = \varphi(v).$$

The involution and antipode are induced by the conjugate data.

Exercise 26.5 (Tannaka–Krein theorem). Prove that $\mathcal{A}_F$ is a Hopf $*$ -algebra and is independent of the chosen bases. For the forgetful fiber functor

$$F : \text{Rep}(G_q) \longrightarrow \text{Hilb}_{\text{fd}},$$

prove that

$$[\varphi \otimes v]_V \longmapsto c_{\varphi, v}$$

is a Hopf $*$ -isomorphism $\mathcal{A}_F \cong \text{Pol}(G_q)$.

Definition 26.7 (Universal R -matrix and braiding). A *universal R -matrix* for a Hopf algebra H is an invertible element $\mathcal{R}$ in a completion of $H \otimes H$ on which the finite-dimensional tensor-product representations under consideration extend, satisfying

$$\mathcal{R}\Delta(x)\mathcal{R}^{-1} = \Delta^{\text{op}}(x), \quad (\Delta \otimes \text{id})(\mathcal{R}) = \mathcal{R}_{13}\mathcal{R}_{23}, \quad (\text{id} \otimes \Delta)(\mathcal{R}) = \mathcal{R}_{13}\mathcal{R}_{12}.$$

For finite-dimensional modules V, W , define

$$c_{V,W} = \tau \circ (\pi_V \otimes \pi_W)(\mathcal{R}) : V \otimes W \longrightarrow W \otimes V.$$

The pair $(H, \mathcal{R})$ is *quasitriangular*.

Exercise 26.6 (Hexagon identities and the Yang–Baxter equation). Prove that $c_{V,W}$ is a natural module intertwiner and that the two coproduct identities for $\mathcal{R}$ give the two hexagon identities. Derive

$$\mathcal{R}_{12}\mathcal{R}_{13}\mathcal{R}_{23} = \mathcal{R}_{23}\mathcal{R}_{13}\mathcal{R}_{12}$$

and, on $V^{\otimes 3}$,

$$c_{12}c_{23}c_{12} = c_{23}c_{12}c_{23}.$$

Prove that the operators $c_{j,j+1}$ define braid-group representations on tensor powers of every finite-dimensional type-1 module.

Definition 26.8 (Ribbon categories). Let $\mathcal{C}$ be a braided rigid $\mathbb{C}$ -linear monoidal category. A *twist* is a natural isomorphism $\theta : \text{id}_{\mathcal{C}} \rightarrow \text{id}_{\mathcal{C}}$ satisfying

$$\theta_1 = \text{id}_1, \quad \theta_{X \otimes Y} = c_{Y,X}c_{X,Y}(\theta_X \otimes \theta_Y).$$

The category is *ribbon* when

$$\theta_{X^\vee} = (\theta_X)^\vee$$

for every object X .

Exercise 26.7 (Ribbon elements). For a quasitriangular Hopf algebra $(H, \mathcal{R})$, set

$$u = m(S \otimes \text{id})(\mathcal{R}_{21}).$$

Let $v \in H$ be central and invertible and satisfy

$$v^2 = uS(u), \quad S(v) = v, \quad \varepsilon(v) = 1, \quad \Delta(v) = (\mathcal{R}_{21}\mathcal{R})^{-1}(v \otimes v).$$

Prove that

$$\theta_V = \pi_V(v^{-1})$$

defines a ribbon twist on the category of finite-dimensional H -modules. Derive invariance of the quantum trace under framed ribbon isotopy.

27 Hecke Algebras and Quantum Schur–Weyl Duality

Definition 27.1 (Root data and affine Weyl groups). Let E be a finite-dimensional Euclidean space. A finite set $R \subset E \setminus \{0\}$ is a *reduced crystallographic root system* if it spans E , is preserved by the reflections

$$s_\alpha(x) = x - \langle x, \alpha^\vee \rangle \alpha, \quad \alpha^\vee = \frac{2\alpha}{\langle \alpha, \alpha \rangle},$$

satisfies $R \cap \mathbb{R}\alpha = \{\alpha, -\alpha\}$, and has $\langle \beta, \alpha^\vee \rangle \in \mathbb{Z}$ for all $\alpha, \beta \in R$. Define

$$W = \langle s_\alpha : \alpha \in R \rangle, \quad Q^\vee = \text{span}_{\mathbb{Z}}\{\alpha^\vee : \alpha \in R\}, \quad W_{\text{aff}} = Q^\vee \rtimes W.$$

The connected components of the complement of the affine reflection hyperplanes

$$H_{\alpha, m} = \{x \in E : \langle x, \alpha \rangle = m\}, \quad \alpha \in R, \quad m \in \mathbb{Z},$$

are the *alcoves*.

Definition 27.2 (Finite Hecke algebras). Let $(W, \{s_i\})$ be a Coxeter system, and let m_{ij} be the order of $s_i s_j$. Its *Hecke algebra* $\mathcal{H}_t(W)$ over $\mathbb{C}(t^{1/2})$ is generated by T_i with the braid relations

$$\underbrace{T_i T_j T_i \cdots}_{m_{ij} \text{ factors}} = \underbrace{T_j T_i T_j \cdots}_{m_{ij} \text{ factors}}$$

and the quadratic relations

$$(T_i - t^{1/2})(T_i + t^{-1/2}) = 0.$$

Definition 27.3 (Quantum Schur–Weyl actions). Let v be generic, let $U_v(\mathfrak{gl}_N)$ act on its vector module $V = \mathbb{C}^N$, and set $t = v^2$. On $V^{\otimes r}$ define

$$\rho_r(T_i) = \text{id}_V^{\otimes(i-1)} \otimes \check{R}_{V,V} \otimes \text{id}_V^{\otimes(r-i-1)}, \quad \check{R}_{V,V} = \tau(\pi_V \otimes \pi_V)(\mathcal{R}),$$

with the normalization for which $\check{R}_{V,V}$ has eigenvalues v and $-v^{-1}$.

Exercise 27.1 (Quantum Schur–Weyl duality). Prove that the coproduct action of $U_v(\mathfrak{gl}_N)$ commutes with $\rho_r(T_i)$. Derive the braid and Hecke relations from the universal R -matrix and prove that ρ_r defines an action of $\mathcal{H}_{v^2}(S_r)$. For generic v and $N \geq r$, prove the double-centralizer identities

$$\rho_r(\mathcal{H}_{v^2}(S_r)) = \text{End}_{U_v(\mathfrak{gl}_N)}(V^{\otimes r}), \quad \pi_r(U_v(\mathfrak{gl}_N)) = \text{End}_{\mathcal{H}_{v^2}(S_r)}(V^{\otimes r}),$$

and the decomposition

$$V^{\otimes r} \cong \bigoplus_{\substack{\lambda \vdash r \\ \ell(\lambda) \leq N}} L_\lambda^{(N)} \otimes S_t^\lambda.$$

Definition 27.4 (Affine Hecke algebras). The extended affine Hecke algebra $\mathcal{H}_t^{\text{aff}}(GL_r)$ is generated by $T_1, \dots, T_{r-1}$ and commuting invertible elements $X_1, \dots, X_r$, subject to the finite Hecke relations and

$$T_i X_i T_i = X_{i+1}, \quad T_i X_j = X_j T_i \quad (j \neq i, i+1).$$

Equivalently, it is the Hecke quotient of the extended affine braid group of $\mathbb{Z}^r \rtimes S_r$.

Definition 27.5 (Affine quantum Schur–Weyl functor). Let $U_v(L\mathfrak{gl}_N)$ be the quantum loop algebra and let

$$\widehat{V} = V \otimes \mathbb{C}[z^{\pm 1}]$$

be the affinization of its vector evaluation module. The tensor power $\widehat{V}^{\otimes r}$ is a $U_v(L\mathfrak{gl}_N)$ – $\mathcal{H}_{v^2}^{\text{aff}}(GL_r)$ bimodule, with the right action determined by the normalized spectral R -matrices and the loop operators. Define

$$\text{SW}_{N,r}^{\text{aff}}(M) = \widehat{V}^{\otimes r} \otimes_{\mathcal{H}_{v^2}^{\text{aff}}(GL_r)} M.$$

Exercise 27.2 (Affine quantum Schur–Weyl duality). Use the spectral Yang–Baxter equation to prove that the two actions on $\widehat{V}^{\otimes r}$ commute and satisfy the affine braid relations. Use the two eigenvalues of the vector R -matrix to derive the affine Hecke quadratic relation. For $N > r$, prove that $\text{SW}_{N,r}^{\text{aff}}$ is an equivalence from finite-dimensional $\mathcal{H}_{v^2}^{\text{aff}}(GL_r)$ -modules to the full subcategory of finite-dimensional polynomial $U_v(L\mathfrak{gl}_N)$ -modules generated by subquotients of tensor products of r vector evaluation modules.

28 DAHA and Macdonald Theory

28.1 Double affine Hecke algebras

Definition 28.1 (Double-affine braid group). Let $T^2 = \mathbb{R}^2/\mathbb{Z}^2$ and let

$$\text{Conf}_r(T^2) = \{(z_1, \dots, z_r) \in (T^2)^r : z_i \neq z_j \text{ for } i \neq j\}.$$

The *elliptic braid group* is

$$B_r^{\text{ell}} = \pi_1(\text{Conf}_r(T^2)/S_r).$$

Its standard generators are adjacent braids T_i and loops X_j, Y_j moving the j th point around the two fundamental cycles of T^2 . The central extension in which the total X – Y commutator is denoted $\mathbf{q}$ is the *double-affine braid group* of type GL_r .

Exercise 28.1 (From affine to double-affine braids). Derive a presentation of B_r^{ell} from the motions T_i, X_j, Y_j . Prove that the T_i satisfy the finite braid relations, that the X_j commute, that the Y_j commute, and that conjugation by T_i permutes the adjacent loop generators. Compute the central commutator of the two total torus motions. Cutting T^2 along one cycle, prove that the remaining generators form an affine braid group.

Definition 28.2 (Double affine Hecke algebras). Let $B^{\text{da}}(R)$ be the double-affine braid group of a root datum R , with braid generators T_i and central parameter $\mathbf{q}$. For nonzero parameters q and t_i , define

$$H_{q,\mathbf{t}}(R) = \mathbb{C}(q, \mathbf{t})[B^{\text{da}}(R)] / \left\langle \mathbf{q} - q, (T_i - t_i^{1/2})(T_i + t_i^{-1/2}) \right\rangle.$$

This is the *double affine Hecke algebra* of R .

Definition 28.3 (Rank-one DAHA and its spherical representation). The universal DAHA $\hat{H}_q(C_1^\vee, C_1)$ is generated by $t_i^{\pm 1}$ for $0 \leq i \leq 3$, subject to

$$t_i t_i^{-1} = 1, \quad t_i + t_i^{-1} \text{ central}, \quad t_0 t_1 t_2 t_3 = q^{-1}.$$

For $k_i \neq 0$, the specialization

$$t_i + t_i^{-1} = k_i + k_i^{-1}$$

is the four-parameter rank-one DAHA. After the standard scalar normalization, let T be its finite Hecke generator with $(T - k)(T + k^{-1}) = 0$, and define

$$e = \frac{T + k^{-1}}{k + k^{-1}}.$$

Its basic polynomial representation is on $\mathcal{L} = \mathbb{C}[z^{\pm 1}]$ by reflection– q -difference operators, and

$$e\mathcal{L} = \mathcal{L}^{z \mapsto z^{-1}} = \mathbb{C}[z + z^{-1}].$$

The action of $e\hat{H}_q e$ on $e\mathcal{L}$ is the *spherical rank-one polynomial representation*.

28.2 Macdonald theory

Definition 28.4 (Type- A polynomial representation). Let

$$\mathcal{P}_r = \mathbb{C}(q, t^{1/2})[x_1^{\pm 1}, \dots, x_r^{\pm 1}],$$

let X_j act by multiplication by x_j , and let s_i exchange x_i and x_{i+1} . Define

$$T_i = t^{1/2}s_i + (t^{1/2} - t^{-1/2}) \frac{s_i - 1}{x_i/x_{i+1} - 1}, \quad 1 \leq i < r,$$

and

$$(\pi f)(x_1, \dots, x_r) = f(x_2, \dots, x_r, qx_1).$$

For generic independent parameters q, t , the operator algebra generated by $X_j^{\pm 1}$, T_i , and $\pi^{\pm 1}$ is the polynomial realization of the GL_r double affine Hecke algebra $H_{q,t}(GL_r)$. The subalgebra generated by T_i and $\pi^{\pm 1}$ is an extended affine Hecke algebra; its translation elements form a commuting family $Y_1, \dots, Y_r$ of q -difference-reflection operators.

Exercise 28.2 (Polynomial representation and the DAHA quotient). Prove that the displayed operators preserve $\mathcal{P}_r$ and satisfy the braid and Hecke relations. Prove

$$\pi X_i \pi^{-1} = X_{i+1} \quad (1 \leq i < r), \quad \pi X_r \pi^{-1} = qX_1,$$

and identify the two affine-Hecke subalgebras generated respectively by T_i, X_j and by T_i, Y_j . Prove that this representation is obtained from the double-affine braid-group algebra by imposing

$$(T_i - t^{1/2})(T_i + t^{-1/2}) = 0$$

and specializing its central parameter to q .

Definition 28.5 (Macdonald operators and polynomials). Let

$$(T_{q,x_i} f)(x_1, \dots, x_r) = f(x_1, \dots, qx_i, \dots, x_r).$$

The first type- A Macdonald operator is

$$D_{q,t} = \sum_{i=1}^r \left(\prod_{j \neq i} \frac{tx_i - x_j}{x_i - x_j} \right) T_{q,x_i}.$$

For generic q, t , the Macdonald polynomial $P_\lambda(x; q, t)$ is the symmetric polynomial triangular in the monomial-symmetric basis and satisfying

$$D_{q,t} P_\lambda = \left(\sum_{i=1}^r q^{\lambda_i} t^{r-i} \right) P_\lambda, \quad \lambda_1 \geq \dots \geq \lambda_r \geq 0.$$

For $0 < q, t < 1$, define

$$\Delta_{q,t}(x) = \prod_{i \neq j} \frac{(x_i/x_j; q)_\infty}{(tx_i/x_j; q)_\infty},$$

and

$$\langle f, g \rangle_{q,t} = \frac{1}{r!} \int_{(S^1)^r} f(x) \overline{g(x)} \Delta_{q,t}(x) \prod_{j=1}^r \frac{dx_j}{2\pi i x_j}.$$

Exercise 28.3 (DAHA Fourier transform and Macdonald spectral theory). Construct the commuting Y -operators and prove that their S_r -invariant Laurent polynomials preserve $\mathcal{P}_r^{S_r}$. Identify the first elementary symmetric polynomial in the Y_j with $D_{q,t}$ up to normalization. Prove existence, uniqueness, orthogonality, and joint diagonalization of the Macdonald polynomials. Construct the DAHA Fourier transform exchanging the X - and Y -polynomial representations and derive its projective $SL_2(\mathbb{Z})$ relations.

Exercise 28.4 (Ordinary and basic hypergeometric degeneration). Set $q = e^{-\varepsilon}$. Prove

$$\lim_{\varepsilon \rightarrow 0^+} \frac{(q^a; q)_m}{(1-q)^m} = (a)_m, \quad \lim_{q \rightarrow 1^-} (1-q)^{1-a} \frac{(q; q)_\infty}{(q^a; q)_\infty} = \Gamma(a).$$

Deduce

$$\lim_{q \rightarrow 1^-} {}_r\phi_{r-1} \left(\begin{matrix} q^{a_1}, \dots, q^{a_r} \\ q^{b_1}, \dots, q^{b_{r-1}} \end{matrix}; q, z \right) = {}_rF_{r-1} \left(\begin{matrix} a_1, \dots, a_r \\ b_1, \dots, b_{r-1} \end{matrix}; z \right).$$

Derive Gauss's summation from the q -Gauss summation. Prove

$$\lim_{q \rightarrow 1^-} (1-q)^{-3n} p_n \left(\frac{q^{ix} + q^{-ix}}{2}; q^a, q^b, q^c, q^d \mid q \right) = W_n(x^2; a, b, c, d).$$

Definition 28.6 (Root-system basic hypergeometric weights). Write $f(z^{\pm m}) = f(z^m)f(z^{-m})$ and

$$f(z_i^{\pm 1} z_j^{\pm 1}) = \prod_{\epsilon, \delta = \pm 1} f(z_i^\epsilon z_j^\delta).$$

For $|q|, |t|, |a_s| < 1$, define

$$\begin{aligned} \Delta_{BC_r}(z) &= \prod_{i=1}^r \frac{(z_i^{\pm 2}; q)_\infty}{\prod_{s=1}^4 (a_s z_i^{\pm 1}; q)_\infty} \\ &\times \prod_{1 \leq i < j \leq r} \frac{(z_i^{\pm 1} z_j^{\pm 1}; q)_\infty}{(t z_i^{\pm 1} z_j^{\pm 1}; q)_\infty}. \end{aligned}$$

The associated *Gustafson integral* uses the normalized torus measure

$$\frac{1}{2^r r!} \prod_{i=1}^r \frac{dz_i}{2\pi i z_i}.$$

Exercise 28.5 (Root systems and Gustafson integrals). Prove that Δ_{BC_r} is invariant under permutations and independent inversions of the z_i . Identify its factors with the roots $\pm e_i$, $\pm 2e_i$, and $\pm e_i \pm e_j$. For $r = 1$, recover the Askey–Wilson weight and its order-two Weyl normalization. Set $t = q^g$ with $g \in \mathbb{N}$, reduce the quotients of q -products to finite products, and compute the limit $q \rightarrow 1^-$ with $z_i = e^{i\theta_i}$.

29 Quantum Spherical Functions and qKZ Transport

Definition 29.1 (Quantum spherical functions). Let $H = U_q(\mathfrak{g})$ and let $B \subseteq H$ be a right coideal subalgebra, so $\Delta(B) \subseteq B \otimes H$. It plays the role of the stabilizer algebra of a quantum homogeneous space. If an H -module V contains a vector v satisfying

$$bv = \varepsilon(b)v \quad (b \in B),$$

then the matrix coefficient

$$\varphi_V(x) = \langle v, \pi_V(x)v \rangle$$

is a *quantum zonal spherical function*. It belongs to the coordinate Hopf dual of H and is invariant under the left and right quantum stabilizer actions.

Exercise 29.1 (Flatness, qKZ transport, and affine Hecke monodromy). Let a vector-valued function satisfy the adjacent exchange rule

$$\Psi(\dots, z_i, z_{i+1}, \dots) = \check{R}_i(z_i/z_{i+1})\Psi(\dots, z_{i+1}, z_i, \dots).$$

Show that consistency of the two routes reversing three adjacent variables is exactly the spectral Yang–Baxter equation. Add a periodic rule that returns z_1 at the end as qz_1 , and explain why translations enlarge the finite braid action to an affine one. If the R -matrices satisfy a Hecke quadratic, prove that the exchange monodromy factors through an affine Hecke algebra. Write the resulting reflection and translation operators in the same generators used in the polynomial representation defined above.

30 Elliptic Hypergeometric Functions and Ruijsenaars Theory

Definition 30.1 (Elliptic shifted factorials and gamma function). For $|p|, |q| < 1$, define

$$\theta(z; p) = (z; p)_\infty (p/z; p)_\infty, \quad (a; q, p)_m = \prod_{j=0}^{m-1} \theta(aq^j; p),$$

$$(a_1, \dots, a_s; q, p)_m = \prod_{i=1}^s (a_i; q, p)_m,$$

and

$$\Gamma(z; p, q) = \prod_{j,k \geq 0} \frac{1 - p^{j+1}q^{k+1}/z}{1 - p^j q^k z}.$$

Definition 30.2 (Hypergeometric deformation hierarchy). For a series $\sum_m c_m$, set $h_m = c_{m+1}/c_m$. It is *ordinary hypergeometric* when h_m is rational in m , and *basic hypergeometric* when h_m is rational in q^m . It is *elliptic hypergeometric* when, after writing $x = q^m$, its term ratio $h(x)$ is multiplicatively p -periodic:

$$h(px) = h(x).$$

Its parameters are *balanced* when the products of the numerator and denominator parameters required by this ellipticity agree. The deformation and degeneration orders are

$$\text{ordinary/additive} \longrightarrow \text{basic/multiplicative} \longrightarrow \text{elliptic},$$

$$\text{elliptic} \xrightarrow{p \rightarrow 0} \text{basic} \xrightarrow{q \rightarrow 1} \text{ordinary}.$$

Exercise 30.1 (Elliptic functional equations and degenerations). Prove

$$\theta(pz; p) = -z^{-1}\theta(z; p), \quad \Gamma(qz; p, q) = \theta(z; p)\Gamma(z; p, q),$$

$$\Gamma(pz; p, q) = \theta(z; q)\Gamma(z; p, q), \quad \Gamma(z; p, q)\Gamma(pq/z; p, q) = 1.$$

Prove

$$\theta(z; 0) = 1 - z, \quad (a; q, 0)_m = (a; q)_m, \quad \Gamma(z; 0, q) = \frac{1}{(z; q)_\infty}.$$

Definition 30.3 (Frenkel–Turaev series and elliptic beta integral). For $a^2q^{n+1} = bcde$, define

$${}_{10}V_9(a; b, c, d, e, q^{-n}; q, p) = \sum_{m=0}^n \frac{\theta(aq^{2m}; p)}{\theta(a; p)} \times \frac{(a, b, c, d, e, q^{-n}; q, p)_m}{(q, aq/b, aq/c, aq/d, aq/e, aq^{n+1}; q, p)_m} q^m.$$

For parameters $t_1, \dots, t_6$ satisfying

$$|t_j| < 1, \quad \prod_{j=1}^6 t_j = pq,$$

the *elliptic beta integral* is

$$\frac{(p; p)_\infty (q; q)_\infty}{4\pi i} \int_{|z|=1} \frac{\prod_{j=1}^6 \Gamma(t_j z^{\pm 1}; p, q)}{\Gamma(z^{\pm 2}; p, q)} \frac{dz}{z}.$$

Exercise 30.2 (Frenkel–Turaev and elliptic beta evaluations). Prove the Frenkel–Turaev summation

$${}_{10}V_9(a; b, c, d, e, q^{-n}; q, p) = \frac{(aq, aq/(bc), aq/(bd), aq/(cd); q, p)_n}{(aq/b, aq/c, aq/d, aq/(bcd); q, p)_n}.$$

Prove the elliptic beta evaluation

$$\frac{(p; p)_\infty (q; q)_\infty}{4\pi i} \int_{|z|=1} \frac{\prod_{j=1}^6 \Gamma(t_j z^{\pm 1}; p, q)}{\Gamma(z^{\pm 2}; p, q)} \frac{dz}{z} = \prod_{1 \leq j < k \leq 6} \Gamma(t_j t_k; p, q).$$

Take $p \rightarrow 0$ in both identities and identify the resulting terminating very-well-poised ${}_8\phi_7$ summation and basic hypergeometric beta integral. Obtain the Askey–Wilson integral by specialization.

Definition 30.4 (Elliptic Ruijsenaars operator). For type A_{r-1} , define

$$D_{p,q,t}^{\text{ell}} = \sum_{i=1}^r \left(\prod_{j \neq i} \frac{\theta(tx_i/x_j; p)}{\theta(x_i/x_j; p)} \right) T_{q,x_i}.$$

Exercise 30.3 (Differential, basic, and elliptic operator hierarchy). Construct the commuting higher elliptic Ruijsenaars operators. Prove

$$D_{p,q,t}^{\text{ell}} \longrightarrow D_{q,t} \quad (p \rightarrow 0).$$

Set $q = e^\varepsilon$ and $t = e^{g\varepsilon}$ and derive the Heckman–Opdam differential operator from the second-order term as $\varepsilon \rightarrow 0$. Take the rational scaling limit of its hyperbolic coefficients and obtain the rational Calogero–Moser operator. Organize the eigenfunction limits as

elliptic Ruijsenaars functions $\longrightarrow$ Macdonald polynomials $\longrightarrow$ Heckman–Opdam functions.

Exercise 30.4 (The Calogero–Sutherland limit). Write $x_i = e^{z_i}$, set $q = e^\epsilon$ and $t = e^{g\epsilon}$, and expand $D_{q,t}$ through order ϵ^2 . After subtracting its scalar value on 1 and removing the center-of-mass drift, obtain

$$\mathcal{L}_g = \sum_i \partial_{z_i}^2 + g \sum_{i < j} \coth\left(\frac{z_i - z_j}{2}\right) (\partial_{z_i} - \partial_{z_j}).$$

On the circle, put $z_i = i\theta_i$ and conjugate by

$$\Psi_0(\theta) = \prod_{i < j} \left| 2 \sin \frac{\theta_i - \theta_j}{2} \right|^g.$$

Recover, up to its ground-state energy, the Sutherland Hamiltonian

$$H_g = - \sum_i \partial_{\theta_i}^2 + \frac{g(g-1)}{2} \sum_{i < j} \frac{1}{\sin^2((\theta_i - \theta_j)/2)}.$$

Compute the eigenfunctions of degrees zero and one and, for $r = 2$, degree two. Identify their Sutherland eigenvalues.

Exercise 30.5 (Selberg measure and the Jacobi beta ensemble). For coupling $g > 0$, set $\beta = 2g$. For $a, b > 0$, take the BC_N Calogero–Sutherland ground state in Jacobi coordinates $x_i \in (0, 1)$,

$$\Psi_0(x) = C \prod_{i=1}^N x_i^{(a-1)/2} (1-x_i)^{(b-1)/2} \prod_{i < j} |x_i - x_j|^{\beta/2}.$$

Prove

$$|\Psi_0(x)|^2 = \frac{1}{Z_{N,\beta}^{\text{Jac}}(a,b)} \prod_{i=1}^N x_i^{a-1} (1-x_i)^{b-1} \prod_{i < j} |x_i - x_j|^\beta,$$

where the Selberg integral gives

$$Z_{N,\beta}^{\text{Jac}}(a,b) = \prod_{j=0}^{N-1} \frac{\Gamma(a + j\beta/2) \Gamma(b + j\beta/2) \Gamma(1 + (j+1)\beta/2)}{\Gamma(a+b + (N+j-1)\beta/2) \Gamma(1 + \beta/2)}.$$

Interpret $|\Psi_0|^2 dx_1 \cdots dx_N$ as the joint law of N random eigenvalues. For the type- A circular ground state, prove instead

$$|\Psi_0(\theta)|^2 \propto \prod_{i < j} |e^{i\theta_i} - e^{i\theta_j}|^\beta.$$

31 Elliptic Star–Triangle and Yang–Baxter Identities

Definition 31.1 (Star–triangle data). A family of edge weights $W_u(x, y)$, one-spin weights $S(x)$, and scalar normalizations $R(u, v, w)$ is *star–triangle data* when

$$\int S(x) W_u(a, x) W_v(b, x) W_w(c, x) dx = R(u, v, w) W_{u'}(b, c) W_{v'}(a, c) W_{w'}(a, b)$$

for its parameter constraint and contour. It is *elliptic* when its weights are finite products and ratios of $\Gamma(\text{cdot}; p, q)$.

Exercise 31.1 (Elliptic beta integral and star–triangle relation). Parametrize the six variables in the elliptic beta integral by three spectral parameters and three external spins. Derive an elliptic star–triangle identity from its evaluation. Apply the identity twice to a three-row lattice strip and prove equality of the two local reorderings that yields commuting transfer matrices. Take $p \rightarrow 0$ and then $q \rightarrow 1$ and derive the corresponding trigonometric and rational star–triangle relations. Compare this integral construction with the $6j$ -symbol/recoupling construction of the Yang–Baxter equation.

32 Root-of-Unity Degeneration and Tensor Categories

Definition 32.1 (Root-of-unity semisimplification). Fix $k \in \mathbb{N}$ and

$$q = e^{\pi i/(k+2)}.$$

Let $\mathcal{T}_k$ be the ribbon category of finite-dimensional type-1 tilting modules for the Lusztig integral form of $U_q(\mathfrak{sl}_2)$ specialized at q . A morphism $f : X \rightarrow Y$ is *negligible* when

$$\mathrm{tr}_q(gf) = 0 \quad \text{for every } g : Y \rightarrow X.$$

Let $\mathcal{N}_k$ be the negligible morphisms and define

$$\mathcal{C}_k = \mathrm{Kar}(\mathcal{T}_k/\mathcal{N}_k),$$

where Kar denotes idempotent completion.

Exercise 32.1 (Semisimplified $SU(2)_k$ fusion). Prove that $\mathcal{N}_k$ is a tensor ideal preserved by duals, braiding, and twist. Prove that $\mathcal{C}_k$ is semisimple with simple objects $V_0, \dots, V_k$ and that

$$V_a \otimes V_b \cong \bigoplus_{\substack{|a-b| \leq c \leq \min(a+b, 2k-a-b) \\ a+b+c \equiv 0 \pmod{2}}} V_c.$$

Compute

$$\dim_q(V_a) = \frac{\sin((a+1)\pi/(k+2))}{\sin(\pi/(k+2))}.$$

Definition 32.2 (Modular tensor categories). A *fusion category* is a semisimple rigid $\mathbb{C}$ -linear monoidal category with simple unit, finite-dimensional morphism spaces, and finitely many simple-object classes. A *modular tensor category* is a ribbon fusion category for which the matrix

$$\tilde{S}_{ij} = \mathrm{tr}_q(c_{X_j, X_i} c_{X_i, X_j})$$

on the simple objects $(X_i)_{i \in I}$ is invertible. Define

$$d_i = \dim_q(X_i), \quad \mathcal{D} = \left(\sum_{i \in I} d_i^2 \right)^{1/2}, \quad S = \mathcal{D}^{-1} \tilde{S}, \quad T_{ij} = \delta_{ij} \theta_i,$$

where $\theta_{X_i} = \theta_i \mathrm{id}_{X_i}$. It is *unitary* when it is a C^* -tensor category and its braiding and twist are unitary.

Exercise 32.2 (Modularity of $SU(2)_k$). Prove that $\mathcal{C}_k$ is a unitary modular tensor category. For $0 \leq a, b \leq k$, compute

$$S_{ab} = \sqrt{\frac{2}{k+2}} \sin\left(\frac{(a+1)(b+1)\pi}{k+2}\right), \quad \theta_a = \exp\left(\frac{\pi i a(a+2)}{2(k+2)}\right).$$

Prove directly that S is unitary and invertible.

Part III

Counting

33 Partitions and Generating Functions

33.1 Partition generating functions and finite boxes

Definition 33.1 (Partitions and q -Pochhammer symbols). An *integer partition* of $n \geq 0$ is a finite nonincreasing sequence $\lambda = (\lambda_1, \dots, \lambda_r)$ of positive integers with $|\lambda| = \sum_i \lambda_i = n$. Let $p(n)$ be the number of partitions, with $p(0) = 1$. For a formal variable q , set

$$(a; q)_n = \prod_{j=0}^{n-1} (1 - aq^j), \quad (a; q)_\infty = \prod_{j=0}^{\infty} (1 - aq^j).$$

Analytically, the infinite product converges for $|q| < 1$. Euler's partition generating function is

$$P(q) = \sum_{n \geq 0} p(n)q^n = \frac{1}{(q; q)_\infty}.$$

Exercise 33.1 (Bosonic and fermionic occupation numbers). Consider oscillator modes with one-particle energies $\hbar\omega, 2\hbar\omega, 3\hbar\omega, \dots$. Show that a bosonic Fock state of excitation energy $n\hbar\omega$ is a partition of n , and derive

$$\prod_{m \geq 1} \frac{1}{1 - q^m} = \sum_{n \geq 0} p(n)q^n.$$

For fermionic occupation numbers in $\{0, 1\}$, show that $\prod_{m \geq 1} (1 + q^m)$ counts partitions into distinct parts. Construct a bijection, by separating powers of two in the multiplicities, between partitions into distinct parts and partitions into odd parts, and conclude

$$\prod_{m \geq 1} (1 + q^m) = \prod_{m \geq 1} \frac{1}{1 - q^{2m-1}}.$$

Interpret the two products as different quasiparticle descriptions with the same exact degeneracies.

Definition 33.2 (Gaussian binomial coefficients). For $0 \leq r \leq n$, the *Gaussian binomial coefficient* is

$$\begin{bmatrix} n \\ r \end{bmatrix}_q = \frac{(q; q)_n}{(q; q)_r (q; q)_{n-r}},$$

and it is zero outside this range.

Exercise 33.2 (Gaussian binomial theorem). Derive the recurrence

$$\begin{bmatrix} n \\ r \end{bmatrix}_q = \begin{bmatrix} n-1 \\ r \end{bmatrix}_q + q^{n-r} \begin{bmatrix} n-1 \\ r-1 \end{bmatrix}_q.$$

Prove that $\begin{bmatrix} n \\ r \end{bmatrix}_q$ is a polynomial in q with nonnegative integer coefficients and that the coefficient of q^m counts partitions of m whose Young diagrams fit inside an $r \times (n-r)$ rectangle.

33.2 Rogers–Ramanujan identities and exclusion rules

Definition 33.3 (Rogers–Ramanujan sum and product series). For $|q| < 1$, define

$$G_{\text{sum}}(q) = \sum_{r \geq 0} \frac{q^{r^2}}{(q; q)_r}, \quad H_{\text{sum}}(q) = \sum_{r \geq 0} \frac{q^{r(r+1)}}{(q; q)_r},$$

$$G_{\text{prod}}(q) = \frac{1}{(q; q^5)_\infty (q^4; q^5)_\infty}, \quad H_{\text{prod}}(q) = \frac{1}{(q^2; q^5)_\infty (q^3; q^5)_\infty}.$$

Exercise 33.3 (Difference conditions and Rogers–Ramanujan counting). Fix the number r of parts. Subtract the staircase $(2r - 1, 2r - 3, \dots, 1)$ to prove that $q^{r^2}/(q; q)_r$ generates partitions with exactly r parts whose successive parts differ by at least two. Use the staircase $(2r, 2r - 2, \dots, 2)$ for the same difference condition with smallest part at least two. Interpret the product sides as partitions into parts congruent to 1 or 4 modulo 5, and to 2 or 3 modulo 5, respectively. Compute all four series through q^{30} using finite products and finite sums, and verify coefficientwise that $G_{\text{sum}} = G_{\text{prod}}$ and $H_{\text{sum}} = H_{\text{prod}}$ to that order. Translate the coefficient equalities into finite tests of the two partition identities and identify which further step would be required for a proof in all degrees.

34 Characters and Exact State Counting

34.1 Virasoro representations and character formulas

Definition 34.1 (The Virasoro algebra and its characters). The *Virasoro algebra* has generators L_n for $n \in \mathbb{Z}$ and a central element acting by c , with

$$[L_m, L_n] = (m - n)L_{m+n} + \frac{c}{12}(m^3 - m)\delta_{m+n,0}.$$

A highest-weight vector $|h\rangle$ satisfies $L_0|h\rangle = h|h\rangle$ and $L_n|h\rangle = 0$ for $n > 0$. For a positive-energy module H_a with finite-dimensional L_0 eigenspaces, its *character* is

$$\chi_a(\tau) = \text{Tr}_{H_a} q^{L_0 - c/24}.$$

The shift by $-c/24$ is the Casimir energy produced by mapping the plane to the cylinder.

Exercise 34.1 (Descendants, partitions, and null states). Use the Virasoro commutation relations to move positive modes to the right and prove directly the triangular Poincaré–Birkhoff–Witt statement that the ordered monomials form a basis of the Verma module:

$$L_{-\lambda_1} \cdots L_{-\lambda_r} |h\rangle, \quad \lambda_1 \geq \cdots \geq \lambda_r \geq 1.$$

Before quotienting by null vectors, deduce the Verma-module character

$$\chi_{c,h}^{\text{Verma}}(\tau) = \frac{q^{h-c/24}}{(q; q)_\infty}.$$

For the vacuum, impose $L_{-1}|0\rangle = 0$ and find the resulting generic vacuum descendant product. Explain why additional null vectors subtract states and turn unrestricted partition numbers into the more structured q -series of minimal models.

34.2 Modular invariance and exact state counting

Definition 34.2 (Rational characters and modular invariants). A finite family of positive-energy sectors H_a is a *rational modular character system* when its characters span a representation of the modular group,

$$\chi_a(-1/\tau) = \sum_b S_{ab} \chi_b(\tau), \quad \chi_a(\tau+1) = \sum_b T_{ab} \chi_b(\tau).$$

A full torus partition function has the form

$$Z(\tau, \bar{\tau}) = \sum_{a,b} M_{ab} \chi_a(\tau) \overline{\chi_b(\tau)}$$

with nonnegative integer multiplicities. It is modular invariant when M commutes with S and T .

Exercise 34.2 (Exact state counting in the Ising CFT). For the Ising theory with $c = 1/2$ and primary weights $0, 1/2, 1/16$, take

$$\begin{aligned} \chi_0 &= \frac{1}{2} \left(\sqrt{\frac{\vartheta_3}{\eta}} + \sqrt{\frac{\vartheta_4}{\eta}} \right), \\ \chi_{1/2} &= \frac{1}{2} \left(\sqrt{\frac{\vartheta_3}{\eta}} - \sqrt{\frac{\vartheta_4}{\eta}} \right), \\ \chi_{1/16} &= \frac{1}{\sqrt{2}} \sqrt{\frac{\vartheta_2}{\eta}}. \end{aligned}$$

Expand each as $q^{h-c/24}$ times a power series and interpret its first coefficients as exact descendant degeneracies. Derive

$$S = \frac{1}{2} \begin{pmatrix} 1 & 1 & \sqrt{2} \\ 1 & 1 & -\sqrt{2} \\ \sqrt{2} & -\sqrt{2} & 0 \end{pmatrix}, \quad T_{aa} = e^{2\pi i(h_a - c/24)}.$$

Verify the modular invariance of $|\chi_0|^2 + |\chi_{1/2}|^2 + |\chi_{1/16}|^2$. Using the Verlinde formula

$$N_{ab}^c = \sum_x \frac{S_{ax} S_{bx} \overline{S_{cx}}}{S_{0x}},$$

recover the Ising fusion rules. Relate these towers to the critical Ising spectroscopy exercise in Quantum Topology.

Exercise 34.3 (The Cardy density of states). Let a modular-invariant unitary CFT have discrete spectrum, unique vacuum, and left and right central charges $c, \bar{c}$. Set $\tau = i\beta/(2\pi)$ and use S -invariance to show that the high-temperature partition function is controlled by the vacuum at inverse temperature $4\pi^2/\beta$. A saddle-point inverse Laplace transform then gives, for large left and right excitation numbers,

$$\log \rho(N, \bar{N}) \sim 2\pi \sqrt{\frac{cN}{6}} + 2\pi \sqrt{\frac{\bar{c}\bar{N}}{6}}.$$

State how c is replaced by $c_{\text{eff}} = c - 24h_{\text{min}}$ in a nonunitary theory. For a nonrotating state with $c = \bar{c}$ and $N = \bar{N}$, recover the thermal entropy used in the BTZ exercise and hence reproduce the horizon entropy from boundary-state counting; see Cardy, *Nuclear Physics B* **270** (1986).

34.3 Integrable lattice models

Definition 34.3 (Transfer matrices and integrability). For a two-dimensional lattice model on a cylinder, a *row transfer matrix* $T_N(u)$ propagates one width- N row to the next and depends on a spectral parameter u . The model is *integrable* when it admits a nontrivial commuting family

$$[T_N(u), T_N(v)] = 0 \quad \text{for every } u, v,$$

usually obtained from a Yang–Baxter or star–triangle relation. Commuting transfer matrices permit simultaneous spectral analysis, but integrability alone does not guarantee that every finite-size degeneracy has a simple closed formula.

Exercise 34.4 (Hard hexagons and exact finite-volume counting). In the hard-hexagon model, occupied vertices of a triangular lattice may not be adjacent. On an $N \times M$ torus define

$$Z_{N,M}(z) = \sum_{I \text{ independent}} z^{|I|}.$$

Index a row transfer matrix by cyclic binary rows and set its matrix element to zero when two consecutive rows violate nearest-neighbor exclusion, and otherwise to $z^{(|a|+|b|)/2}$. Prove $Z_{N,M}(z) = \text{Tr}(T_N(z)^M)$ and compute the polynomial for several small sizes. Show that the coefficients are exact state counts, whereas the largest eigenvalue controls the thermodynamic free energy.

Verify the algebraic identity

$$z_c = \frac{11 + 5\sqrt{5}}{2} = \left(\frac{1 + \sqrt{5}}{2} \right)^5,$$

and, for the finite matrices already constructed, evaluate the largest eigenvalue and its first two z -derivatives near z_c . Plot their change with N and explain why finite matrices remain analytic even when their large-size limit may develop a singularity.

35 Probability Spaces and Convergence

Definition 35.1 (Probability spaces and convergence). A *probability space* is a measure space $(\Omega, \mathcal{F}, \mathbb{P})$ with $\mathbb{P}(\Omega) = 1$. A *random variable* is a measurable map $X : \Omega \rightarrow Y$, its *law* is $\mathbb{P}_X(B) = \mathbb{P}(X^{-1}(B))$, and, when it is scalar-valued and integrable, its *expectation* is its integral. A random vector $X = (X_1, \dots, X_N)$ has *joint density* p with respect to a measure μ when

$$\mathbb{P}(X \in B) = \int_B p d\mu.$$

The Gaussian law $N(m, v)$ on $\mathbb{R}$ has density

$$\frac{1}{\sqrt{2\pi v}} \exp \left[-\frac{(x - m)^2}{2v} \right].$$

A sequence of random variables in a metric space (Y, d) *converges in probability* to X when

$$\mathbb{P}(d(X_N, X) > \varepsilon) \longrightarrow 0 \quad (\varepsilon > 0).$$

It *converges in distribution*, written $X_N \implies X$, when

$$\int f d\mathbb{P}_{X_N} \longrightarrow \int f d\mathbb{P}_X$$

for every bounded continuous f .

36 Random Matrices and Statistical Universality

Definition 36.1 (Scalar determinants, Vandermonde determinant, and Dyson classes). For $A = (a_{ij}) \in M_N(\mathbb{C})$, define

$$\det A = \sum_{\sigma \in S_N} \operatorname{sgn}(\sigma) \prod_{i=1}^N a_{i, \sigma(i)}.$$

For $\lambda = (\lambda_1, \dots, \lambda_N)$, define

$$\Delta(\lambda) = \det[\lambda_i^{j-1}]_{i,j=1}^N = \prod_{i < j} (\lambda_j - \lambda_i).$$

Let $\mathbb{F} = \mathbb{R}, \mathbb{C}$, or $\mathbb{H}$ and set

$$\beta = \dim_{\mathbb{R}} \mathbb{F} \in \{1, 2, 4\}.$$

The Gaussian orthogonal, unitary, and symplectic ensembles, denoted GOE, GUE, and GSE, consist of the self-adjoint matrices $H = H^*$ over $\mathbb{R}, \mathbb{C}$, and $\mathbb{H}$, respectively, with density

$$d\mathbb{P}_{N,\beta}(H) = C_{N,\beta} \exp\left[-\frac{\beta N}{4\sigma^2} \operatorname{Tr}(H^2)\right] dH.$$

Here H^* is conjugate transpose and dH is Lebesgue measure on the independent real coordinates.

Exercise 36.1 (Eigenvalue–eigenvector Jacobian). Diagonalize $H = U \operatorname{diag}(\lambda) U^{-1}$ in each Dyson class. Use the invariant metric $\operatorname{Tr}(dH^2)$ to prove

$$dH = C |\Delta(\lambda)|^\beta \prod_{i=1}^N d\lambda_i d\mu(U),$$

with the stabilizer and permutation factors included in C . Integrate over the eigenvectors and derive

$$p_{N,\beta}(\lambda) = \frac{1}{Z_{N,\beta}} \exp\left[-\frac{\beta N}{4\sigma^2} \sum_{i=1}^N \lambda_i^2\right] \prod_{i < j} |\lambda_i - \lambda_j|^\beta.$$

Definition 36.2 (Beta ensembles and Coulomb gases). For an interval $I \subseteq \mathbb{R}$, a confining potential V , and $\beta > 0$, the *beta ensemble* is

$$d\mathbb{P}_{N,\beta,V}(\lambda) = \frac{1}{Z_{N,\beta,V}} \prod_{i < j} |\lambda_i - \lambda_j|^\beta \prod_{i=1}^N e^{-\beta N V(\lambda_i)/2} d\lambda_i.$$

Its *Coulomb-gas Hamiltonian* is

$$\mathcal{H}_N(\lambda) = \frac{N}{2} \sum_{i=1}^N V(\lambda_i) - \sum_{i < j} \log |\lambda_i - \lambda_j|, \quad d\mathbb{P}_{N,\beta,V} = Z^{-1} e^{-\beta \mathcal{H}_N} d\lambda.$$

The Jacobi beta ensemble is the Selberg law on $[0, 1]^N$, and the circular beta ensemble is

$$\frac{1}{Z_{N,\beta}^{\text{circ}}} \prod_{i < j} |e^{i\theta_i} - e^{i\theta_j}|^\beta \prod_{i=1}^N \frac{d\theta_i}{2\pi}.$$

Exercise 36.2 (Logarithmic equilibrium and the semicircle law). For the empirical measure $L_N = N^{-1} \sum_i \delta_{\lambda_i}$, derive the mean-field energy

$$\mathcal{I}_V(\mu) = \int V(x) d\mu(x) - \iint \log |x - y| d\mu(x) d\mu(y).$$

Prove that its minimizer satisfies

$$V(x) - 2 \int \log |x - y| d\mu_V(y) = \ell$$

on its support. For $V(x) = x^2/(2\sigma^2)$, solve the singular integral equation and obtain

$$\rho_{\text{sc}}(x) = \frac{1}{2\pi\sigma^2} \sqrt{4\sigma^2 - x^2} \mathbf{1}_{[-2\sigma, 2\sigma]}(x).$$

Definition 36.3 (Wigner matrices and empirical spectral measures). A *Wigner matrix* H_N is an $N \times N$ Hermitian random matrix whose upper-triangular entries are independent and centered, with $\mathbb{E}|H_{ij}|^2 = \sigma^2/N$ off the diagonal and, for every fixed m ,

$$\sup_{N,i,j} \mathbb{E} |\sqrt{N} H_{ij}|^m < \infty.$$

Its *empirical spectral measure* is

$$L_{H_N} = \frac{1}{N} \sum_{j=1}^N \delta_{\lambda_j(H_N)}, \quad \int x^k dL_{H_N}(x) = \frac{1}{N} \text{Tr}(H_N^k).$$

Exercise 36.3 (Wigner semicircle law). Expand $N^{-1} \mathbb{E} \text{Tr}(H_N^k)$ as a sum over closed index walks. Prove that only noncrossing pair-matched walks survive as $N \rightarrow \infty$ and that their number is the Catalan number. Prove that the variance tends to zero and conclude that L_{H_N} converges in probability, in moments, to $\rho_{\text{sc}}(x)dx$.

Definition 36.4 (Unfolded spectra and level repulsion). For a smooth approximation $\bar{N}(E)$ to the cumulative level count, define

$$x_j = \bar{N}(E_j), \quad s_j = x_{j+1} - x_j.$$

The variables x_j are the *unfolded levels*. Their local mean spacing is one. Exact symmetry sectors are unfolded separately. The factor $|\Delta(\lambda)|^\beta$ defines *level repulsion* of order β .

Exercise 36.4 (Wigner surmises and repulsion exponents). For 2×2 GOE and GUE matrices, separate the trace from the traceless coordinates and compute the radial distribution of the eigenvalue gap. After rescaling to unit mean spacing, derive

$$P_1(s) = \frac{\pi}{2} s e^{-\pi s^2/4}, \quad P_2(s) = \frac{32}{\pi^2} s^2 e^{-4s^2/\pi}.$$

Use the beta-ensemble density to prove

$$P_\beta(s) \sim c_\beta s^\beta \quad (s \downarrow 0),$$

and compare it with the Poisson spacing law e^{-s} .

Definition 36.5 (Global, bulk, and edge scales). Order the eigenvalues as $\lambda_1 \leq \dots \leq \lambda_N$. Define the classical location γ_j by

$$\frac{j}{N} = \int_{-2\sigma}^{\gamma_j} \rho_{\text{sc}}(x) dx.$$

At $E \in (-2\sigma, 2\sigma)$, the bulk coordinate is

$$u = N\rho_{\text{sc}}(E)(\lambda - E),$$

and at the upper soft edge it is

$$\xi = \frac{N^{2/3}}{\sigma}(\lambda - 2\sigma).$$

Exercise 36.5 (Bulk and edge scales). Use the expected count $N\rho_{\text{sc}}(E)dE$ to derive the bulk spacing scale $[N\rho_{\text{sc}}(E)]^{-1}$. Prove

$$\rho_{\text{sc}}(2\sigma - s) \sim \frac{1}{\pi\sigma^{3/2}}\sqrt{s} \quad (s \downarrow 0),$$

and derive the edge scale $s \asymp \sigma N^{-2/3}$.

Definition 36.6 (Airy kernel and the Tracy–Widom law). Let Ai be the solution of $y'' = xy$ decaying at $+\infty$. Define

$$K_{\text{Ai}}(x, y) = \frac{\text{Ai}(x)\text{Ai}'(y) - \text{Ai}'(x)\text{Ai}(y)}{x - y}$$

by continuity on the diagonal. The GUE Tracy–Widom distribution is

$$F_2(s) = \det(I - K_{\text{Ai}})_{L^2(s, \infty)}.$$

36.1 Correlation functions and point processes

Definition 36.7 (Point processes and correlation functions). Let (X, μ) be a locally compact measure space. A *simple point process* is a random locally finite counting measure

$$\Xi = \sum_i \delta_{x_i}$$

with distinct points almost surely. Its k -point correlation function ρ_k is defined by

$$\mathbb{E} \sum_{i_1, \dots, i_k}^{\neq} f(x_{i_1}, \dots, x_{i_k}) = \int_{X^k} f(x_1, \dots, x_k) \rho_k(x_1, \dots, x_k) \prod_{j=1}^k d\mu(x_j).$$

For a symmetric joint density p_N of exactly N points,

$$\rho_k(x_1, \dots, x_k) = \frac{N!}{(N-k)!} \int_{X^{N-k}} p_N(x_1, \dots, x_N) \prod_{j=k+1}^N d\mu(x_j).$$

Exercise 36.6 (Factorial moments and gap probabilities). For a relatively compact measurable set B and $N_B = \Xi(B)$, prove, whenever either side is finite,

$$\mathbb{E}(N_B)_k = \int_{B^k} \rho_k d\mu^{\otimes k}, \quad (N_B)_k = N_B(N_B - 1) \cdots (N_B - k + 1).$$

If $\mathbb{E}[2^{N_B}] < \infty$, use inclusion–exclusion and justify the exchange of expectation and summation to prove

$$\mathbb{P}(N_B = 0) = \sum_{k=0}^{\infty} \frac{(-1)^k}{k!} \int_{B^k} \rho_k d\mu^{\otimes k}.$$

36.2 Determinantal/Pfaffian processes and Fredholm determinants

Definition 36.8 (Determinantal and Pfaffian point processes). For an antisymmetric matrix $A = (a_{ij}) \in M_{2m}(\mathbb{C})$, define

$$\text{Pf}(A) = \frac{1}{2^m m!} \sum_{\sigma \in S_{2m}} \text{sgn}(\sigma) \prod_{j=1}^m a_{\sigma(2j-1), \sigma(2j)}.$$

A point process is *determinantal* with kernel K when

$$\rho_k(x_1, \dots, x_k) = \det[K(x_i, x_j)]_{i,j=1}^k.$$

It is *Pfaffian* with antisymmetric 2×2 matrix kernel $\mathbb{K}$ when

$$\rho_k(x_1, \dots, x_k) = \text{Pf}[\mathbb{K}(x_i, x_j)]_{i,j=1}^k.$$

For a trace-class integral operator K_B on $L^2(B, \mu)$, its *Fredholm determinant* is

$$\det(I + zK_B) = \sum_{k=0}^{\infty} \frac{z^k}{k!} \int_{B^k} \det[K(x_i, x_j)]_{i,j=1}^k d\mu^{\otimes k}.$$

The Fredholm Pfaffian is defined by the analogous series with Pfaffians of the matrix kernel. A *positive contraction* is an operator K satisfying $0 \leq K \leq I$.

Exercise 36.7 (Counting statistics and Fredholm formulas). For a determinantal process, prove

$$\mathbb{E}[z^{N_B}] = \det(I + (z-1)K_B), \quad \mathbb{P}(N_B = 0) = \det(I - K_B).$$

Derive the corresponding Fredholm-Pfaffian formula for a Pfaffian process. If K_B is a positive trace-class contraction with eigenvalues (λ_j) , prove

$$\det(I + zK_B) = \prod_j (1 + z\lambda_j)$$

and that N_B is distributed as a sum of independent Bernoulli variables with parameters λ_j .

36.3 Orthogonal-polynomial ensembles

Definition 36.9 (Orthogonal-polynomial ensembles). Let $w(x)d\mu(x)$ be a measure with finite moments and let p_j be its monic orthogonal polynomials,

$$\int p_j(x)p_k(x)w(x)d\mu(x) = h_j\delta_{jk}.$$

The associated *orthogonal-polynomial ensemble* has density

$$p_N(x_1, \dots, x_N) = \frac{1}{Z_N} \Delta(x)^2 \prod_{i=1}^N w(x_i), \quad Z_N = N! \prod_{j=0}^{N-1} h_j.$$

Its correlation kernel is

$$K_N(x, y) = \sqrt{w(x)w(y)} \sum_{j=0}^{N-1} \frac{p_j(x)p_j(y)}{h_j}.$$

Exercise 36.8 (Andréief identity and determinantal correlations). Prove Andréief's identity

$$\int_{X^N} \det[f_i(x_j)] \det[g_i(x_j)] \prod_{j=1}^N d\mu(x_j) = N! \det \left[\int_X f_i(x) g_j(x) d\mu(x) \right].$$

Use the Vandermonde determinant to prove

$$p_N(x_1, \dots, x_N) = \frac{1}{N!} \det[K_N(x_i, x_j)]_{i,j=1}^N$$

and

$$\rho_k(x_1, \dots, x_k) = \det[K_N(x_i, x_j)]_{i,j=1}^k.$$

Exercise 36.9 (Skew-orthogonal ensembles). For an antisymmetric bilinear form $\langle \cdot, \cdot \rangle_s$, construct skew-orthogonal polynomials satisfying

$$\langle p_{2j}, p_{2k+1} \rangle_s = r_j \delta_{jk}, \quad \langle p_{2j}, p_{2k} \rangle_s = \langle p_{2j+1}, p_{2k+1} \rangle_s = 0.$$

Prove de Bruijn's Pfaffian integration identity and use it to derive the matrix correlation kernels for the GOE and GSE.

36.4 Christoffel–Darboux and integrable kernels

Definition 36.10 (Christoffel–Darboux and integrable kernels). Let $q_j = p_j / \sqrt{h_j}$ and let κ_j be the leading coefficient of q_j . The Christoffel–Darboux kernel is

$$K_N(x, y) = \sqrt{w(x)w(y)} \frac{\kappa_{N-1}}{\kappa_N} \frac{q_N(x)q_{N-1}(y) - q_{N-1}(x)q_N(y)}{x - y}.$$

An *integrable kernel* is a kernel of the form

$$K(x, y) = \frac{f(x)^\top g(y)}{x - y}, \quad f(x)^\top g(x) = 0.$$

Exercise 36.10 (Projection and resolvent identities). Derive the Christoffel–Darboux formula from the three-term recurrence. Prove

$$\int K_N(x, z) K_N(z, y) d\mu(z) = K_N(x, y), \quad \int K_N(x, x) d\mu(x) = N.$$

For an integrable trace-class kernel K , prove that the resolvent $R = K(I - K)^{-1}$ is again integrable and express its numerator using $(I - K)^{-1}f$ and $(I - K^\top)^{-1}g$.

36.5 Bulk, soft-edge, and hard-edge universality

Definition 36.11 (Universal limiting kernels). The bulk, soft-edge, and hard-edge kernels are respectively

$$K_{\sin}(u, v) = \frac{\sin \pi(u - v)}{\pi(u - v)},$$

$$K_{\text{Ai}}(u, v) = \frac{\text{Ai}(u) \text{Ai}'(v) - \text{Ai}'(u) \text{Ai}(v)}{u - v},$$

and, for $\alpha > -1$,

$$K_{\text{Bes}}^{(\alpha)}(u, v) = \frac{J_\alpha(\sqrt{u})\sqrt{v} J'_\alpha(\sqrt{v}) - \sqrt{u} J'_\alpha(\sqrt{u}) J_\alpha(\sqrt{v})}{2(u - v)}.$$

The values on the diagonal are defined by continuity.

Exercise 36.11 (Classical kernel universality). Use the Christoffel–Darboux formula and the asymptotics of the Hermite, Laguerre, and Jacobi polynomials to prove convergence of their rescaled kernels, where applicable, to K_{sin} in the bulk, K_{Ai} at a soft edge, and $K_{\text{Bes}}^{(\alpha)}$ at a hard edge. Deduce convergence of every correlation function and of the associated Fredholm gap probabilities. Recover the Tracy–Widom law for the largest GUE eigenvalue.

37 Integrable Probability and KPZ Universality

37.1 RSK and last-passage percolation

Definition 37.1 (Partitions, tableaux, and Schur functions). A partition is a sequence $\lambda_1 \geq \lambda_2 \geq \dots \geq 0$ with finite sum $|\lambda| = \sum_i \lambda_i$. A semistandard Young tableau of shape λ has weakly increasing rows and strictly increasing columns. For variables $a = (a_1, \dots, a_m)$, define

$$s_\lambda(a) = \sum_T \prod_{i=1}^m a_i^{m_i(T)},$$

where $m_i(T)$ is the number of entries equal to i .

Definition 37.2 (RSK correspondence and last-passage time). The Robinson–Schensted–Knuth construction associates

$$A = (a_{ij}) \mapsto (P, Q)$$

to a matrix of nonnegative integers a pair of semistandard Young tableaux of the same shape λ , with the row and column sums of A giving the contents of P and Q . For weights $a_{ij} \geq 0$, define

$$G(m, n) = \max_{\pi: (1,1) \nearrow (m,n)} \sum_{(i,j) \in \pi} a_{ij}.$$

Exercise 37.1 (Schensted and Greene theorems). Prove the RSK bijection and Greene’s theorem. For $A \in M_{m,n}(\mathbb{Z}_{\geq 0})$, deduce

$$\lambda_1 = G(m, n)$$

and identify $\lambda_1 + \dots + \lambda_r$ with the maximal weight of r disjoint up-right paths. For independent geometric weights with parameters $a_i b_j$, use RSK to derive

$$\mathbb{P}(\lambda) = \prod_{i,j} (1 - a_i b_j) s_\lambda(a) s_\lambda(b)$$

and prove the Cauchy identity

$$\sum_{\lambda} s_\lambda(a) s_\lambda(b) = \prod_{i,j} (1 - a_i b_j)^{-1}.$$

37.2 Schur and Macdonald processes

Definition 37.3 (Schur measures and processes). For $\mu \subseteq \lambda$, the *skew Schur function* $s_{\lambda/\mu}$ is the tableau sum over the skew diagram $\lambda \setminus \mu$. A *specialization* is an algebra homomorphism from the

algebra of symmetric functions to $\mathbb{C}$; it is *Schur-positive* when $s_{\lambda/\mu}(\rho) \geq 0$ for all $\mu \subseteq \lambda$. The *Schur measure* associated to two Schur-positive specializations ρ^+, ρ^- is

$$\mathbb{P}(\lambda) = \frac{s_\lambda(\rho^+)s_\lambda(\rho^-)}{\sum_\nu s_\nu(\rho^+)s_\nu(\rho^-)}.$$

An ascending *Schur process* is a probability measure on $\lambda^{(1)} \subseteq \dots \subseteq \lambda^{(m)}$ proportional to

$$s_{\lambda^{(1)}}(\rho_1^+) \prod_{r=2}^m s_{\lambda^{(r)}/\lambda^{(r-1)}}(\rho_r^+) s_{\lambda^{(m)}}(\rho^-).$$

Exercise 37.2 (Schur processes as determinantal processes). Encode a partition by the Maya diagram

$$\mathcal{X}(\lambda) = \{\lambda_i - i + \frac{1}{2} : i \geq 1\}.$$

Use the Cauchy and skew-Cauchy identities to prove that the induced space-time point process is determinantal. Derive its double-contour correlation kernel and express the distribution of λ_1 as a Fredholm determinant. Recover geometric last-passage percolation as a one-time marginal.

Definition 37.4 (Macdonald measures and processes). Let $P_\lambda(\cdot; q, t)$ be the Macdonald polynomials and let $Q_\lambda = b_\lambda(q, t)P_\lambda$ be their dual basis. Define the skew functions by

$$P_\lambda(x, y) = \sum_\mu P_{\lambda/\mu}(x)P_\mu(y), \quad Q_\lambda(x, y) = \sum_\mu Q_{\lambda/\mu}(x)Q_\mu(y).$$

For finite specializations a, b , define

$$\Pi(a; b) = \prod_{i,j} \frac{(ta_i b_j; q)_\infty}{(a_i b_j; q)_\infty}.$$

The *Macdonald measure* is

$$\mathbb{P}_{q,t}(\lambda) = \frac{P_\lambda(a; q, t)Q_\lambda(b; q, t)}{\Pi(a; b)}.$$

The ascending *Macdonald process* is

$$\frac{P_{\lambda^{(1)}}(a^{(1)}) \prod_{r=2}^m P_{\lambda^{(r)}/\lambda^{(r-1)}}(a^{(r)}) Q_{\lambda^{(m)}}(b)}{\Pi(a^{(1)}, \dots, a^{(m)}; b)}.$$

Exercise 37.3 (Macdonald observables and degenerations). Apply the commuting Macdonald difference operators to the Cauchy identity and derive contour-integral formulas for the joint moments of

$$\sum_i q^{\lambda_i} t^{N-i}.$$

Prove that $q = t$ gives the Schur process, $q = 0$ gives the Hall–Littlewood process, and $t = 0$ gives the q -Whittaker process. Obtain Jack processes from the limit $q = t^\alpha$, $t \uparrow 1$.

37.3 Dyson Brownian motion and KPZ-type dynamics

Definition 37.5 (Brownian motion and space-time white noise). A *standard Brownian motion* is a continuous real process $(B_t)_{t \geq 0}$ with $B_0 = 0$ and independent increments satisfying

$$B_t - B_s \sim N(0, t - s) \quad (0 \leq s < t).$$

Itô differentials satisfy

$$dB_i dB_j = \delta_{ij} dt, \quad dB_i dt = dt^2 = 0.$$

Space-time white noise on $\mathbb{R}_+ \times \mathbb{R}$ is the centered Gaussian generalized process ξ satisfying

$$\mathbb{E}[\xi(\varphi)\xi(\psi)] = \int_{\mathbb{R}_+ \times \mathbb{R}} \varphi(t, x)\psi(t, x) dt dx$$

for test functions φ, ψ .

Definition 37.6 (Dyson Brownian motion). For $\beta \geq 1$ and a confining potential V , *Dyson Brownian motion* is the interacting diffusion

$$d\lambda_i = \sqrt{\frac{2}{\beta N}} dB_i + \left(\frac{1}{N} \sum_{j \neq i} \frac{1}{\lambda_i - \lambda_j} - \frac{1}{2} V'(\lambda_i) \right) dt.$$

Its reversible density is

$$Z^{-1} \prod_{i < j} |\lambda_i - \lambda_j|^\beta \prod_i e^{-\beta N V(\lambda_i)/2}.$$

Exercise 37.4 (Matrix diffusion and eigenvalue dynamics). Apply Itô calculus to Hermitian and real-symmetric Ornstein–Uhlenbeck matrix processes and derive Dyson Brownian motion for $\beta = 2$ and $\beta = 1$. Prove noncollision, compute the generator in divergence form, and verify the stated reversible measure. Derive the complex Burgers equation for the large- N Stieltjes transform and, in the unitary case, the sine- and Airy-kernel local equilibria.

Definition 37.7 (TASEP and the KPZ equation). In the totally asymmetric simple exclusion process, particles on $\mathbb{Z}$ attempt rate-one jumps from x to $x + 1$, and a jump is suppressed when $x + 1$ is occupied. Its integrated current defines a height function. The one-dimensional KPZ equation is

$$\partial_t h = \nu \partial_x^2 h + \frac{\lambda}{2} (\partial_x h)^2 + \sqrt{D} \xi,$$

where ξ is space-time white noise. Its Cole–Hopf solution is $h = (2\nu/\lambda) \log Z$, where Z solves the corresponding stochastic heat equation with multiplicative noise.

Exercise 37.5 (Last passage, particle dynamics, and KPZ scaling). Use the recursion

$$G(m, n) = \max\{G(m-1, n), G(m, n-1)\} + a_{mn}$$

to identify exponential last-passage times with TASEP currents. For i.i.d. exponential weights, prove

$$\frac{G(n, n) - 4n}{2^{4/3} n^{1/3}} \Longrightarrow \chi_2, \quad \mathbb{P}(\chi_2 \leq s) = F_2(s).$$

Derive the 1:2:3 KPZ fluctuation scaling and its spatial Airy process limit.

Exercise 37.6 (Macdonald dynamics and KPZ degenerations). At $t = 0$, intertwine the q -Whittaker difference operators with the q -TASEP generator. Derive a Fredholm-determinant formula for the q -Laplace transform of a particle position. Take the $q \uparrow 1$ limit to the O’Connell–Yor semi-discrete polymer and then its intermediate-disorder limit to the stochastic heat equation and the KPZ equation. Identify the corresponding degenerations of the Macdonald-process observables.

38 Free Probability and Asymptotic Freeness

38.1 Free independence and additive convolution

Definition 38.1 (Free independence and free cumulants). Unital subalgebras $(A_i)_i$ of (A, φ) are *freely independent* if

$$\varphi(a_1 \cdots a_m) = 0$$

whenever $a_j \in A_{i_j}$, $\varphi(a_j) = 0$, and neighboring indices are distinct. Variables are free when their generated unital subalgebras are.

Let $NC(n)$ be the lattice of noncrossing partitions of $\{1, \dots, n\}$. The *free cumulants* κ_n are the multilinear functionals determined recursively by

$$\varphi(a_1 \cdots a_n) = \sum_{\pi \in NC(n)} \kappa_\pi(a_1, \dots, a_n),$$

where κ_π is the product of the cumulants associated to the blocks of π . Freeness is equivalently the vanishing of every mixed free cumulant.

Definition 38.2 (Semicircular variables). A self-adjoint variable S is *semicircular* with variance σ^2 if its distribution is

$$d\mu_{\text{sc}, \sigma}(x) = \frac{1}{2\pi\sigma^2} \sqrt{4\sigma^2 - x^2} \mathbf{1}_{[-2\sigma, 2\sigma]}(x) dx.$$

Equivalently, its only nonzero free cumulant is $\kappa_2(S, S) = \sigma^2$.

Exercise 38.1 (Free central limit theorem). Use noncrossing pairings to prove

$$\varphi(S^{2m+1}) = 0, \quad \varphi(S^{2m}) = C_m \sigma^{2m}, \quad C_m = \frac{1}{m+1} \binom{2m}{m}.$$

Let $X_1, X_2, \dots$ be freely independent, identically distributed, centered self-adjoint variables with $\varphi(X_i^2) = \sigma^2$ and moments of every order. Use multilinearity and vanishing mixed cumulants to prove that the moments of $n^{-1/2}(X_1 + \cdots + X_n)$ converge to these semicircular moments. Compare this free central limit theorem with the Gaussian limit for classically independent variables.

Definition 38.3 (Cauchy and R -transforms). For a compactly supported probability measure μ on $\mathbb{R}$, its *Cauchy transform* is

$$G_\mu(z) = \int_{\mathbb{R}} \frac{d\mu(t)}{z - t}, \quad z \in \mathbb{C} \setminus \text{supp } \mu.$$

Near infinity it has a local compositional inverse $K_\mu(w) = G_\mu^{(-1)}(w)$ near $w = 0$. The *R -transform* is defined by

$$K_\mu(w) = \frac{1}{w} + R_\mu(w), \quad R_\mu(w) = \sum_{n \geq 1} \kappa_n(\mu) w^{n-1}.$$

If freely independent self-adjoint variables X, Y have laws μ, ν , then the law of $X + Y$ is their *free additive convolution* $\mu \boxplus \nu$, characterized by

$$R_{\mu \boxplus \nu} = R_\mu + R_\nu.$$

Exercise 38.2 (Free convolution and the semicircle equation). Derive the additivity of the R -transform from the vanishing of mixed free cumulants. Show that a centered semicircular law of variance σ^2 has $R(w) = \sigma^2 w$ and hence

$$\sigma^2 G(z)^2 - zG(z) + 1 = 0.$$

Select the solution satisfying $zG(z) \rightarrow 1$ at infinity and recover the semicircle density from the Stieltjes inversion formula. Deduce that the free sum of semicircular variables of variances σ_1^2 and σ_2^2 is semicircular of variance $\sigma_1^2 + \sigma_2^2$, and rederive the free central limit theorem at the level of transforms.

38.2 Random matrices and asymptotic freeness

Definition 38.4 (Asymptotic freeness). Random matrix families $A_N^{(1)}, \dots, A_N^{(r)}$ are *asymptotically free* when their normalized expected joint trace moments converge to the moments of freely independent variables:

$$\lim_{N \rightarrow \infty} \frac{1}{N} \mathbb{E} \operatorname{Tr} P(A_N^{(1)}, \dots, A_N^{(r)}) = \varphi(P(a_1, \dots, a_r))$$

for every noncommutative polynomial P .

Exercise 38.3 (Asymptotic freeness of independent GUE matrices). For independent GUE matrices $H_N^{(1)}, \dots, H_N^{(r)}$, apply Wick's rule to normalized mixed traces of centered polynomials. Prove that crossing pairings are lower order and that mixed free cumulants vanish in the limit. Conclude that the family converges in joint moments to freely independent semicircular variables.