

Coordinate-free Mathematics

Paul Gustafson

May 2025

1 Linear Algebra

Linear algebra studies linear maps between vector spaces. A **vector space** V over a field k is a set equipped with vector addition and scalar multiplication operations satisfying straightforward compatibility conditions. A **linear map** $f : V \rightarrow U$ between k -vector spaces is a map of underlying sets such that $f(\alpha v + \beta w) = \alpha f(v) + \beta f(w)$ for all $\alpha, \beta \in k$ and $v, w \in V$. Examples include differential operators, group representations, coefficient matrices of linear systems of equations, and local approximations of globally nonlinear functions.

The **free k -vector space on a set X** , denoted kX , is the set of finitely supported set-theoretical functions $X \rightarrow k$ with pointwise addition and scalar multiplication. The **direct sum** vector space $V \oplus W$ is the set $V \times W$ with pointwise addition and scalar multiplication defined by $\alpha(v \oplus w) = \alpha v \oplus \alpha w$. The **quotient** of V by a subspace W , denoted V/W , is the set of equivalence classes of V modulo the relation $x \sim y$ iff $x - y \in W$. The **tensor product** $V \otimes W$ is the vector space $k(V \times W)/N$ where N is the subspace generated by the obvious linearity relations on V and W .

Exercise 1 An **isomorphism** of vector spaces $V \cong W$ consists of a pair of linear maps $f : V \rightarrow W$ and $g : W \rightarrow V$ such that $f \circ g = 1_W$ and $g \circ f = 1_V$. Prove the following:

- $k(X \sqcup Y) \cong kX \oplus kY$
- $k(X \times Y) \cong kX \otimes kY$
- The first isomorphism theorem for vector spaces (and second and third if you want).
- Find universal properties determining up to canonical isomorphism the free k -vector space, direct sum, quotient, tensor product, and dual space (defined below), respectively.

The **dual space** V^* is the space of linear maps from a vector space V to its ground field k . The evaluation map $\text{ev}_V : V^* \otimes V \rightarrow k$ is the linear extension of the evaluation morphism $f \otimes x \mapsto f(x)$. We say that V is **finite dimensional**

if there is a co-evaluation morphism $\text{coev}_V : k \rightarrow V \otimes V^*$ satisfying the snake equations $(1_V \otimes \text{ev}_V) \circ (\text{coev}_V \otimes 1_V) = 1_V$ and $(\text{ev}_V \otimes 1_{V^*}) \circ (1_{V^*} \otimes \text{coev}_V) = 1_{V^*}$. We define the **dimension** $\dim(V) = \text{tr}(1_V)$, where $\text{tr}(f) : \text{End}(V) \rightarrow k$ is given by $f \mapsto (\text{ev}_V \circ s_{V,V^*} \circ (f \otimes 1_{V^*}) \circ \text{coev}_V)(1)$ where $s_{X,Y} : X \otimes Y \rightarrow Y \otimes X$ is the symmetric braiding $x \otimes y \mapsto y \otimes x$.

Exercise 2 Prove the following for finite-dimensional vector spaces V, W and linear maps $f, g : V \rightarrow V$.

- $\text{tr}(fg) = \text{tr}(gf)$
- $\text{tr}(f \oplus g) = \text{tr}(f) + \text{tr}(g)$
- $\text{tr}(f \otimes g) = \text{tr}(f) \text{tr}(g)$
- $\dim(V) \in \mathbf{N}$
- $V \cong W$ iff $\dim(V) = \dim(W)$

The simplest linear maps are scalar multiples of the identity. Although most linear maps are not of this form, it is useful to decompose them into sums of scalar maps when possible. One condition under which we can decompose a linear map into a direct sum of scalars is if the map is self-adjoint. Suppose V is finite dimensional over $\mathbb{C}$. A Hermitian inner product $\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{C}$ is a positive-definite sesquilinear form (linear in the first variable, anti-linear in the second variable, and conjugate symmetric). Write $|v| = \sqrt{\langle v, v \rangle}$. For a linear map $f : V \rightarrow W$ between Hermitian spaces, the adjoint $f^* : W \rightarrow V$ is characterized by $\langle fv, w \rangle = \langle v, f^*w \rangle$. An endomorphism f is **self-adjoint** if $f = f^*$. An isomorphism f is **unitary** if it preserves inner products, equivalently if $f^* = f^{-1}$.

For maps $f : V \rightarrow V'$ and $g : W \rightarrow W'$, define $(f \otimes g)(v \otimes w) = f(v) \otimes g(w)$. The tensor product of Hermitian spaces has inner product

$$\langle v_1 \otimes w_1, v_2 \otimes w_2 \rangle = \langle v_1, v_2 \rangle \langle w_1, w_2 \rangle.$$

The Cauchy–Schwarz inequality is $|\langle v, w \rangle| \leq |v||w|$, with equality iff v and w are linearly dependent.

Exercise 3 (No-cloning theorem) Let V be a finite-dimensional complex inner-product space, let $b \in V$ be a unit vector, and suppose that $C : V \otimes V \rightarrow V \otimes V$ is unitary. Prove that if

$$C(v \otimes b) = v \otimes v, \quad C(w \otimes b) = w \otimes w$$

for unit vectors $v, w \in V$, then

$$\langle v, w \rangle = \langle v, w \rangle^2.$$

Deduce that either $\langle v, w \rangle = 0$ or $v = w$. Conclude that no unitary map can satisfy $C(v \otimes b) = v \otimes v$ for every unit vector $v \in V$.

A scalar $\lambda \in k$ is an **eigenvalue** for a linear map $f : V \rightarrow V$ if there exists a nonzero vector $v \in V$ for which $fv = \lambda v$. The **eigenspace** of λ is

$$\ker(f - \lambda 1_V),$$

whose nonzero elements are eigenvectors; its dimension is called the **geometric multiplicity** of λ .

Exercise 4 *Prove that any self-adjoint linear map f on a finite-dimensional complex vector space V is diagonalizable:*

$$f = \sum \lambda_i p_i,$$

where $\sum p_i = 1_V$ and $p_i p_j = \delta_{ij} p_i$. Use this fact to prove the existence of the singular value decomposition for an arbitrary linear map between complex finite-dimensional vector spaces.

For $\psi \in V \otimes W$, define

$$T_\psi : V^* \rightarrow W, \quad T_\psi(\phi) = (\phi \otimes 1_W)(\psi).$$

The map $j_V : V \rightarrow V^*$ given by $j_V(v) = \langle \cdot, v \rangle$ is conjugate-linear. Equip V^* with the Hermitian inner product determined by $\langle j_V(v), j_V(w) \rangle = \langle w, v \rangle$. With $T_\psi^* : W \rightarrow V^*$ denoting the adjoint, define

$$\rho_V = j_V^{-1} T_\psi^* T_\psi j_V, \quad \rho_W = T_\psi T_\psi^*.$$

Write $U(V)$ for the group of unitary endomorphisms of V , acting with $U(W)$ on $V \otimes W$ by tensor products. A nonzero tensor is **decomposable** if it has the form $v \otimes w$. The rank of an endomorphism is the dimension of its image.

Exercise 5 (Schmidt decomposition and reduced endomorphisms)

Let V and W be finite-dimensional Hermitian vector spaces and let $0 \neq \psi \in V \otimes W$. Prove that there exist orthonormal families (v_i) in V and (w_i) in W , and positive scalars s_i , such that

$$\psi = \sum_{i=1}^r s_i v_i \otimes w_i.$$

Prove the following:

- The nonzero eigenvalues of ρ_V and ρ_W are the numbers s_i^2 , with the same multiplicities.
- ψ is decomposable iff exactly one s_i is nonzero.
- ρ_V has rank one iff ρ_W has rank one iff ψ is decomposable.
- The multiset (s_i) is invariant under the action of $U(V) \times U(W)$.

For a unit vector $\psi \in V$ and a self-adjoint $A \in \text{End}(V)$, define

$$m_\psi(A) = \langle \psi, A\psi \rangle, \quad \Delta_\psi(A) = |(A - m_\psi(A)1_V)\psi|.$$

Exercise 6 (Uncertainty principle) Let $A, B \in \text{End}(V)$ be self-adjoint and let $\psi \in V$ be a unit vector. Prove that

$$\Delta_\psi(A)\Delta_\psi(B) \geq \frac{1}{2} |\langle \psi, [A, B]\psi \rangle|, \quad [A, B] = AB - BA.$$

Determine the condition for equality from the equality condition in the Cauchy-Schwarz inequality.

For $K \in \text{End}(V)$, define $e^K = \sum_{n \geq 0} K^n/n!$. If H is self-adjoint, write $U_t = e^{-itH}$.

Exercise 7 (Conservation from commutators)

Let $A, H \in \text{End}(V)$ be self-adjoint, let $U_t = e^{-itH}$, and let $\psi \in V$. Prove that

$$\frac{d}{dt} \langle U_t\psi, AU_t\psi \rangle = -i \langle U_t\psi, [H, A]U_t\psi \rangle.$$

Deduce that $[H, A] = 0$ implies that $\langle U_t\psi, AU_t\psi \rangle$ is independent of t for every $\psi \in V$. Prove the converse.

We can solve for the eigenvalues of a linear operator on a finite-dimensional vector space via the determinant. Define the tensor algebra and **exterior algebra** of V by

$$T(V) = \bigoplus_{k \geq 0} V^{\otimes k}, \quad \Lambda V = T(V)/\langle v \otimes v : v \in V \rangle.$$

The space $\Lambda^k V$ is the degree- k component of ΛV .

Exercise 8 Show that the highest graded component of ΛV is one-dimensional.

The **determinant** $\det(f)$ is the coefficient of the linear map on the highest-graded component of ΛV given by $v_1 \wedge \cdots \wedge v_{\dim(V)} \mapsto f(v_1) \wedge \cdots \wedge f(v_{\dim(V)})$.

Exercise 9 Prove the following:

- $\det(f \oplus g) = \det(f)\det(g)$
- If $V = \mathbb{R}^n$, then $\det(f)$ is the signed volume of $f(I^n)$ where $I = [0, 1]$.
- Cramer's rule

For $f \in \text{End}(V)$, the **characteristic polynomial** is the polynomial $\lambda \mapsto \det(f - \lambda 1_V)$.

Exercise 10 Assume that the characteristic polynomial of f splits over k . Prove the following for a linear operator f on a finite-dimensional vector space V :

- Every eigenvalue of f is a root of its characteristic polynomial.
- The geometric multiplicity of each eigenvalue is at most equal to its **algebraic multiplicity** (i.e. multiplicity as a root of the characteristic polynomial)
- $\text{tr}(f) = \sum_i \lambda_i$ and $\det(f) = \prod_i \lambda_i$, where each eigenvalue λ_i is repeated according to its algebraic multiplicity
- The operator f satisfies its own characteristic polynomial (replacing λ with f)
- The existence of the Jordan normal form for a linear operator on a finite-dimensional complex vector space

2 Calculus

A **smooth manifold** is a topological space that is locally diffeomorphic to Euclidean space $\mathbb{R}^n$. A **smooth vector bundle** $p : E \rightarrow M$ consists of a smooth surjection p between smooth manifolds E (the total space) and M (the base space) such that all fibers are isomorphic to a single vector space V , and smooth local trivializations exist (locally, the fiber bundle looks like a product space). A prototypical example is the **tangent bundle** TM which consists of all tangent spaces of points of a given manifold. A **vector field** $X \in \Gamma(TM)$ is a smooth section of the tangent bundle (i.e. a smooth map $s : M \rightarrow TM$ such that $p \circ s = 1_M$).

Exercise 11 Write down precise definitions for the following:

- smooth manifold, smooth map, vector bundle, tangent bundle.
- bundle map, direct sum, tensor product, and dual bundle
- derivative of a smooth function between manifolds

To define integration in a coordinate-free way, we will integrate differential forms over chains. The space of **differential n -forms** is $\Omega^n(M) = \Gamma(\Lambda^n T^*M)$. The exterior derivative $d : \Omega^n(M) \rightarrow \Omega^{n+1}(M)$ extends the differential of smooth functions and satisfies

$$d(\alpha \wedge \beta) = d\alpha \wedge \beta + (-1)^p \alpha \wedge d\beta$$

for $\alpha \in \Omega^p(M)$. A **singular n -simplex** is a smooth map $\sigma : \Delta^n \rightarrow M$ where Δ^n is the standard n -simplex. The collection of singular n -simplices in M freely generates the abelian group $C_n(M)$ of **singular n -chains** in M . The **boundary map** $\partial : C_n(M) \rightarrow C_{n-1}(M)$ is defined on each simplex $\sigma : \Delta^n \rightarrow M$ by

$$\partial\sigma = \sum_{j=0}^n (-1)^j \sigma \circ f_j,$$

where $f_j : \Delta^{n-1} \rightarrow \Delta^n$ is the inclusion of the face opposite the j -th vertex of Δ^n .

A **chain complex** $(A_n, d_n)_n$ is a sequence of abelian groups A_n and homomorphisms $d_n : A_n \rightarrow A_{n-1}$ such that $d_{n-1} \circ d_n = 0$ for all n . The n -th **homology group** H_n of the chain complex is

$$H_n = \ker(d_n) / \operatorname{im}(d_{n+1}).$$

The prototypical example is singular homology which corresponds to the chain complex of singular chains with the boundary map defined above.

Cohomology and cochains are defined similarly, using superscripts instead of subscripts and boundary maps $d^n : A^n \rightarrow A^{n+1}$ increase degree instead of decrease it. Singular cohomology corresponds to the cochain complex given by dualizing the singular chain complex above.

The **de Rham cohomology** of M is

$$H_{\text{dR}}^n(M) = \ker(d : \Omega^n(M) \rightarrow \Omega^{n+1}(M)) / \operatorname{im}(d : \Omega^{n-1}(M) \rightarrow \Omega^n(M)).$$

Exercise 12 *Prove the following:*

- Show that $d^2 = 0$, so differential forms on a manifold M form a cochain complex.
- Define the integral of an n -form on an n -chain. Prove that $\int_c d\omega = \int_{\partial c} \omega$ for any n -chain c and differential $n-1$ -form ω .
- Singular cohomology with real coefficients and de Rham cohomology are naturally isomorphic.
- Bounded chain complexes of finite-dimensional vector spaces form a symmetric monoidal abelian category with duals, the trace of an endomorphism is its Lefschetz number, and the dimension of a chain complex is its Euler characteristic.

For a vector field X , contraction is the map $\iota_X : \Omega^p(M) \rightarrow \Omega^{p-1}(M)$ defined by

$$(\iota_X \alpha)(X_2, \dots, X_p) = \alpha(X, X_2, \dots, X_p).$$

The Lie derivative of forms is determined by Cartan's formula $\mathcal{L}_X = d\iota_X + \iota_X d$.

A **symplectic form** is a two-form $\omega \in \Omega^2(M)$ such that $d\omega = 0$ and $X \mapsto \iota_X \omega$ is a bundle isomorphism $TM \rightarrow T^*M$. Write $C^\infty(M)$ for the smooth real-valued functions on M . For $H \in C^\infty(M)$, define X_H by

$$\iota_{X_H} \omega = dH.$$

Exercise 13 (Liouville's theorem)

Let (M, ω) be $2n$ -dimensional with ω symplectic. For $H \in C^\infty(M)$, prove using Cartan's formula that

$$\mathcal{L}_{X_H} \omega = 0.$$

Deduce that $\mathcal{L}_{X_H}\omega^n = 0$ and hence that the local flow of X_H preserves the volume form

$$\frac{\omega^n}{n!}.$$

For $f, g \in C^\infty(M)$, define

$$\{f, g\} = \omega(X_f, X_g).$$

Exercise 14 (Hamiltonian Noether theorem)

Let (M, ω) be symplectic. Prove that

$$X_g(f) = \{f, g\}.$$

Let Φ_t be the local flow of X_f . Prove that the following are equivalent:

- $H \circ \Phi_t = H$ whenever both sides are defined.
- $X_f(H) = 0$.
- $\{H, f\} = 0$.
- f is constant along the integral curves of X_H .

Write vol_g for the metric volume form and $\langle \cdot, \cdot \rangle_g$ for the induced pairing of forms. On an oriented pseudo-Riemannian manifold, the Hodge star $\star$ is determined by

$$\alpha \wedge \star \beta = \langle \alpha, \beta \rangle_g \text{vol}_g.$$

Exercise 15 (Charge conservation from Maxwell's equations)

Let M be an oriented four-dimensional pseudo-Riemannian manifold. Suppose that $F \in \Omega^2(M)$ and $J \in \Omega^3(M)$ satisfy

$$dF = 0, \quad d \star F = J.$$

Prove that $dJ = 0$. If C is an oriented compact four-dimensional chain, deduce that

$$\int_{\partial C} J = 0.$$

If the three-dimensional pieces of its boundary satisfy

$$\partial C = \Sigma_1 - \Sigma_0 + B,$$

prove that

$$\int_{\Sigma_1} J - \int_{\Sigma_0} J = - \int_B J.$$

For a closed form A , write $[A]$ for its de Rham cohomology class.

Exercise 16 (Aharonov–Bohm cohomological obstruction) Let M be a compact smooth manifold and let $A \in \Omega^1(M)$ satisfy $dA = 0$. Prove the following:

- For every contractible open set $U \subseteq M$, there exists $\lambda \in C^\infty(U)$ such that $A|_U = d\lambda$.
- For every smooth closed curve γ , the number $\int_\gamma A$ depends only on the homology class of γ .
- There exists a global $\lambda \in C^\infty(M)$ with $A = d\lambda$ iff $[A] = 0 \in H_{\text{dR}}^1(M)$.
- If $[A] \neq 0$, then there exists a smooth closed curve γ such that $\int_\gamma A \neq 0$.

An **affine connection** is a map $\nabla : \Gamma(TM) \times \Gamma(TM) \rightarrow \Gamma(TM)$ such that $\nabla_f X Y = f \nabla_X Y$ and $\nabla_X(fY) = (\partial_X f)Y + f \nabla_X Y$ for any scalar field f on M .

References

- [1] Paul R. Halmos, *Finite-Dimensional Vector Spaces*, 2nd ed., Springer, 1974.
- [2] Kenneth Hoffman and Ray Kunze, *Linear Algebra*, 2nd ed., Prentice Hall, 1971.
- [3] Michael Spivak, *Calculus on Manifolds: A Modern Approach to Classical Theorems of Advanced Calculus*, Addison-Wesley, 1965.
- [4] Serge Lang, *Algebra*, 3rd ed., Springer, 2002.
- [5] Kate Ponto and Michael Shulman, *Traces in symmetric monoidal categories*, *Expositiones Mathematicae*, 32(3):248–273, 2014.
- [6] Raoul Bott and Loring W. Tu, *Differential Forms in Algebraic Topology*, Graduate Texts in Mathematics, Vol. 82, Springer, 1982.