



Getting started with TEOS-10 and the **Gibbs Seawater (GSW)** Oceanographic Toolbox

VERSION 3.0

May 2011



The International Thermodynamic Equation Of Seawater – 2010 (TEOS-10) was developed by SCOR/IAPSO Working Group 127 and was adopted by the Intergovernmental Oceanographic Commission (IOC). The members of SCOR/IAPSO Working Group 127 are as follows.

Trevor J. McDougall, (chair), CSIRO, Hobart, Australia
Rainer Feistel, Leibniz-Institut fuer Ostseeforschung, Warnemuende, Germany
Daniel G. Wright⁺, Bedford Institute of Oceanography, Dartmouth, Canada
Rich Pawlowicz, University of British Columbia, Vancouver, Canada
Frank J. Millero, University of Miami, Florida, USA
David R. Jackett, formerly of CSIRO, Hobart, Australia
Brian A. King, National Oceanography Centre, Southampton, UK
Giles M. Marion, Desert Research Institute, Reno, USA
Steffen Seitz, Physikalisch-Technische Bundesanstalt (PTB), Braunschweig, Germany
Petra Spitzer, Physikalisch-Technische Bundesanstalt (PTB), Braunschweig, Germany
C-T. Arthur Chen, National Sun Yat-Sen University, Taiwan, R.O.C.

⁺ deceased, 8th July 2010.

The photograph on the front cover of a CTD and lowered ADCP hovering just below the sea surface was taken south of Timor from the *Southern Surveyor* in August 2003 by Ann Gronell Thresher. Document cover by Louise Bell.

For bibliographic purposes, this document should be cited as follows:

McDougall T. J. and P. M. Barker, 2011: Getting started with TEOS-10 and the Gibbs Seawater (GSW) Oceanographic Toolbox, 28pp., SCOR/IAPSO WG127, ISBN 978-0-646-55621-5.

This document is available from www.TEOS-10.org.

May 2011

Author: McDougall, Trevor J.
Title: Getting started with TEOS-10 and the Gibbs Seawater (GSW) Oceanographic Toolbox / Trevor J. McDougall and Paul M. Barker.
ISBN: 9780646556215 (pbk.)
Subjects: Gibbs Seawater (GSW) Oceanographic Toolbox (Computer program)
Seawater--Thermodynamics--Computer programs
Other Authors/Contributors: Barker, Paul M.
Dewey Number: 551.46

Getting started with TEOS-10 and the Gibbs Seawater (GSW) Oceanographic Toolbox version 3.0

Trevor J. McDougall¹ and Paul M. Barker¹

May 2011

Table of Contents

	page
1. Preamble	2
2. Installing the GSW Oceanographic Toolbox in MATLAB	4
3. Absolute Salinity S_A	5
4. Preformed Salinity S_*	8
5. Conservative Temperature Θ	10
6. Which types of salinity and temperature should be archived?	12
7. The 48-term expression $\hat{\rho}(S_A, \Theta, p)$ for density	13
8. Changes to oceanographic practice under TEOS-10	16
9. Ocean modelling using TEOS-10	17
10. A guide to the GSW Oceanographic Toolbox	19
11. References	23
12. Recommended nomenclature, symbols and units in oceanography.	24

¹CSIRO Marine and Atmospheric Research, Hobart, Australia.
email: Trevor.McDougall@TEOS-10.org

1. Preamble

The International Thermodynamic Equation Of Seawater – 2010 (TEOS-10) allows all the thermodynamic properties of pure water, ice, seawater and moist air to be evaluated in a self-consistent manner. For the first time the effects of the variations in seawater composition around the world ocean are accounted for; these spatial variations of seawater composition cause density differences that are equivalent to ten times the precision of our Practical Salinity measurements at sea.

The GSW Oceanographic Toolbox of TEOS-10 is concerned primarily with the properties of pure liquid water and of seawater; the TEOS-10 software for evaluating the properties of ice and of humid air is available in the SIA (Seawater-Ice-Air) software library from the TEOS-10 web site, <http://www.TEOS-10.org>.

TEOS-10 has introduced several new variables into oceanography, including Absolute Salinity S_A , Preformed Salinity S_* , and Conservative Temperature Θ . These variables are introduced in this document, and then the use of these variables is discussed, followed by the complete listing and description of the functions available in the GSW toolbox.

Absolute Salinity is the salinity argument of the TEOS-10 algorithms which give the various thermodynamic properties of seawater, and under TEOS-10 Absolute Salinity S_A is the salinity variable to be used in scientific publications. Note, however, it is Practical Salinity S_p which must be reported to and stored in national databases. The practice of storing one type of salinity in national databases (Practical Salinity), but using a different type of salinity in publications (Absolute Salinity), is exactly analogous to our present practice with temperature; *in situ* temperature is stored in databases (since it is the measured quantity), but the temperature variable that is used in publications is a calculated quantity, being potential temperature to date, and from now, Conservative Temperature.

For the past thirty years, under EOS-80 we have taken the “raw” data of Practical Salinity S_p (PSS-78), *in situ* temperature t (now ITS-90) and pressure p and we have used an algorithm to calculate potential temperature θ in order to analyze and publish water-mass characteristics on the $S_p - \theta$ diagram. On this $S_p - \theta$ diagram we have been able to draw curved contours of potential density using EOS-80. Under TEOS-10 this practice has now changed. Density and potential density (and all types of geostrophic streamfunction including dynamic height anomaly) are now not functions of Practical Salinity S_p but rather are functions of Absolute Salinity S_A . TEOS-10 also defines a new temperature variable, Conservative Temperature Θ , which takes the place of potential temperature θ . Conservative Temperature Θ has the advantage over θ of more accurately representing the “heat content” of seawater. Under TEOS-10 is not possible to draw isolines of potential density on a $S_p - \theta$ diagram. Rather, because of the spatial variations of seawater composition, a given value of potential density defines an area on the $S_p - \theta$ diagram, not a curved line. Hence for the analysis and publication of ocean data under TEOS-10 we need to change from using the $S_p - \theta$ diagram which was appropriate under EOS-80, to using the $S_A - \Theta$ diagram. It is on this $S_A - \Theta$ diagram that the isolines of potential density can be drawn under TEOS-10.

As a fast-track precursor to the rest of this document, we note that these calculations can be performed using the functions of the GSW Oceanographic Toolbox as follows. The observed variables (S_p, t, p) , together with longitude and latitude, are used to first form Absolute Salinity S_A using `gsw_SA_from_SP`, and then Conservative Temperature Θ is calculated using `gsw_CT_from_t`. Oceanographic water masses are then analyzed on the $S_A - \Theta$ diagram (using `gsw_SA_CT_plot`), and potential density contours can be drawn on this $S_A - \Theta$ diagram using `gsw_rho(SA,CT,p_ref)`.

The more prominent advantages of TEOS-10 compared with EOS-80 are

- For the first time the influence of the spatially varying composition of seawater is systematically taken into account through the use of Absolute Salinity S_A . In the open ocean, this has a non-trivial effect on the horizontal density gradient, and thereby on ocean velocities and “heat” transports calculated via the “thermal wind” relation.
- The new salinity variable, Absolute Salinity S_A , is measured in SI units (e.g. g kg^{-1}).
- The Gibbs function approach of TEOS-10 allows the calculation of internal energy, entropy, enthalpy, potential enthalpy and the chemical potentials of seawater as well as the freezing temperature, and the latent heats of melting and of evaporation. These quantities were not available from EOS-80 but are essential for the accurate accounting of “heat” in the ocean and for the consistent and accurate treatment of air-sea and ice-sea heat fluxes in coupled climate models.
- In particular, Conservative Temperature Θ accurately represents the “heat content” per unit mass of seawater, and is to be used in place of potential temperature θ in oceanography.
- The thermodynamic quantities available from TEOS-10 are totally consistent with each other, while this was not the case with EOS-80.
- A single algorithm for seawater density (the 48-term computationally-efficient expression $\hat{\rho}(S_A, \Theta, p)$) can now be used for ocean modelling, for observational oceanography, and for theoretical studies. By contrast, for the past 30 years we have used different algorithms for density in ocean modelling and in observational oceanography and inverse modelling.

The present document (McDougall and Barker, 2011) provides a short description of the three new oceanographic variables S_A , S_* and Θ , leading into a discussion of the changes to observational oceanography and ocean modelling under TEOS-10 (compared with EOS-80), and then we list and describe the functions in the GSW Oceanographic Toolbox. The present document ends with the recommendations of SCOR/IAPSO Working Group 127, as endorsed by the Intergovernmental Oceanographic Commission, for the nomenclature, symbols and units to be used in physical oceanography, repeated from appendix L of IOC *et al.* (2010). Another document “What every oceanographer needs to know about TEOS-10 (The TEOS-10 Primer)” (Pawlowicz, 2010) provides a succinct introduction to the thermodynamic theory underlying TEOS-10 and is available from www.TEOS-10.org.

Note that when referring to the use of TEOS-10, it is the TEOS-10 Manual which should be referenced as IOC *et al.* (2010) [IOC, SCOR and IAPSO, 2010: *The international thermodynamic equation of seawater – 2010: Calculation and use of thermodynamic properties*. Intergovernmental Oceanographic Commission, Manuals and Guides No. 56, UNESCO (English), 196 pp.].

2. Installing the GSW Oceanographic Toolbox in MATLAB

Step 1

Download the GSW Oceanographic Toolbox in MATLAB from www.TEOS-10.org.

Step 2

Unzip the Toolbox to a directory you name "GSW".

**ENSURE THAT THE THREE SUBFOLDERS (html, library, pdf)
HAVE ALSO BEEN EXTRACTED.**

Step 3 (within MATLAB)

Add the "GSW" directory to your MATLAB path using "Add with subfolders ..."
That is, use the menus as follows "File" → "Set Path..." → "Add with subfolders ...".
(Alternatively, the "addpath" command could be used).

**ENSURE THAT THE THREE SUBFOLDERS (html, library, pdf)
HAVE ALSO BEEN ADDED TO THE PATH.**

Step 4

Run **gsw_check_functions** to check that the Toolbox is correctly installed and that there are no conflicts. (This function runs three stored vertical profiles through all of the GSW functions, and checks that the outputs are within pre-defined limits of the correct values. These pre-defined limits are a factor of approximately a hundred larger than the errors expected from the numerical precision of different computers, at the standard double precision of MATLAB).

If the MATLAB Desktop is running,

Step 5

Run **gsw_front_page** to gain access to the front page of the GSW Oceanographic Toolbox, which describes all aspects of the Toolbox.

Having installed the GSW Oceanographic Toolbox, the command **gsw_contents** will show the contents list of the software functions. The software descriptions and the help files for the GSW functions can be accessed by clicking on the function names on this list.

In addition, we have included a short demonstration function, **gsw_demo**, to introduce the user to the GSW Oceanographic Toolbox. **gsw_demo** uses two stored (S_p, t, p) profiles from the North Pacific and demonstrates, in a step-by-step manner, how to convert these into (S_A, Θ, p) profiles. **gsw_demo** then demonstrates how to evaluate several water-column properties such as dynamic height, geostrophic streamfunction and geostrophic velocity, as well as forming potential density contours on the $S_A - \Theta$ diagram.

A user may want to run **gsw_check_functions** periodically to confirm that the software remains uncorrupted.

3. Absolute Salinity S_A

Perhaps the most apparent change in using TEOS-10 compared with using the International Equation of State of seawater (EOS-80) is the adoption of Absolute Salinity S_A instead of Practical Salinity S_p (PSS-78) as the salinity argument for evaluating the thermodynamic properties of seawater. Importantly, Practical Salinity is retained as the salinity variable that is stored in national databases. This is done to maintain continuity in the archived salinity variable, and also because Practical Salinity is virtually the measured variable (whereas Absolute Salinity is a calculated variable).

The “raw” physical oceanographic data, as collected from ships and from autonomous platforms (e. g. Argo), and as stored in national oceanographic data bases, are

- Practical Salinity (S_p , unitless, PSS-78) and
- *in situ* temperature (t , °C, ITS-90) as functions of
- sea pressure (p , dbar), at a series of
- longitudes and latitudes.

Under TEOS-10 all the thermodynamic properties are functions of Absolute Salinity S_A (rather than of Practical Salinity), hence the first step in processing oceanographic data is to calculate Absolute Salinity, and this is accomplished by the GSW function **gsw_SA_from_SP**. Hence the function **gsw_SA_from_SP** is perhaps the most fundamental of the GSW functions as it is the gateway leading from oceanographic measurements to all the thermodynamic properties of seawater under TEOS-10. A call to this function can be avoided only if one is willing to ignore the influence of the spatial variations in the composition of seawater on seawater properties (such as density and specific volume). If this is indeed the intention, then the remaining GSW functions must be called with the salinity argument being Reference Salinity S_R , and most definitely, not with Practical Salinity S_p . Reference Salinity S_R can be obtained from the function **gsw_SR_from_SP**.

The **gsw_SA_from_SP**($S_p, p, \text{long}, \text{lat}$) function first interpolates the global Absolute Salinity Anomaly Ratio (R^δ) data set using the internal GSW library function **gsw_SAAR** to the ($p, \text{long}, \text{lat}$) location. **gsw_SA_from_SP** then uses this interpolated value of R^δ to calculate Absolute Salinity S_A according to (see Eqn. (A.5.10) of appendix A.5 of the TEOS-10 Manual, IOC *et al.* (2010))

$$S_A = \frac{35.165\,04\,\text{g kg}^{-1}}{35} S_p (1 + R^\delta). \quad \text{Non-Baltic} \quad (1)$$

In this expression $(35.165\,04\,\text{g kg}^{-1}/35) S_p$ is the Reference Salinity S_R , which is the best estimate of Absolute Salinity of a Standard Seawater sample.

Eqn. (1) is the value of Absolute Salinity returned by **gsw_SA_from_SP** unless the function detects that the location is in the Baltic Sea (where incidentally the internal GSW library function **gsw_SAAR** returns a value of R^δ of zero). If the observation is from the Baltic Sea, the Absolute Salinity Anomaly δS_A is calculated according to $S_A - S_R = 0.087\,\text{g kg}^{-1} \times (1 - S_p/35)$ (from Eqn. (A.5.16) of IOC *et al.* (2010), following Feistel *et al.* (2010)), so that Absolute Salinity S_A is given by

$$S_A = \frac{(35.165\,04 - 0.087)\,\text{g kg}^{-1}}{35} S_p + 0.087\,\text{g kg}^{-1}. \quad \text{Baltic Sea} \quad (2)$$

In summary, the **gsw_SA_from_SP** function returns either Eqn. (1) or Eqn. (2) depending on whether the longitude and latitude of the sample put the observation outside or inside the Baltic Sea. Since Practical Salinity should always be positive but there are sometimes a few negative values from a CTD, any negative input values of S_p to this function **gsw_SA_from_SP** are set to zero.

If the latitude and longitude are such as to place the observation well away from the ocean, a flag ‘in_ocean’ is set to zero as a warning, otherwise it is 1. This flag is only set when the observation is well and truly on dry land; often the warning flag is not set until

one is several hundred kilometers inland from the coast. When the function detects that the observation is not from the ocean, R^δ is set equal to zero and `gsw_SA_from_SP` returns $S_A = S_R = (35.165\,04\text{ g kg}^{-1}/35) S_p$ in accordance with Eqn. (1).

The largest influence of the variable seawater composition occurs in the northern North Pacific where $S_A - S_R = \delta S_A$ is as large as 0.027 g kg^{-1} (see Figure 2 of IOC *et al.* (2010) which is reproduced below), this being the difference between Absolute Salinity and the estimate of Absolute Salinity which can be made on the basis of Practical Salinity alone. This increment of salinity equates to an increment of density of approximately 0.020 kg m^{-3} .




Figure 2 (a). Absolute Salinity Anomaly δS_A at $p = 2000\text{ dbar}$.




Figure 2 (b). A vertical section of Absolute Salinity Anomaly δS_A along 180°E in the Pacific Ocean.

In order to gauge the importance of the spatial variation of seawater composition, the northward gradient of density at constant pressure is reproduced below from Fig. A.5.1 of IOC *et al.* (2010) for the data in a world ocean hydrographic atlas deeper than 1000m. The vertical axis in this figure is the magnitude of the difference between the northward density gradient at constant pressure when the TEOS-10 algorithm for density is called with S_A (as it should be) compared with calling the same TEOS-10 density algorithm with S_R as the salinity argument. Figure A.5.1 shows that the “thermal wind” is misestimated by more than 2% for 58% of the data in the world ocean below a depth of 1000m if the effects of the variable seawater composition are ignored. When this same comparison is done for only the North Pacific, it is found that 60% of the data deeper than 1000m has “thermal wind” misestimated by more than 10% if S_R is used in place of S_A .

The first version of **gsw_SA_from_SP** was made available in January 2009 (then called **gsw_ASsal**). The second version (version 2.0) was released in October 2010 and superseded version 1. The third version (version 3.0) was released in May 2011 and supersedes version 2.0.




Figure A.5.1. The northward density gradient at constant pressure (the horizontal axis) for data in the global ocean atlas of Gouretski and Koltermann (2004) for $p > 1000$ dbar. The vertical axis is the magnitude of the difference between evaluating the density gradient using S_A versus S_R as the salinity argument in the TEOS-10 expression for density.

As discussed in Pawlowicz (2010), Wright *et al.* (2011) and IOC *et al.* (2010), there are actually several contenders for the title of the “absolute salinity” of seawater, namely “Solution Salinity”, “Added-Mass Salinity”, and “Density Salinity”. The nuances surrounding these different definitions need not concern most physical oceanographers. Under TEOS-10 the words Absolute Salinity and symbol S_A are reserved for “Density Salinity” such as can be deduced using laboratory measurements with a vibrating beam densimeter.

4. Preformed Salinity S_*

Absolute Salinity S_A , Reference Salinity S_R and Practical Salinity S_p are all conservative salinity variables under the processes of (i) adiabatic pressure changes, and (ii) turbulent mixing, but none of these three salinity variables are conservative in the presence of (iii) biogeochemical processes. Preformed Salinity S_* is designed to be a conservative salinity variable which is unaffected by biogeochemical activity in the ocean; it is defined as Absolute Salinity less the contributions of biogeochemical processes to Absolute Salinity.

The **gsw_Sstar_from_SP**(SP,p,long,lat) function first interpolates the global Absolute Salinity Anomaly Ratio (R^δ) data set using the internal GSW library function **gsw_SAAR** to the (p,long,lat) location. **gsw_Sstar_from_SP** then uses this interpolated value of R^δ to calculate Preformed Salinity S_* according to (see Eqn. (A.5.11) of appendix A.5 of the TEOS-10 Manual, IOC *et al.* (2010))

$$S_* = \frac{35.165\,04\,\text{g kg}^{-1}}{35} S_p (1 - r_1 R^\delta), \quad \text{Non-Baltic} \quad (3)$$

where r_1 is taken to be the constant 0.35 based on the work of Pawlowicz *et al.* (2011). Note that $(35.165\,04\,\text{g kg}^{-1}/35) S_p$ is Reference Salinity S_R which is the best estimate of Absolute Salinity for a Standard Seawater sample. The figure below (from Figure A.4.1 of the TEOS-10 Manual, IOC *et al.*, 2010) illustrates the relationships between Preformed Salinity S_* , Reference Salinity S_R and Absolute Salinity S_A .




Figure A.4.1. Number line of salinity, illustrating the differences between Preformed Salinity S_* , Reference Salinity S_R , and Absolute Salinity S_A for seawater whose composition differs from that of Standard Seawater.

Equation (3) is the value of Preformed Salinity S_* returned by **gsw_Sstar_from_SP** unless the function detects that the location is in the Baltic Sea (where incidentally the internal GSW library function **gsw_SAAR** returns a value of R^δ of zero). In the Baltic Sea the deviations of Absolute Salinity from Reference Salinity are not due to non-conservative biogeochemical processes but rather are due to the anomalous composition entering the Baltic from rivers. Since these anomalous constituents are conservative, Preformed Salinity S_* in the Baltic Sea is Absolute Salinity S_A . Hence, if the observation is from the Baltic Sea, Preformed Salinity S_* is calculated using the relation $S_A - S_R = 0.087\,\text{g kg}^{-1} \times (1 - S_p/35)$ (from Eqn. (A.5.6) of IOC *et al.* (2010), following Feistel *et al.* (2010)), so that

$$S_* = S_A = \frac{(35.165\,04 - 0.087)\,\text{g kg}^{-1}}{35} S_p + 0.087\,\text{g kg}^{-1}. \quad \text{Baltic Sea} \quad (4)$$

In summary, the **gsw_Sstar_from_SP** function returns either Eqn. (3) or Eqn. (4) depending on whether the longitude and latitude of the sample put the observation outside or inside the Baltic Sea. Since Practical Salinity should always be positive but there are sometimes be a few negative values from a CTD, any negative input values of S_p to this function **gsw_Sstar_from_SP** are set to zero.

If the latitude and longitude are such as to place the observation well away from the ocean, a flag ‘in_ocean’ is set to zero as a warning, otherwise it is 1. This flag is only set when the observation is well and truly on dry land; often the warning flag is not set until one is several hundred kilometers inland from the coast. When the function detects that the observation is not from the ocean, R^δ is set equal to zero and **gsw_Sstar_from_SP** returns $S_* = S_R = (35.165\,04 \text{ g kg}^{-1}/35) S_P$ in accordance with Eqn. (3).

The largest influence of the variable seawater composition occurs in the northern North Pacific where $S_R - S_*$ is almost 0.01 g kg^{-1} , and the difference between Absolute Salinity and the conservative Preformed Salinity, $S_A - S_*$, is as large as 0.036 g kg^{-1} , equivalent to an increment of density of approximately 0.028 kg m^{-3} .

Continuing to concentrate on the mid-depth northern North Pacific, recall that the salinity difference $S_A - S_R = \delta S_A$ of 0.027 g kg^{-1} represents the difference between Absolute Salinity and the estimate of it using only Practical Salinity, while the value of $S_A - S_*$ of 0.036 g kg^{-1} is the total influence of biogeochemical processes on Absolute Salinity. An ocean model which treats its salinity variable as being conservative needs to account for this salinity difference, $S_A - S_*$, before density and the “thermal wind” can be accurately calculated. The reason why the salinity differences $S_A - S_R$ and $S_A - S_*$ are not equal is that biogeochemical processes have an effect on the conductivity of seawater and therefore on Practical Salinity and Reference Salinity.

What then is the appropriate use of Preformed Salinity S_* ? This salinity variable is the one which can be treated as being conservative. Hence, in contrast to the evolution equation of Absolute Salinity S_A , the evolution equation for Preformed Salinity S_* does not contain non-conservative source terms caused by biogeochemistry (see appendix A.20 of IOC *et al.* (2010)). This means that Preformed Salinity S_* is ideal for use as

- (i) the salinity variable that is advected and diffused in forward ocean models,
- (ii) the salinity variable that is advected and diffused in inverse ocean models, and
- (iii) the salinity variable that is averaged when forming a hydrographic atlas.

In these applications, the salinity difference $S_A - S_*$ is added to the averaged atlas or model salinity variable to form Absolute Salinity S_A before other thermodynamic properties (such as density etc.) are calculated. Because of its conservative nature, there is an argument that suggests that Preformed Salinity S_* should be used in place of Absolute Salinity S_A as the salinity axis of the $S_A - \Theta$ diagram, since water masses are notionally mixed on this diagram. However, to do so would take away the ability to draw isolines of potential density on this diagram, so the $S_A - \Theta$ diagram (as opposed to the $S_* - \Theta$ diagram) is recommended for water-mass analysis.

5. Conservative Temperature Θ

Because the TEOS-10 properties of seawater are all derived from a Gibbs function, it is possible to find thermodynamic properties such as enthalpy, internal energy and entropy. Hence potential enthalpy and Conservative Temperature (which is simply proportional to potential enthalpy) are readily available.

Conservative Temperature is in some respects quite similar to potential temperature in that the same artificial thought experiment is involved with their definitions. In both cases one takes a seawater sample at an arbitrary pressure in the ocean and one imagines decreasing the pressure on the seawater parcel in an adiabatic and isohaline manner until the sea pressure $p = 0$ dbar is reached. The temperature of the fluid parcel at the end of this artificial thought experiment is defined to be the potential temperature θ . Similarly, the enthalpy at the end of this artificial thought experiment is defined to be the potential enthalpy h^0 , and Conservative Temperature Θ is simply potential enthalpy divided by the fixed “heat capacity” $c_p^0 \equiv 3991.867\,957\,119\,63 \text{ J kg}^{-1} \text{ K}^{-1}$.

Conservative Temperature Θ represents the “heat content” of seawater much more accurately than does potential temperature θ . Θ can be evaluated from *in situ* temperature t from the function **gsw_CT_from_t**. The difference between potential temperature and Conservative Temperature can be as large as $\theta - \Theta = -1.4^\circ\text{C}$ but is more typically no more than $\pm 0.1^\circ\text{C}$ (see Figure A.17.1 of IOC *et al.* (2010) which is reproduced below). To put a temperature difference of 0.1°C in context, this is the typical difference between *in situ* and potential temperatures for a pressure difference of 1000 dbar, and it is approximately 40 times as large as the typical differences between t_{90} and t_{68} in the ocean.




Figure A.17.1. Contours (in $^\circ\text{C}$) of the difference between potential temperature and Conservative Temperature $\theta - \Theta$. This plot illustrates the non-conservative production of potential temperature θ in the ocean.

The air-sea heat flux is exactly proportional to the flux of Conservative Temperature, and because Θ is almost a perfectly conservative variable, the meridional “heat” flux is very accurately given by the meridional flux of Θ (as opposed to the meridional flux of potential temperature). Also, the parameterized lateral diffusion of “heat” along neutral tangent planes can be more than 1% different when such lateral diffusive heat fluxes are estimated using gradients of potential temperature rather than gradients of Conservative Temperature (see Figure A.14.1 of IOC *et al.* (2010) which is reproduced below).

For these reasons Conservative Temperature Θ is the appropriate temperature variable to be used in ocean analyses. Just as Absolute Salinity S_A is now to be used in oceanographic publications instead of Practical Salinity, so too Conservative Temperature Θ takes the place of potential temperature θ under TEOS-10.




Figure A.14.1. Contours of $(|\nabla_n \theta|/|\nabla_n \Theta| - 1) \times 100\%$ at $p = 0$, showing the percentage difference between the epineutral gradients of θ and Θ . The red dots are from the global ocean atlas of Gouretski and Koltermann (2004) at $p = 0$.

6. Which types of salinity and temperature should be archived?

Since 1978 we have gone to sea and collected and then archived Practical Salinity S_p (PSS-78), *in situ* temperature t (now ITS-90) and pressure p at known values of longitude, latitude and time. Under TEOS-10, nothing about this data collection and storage changes. Our instruments still need to measure properly calibrated data of (S_p, t, p) at known longitude, latitude and time. These are also exactly the same data that should continue to be archived in national oceanographic archives.

In particular, it cannot be over-emphasized that NONE of Reference Salinity S_R , Absolute Salinity S_A or Preformed Salinity S_* should EVER be submitted to or stored in national oceanographic databases. This resonates with present practice regarding temperature; we archive the measured quantity, *in situ* temperature t , not the calculated quantity, potential temperature θ . Similarly, just as potential temperature is not stored in national databases, so too Conservative Temperature should NEVER be submitted to or stored in such national databases. In short, such databases should store measured (not calculated) quantities, and for this purpose, Practical Salinity is regarded as a measured quantity.

7. The 48-term expression $\hat{\rho}(S_A, \Theta, p)$ for density

The computationally efficient 48-term rational function expression for density, $\hat{\rho}(S_A, \Theta, p)$, is described in McDougall *et al.* (2011b), in appendix A.30 and appendix K of the TEOS-10 Manual (IOC *et al.* (2010)), and is the function **gsw_rho**(SA,CT,p) in the GSW Oceanographic Toolbox. Seawater density data was fitted in a “funnel” of data points in (S_A, Θ, p) space which is described in more detail in McDougall *et al.* (2011b). The “funnel” extends to a pressure of 8000 dbar. At the sea surface the “funnel” covers the full range of temperature and salinity while for pressures greater than 6500 dbar, the maximum Conservative Temperature of the fitted data is 10°C and the minimum Absolute Salinity is 30 g kg⁻¹. That is, the 48-term fit has been performed over a region of parameter space which includes water that is approximately 10°C warmer and 5 g kg⁻¹ fresher in the deep ocean than the seawater which exists in the present ocean.

The rms error of this 48-term approximation to the TEOS-10 density over the oceanographic “funnel” is 0.00046 kg m⁻³; this can be compared with the rms uncertainty of 0.004 kg m⁻³ of the underlying laboratory density data to which the TEOS-10 Gibbs function was fitted. Similarly, the appropriate thermal expansion coefficient,

$$\alpha^\Theta = - \frac{1}{\rho} \left. \frac{\partial \rho}{\partial \Theta} \right|_{S_A, p},$$

of the 48-term equation of state is different from the same thermal expansion coefficient evaluated directly from TEOS-10 with an rms error in the “funnel” of $0.069 \times 10^{-6} \text{ K}^{-1}$, compared with the rms error of the thermal expansion coefficient of the laboratory data to which the Feistel (2008) Gibbs function was fitted of $0.73 \times 10^{-6} \text{ K}^{-1}$. In terms of the evaluation of density gradients, the haline contraction coefficient evaluated from the 48-term equation is many times more accurate than the thermal expansion coefficient.

In dynamical oceanography it is the thermal expansion and haline contraction coefficients α^Θ and β^Θ which are the most important aspects of the equation of state since the “thermal wind” is proportional to $\alpha^\Theta \nabla_p \Theta - \beta^\Theta \nabla_p S_A$ and the vertical static stability is given in terms of the buoyancy frequency N by $g^{-1} N^2 = \alpha^\Theta \Theta_z - \beta^\Theta (S_A)_z$. Hence for dynamical oceanography the 48-term rational function expression for density retains essentially the full accuracy of TEOS-10. The use of the 48-term expression for density has several advantages over using the exact formulation, namely

- it is a function of Conservative Temperature, so eliminating the need to be continually converting between Conservative and *in-situ* temperatures in order to evaluate density,
- it is computationally faster (by a factor of 6.5) to use the 48-term expression, **gsw_rho**(SA,CT,p) (or equivalently **gsw_rho_CT**(SA,CT,p)), rather than using **gsw_rho_CT_exact**(SA,CT,p) which is based on the sum of the Gibbs functions of pure water (IAPWS-09) and of sea salt (IAPWS-08),
- ocean models will use this 48-term equation of state, and it is advantageous for the fields of observational and theoretical oceanography to use the same equation of state as ocean models.

The functions of the GSW Oceanographic Toolbox are listed on the next two pages, after which section 8 begins on page 16.



documentation set

gsw_front_page
gsw_contents
gsw_check_functions
gsw_demo

front page to the GSW Oceanographic Toolbox
contents of the GSW Oceanographic Toolbox
checks that all the GSW functions work correctly
demonstrates many GSW functions and features

Practical Salinity (SP), PSS-78

gsw_SP_from_C
gsw_C_from_SP
gsw_SP_from_R
gsw_R_from_SP
gsw_SP_salinometer

Practical Salinity from conductivity, C (incl. for SP < 2)
conductivity, C, from Practical Salinity (incl. for SP < 2)
Practical Salinity from conductivity ratio, R (incl. for SP < 2)
conductivity ratio, R, from Practical Salinity (incl. for SP < 2)
Practical Salinity from a laboratory salinometer (incl. for SP < 2)

Absolute Salinity (SA), Preformed Salinity (Sstar) and Conservative Temperature (CT)

gsw_SA_from_SP
gsw_Sstar_from_SP
gsw_CT_from_t

Absolute Salinity from Practical Salinity
Preformed Salinity from Practical Salinity
Conservative Temperature from in-situ temperature

Absolute Salinity – Conservative Temperature plotting function

function to plot Absolute Salinity – Conservative Temperature profiles on the SA-CT diagram, including the freezing line and selected potential density contours

other conversions between temperatures, salinities, entropy, pressure and height

gsw_deltaSA_from_SP
gsw_SA_Sstar_from_SP
gsw_SR_from_SP
gsw_SP_from_SR
gsw_SP_from_SA
gsw_Sstar_from_SA
gsw_SA_from_Sstar
gsw_SP_from_Sstar
gsw_pt_from_CT
gsw_t_from_CT
gsw_CT_from_pt
gsw_pot_enthalpy_from_pt
gsw_pt0_from_t
gsw_pt_from_t
gsw_t00_from_t48
gsw_t00_from_t68
gsw_z_from_p
gsw_p_from_z
gsw_z_from_depth
gsw_depth_from_z
gsw_Abs_Pressure_from_p
gsw_p_from_Abs_Pressure
gsw_entropy_from_CT
gsw_CT_from_entropy
gsw_entropy_from_pt
gsw_pt_from_entropy
gsw_molality_from_SA
gsw_ionic_strength_from_SA

Absolute Salinity Anomaly from Practical Salinity
Absolute Salinity & Preformed Salinity from Practical Salinity
Reference Salinity from Practical Salinity
Practical Salinity from Reference Salinity
Practical Salinity from Absolute Salinity
Preformed Salinity from Absolute Salinity
Absolute Salinity from Preformed Salinity
Practical Salinity from Preformed Salinity
potential temperature from Conservative Temperature
in-situ temperature from Conservative Temperature
Conservative Temperature from potential temperature
potential enthalpy from potential temperature
potential temperature with reference pressure of 0 dbar
potential temperature
ITS-90 temperature from IPTS-48 temperature
ITS-90 temperature from IPTS-68 temperature
height from pressure
pressure from height
height from depth
depth from height
Absolute Pressure, P, from sea pressure, p
sea pressure, p, from Absolute Pressure, P
entropy from Conservative Temperature
Conservative Temperature from entropy
entropy from potential temperature
potential temperature from entropy
molality of seawater
ionic strength of seawater

density and enthalpy, based on the 48-term expression for density, $\hat{\rho}(S_A, \Theta, p)$

The functions in this group ending in “_CT” may also be called without “_CT”.

gsw_rho_CT
gsw_alpha_CT
gsw_beta_CT
gsw_rho_alpha_beta_CT
gsw_specvol_CT
gsw_specvol_anom_CT
gsw_sigma0_CT
gsw_sigma1_CT
gsw_sigma2_CT
gsw_sigma3_CT
gsw_sigma4_CT
gsw_sound_speed_CT
gsw_internal_energy_CT
gsw_enthalpy_CT
gsw_enthalpy_diff_CT
gsw_dynamic_enthalpy_CT
gsw_SA_from_rho_CT
gsw_CT_from_rho
gsw_CT_maxdensity

in-situ density, and potential density
thermal expansion coefficient with respect to CT
saline contraction coefficient at constant CT
in-situ density, thermal expansion & saline contraction coefficients
specific volume
specific volume anomaly
sigma0 from CT with reference pressure of 0 dbar
sigma1 from CT with reference pressure of 1000 dbar
sigma2 from CT with reference pressure of 2000 dbar
sigma3 from CT with reference pressure of 3000 dbar
sigma4 from CT with reference pressure of 4000 dbar
sound speed (approximate, with r.m.s. error of 0.067 m/s)
internal energy
enthalpy
difference of enthalpy between two pressures
dynamic enthalpy
Absolute Salinity from density
Conservative Temperature from density
Conservative Temperature of maximum density of seawater

water column properties, based on the 48-term expression for density, $\hat{\rho}(S_A, \Theta, p)$

gsw_Nsquared
gsw_Turner_Rsubrho
gsw_IPV_vs_fNsqared_ratio

buoyancy (Brunt-Väisälä) frequency squared (N²)
Turner angle & Rsubrho
ratio of the vertical gradient of potential density (with reference pressure, p_ref), to the vertical gradient of locally-referenced potential density

neutral and non-linear properties, based on the 48-term expression for density, $\hat{\rho}(S_A, \Theta, p)$

gsw_cabbeling
gsw_thermobaric
gsw_isopycnal_slope_ratio
gsw_ntp_pt_vs_CT_ratio
gsw_isopycnal_vs_ntp_CT_ratio

cabbeling coefficient
thermobaric coefficient
ratio of the slopes of isopycnals on the SA-CT diagram for p & p_ref
ratio of gradients of pt & CT in a neutral tangent plane
ratio of the gradient of CT in a potential density surface to that in the neutral tangent plane

geostrophic streamfunctions, based on the 48-term expression for density, $\hat{\rho}(S_A, \Theta, p)$

gsw_geo_stfr_dyn_height
gsw_geo_stfr_dyn_height_pc
gsw_geo_stfr_isopycnal
gsw_geo_stfr_isopycnal_pc
gsw_geo_stfr_Cunningham
gsw_geo_stfr_Montgomery

dynamic height anomaly
dynamic height anomaly for piecewise constant profiles
approximate isopycnal geostrophic streamfunction
approximate isopycnal geostrophic streamfunction for piecewise constant profiles
Cunningham geostrophic streamfunction
Montgomery geostrophic streamfunction

geostrophic velocity

gsw_geostrophic_velocity

geostrophic velocity

derivatives of enthalpy, entropy, CT and pt

gsw_CT_first_derivatives
gsw_CT_second_derivatives
gsw_enthalpy_first_derivatives
gsw_enthalpy_second_derivatives
gsw_entropy_first_derivatives
gsw_entropy_second_derivatives
gsw_pt_first_derivatives
gsw_pt_second_derivatives

freezing temperatures

gsw_CT_freezing
gsw_t_freezing
gsw_brineSA_CT
gsw_brineSA_t

Conservative Temperature freezing temperature of seawater
in-situ freezing temperature of seawater
Absolute Salinity of seawater at the freezing point (for given CT)
Absolute Salinity of seawater at the freezing point (for given t)

isobaric melting enthalpy and isobaric evaporation enthalpy

gsw_latentheat_melting
gsw_latentheat_evap_CT

latent heat of melting of ice into seawater (isobaric melting enthalpy)
latent heat of evaporation of water from seawater (isobaric evaporation enthalpy) with CT as input temperature
latent heat of evaporation of water from seawater (isobaric evaporation enthalpy) with in-situ temperature, t, as input

planet Earth properties

gsw_f
gsw_grav
gsw_distance

Coriolis parameter
gravitational acceleration
spherical earth distance between points in the ocean

steric height

gsw_steric_height

dynamic height anomaly divided by 9.7963 m s^{-2}

TEOS–10 constants

gsw_T0
gsw_P0
gsw_SSO
gsw_uPS
gsw_cp0
gsw_C3515
gsw_SonCI
gsw_valence_factor
gsw_atomic_weight

Celsius zero point; 273.15 K
one standard atmosphere; 101 325 Pa
Standard Ocean Reference Salinity; 35.165 04 g/kg
unit conversion factor for salinities; (35.165 04/35) g/kg
the “specific heat” for use with CT; 3991.867 957 119 63 (J/kg)/K
conductivity of SSW at SP=35, t_68=15, p=0; 42.9140 mS/cm
ratio of SP to Chlorinity; 1.80655 (g/kg)⁻¹
valence factor of sea salt; 1.2452898
mole-weighted atomic weight of sea salt; 31.4038218... g/mol

density and enthalpy in terms of CT, based on the exact Gibbs function

gsw_rho_CT_exact
gsw_alpha_CT_exact
gsw_beta_CT_exact
gsw_rho_alpha_beta_CT_exact
gsw_specvol_CT_exact
gsw_specvol_anom_CT_exact
gsw_sigma0_CT_exact
gsw_sigma1_CT_exact
gsw_sigma2_CT_exact
gsw_sigma3_CT_exact
gsw_sigma4_CT_exact
gsw_sound_speed_CT_exact
gsw_internal_energy_CT_exact
gsw_enthalpy_CT_exact
gsw_enthalpy_diff_CT_exact
gsw_dynamic_enthalpy_CT_exact
gsw_SA_from_rho_CT_exact
gsw_CT_from_rho_exact
gsw_CT_maxdensity_exact

in-situ density from CT, and potential density from CT
thermal expansion coefficient with respect to CT
saline contraction coefficient at constant CT
density, thermal expansion & saline contraction coefficients from CT
specific volume from CT
specific volume anomaly from CT
sigma0 from CT with reference pressure of 0 dbar
sigma1 from CT with reference pressure of 1000 dbar
sigma2 from CT with reference pressure of 2000 dbar
sigma3 from CT with reference pressure of 3000 dbar
sigma4 from CT with reference pressure of 4000 dbar
sound speed from CT
internal energy from CT
enthalpy from CT
difference of enthalpy from CT between two pressures
dynamic enthalpy from CT
Absolute Salinity from density & CT
Conservative Temperature from density
Conservative Temperature of maximum density of seawater

basic thermodynamic properties in terms of in-situ t, based on the exact Gibbs function

gsw_rho_t_exact
gsw_pot_rho_t_exact
gsw_sigma0_pt0_exact
gsw_alpha_wrt_CT_t_exact
gsw_alpha_wrt_pt_t_exact
gsw_alpha_wrt_t_exact
gsw_beta_const_CT_t_exact
gsw_beta_const_pt_t_exact
gsw_beta_const_t_exact
gsw_specvol_t_exact
gsw_specvol_anom_t_exact
gsw_sound_speed_t_exact
gsw_kappa_t_exact
gsw_kappa_const_t_exact
gsw_internal_energy_t_exact
gsw_enthalpy_t_exact
gsw_dynamic_enthalpy_t_exact
gsw_SA_from_rho_t_exact
gsw_t_from_rho_exact
gsw_t_maxdensity_exact
gsw_entropy_t_exact
gsw_cp_t_exact
gsw_isochoric_heat_cap_t_exact
gsw_chem_potential_relative_t_exact
gsw_chem_potential_water_t_exact
gsw_chem_potential_salt_t_exact
gsw_Helmholtz_energy_t_exact
gsw_adiabatic_lapse_rate_t_exact
gsw_osmotic_coefficient_t_exact
gsw_osmotic_pressure_t_exact

in-situ density
potential density
sigma0 from pt0 with reference pressure of 0 dbar
thermal expansion coefficient with respect to CT
thermal expansion coefficient with respect to pt
thermal expansion coefficient with respect to t
saline contraction coefficient at constant CT
saline contraction coefficient at constant pt
saline contraction coefficient at constant t
specific volume
specific volume anomaly
sound speed
isentropic compressibility
isothermal compressibility
internal energy
enthalpy
dynamic enthalpy
Absolute Salinity from density
in-situ temperature from density
in-situ temperature of maximum density of seawater
entropy
isobaric heat capacity
isochoric heat capacity of seawater
relative chemical potential
chemical potential of water in seawater
chemical potential of salt in seawater
Helmholtz energy
adiabatic lapse rate
osmotic coefficient of seawater
osmotic pressure of seawater

The GSW Toolbox is available from

www.TEOS-10.org



8. Changes to oceanographic practice under TEOS-10

For the past thirty years we have taken the “raw” data of Practical Salinity S_p (PSS-78), *in situ* temperature t (now ITS-90) and pressure p and we have used an algorithm to calculate potential temperature θ in order to analyze and publish water-mass characteristics on the $S_p - \theta$ diagram. On this $S_p - \theta$ diagram we have been able to draw curved contours of potential density using EOS-80.

Under TEOS-10 this practice has now changed:- density and potential density (and all types of geostrophic streamfunction including dynamic height anomaly) are now not functions of Practical Salinity S_p but rather are functions of Absolute Salinity S_A .

TEOS-10 also defines a new temperature variable, Conservative Temperature Θ , which takes the place of potential temperature θ (see section 5 above). Operationally, the calculation of Conservative Temperature Θ as a function of (S_A, t, p) under TEOS-10 is no different in principle from the way potential temperature was calculated from (S_p, t, p) under EOS-80; in both cases a simple computer algorithm is called. Conservative Temperature Θ has the advantage over θ of more accurately representing the “heat content” of seawater, and is also much closer (by a factor of a hundred) to being a conservative variable than is potential temperature. Heat is exchanged between the ocean and its atmosphere and ice boundaries as a flux of potential enthalpy which is exactly $c_p^0 \equiv 3991.867\,957\,119\,63\text{ J kg}^{-1}\text{ K}^{-1}$ times the density times the corresponding flux of Θ . The transport of potential enthalpy $c_p^0 \Theta$ in the ocean, and in particular across ocean sections, can be regarded as the transport of “heat” irrespective of whether there are non-zero fluxes of mass and/or of salt across such ocean sections (IOC *et al.*, 2010).

Under TEOS-10 is not possible to draw isolines of potential density on a $S_p - \theta$ diagram. Rather, because of the spatial variations of seawater composition, a given value of potential density defines an area on the $S_p - \theta$ diagram, not a curved line. Hence for the analysis and publication of ocean data under TEOS-10 we need to change from using the $S_p - \theta$ diagram which was appropriate under EOS-80, to using the $S_A - \Theta$ diagram. It is on this $S_A - \Theta$ diagram that the isolines of potential density can be drawn under TEOS-10.

Density may be calculated from the sum of the Gibbs functions of pure water (IAPWS-09, Feistel (2003)) and of salt (IAPWS-08, Feistel (2008)) using `gsw_rho_CT_exact(SA,CT,p)` or from the 48-term rational function expression `gsw_rho(SA,CT,p)`. The errors involved with using the 48-term expression for density are much less than the uncertainty in the effect of seawater composition on density, and are also much less than the uncertainty of the underlying laboratory density data to which the TEOS-10 Gibbs function was fitted (IAPWS-08, Feistel (2008)). This computationally efficient 48-term expression for density is the obvious choice for use in ocean models since it is a function of the model’s temperature variable, Conservative Temperature. The highly accurate nature of the 48-term expression means that theoretical studies, observational oceanography and ocean modeling can all be performed using the same equation of state which is conveniently expressed in terms of Conservative Temperature. This eliminates the need to continually transform from Conservative Temperature back to *in situ* temperature in order to calculate density and its derivatives.

These advantages lead us to recommend the 48-term expression for general use by oceanographers, including for observational studies, for ocean modelling and for theoretical studies, thus ensuring consistency between these different branches of oceanography. The GSW Oceanographic Toolbox provides many functions based on this 48-term equation of state, including `gsw_Nsquared(SA,CT,p)` to evaluate the square of the buoyancy frequency, `gsw_enthalpy(SA,CT,p)` to evaluate the specific enthalpy of seawater, and several functions to evaluate various geostrophic streamfunctions. The geostrophic streamfunction to be used for flow in an isobaric surface is `gsw_geo_strf_dyn_height` while that to be used in approximately neutral surfaces (including potential density surfaces, ω -surfaces and

Neutral Density (γ^n) surfaces) is **gsw_geo_strf_isopycnal**. Also, it is this 48-term expression for density that will be the basis for updated algorithms for ω -surfaces (Klocker *et al.* (2010)) and Neutral Density γ^n (Jackett and McDougall (1997)).

In summary, under EOS-80 we have to date used the observed variables (S_p, t, p) to first form potential temperature θ and then we have analyzed water masses on the $S_p - \theta$ diagram, and we have been able to draw curved contours of potential density on this same $S_p - \theta$ diagram. Under TEOS-10, the observed variables (S_p, t, p), together with longitude and latitude, are used to first form Absolute Salinity S_A using **gsw_SA_from_SP**, and then Conservative Temperature Θ is calculated using **gsw_CT_from_t**. Oceanographic water masses are then analyzed on the $S_A - \Theta$ diagram (using **gsw_SA_CT_plot**), and potential density contours can be drawn on this $S_A - \Theta$ diagram using **gsw_rho**(SA,CT,p_ref).

The various oceanographic properties that rely on the equation of state have been written in terms of S_A and Θ in the GSW Oceanographic Toolbox, and all of the oceanographic variables in common use (including geostrophic streamfunctions) have been written using the 48-term expression for density to ensure consistency between ocean models, observational studies and theoretical work. The use of many of these functions can be seen by running **gsw_demo**.

9. Ocean modelling using TEOS-10

Ocean models treat their salinity and temperature variables as being conservative, with the choice of variables to date being Practical Salinity and potential temperature. Converting ocean models to be TEOS-10 compatible requires several changes. The model's temperature variable needs to

- (i) accurately represent the “heat content” per unit mass of seawater and
- (ii) to be as conservative as possible under ocean mixing processes.

Conservative Temperature Θ has these properties whereas potential temperature θ does not. Fortunately it is relatively easy to change ocean models to have Conservative Temperature as their temperature variable. With the expression for density being cast in terms of Absolute Salinity S_A and Conservative Temperature Θ as $\hat{\rho}(S_A, \Theta, p)$, the interior of an ocean model can be written totally in terms of this one temperature variable, Θ . In the air-sea interaction module of an ocean model the sea-surface-temperature (SST) needs to be evaluated for use in bulk air-sea flux formulae, and this is done by calling the function **gsw_pt_from_CT**. This conversion from Θ to SST needs to be done just at the sea surface in the air-sea interaction module.

The current practice in numerical models is to treat salinity as a perfectly conserved quantity in the interior of the ocean. In order to continue this practice the appropriate model salinity variable is Preformed Salinity S_* . Preformed Salinity and Absolute Salinity are related to S_R and S_* respectively by Eqns. (A.20.1) and (A.20.2) of the TEOS-10 Manual, repeated here

$$S_* = S_R (1 - r_1 R^\delta), \quad (5)$$

$$S_A = S_* (1 + F^\delta), \quad (6)$$

where

$$R^\delta \equiv \frac{\delta S_A^{\text{ref}}}{S_R^{\text{ref}}} \quad \text{and} \quad F^\delta = \frac{[1 + r_1] R^\delta}{(1 - r_1 R^\delta)}. \quad (7a, b)$$

Recall that $R^\delta \equiv \delta S_A^{\text{ref}} / S_R^{\text{ref}}$ is the ratio of the reference (ref) values of Absolute Salinity Anomaly and Reference Salinity.

Because Preformed Salinity S_* is designed to be a conservative salinity variable, blind to the effects of biogeochemical processes, its evolution equation is in the conservative form (see appendix A.21 of IOC *et al.* (2011)),

$$\frac{d\hat{S}_*}{dt} = h^{-1}\nabla_n \cdot (hK\nabla_n \hat{S}_*) + \left(D \frac{\partial \hat{S}_*}{\partial z} \right)_z. \quad (8)$$

Here the over-tilde of \hat{S}_* indicates that this variable is the thickness-weighted average Preformed Salinity, having been averaged between a pair of closely-spaced neutral tangent planes. The material derivative on the left-hand side of Eqn. (8) is with respect to the sum of the Eulerian and quasi-Stokes velocities of height coordinates (equivalent to the description in appendix A.21 of IOC *et al.* (2010) in terms of the thickness-weighted mean horizontal velocity and the mean dianeutral velocity), while the right-hand side of this equation is the standard notation indicating that \hat{S}_* is being diffused along neutral tangent planes with the diffusivity K and in the vertical direction with the diapycnal diffusivity D (and h here is the average thickness between two closely spaced neutral tangent planes). The model is initialized with values of Preformed Salinity using Eqn. (5) based on observations of Practical Salinity and on the interpolated global observed data base of R^δ ; this is best done by calling **gsw_Sstar_from_SP**.

In order to evaluate density during the running of an ocean model, Absolute Salinity must be evaluated based on the model's primary salinity variable, Preformed Salinity, and Eqn. (6). This can be done by carrying the following evolution equation for F^δ

$$\frac{dF^\delta}{dt} = h^{-1}\nabla_n \cdot (hK\nabla_n F^\delta) + \left(D \frac{\partial F^\delta}{\partial z} \right)_z + \tau^{-1}(F^{\delta\text{obs}} - F^\delta). \quad (9)$$

The model variable F^δ (note that $F^\delta = S_A/S_* - 1$) is initialized based on observations of $R^\delta \equiv \delta S_A^{\text{ref}}/S_R^{\text{ref}}$ and the use of Eqn. (7b); this is best done by calling **gsw_Fdelta**. Equation (9) shows that F^δ is advected and diffused like any other tracer, but in addition, there is a non-conservative source term $\tau^{-1}(F^{\delta\text{obs}} - F^\delta)$ which serves to restore the model variable F^δ towards the observed value (found from **gsw_Fdelta**) with a restoring time τ that can be chosen to suit particular modeling needs (see the discussion in appendix A.20 of the TEOS-10 Manual, IOC *et al.* (2010)).

In summary, the approach for handling salinity in ocean models suggested in IOC *et al.* (2010) and summarized here carries the evolution Eqns. (8) and (9) for \hat{S}_* and F^δ , while \hat{S}_A is calculated from these two model variables at each time step according to

$$\hat{S}_A = \hat{S}_*(1 + F^\delta). \quad (10)$$

It is this salinity, \hat{S}_A , which is used as the argument for the model's expression for density at each time step of the model.

The Baltic Sea is somewhat of an exception because its compositional variations are not due to biogeochemistry but to anomalous riverine input of dissolved salts which behave conservatively. Preformed Salinity S_* in the Baltic is equal to Absolute Salinity S_A , which implies that $r_1 = -1$ in the Baltic Sea. Hence in the Baltic, an ocean model simply puts $S_A = S_*$ and the value of Absolute Salinity Anomaly δS_A is immaterial during the running of the model. Of course the values of δS_A in the Baltic are important for relating Absolute Salinity and Preformed Salinity to measured values of Practical Salinity there.

If an ocean model is to be run for only a short time (less than a century) then it may be sufficiently accurate to carry only one salinity variable, namely Absolute Salinity S_A . For longer integrations the neglect of the non-conservative biogeochemical source term means that the model's salinity variable S_A will depart from reality. A more detailed discussion of these points is available in appendix A.20 of the TEOS-10 Manual (IOC *et al.* (2010)).

In summary, the changes needed to make ocean models TEOS-10 compatible are

- (i) use an equation of state in terms of S_A and Θ , $\hat{\rho}(S_A, \Theta, p)$, such as the 48-term expression to be found in **gsw_rho**(S_A, CT, p),
- (ii) have Conservative Temperature Θ as the model's temperature variable (note that SST needs to be evaluated in the model's air-sea flux module using **gsw_pt_from_CT** at the sea surface only),
- (iii) incorporate the effects of the spatially variable seawater composition using the techniques of appendix A.20 of IOC *et al.* (2010) as summarized above,
- (iv) restoring boundary conditions for ocean-only models can be imposed on the model variables S_* and Θ ,
- (v) model output salinities and temperatures are best made as Absolute Salinity S_A and Conservative Temperature Θ , consistent with the variables which will be published in oceanographic journals.

10. A guide to the GSW Oceanographic Toolbox

The key attributes of the three new oceanographic variables S_A , S_* and Θ may be summarized as follows. Preformed Salinity S_* and Conservative Temperature Θ are the ideal variables for representing the “salt content” and “heat content” of seawater in the standard conservation equations of physical oceanography. However, the thermodynamic properties of seawater (in particular, density) depend not on Preformed Salinity S_* , but rather on Absolute Salinity S_A . While Practical Salinity S_p is relatively easy to measure accurately, it should now be regarded as a stepping stone on the way to calculating the two more attractive salinity variables, S_A and S_* .

The GSW functions are listed on the central two pages of this document. These two pages are also available as a separate double-sided laminated A4 page for ready reference. Printed versions of the present document and of the laminated list of GSW functions are available from the authors. The GSW functions are grouped under several headings of functions with similar characteristics. The first group, called “documentation set” contains tools to assist the user with the GSW Toolbox. By running **gsw_front_page** the user gains access to many of the introductory documents that are found on the TEOS-10 web site, www.TEOS-10.org. Essentially **gsw_front_page** is the front page to the GSW Oceanographic Toolbox, and from this front page much of the TEOS-10 documentation is available (including the present document). The function **gsw_contents** displays all the GSW functions as a list from which the help files can be read by clicking on their function names. The function **gsw_check_functions** confirms that the GSW Oceanographic Toolbox is correctly installed and that there are no conflicts. This function runs three stored vertical profiles through all the other GSW functions, and checks that the outputs are within predefined limits of the correct answers. These pre-defined limits are a factor of approximately a hundred larger than the errors expected from numerical round-off (at the standard double precision of MATLAB). The user may want to run **gsw_check_functions** periodically to confirm that the software remains uncorrupted. **gsw_demo** runs and displays results from several of the GSW functions, so introducing the user to some of the features of the Toolbox.

The second group of functions are the PSS-78 routines for Practical Salinity in terms of either conductivity C or conductivity ratio R , as well as their inverse functions. The input temperature to these functions is *in situ* temperature (ITS-90), and the inverse algorithms are iterated until the Practical Salinity is equal to the input value to within 2×10^{-14} , that is, to machine precision. These functions incorporate a modified form of the extension of Hill *et al.* (1986) to Practical Salinities between zero and 2. The modification ensures that the algorithm is exactly PSS-78 for $S_p \geq 2$ and is continuous at $S_p = 2$. The last function in this

group, **gsw_SP_salinometer**, calculates Practical Salinity from the two outputs of a laboratory salinometer, namely R_t and the bath temperature.

The third group delivers the three new oceanographic variables, Absolute Salinity S_A , Preformed Salinity S_* , and Conservative Temperature Θ . The first two functions have Practical Salinity S_p , pressure, longitude and latitude as input variables. Note that virtually all of the functions which follow this third group require Absolute Salinity S_A as an input. Hence it is clear that when analyzing oceanic data, the very first function call must be to **gsw_SA_from_SP**. Hence this function is the most fundamental in the GSW toolbox. This function can be avoided only by ignoring the influence of the spatial variations of seawater composition, in which case the remaining GSW functions would be called with Reference Salinity S_R (given by calling **gsw_SR_from_SP**) in place of S_A . The function **gsw_CT_from_t** evaluates Conservative Temperature Θ , as a function of Absolute Salinity S_A , *in situ* temperature t and pressure p .

The fourth group contains just the function **gsw_SA_CT_plot** which plots the TEOS-10 version of the “ T - S ” diagram for a series of vertical profiles. The Conservative Temperature at the freezing point for $p = 0$ dbar, and user-selected potential density contours are also displayed on this $S_A - \Theta$ diagram using the 48-term expression for the density of seawater, **gsw_rho_CT**(S_A, CT, p).

The fifth grouping of functions has the heading “other conversions between temperatures, salinities, entropy, pressure and height”. Some of these functions are the reverse of those in the previous groups (namely **gsw_SP_from_SA**, **gsw_SP_from_Sstar** and **gsw_t_from_CT**) while others perform familiar functions such as **gsw_pt_from_t**(S_A, t, p, p_{ref}) which evaluates the potential temperature of the “bottle” (S_A, t, p) referenced to the pressure p_{ref} .

The next four groups of functions are all derived from the computationally-efficient 48-term expression for density, $\hat{\rho}(S_A, \Theta, p)$ of McDougall *et al.* (2011b) (as summarized in appendices A.30 and K of IOC *et al.* (2010)). The first group called “density and enthalpy, based on the 48-term expression for density, $\hat{\rho}(S_A, \Theta, p)$ ” contains the function **gsw_rho_CT** to evaluate both density and potential density, and **gsw_alpha_CT** to evaluate the relevant thermal expansion coefficient. This 48-term expression for density is essentially as accurate as the full TEOS-10 expression for density, and this 48-term expression has the advantage that its temperature argument is Conservative Temperature. The functions **gsw_enthalpy_CT** and **gsw_enthalpy_diff_CT** are used when evaluating various geostrophic streamfunctions, since under isentropic and isohaline conditions, enthalpy is the pressure integral of specific volume. The functions **gsw_SA_from_rho_CT** and **gsw_CT_from_rho** are essential the inverse functions of the equation of state in that they return the Absolute Salinity (or Conservative Temperature respectively) for given values of density, pressure and either Θ or S_A respectively. Note that all of the functions in this group which end in “_CT” can also be called without the last “_CT” of each function name. For example, the functions **gsw_rho**(S_A, CT, p) and **gsw_rho_CT**(S_A, CT, p) are identical. Both forms of the function names have been retained so that the functional dependence on Θ rather than on t can be emphasized in this table.

The next group of three functions delivers variables which are defined in terms of the vertical gradients of S_A and Θ on an individual vertical profile, and so are inherently water column properties. These functions deliver the square of the buoyancy frequency (**gsw_Nsquared**), the Turner angle, and the ratio of the vertical gradient of potential density to the vertical gradient of locally-referenced potential density.

The next group of functions, concerned with various neutral and non-linear attributes of the seawater equation of state, returns properties such as the cabbeling coefficient (**gsw_cabbeling**) and the thermobaric coefficient (**gsw_thermobaric**) which are concerned with how the non-linear nature of the equation of state causes mean diapycnal advection in the ocean, even in the absence of small-scale diapycnal mixing.

The following group is for calculating four different geostrophic streamfunctions. All of these GSW geostrophic streamfunction functions have S_A and Θ as their input salinity and temperature. It is important to realize that a particular geostrophic streamfunction is only accurate when used in the surface for which it is derived. For example, dynamic height anomaly is the geostrophic streamfunction in an isobaric surface while the Montgomery streamfunction is the geostrophic streamfunction in a specific volume anomaly surface. When one is working in some type of approximately neutral surface, the Cunningham geostrophic streamfunction is more accurate than the Montgomery streamfunction, while the “isopycnal” geostrophic streamfunction **gsw_geo_strf_isopycnal** of McDougall and Klocker (2010) is the most accurate (see Figures 1, 2 and 3 of McDougall and Klocker (2010)).

The next group contains just the one function, **gsw_geostrophic_velocity**, which calculates the geostrophic velocity in a given surface with respect to the velocity in a reference surface. This function should be called with dynamic height anomaly if the surface in which the geostrophic velocity is required is an isobaric surface. Similarly, **gsw_geostrophic_velocity** should be called with the “isopycnal” geostrophic streamfunction **gsw_geo_strf_isopycnal** if the surface in which the geostrophic velocity is evaluated is an approximately neutral surface (such as a Neutral Density surface (Jackett and McDougall (1997)), an ω -surface (Klocker *et al.* (2010)) or a potential density surface).

The following group “derivatives of enthalpy, entropy, CT and pt” contains functions which use the full TEOS-10 Gibbs function and have a variety of input temperatures, appropriate to the variable being differentiated. The outputs of these functions are used, for example, in evaluating the amount of non-conservative production associated with each variable (enthalpy, entropy, CT and pt) when two seawater parcels are mixed.

The following group gives the freezing temperatures of seawater (both the Conservative Temperature and the *in situ* temperature at freezing). These freezing temperatures are functions of Absolute Salinity, pressure and the air saturation_fraction (which must be between 0 and 1). The functions **gsw_brineSA_CT** and **gsw_brineSA_t** return the Absolute Salinity at which seawater freezes at given values of Conservative Temperature (or *in situ* temperature, respectively), pressure and saturation_fraction. This group is followed by a group of functions which gives the latent heats of melting and of evaporation.

The next group, “planet Earth properties”, delivers straightforward properties of the rotating planet of the solar system on which we presently reside.

The following group, “Absolute Pressure P and sea pressure p” contains two functions for transforming between these two different types of pressure.

The group “TEOS-10 constants” simply returns various constants which are basic to TEOS-10. Note that the constant **gsw_C3515** is not a fundamental constant of either PSS-78 or TEOS-10 but is required to convert a measured conductivity value C into conductivity ratio R (which is a fundamental property of PSS-78).

The final two groups contain only functions evaluated using the full TEOS-10 Gibbs function (being the sum of the IAPWS-09 and IAPWS-08 Gibbs functions). The group of GSW functions, headed “density and enthalpy in terms of CT, based on the exact Gibbs function” delivers exactly the same outputs as the corresponding group based on the 48-term expression for density, $\hat{\rho}(S_A, \Theta, p)$, having also the same inputs as those functions. The functions in this group can be used to confirm that the use of the 48-term computationally efficient equation of state does not noticeably degrade any output property.

The final group of GSW functions is headed “basic thermodynamic properties in terms of in-situ t , based on the exact Gibbs function”. These functions have *in situ* temperature t as their input temperature variable. All the functions in this group use the full TEOS-10 Gibbs function, namely the sum of the Gibbs functions of IAPWS-09 and IAPWS-08 (rather than the 48-term expression for density). Many of the functions in this group are called by the functions in the previous group.

The GSW Oceanographic Toolbox is designed to be comprehensive and to be installed in its entirety, even though most users may use relatively few of the functions for routine oceanographic analyses. For example, the most basic use of the GSW Oceanographic Toolbox would begin with a data set of (S_p, t, p) at known longitudes and latitudes. The first steps are to call **gsw_SA_from_SP** and then **gsw_CT_from_t** to convert to a data set of (S_A, Θ, p) . With the data set in this form, water masses may be analyzed accurately on the $S_A - \Theta$ diagram, and *in situ* density and potential density are available by calling the computationally-efficient 48-term expression for density, **gsw_rho**, with the pressure input being the *in situ* sea pressure p , and the reference sea pressure p_{ref} , respectively. That is, *in situ* density is evaluated as **gsw_rho**(SA,CT,p) and potential density with respect to the reference pressure p_{ref} is given by **gsw_rho**(SA,CT, p_{ref}).

In addition to the functions discussed above and listed on the central two pages of this document, there is an additional group of library functions which are listed below. These are internal functions which are not intended to be called by users. There is nothing stopping a skilled operator using these programs, but unless the user is confident, it is safer to access these library routines via one of the public functions; for example, there is little or no checking on the array sizes of the input variables in these internal library functions. The data set gsw_data_v3_0 must not be tampered with.

library functions of the GSW Toolbox (internal functions; not intended to be called by users)

The GSW functions on pages 14 and 15 call the following library functions,

gsw_gibbs	the TEOS-10 Gibbs function and its derivatives
gsw_SAAR	Absolute Salinity Anomaly Ratio (excluding the Baltic Sea)
gsw_Fdelta	ratio of Absolute to Preformed Salinity, minus 1
gsw_delta_SA_ref	Absolute Salinity Anomaly ref. value (excluding the Baltic Sea)
gsw_SA_from_SP_Baltic	Absolute Salinity from SP in the Baltic Sea
gsw_SP_from_SA_Baltic	Practical Salinity from SA in the Baltic Sea
gsw_infunnel	"oceanographic funnel" check for the 48-term density equation
gsw_entropy_part	entropy minus the terms that are a function of only SA
gsw_entropy_part_zerop	entropy_part evaluated at 0 dbar
gsw_interp_ref_cast	linearly interpolates the reference cast of the isopycnal streamfunction
gsw_interp_SA_CT	linearly interpolates (SA,CT,p) to the desired p
gsw_gibbs_pt0_pt0	$\text{gibbs}(0,2,0,SA,t,0)$
gsw_specvol_SSO_0_p	$\text{specvol_CT}(35.16504,0,p)$
gsw_enthalpy_SSO_0_p	$\text{enthalpy_CT}(35.16504,0,p)$
gsw_Hill_ratio_at_SP2	Hill ratio at a Practical Salinity of 2

The GSW data set

gsw_data_v3_0	contains
	(1) the global data set of Absolute Salinity Anomaly Ratio, R^δ ,
	(2) the global data set of Absolute Salinity Anomaly ref., δS_A^{ref} ,
	(3) a reference cast (for the isopycnal streamfunction),
	(4) two reference casts that are used by gsw_demo, and
	(5) three vertical profiles of (SP, t, p) at known long & lat, plus
	the outputs of all the GSW functions for these 3 profiles, and
	the required accuracy of all these outputs.

licences and 3rd party software

gsw_licence	licence for the GSW Oceanographic Toolbox
cprintf	prints colour text to the screen (3 rd party software)
cprintf_licence	licence for cprintf software

11. References

- Feistel, R., 2003: A new extended Gibbs thermodynamic potential of seawater, *Progr. Oceanogr.*, **58**, 43-114.
- Feistel, R., 2008: A Gibbs function for seawater thermodynamics for -6 to 80 °C and salinity up to 120 g kg^{-1} , *Deep-Sea Res. I*, **55**, 1639-1671.
- Feistel, R., S. Weinreben, H. Wolf, S. Seitz, P. Spitzer, B. Adel, G. Nausch, B. Schneider and D. G. Wright, 2010: Density and Absolute Salinity of the Baltic Sea 2006–2009. *Ocean Science*, **6**, 3–24. <http://www.ocean-sci.net/6/3/2010/os-6-3-2010.pdf>
- Hill, K. D., T. M. Dauphinee and D. J. Woods, 1986: The extension of the Practical Salinity Scale 1978 to low salinities, *IEEE J. Oceanic Eng.*, **11**, 109–112.
- IAPWS, 2008: Release on the IAPWS Formulation 2008 for the Thermodynamic Properties of Seawater. The International Association for the Properties of Water and Steam. Berlin, Germany, September 2008, available from www.iapws.org. This Release is referred to in the text as **IAPWS-08**.
- IAPWS, 2009: Supplementary Release on a Computationally Efficient Thermodynamic Formulation for Liquid Water for Oceanographic Use. The International Association for the Properties of Water and Steam. Doorwerth, The Netherlands, September 2009, available from <http://www.iapws.org>. This Release is referred to in the text as **IAPWS-09**.
- IOC, SCOR and IAPSO, 2010: *The international thermodynamic equation of seawater – 2010: Calculation and use of thermodynamic properties*. Intergovernmental Oceanographic Commission, Manuals and Guides No. 56, UNESCO (English), 196 pp. Available from <http://www.TEOS-10.org>
- Jackett, D. R. and T. J. McDougall, 1997: A neutral density variable for the world's oceans. *Journal of Physical Oceanography*, **27**, 237-263.
- Klocker, A., T. J. McDougall and D. R. Jackett, 2009: A new method for forming approximately neutral surfaces. *Ocean Sci.*, **5**, 155-172.
- McDougall T. J. and P. M. Barker, 2011: Getting started with TEOS-10 and the Gibbs Seawater (GSW) Oceanographic Toolbox, 28pp., SCOR/IAPSO WG127, ISBN 978-0-646-55621-5, available from www.TEOS-10.org (the present document)
- McDougall T. J., P. M. Barker, R. Feistel and D. R. Jackett, 2011b: A computationally efficient 48-term expression for the density of seawater in terms of Conservative Temperature, and related properties of seawater. To be submitted to *J. Atm. Ocean. Technol.*, **xx**, yyy-zzz.
- McDougall, T. J., D. R. Jackett and F. J. Millero, 2011a: An algorithm for estimating Absolute Salinity in the global ocean. submitted to *Ocean Science*, a preliminary version is available at *Ocean Sci. Discuss.*, **6**, 215-242.
- McDougall, T. J. and A. Klocker, 2010: An approximate geostrophic streamfunction for use in density surfaces. *Ocean Modelling*, **32**, 105-117.
- Millero, F. J., R. Feistel, D. G. Wright, and T. J. McDougall, 2008: The composition of Standard Seawater and the definition of the Reference-Composition Salinity Scale, *Deep-Sea Res. I*, **55**, 50-72.
- Pawlowicz, R., 2010: What every oceanographer needs to know about TEOS-10 (The TEOS-10 Primer), unpublished manuscript, available from www.TEOS-10.org.
- Pawlowicz, R., D. G. Wright and F. J. Millero, 2011: The effects of biogeochemical processes on oceanic conductivity/salinity/density relationships and the characterization of real seawater. *Ocean Science*, **7**, 363–387. <http://www.ocean-sci.net/7/363/2011/os-7-363-2011.pdf>
- Wright, D. G., R. Pawlowicz, T. J. McDougall, R. Feistel and G. M. Marion, 2011: Absolute Salinity, “Density Salinity” and the Reference-Composition Salinity Scale: present and future use in the seawater standard TEOS-10. *Ocean Sci.*, **7**, 1-26. <http://www.ocean-sci.net/7/1/2011/os-7-1-2011.pdf>

12. Recommended nomenclature, symbols and units in oceanography

This section 12 is from appendix L of the TEOS-10 Manual, IOC *et al.* (2010).

L.1 Recommended nomenclature

The strict SI units of Absolute Salinity, temperature and pressure are kg kg^{-1} , Absolute Temperature in K and Absolute Pressure P in Pa. These are the units predominantly adopted in the SIA computer software for the input and output variables. If oceanographers were to adopt this practice of using strictly SI quantities it would simplify many thermodynamic expressions at the cost of using unfamiliar units.

The GSW Oceanographic Toolbox (appendix N) adopts as far as possible the currently used oceanographic units, so that the input variables for all the computer algorithms are Absolute Salinity in S_A in g kg^{-1} , temperature in $^{\circ}\text{C}$ and pressure as sea pressure in dbar. The outputs of the functions are also generally consistent with this choice of units, but some variables are more naturally expressed in SI units.

It seems impractical to recommend that the field of oceanography fully adopt strict basic SI units. It is however very valuable to have the field adopt uniform symbols and units, and in the interests of achieving this uniformity we recommend the following symbols and units. These are the symbols and units we have adopted in the GSW Oceanographic Toolbox.

Table L.1. Recommended Symbols and Units in Oceanography

Quantity	Symbol	Units	Comments
Chlorinity	Cl	g kg^{-1}	<i>Chlorinity</i> is defined as the following mass fraction; it is 0.328 523 4 times the ratio of the mass of pure silver required to precipitate all dissolved chloride, bromide and iodide in seawater to the mass of seawater.
Standard Ocean Reference Salinity	S_{SO}	g kg^{-1}	35.165 04 g kg^{-1} being exactly 35 u_{PS} , corresponding to the standard ocean Practical Salinity of 35.
freezing temperatures	t^f, Θ^f	$^{\circ}\text{C}$	<i>in situ</i> and conservative values, each as a function of S_A and p .
absolute pressure	P	Pa	When absolute pressure is used it should always be in Pa, not in Mpa nor in dbar.
sea pressure. Sea pressure is the pressure argument to all the GSW Toolbox functions.	p	dbar	Equal to $P - P^0$ and usually expressed in dbar not Pa.

gauge pressure. Gauge pressure (also called applied pressure) is sometimes reported from ship-born instruments.	p^{gauge}	dbar	Equal to the absolute pressure P minus the local atmospheric pressure at the time of the instrument calibration, and expressed in dbar not Pa. Sea pressure p is preferred over gauge pressure p^{gauge} , as p is the argument to the seawater Gibbs function.
reference pressure	p_r	dbar	The value of the sea pressure p to which potential temperature and/or potential density are referenced.
one standard atmosphere	P_0	Pa	exactly 101 325 Pa (= 10.1325 dbar)
isopycnal slope ratio	r	1	$r = \frac{\alpha^\Theta(p)/\beta^\Theta(p)}{\alpha^\Theta(p_r)/\beta^\Theta(p_r)}$
Stability Ratio	R_ρ	1	$R_\rho = \alpha^\Theta \Theta_z / \beta^\Theta (S_A)_z \approx \alpha^\Theta \theta_z / \beta^\Theta (S_A)_z .$
isopycnal temperature gradient ratio	G^Θ	1	$G^\Theta = r[R_\rho - 1] / [R_\rho - r]; \quad \nabla_\sigma \Theta = G^\Theta \nabla_n \Theta$
Practical Salinity	S_p	1	Defined in the range $2 < S_p < 42$ by PSS-78 based on measured conductivity ratios.
Reference Salinity	S_R	g kg^{-1}	Reference-Composition Salinity (or Reference Salinity for short) is the Absolute Salinity of seawater samples that have Reference Composition. At $S_p = 35$, S_R is exactly $u_{\text{PS}} S_p$. while in the range $2 < S_p < 42$ $S_R \approx u_{\text{PS}} S_p$.
Absolute Salinity (This is the salinity argument to all the GSW Toolbox functions.)	$S_A = S_A^{\text{dens}}$	g kg^{-1}	$S_A = S_R + \delta S_A \approx u_{\text{PS}} S_p + \delta S_A$ Absolute Salinity is the sum of S_R on the Millero <i>et al.</i> (2008a) Reference-Salinity Scale and the Absolute Salinity Anomaly. The full symbol for S_A is S_A^{dens} as it is the type of absolute salinity which delivers the best estimate of density when used as the salinity argument of the TEOS-10 Gibbs function. Another name for $S_A = S_A^{\text{dens}}$ is “Density Salinity”.
Absolute Salinity Anomaly	δS_A	g kg^{-1}	$\delta S_A = S_A - S_R$, the difference between Absolute Salinity, $S_A = S_A^{\text{dens}}$, and Reference-Composition Salinity. In terms of the full nomenclature of Pawlowicz <i>et al.</i> (2010), Wright <i>et al.</i> (2010b) and appendix A.4 herein, the Absolute Salinity Anomaly δS_A is δS_R^{dens} .
“Preformed Absolute Salinity”, often shortened to “Preformed Salinity”	S_*	g kg^{-1}	Preformed Absolute Salinity S_* is a salinity variable that is designed to be as conservative as possible, by removing the estimated biogeochemical influences on the seawater composition from other forms of salinity (see Pawlowicz <i>et al.</i> (2010), Wright <i>et al.</i> (2010b) and appendix A.4 herein).
“Solution Absolute Salinity”, often shortened to “Solution Salinity”	S_A^{soln}	g kg^{-1}	The mass fraction of non-H ₂ O constituents in seawater after it has been brought to chemical equilibrium at $t = 25^\circ\text{C}$ and $p = 0$ dbar (see Pawlowicz <i>et al.</i> (2010), Wright <i>et al.</i> (2010b) and appendix A.4 herein).
“Added-Mass Salinity”	S_A^{add}	g kg^{-1}	$S_A^{\text{add}} - S_R$ is the estimated mass fraction of non-H ₂ O constituents needed as ingredients to be added to Standard Seawater which when mixed and brought to chemical equilibrium at $t = 25^\circ\text{C}$ and $p = 0$ dbar results in the observed seawater composition.

temperature	t	°C	
Absolute Temperature	T	K	$T / \text{K} \equiv T_0 / \text{K} + t / (^\circ\text{C}) = 273.15 + t / (^\circ\text{C})$
temperature derivatives	T	K	When a quantity is differentiated with respect to <i>in situ</i> temperature, the symbol T is used in order to distinguish this variable from time.
Celsius zero point	T_0	K	$T_0 \equiv 273.15 \text{ K}$
potential temperature	θ	°C	Defined implicitly by Eqn. (3.1.3)
Conservative Temperature	Θ	°C	Defined in Eqn. (3.3.1) as exactly potential enthalpy divided by c_p^0 .
the “specific heat”, for use with Conservative Temperature	c_p^0	$\text{J kg}^{-1} \text{K}^{-1}$	$c_p^0 \equiv 3991.867\,957\,119\,63 \text{ J kg}^{-1} \text{K}^{-1}$. This 15-digit number is defined to be the exact value of c_p^0 . c_p^0 is the ratio of potential enthalpy h^0 to Θ .
combined standard uncertainty	u_c	Varies	
enthalpy	H	J	
specific enthalpy	h	J kg^{-1}	$h = u + (p + P_0)v$. Here p and P_0 must be in Pa not dbar.
specific potential enthalpy	h^0	J kg^{-1}	specific enthalpy referenced to zero sea pressure, $h^0 = h(S_A, \theta[S_A, t, p, p_r = 0], p_r = 0)$
specific isobaric heat capacity	c_p	$\text{J kg}^{-1} \text{K}^{-1}$	$c_p = \partial h / \partial T _{S_A, p}$
internal energy	U	J	
specific internal energy	u	J kg^{-1}	
specific isochoric heat capacity	c_v	$\text{J kg}^{-1} \text{K}^{-1}$	$c_v = \partial u / \partial T _{S_A, v}$
Gibbs function (Gibbs energy)	G	J	
specific Gibbs function (Gibbs energy)	g	J kg^{-1}	
specific Helmholtz energy	f	J kg^{-1}	
unit conversion factor for salinities	u_{ps}	g kg^{-1}	$u_{\text{ps}} \equiv (35.16504/35) \text{ g kg}^{-1} \approx 1.004\,715\dots \text{g kg}^{-1}$ The first part of this expression is exact. This conversion factor is an important and invariant constant of the 2008 Reference-Salinity Scale (Millero <i>et al.</i> (2008a)).
entropy	Σ	J K^{-1}	
specific entropy	η	$\text{J kg}^{-1} \text{K}^{-1}$	In many other publications the symbol s is used for specific entropy.
density	ρ	kg m^{-3}	
density anomaly	σ^t	kg m^{-3}	$\rho(S_A, t, 0) - 1000 \text{ kg m}^{-3}$
potential density anomaly referenced to a sea pressure of 2000 dbar	σ_2	kg m^{-3}	$\rho(S_A, \theta[S_A, t, p, p_r], p_r) - 1000 \text{ kg m}^{-3}$ where $p_r = 2000 \text{ dbar}$
potential density anomaly referenced to a sea pressure of 4000 dbar	σ_4	kg m^{-3}	$\rho(S_A, \theta[S_A, t, p, p_r], p_r) - 1000 \text{ kg m}^{-3}$ where $p_r = 4000 \text{ dbar}$
thermal expansion coefficient with respect to <i>in situ</i> temperature	α^t	K^{-1}	$v^{-1} \partial v / \partial T _{S_A, p} = -\rho^{-1} \partial \rho / \partial T _{S_A, p}$
thermal expansion coefficient with respect to potential temperature θ	α^θ	K^{-1}	$v^{-1} \partial v / \partial \theta _{S_A, p} = -\rho^{-1} \partial \rho / \partial \theta _{S_A, p}$

thermal expansion coefficient with respect to Conservative Temperature Θ	α^Θ	K^{-1}	$v^{-1}\partial v / \partial \Theta _{S_A, p} = -\rho^{-1}\partial \rho / \partial \Theta _{S_A, p}$
saline contraction coefficient at constant <i>in situ</i> temperature	β^t	kg g^{-1}	$-v^{-1}\partial v / \partial S_A _{T, p} = \rho^{-1}\partial \rho / \partial S_A _{T, p}$ Note that the units for β^t are consistent with S_A being in g kg^{-1} .
saline contraction coefficient at constant potential temperature	β^θ	kg g^{-1}	$-v^{-1}\partial v / \partial S_A _{\theta, p} = \rho^{-1}\partial \rho / \partial S_A _{\theta, p}$ Note that the units for β^θ are consistent with S_A being in g kg^{-1} .
saline contraction coefficient at constant Conservative Temperature	β^Θ	kg g^{-1}	$-v^{-1}\partial v / \partial S_A _{\Theta, p} = +\rho^{-1}\partial \rho / \partial S_A _{\Theta, p}$ Note that the units for β^Θ are consistent with S_A being in g kg^{-1} .
isothermal compressibility	κ^t	Pa^{-1}	
isentropic and isohaline compressibility	κ	Pa^{-1}	
chemical potential of water in seawater	μ^w	J g^{-1}	
chemical potential of sea salt in seawater	μ^s	J g^{-1}	
relative chemical potential of (sea salt and water in) seawater	μ	J g^{-1}	$(\partial g / \partial S_A)_{t, p} = \mu^s - \mu^w$
dissipation rate of kinetic energy per unit mass	ε	$\text{J kg}^{-1} \text{s}^{-1} = \text{m}^2 \text{s}^{-3}$	
adiabatic lapse rate	Γ	K Pa^{-1}	$\Gamma = \frac{\partial t}{\partial P} _{S_A, \theta} = \frac{\partial t}{\partial P} _{S_A, \Theta} = \frac{\partial t}{\partial P} _{S_A, \eta}$
sound speed	c	m s^{-1}	
specific volume	v	$\text{m}^3 \text{kg}^{-1}$	$v = \rho^{-1}$
specific volume anomaly	δ	$\text{m}^3 \text{kg}^{-1}$	
thermobaric coefficient based on θ	T_b^θ	$\text{K}^{-1} \text{Pa}^{-1}$	$T_b^\theta = \beta^\theta \partial (\alpha^\theta / \beta^\theta) / \partial P _{S_A, \theta}$
thermobaric coefficient based on Θ	T_b^Θ	$\text{K}^{-1} \text{Pa}^{-1}$	$T_b^\Theta = \beta^\Theta \partial (\alpha^\Theta / \beta^\Theta) / \partial P _{S_A, \Theta}$
cabbeling coefficient based on θ	C_b^θ	K^{-2}	$C_b^\theta = \partial \alpha^\theta / \partial \theta _{S_A, p} + 2 \frac{\alpha^\theta}{\beta^\theta} \partial \alpha^\theta / \partial S_A _{\theta, p} - \left(\frac{\alpha^\theta}{\beta^\theta}\right)^2 \partial \beta^\theta / \partial S_A _{\theta, p}$
cabbeling coefficient based on Θ	C_b^Θ	K^{-2}	$C_b^\Theta = \partial \alpha^\Theta / \partial \Theta _{S_A, p} + 2 \frac{\alpha^\Theta}{\beta^\Theta} \partial \alpha^\Theta / \partial S_A _{\Theta, p} - \left(\frac{\alpha^\Theta}{\beta^\Theta}\right)^2 \partial \beta^\Theta / \partial S_A _{\Theta, p}$
buoyancy frequency	N	s^{-1}	$N^2 = g(\alpha^\Theta \Theta_z - \beta^\Theta S_{A_z}) = g(\alpha^\theta \theta_z - \beta^\theta S_{A_z})$
neutral helicity	H^n	m^{-3}	defined by Eqns. (3.13.1) and (3.13.2)
Neutral Density	γ^n	kg m^{-3}	a density variable whose iso-surfaces are designed to be approximately neutral, i. e. $\alpha^\Theta \nabla_\gamma \Theta \approx \beta^\Theta \nabla_\gamma S_A$.
Neutral-Surface-Potential-Vorticity	$NSPV$	s^{-3}	$NSPV = -g\rho^{-1}f\gamma_z^n$ where f is the Coriolis parameter.
dynamic height anomaly	Ψ	$\text{m}^2 \text{s}^{-2}$	$\text{Pa m}^3 \text{kg}^{-1} = \text{m}^2 \text{s}^{-2}$
Montgomery geostrophic streamfunction	Ψ^M	$\text{m}^2 \text{s}^{-2}$	$\text{Pa m}^3 \text{kg}^{-1} = \text{m}^2 \text{s}^{-2}$

PISH (Pressure-Integrated Steric Height)	Ψ'	kg s^{-2}	streamfunction for f times the depth-integrated relative mass flux, see Eqns. (3.31.1) – (3.31.5).
Coriolis parameter	f	s^{-1}	$1.458\,42 \times 10^{-4} \sin \phi \text{ s}^{-1}$, where ϕ is latitude
molality	m_{sw}	mol kg^{-1}	$m_{\text{sw}} = \sum_i m_i = \frac{1}{M_S} \frac{S_A}{(1-S_A)}$ where M_S is the mole-weighted average atomic weight of the elements of sea salt, $M_S = 0.031\,403\,821\,8 \dots \text{kg mol}^{-1}$
ionic strength	I	mol kg^{-1}	$I = \frac{1}{2} m_{\text{sw}} \langle Z^2 \rangle = \frac{1}{2} \sum_i m_i z_i^2$ $= 0.622\,644\,9 m_{\text{sw}}$ $\approx \frac{622.644\,9}{31.403\,821\,8} \text{mol kg}^{-1} \frac{S_A}{(1-S_A)}$
osmotic coefficient	ϕ	1	$\phi(S_A, T, p) = \frac{g(0, t, p) - \mu^{\text{w}}(S_A, t, p)}{m_{\text{sw}} R (T_0 + t)}$ where the molar gas constant, $R = 8.314\,472 \text{ J mol}^{-1} \text{ K}^{-1}$. See also Eqns. (2.14.1) and (3.40.9) for an equivalent definition of ϕ .

Acknowledgements

The authors of the present document acknowledge partial financial support from CSIRO and the Wealth from Oceans National Flagship during the development of the GSW Oceanographic Toolbox.


TEOS-10 and GSW in a nutshell

In order to analyse oceanographic data under TEOS-10, the observed values of Practical Salinity S_p and *in situ* temperature t need to be converted into Absolute Salinity S_A and Conservative Temperature Θ , as follows,


Step 1. calculate Absolute Salinity, $S_A = \text{gsw_SA_from_SP}(S_p, p, \text{long}, \text{lat})$,

Step 2. calculate Conservative Temperature, $\Theta = \text{gsw_CT_from_t}(S_A, t, p)$.

Having converted (S_p, t, p) to (S_A, Θ, p) , the GSW functions listed on page 14 of this document are then used for analysing the data. The use of these GSW functions ensures consistency between theoretical oceanography, observational oceanography and ocean modelling.



Under TEOS-10, the $S_A - \Theta$ diagram is the new “T-S” diagram. The above $S_A - \Theta$ diagram was plotted using `gsw_SA_CT_plot`. The σ_2 potential density anomaly contours were evaluated inside this function using `gsw_rho_CT(SA,CT,2000)`. This same function can be used to evaluate *in situ* density via `gsw_rho_CT(SA,CT,p)`.



www.TEOS-10.org