

Task Solution Chapter 3

Witold Wolski

December 3, 2016

Task 1

NO Idea where to start.

verify formula for $var(Y)$

$$var(Y) = \frac{\phi}{w} b''(\theta)$$

For mean:

$$\begin{aligned} 0 &= E\left(\frac{\partial l}{\partial \theta}\right) \\ &= E\left(\frac{y - b'(\theta)}{\phi/w}\right) = w/\phi E(y - b'(\theta)) = w/\phi (E(y) - b'(\theta)) \\ &\rightarrow E(Y) = b'(\theta) \end{aligned}$$

According to Result 2 on page 64

with:

$$\begin{aligned} l(\theta, \phi, y) &= \frac{y\theta - b(\theta)}{\phi/w} + c(y, \phi) \\ -E\left(\frac{\partial^2 l}{\partial \theta^2}\right) &= E\left(\frac{\partial^2 l}{\partial \theta^2}\right) \\ -E(-wb''(\theta)/\phi) &= E\left(\left(\frac{y - b'(\theta)}{\phi/w}\right)^2\right) = (w/\phi)^2 E((y - E(Y))^2) \\ \frac{\phi}{w} b''(\theta) &= Var(Y) \end{aligned}$$

Task 2

Use equations (3.8) and (3.9) to verify the entries in the table above.

Equation (3.8) is:

$$\mu = E(Y) = b'(\theta)$$

Equation (3.9) is:

$$var(Y) = \frac{\phi}{w} b''(\theta)$$

i

For normal distribution:

$$b(\theta) = \theta^2/2 \rightarrow b'(\theta) = \theta \rightarrow b''(\theta) = 1$$

ii

For Poisson distribution

$$b(\theta) = e^\theta \rightarrow b'(\theta) = e^\theta \rightarrow b''(\theta) = e^\theta$$

iii

For Binomial

$$b(\theta) = \log(1 + e^\theta) \rightarrow b'(\theta) = \frac{e^\theta}{1 + e^\theta} \rightarrow b''(\theta) = \frac{e^\theta}{e^\theta + 1} - \frac{e^{2\theta}}{(e^\theta + 1)^2}$$

Need to check how to differentiate this.

iv

For Gamma

$$b(\theta) = -\log(-\theta) \rightarrow b'(\theta) = +1/\theta \rightarrow b''(\theta) = 1/\theta^2 = (-1\theta)^2 = \mu^2$$

Task 3

Task 3 Verify the canonical links for the Normal, Binomial, Poisson and Gamma.

The link function g for which $\theta_i = x_i^T \beta$ (that is, for which $g = (b')^{-1}$ is called the canonical link.

Hence, the task is to find the inverse of $g = (b')^{-1}$.

for Normal

$$b'(\theta) = \theta \rightarrow \theta = g(\mu) = 1(\mu)$$

for Poisson

$$b'(\theta) = e^\theta \rightarrow \mu = e^\theta \rightarrow \theta = \log(\mu)$$

for Binomial

$$b'(\theta) = \log(1 + e^\theta) \rightarrow \mu = \log(1 + e^\theta) \rightarrow \theta = \log(\mu_i / (1 - \mu_i))$$

For Gamma

$$b'(\theta) = -1/\theta \rightarrow \mu = -1/\theta \rightarrow \theta = g(\mu) = -1/\mu$$

Task 4

For the canonical link (see, in particular, equation (3.10)) show that

$$\begin{aligned}\sum w_i y_i \frac{\partial \theta}{\partial \beta} &= \sum w_i \frac{\partial b(\theta)}{\partial \beta} \\ \frac{\partial^2 l}{\partial \beta_j \partial \beta_k} &= \\ &= \frac{1}{\phi} \frac{\partial^2 l}{\partial \beta_j \partial \beta_k} \left(\sum_i \{w_i [y_i \theta_i - b(\theta_i)]\} \right) \\ &= \frac{1}{\phi} \sum_i \{-x_{ij} w_i h'(x_i^T \beta) x_{ik}\}\end{aligned}$$

Hint :

$$b'(\theta) = h(x^T \beta) \rightarrow b(\theta) = \int h(x^T \beta) + C$$

than:

$$\begin{aligned}&= \frac{1}{\phi} \frac{\partial^2 l}{\partial \beta_j \partial \beta_k} \left(\sum_i \{w_i [y_i \theta_i - b(\theta_i)]\} \right) \\ &= \frac{1}{\phi} \frac{\partial^2 l}{\partial \beta_j \partial \beta_k} \left(\sum_i \{w_i [y_i \theta_i - (\int h(x^T \beta) + C)]\} \right)\end{aligned}$$

Task 5

Show that, for the maximal model, y_i is the maximum likelihood estimate of μ_i . (Hint: the likelihood equation for θ_i is $y_i = b'(\theta_i)$, and we already know that $b'(\theta_i) = \mu_i$.)

$$y_i = b'(\theta_i) = \mu_i$$

$$\begin{aligned}l(\mu; y, \phi) &= \sum_i \frac{y_i \theta - b(\theta)}{\phi/w_i} + c(y, \phi) \\ \partial l / \partial \theta_i &= w_i / \phi \{y_i - b'(\theta_i)\} = 0 \\ \rightarrow y_i &= b'(\theta_i) = \mu_i\end{aligned}$$

The model that allows the mean of each observation to be a separate parameter is called the **full** or **saturated** or maximal model.

The maximum possible value of L as the components of μ vary without any restrictions (or, essentially equivalently, when there is an element of β for each observation, so that $\mu_i^\diamond = h(\eta_i) = h(\beta_i)$ occurs when $\mu_i^\diamond = y_i$, hence $\mu_i^\diamond = y_i$ for all i).

Task 6

Deviances for common distributions.

Confirm results in table these using the table in 3.3.4 and equation (3.12).

Eq 3.12 is:

$$D(y, \hat{\mu}) = \phi S(y, \hat{\mu}) = 2 \sum_i w_i [y_i(\hat{\theta}_i^\diamond - \hat{\theta}_i) - b(\hat{\theta}_i^\diamond) + b(\hat{\theta}_i)]$$

For Normal

$\hat{\theta}^\diamond = y_i$ and $\hat{\theta} = \mu$; $b(\theta) = \theta^2/2$ than $b(\hat{\theta}^\diamond) = y_i^2/2$ and $b(\hat{\theta}) = \mu^2/2$; and $w_i = 1$.

$$\begin{aligned} 2 \sum_i [y_i(y_i - \mu) - y_i^2/2 + \mu^2/2] \\ &= 2 \sum_i [y_i^2 - y_i\mu - y_i^2/2 + \mu^2/2] \\ &= \sum_i [y_i^2 - 2y_i\mu + \mu^2] \\ &= \sum_i (y_i - \mu)^2 \end{aligned}$$

For Poisson

$\mu = e^\theta \rightarrow \theta = \log(\mu)$. Than

$\hat{\theta} = \log(\mu)$ and $\hat{\theta}^\diamond = \log y_i$; $b(\theta) = e^\theta$ than

$b(\hat{\theta}^\diamond) = e^{\log(y_i)} = y_i$ and $b(\hat{\theta}) = e^{\log(\mu)} = \mu$ and $\phi = 1$, $w_i = 1$

$$2 \sum_i \{y_i(\log(y_i) - \log(\mu)) - y_i + \mu\} = 2 \sum_i \{y_i \log(y_i/\mu) - y_i + \mu\}$$

For Binomial

Task 7

Use the results in 3.6.4 to obtain simple expressions for the null deviances for Poisson (weights must be one) and Gamma (assume weights are one).

If $W = \sum w_i$

$$\mu = \sum w_i y_i / W$$

For Poisson and Gamma model $w_i = 1$, than $\mu = 1/N \sum y_i$

For Poisson

$$\begin{aligned}
2 \sum y_i \log(y_i/\hat{\mu}_i) - (y_i - \hat{\mu}_i) &= \\
&= 2 \sum (y_i \log(\frac{y_i}{1/N \sum y_i}) - (y_i - 1/N \sum y_i)) \\
&= 2 \sum (y_i \log(y_i) - \log(1/N \sum y_i) - (y_i) + (1/N \sum y_i)) \\
&= 2 \sum y_i (\log(y_i) - \log(1/N \sum y_i))
\end{aligned}$$

For Gamma

Similarly

Task 8

For the Poisson distribution, verify that D and χ^2 are asymptotically equivalent. (Hint: use that for small $|x - y|$, $x \log(x/y) \approx (x - y) + (x - y)^2/(2y)$.)

$$\chi^2 = \sum e_{P,i}^2 = \sum (\sqrt{w_i} \frac{y_i - \hat{\mu}_i}{\sqrt{V(\hat{\mu}_i)}})^2$$

for Poisson $w_i = 1$ and $Var(Y_i) = \frac{\phi}{w_i} V(\mu_i) = \frac{\phi}{w_i} b''(\theta)$ and $b''(\theta) = V(\mu_i) = \mu$,
than:

$$\chi^2 = \sum e_{P,i}^2 = \sum (\frac{y_i - \hat{\mu}_i}{\sqrt{\mu_i}})^2$$

$$\begin{aligned}
D(y_i, \mu_i) &= 2 \sum y_i \log(y_i/\hat{\mu}_i) - (y_i - \hat{\mu}_i) = \\
&= 2 \sum (y_i - \hat{\mu}_i) + (y_i - \hat{\mu}_i)^2/(2\hat{\mu}_i) - (y_i - \hat{\mu}_i) = \\
&= \sum ((y_i - \hat{\mu}_i)^2/\hat{\mu}_i)
\end{aligned}$$

Task 9

Obtain the form of the **Pearson residual** for the Binomial, Poisson and Gamma. (These are given in Chapters 4 and 5)

For Binomial

$$Var(\mu_i) = \phi/w_i b''(\theta) = (1/n)\mu(1 - \mu)$$

than

$$e_{p,i} = \frac{y_i - \mu_i}{Var(\mu_i)} \stackrel{with1}{=} \frac{y_i - \mu_i}{\mu_i(1 - \mu_i)/n_i}$$

For Poisson

$$Var(\mu_i) = \phi/w_i b''(\theta) = (1/1)\mu$$

than

$$e_{p,i} = \frac{y_i - \mu_i}{\mu_i}$$

For Gamma

$$Var(\mu_i) = \phi/w_i b''(\theta) = (\phi/1)\mu^2$$

than

$$e_{p,i} = \frac{y_i - \mu_i}{\phi\mu_i^2}$$

Task 10

- How does a GLM with identity link relate to a linear model?

It's the same as a linear model.

- Under which situations does it make sense to fit a linear model instead of a GLM with identity link?

The response y is continuous and normally distributed.

Task 11

Confirm that:

$$\sum e_{P,i}^2 = \sum \left\{ \frac{[s_i - n_i \hat{\mu}_i]^2}{n_i \hat{\mu}_i} + \frac{[(n_i - s_i) - n_i(1 - \hat{\mu}_i)]^2}{n_i(1 - \hat{\mu}_i)} \right\}$$

With

$$e_{P,i} = \frac{y_i - \hat{\mu}_i}{\sqrt{\hat{\mu}_i(1 - \hat{\mu}_i)/n_i}}$$

and

$$\chi^2 = \sum e_{P,i}^2$$