

Quiz 3

Chemistry 3BB3; Winter 2004

1. Write the electronic Schrödinger Equation for the Beryllium ($Z=4$, $N=4$) atom, showing the dependence on \hbar , e , ε_0 , and m_e . (Do not use atomic units. You may use the Born-Oppenheimer approximation.)
2. Write a Slater determinant wave function for the $1s^2 4s^1$ excited state of the Lithium atom. Do not use the “shorthand” notation. Denote the spatial part of the 1s orbital with $\psi_{1s}(\mathbf{r})$ and the spatial part of the 2s orbital with $\psi_{4s}(\mathbf{r})$.

Using separation of variables, the wave function for the hydrogen atom can be written as

$$\Psi_{n,l,m}(\mathbf{r}) = R_{n,l}(r) Y_l^m(\theta, \phi).$$

3. Write the Schrödinger equation for radial wave function, $R_{n,l}(r)$, in the hydrogen atom. You may use atomic units.
4. Is $\Psi(\mathbf{r}_1, \sigma(1), \mathbf{r}_2, \sigma(2)) \propto e^{-\zeta_1 r_1} e^{-\zeta_2 r_2} (1 + b|\mathbf{r}_1 - \mathbf{r}_2| + c|\mathbf{r}_1 - \mathbf{r}_2|^2)(\alpha(1)\beta(2) - \alpha(2)\beta(1))$ an acceptable approximate wave function for the Helium atom?
 (a) yes (b) no

Name:

5. Which of the following operators commute with the electronic Hamiltonian for an atom. (There may be more than one answer.)

- | | | |
|-----------------|-----------------|-----------------|
| (a) \hat{L}_x | (d) \hat{L}^2 | (g) \hat{J}_x |
| (b) \hat{L}_y | (e) \hat{S}^2 | (h) \hat{S}_y |
| (c) \hat{L}_z | (f) \hat{J}^2 | (i) \hat{S}_x |

6. Which of the following operators commute with \hat{L}_x . (There may be more than one answer.)

- | | | |
|-----------------|-----------------|-----------------|
| (a) \hat{L}_x | (e) \hat{J}_y | (g) \hat{S}_x |
| (b) \hat{L}_y | (f) \hat{L}^2 | (h) \hat{S}_y |
| (c) \hat{L}_z | (g) \hat{S}^2 | (i) \hat{S}_z |
| (d) \hat{J}_x | (h) \hat{J}^2 | |

7. Which of the following operators commute with \hat{S}_x . (There may be more than one answer.)

- | | | |
|-----------------|-----------------|-----------------|
| (a) \hat{L}_x | (e) \hat{J}_y | (g) \hat{S}_x |
| (b) \hat{L}_y | (f) \hat{L}^2 | (h) \hat{S}_y |
| (c) \hat{L}_z | (g) \hat{S}^2 | (i) \hat{S}_z |
| (d) \hat{J}_x | (h) \hat{J}^2 | |

Below, there is a table with three term symbols.

8. What are the possible values of J ?

9. What are the possible values for M_S ?

10. What are the possible values of M_L ?

	Term Symbol		
Values of	6D	3G	1S
J			
M_S			
M_J			

Quiz 3

Chemistry 3BB3; Winter 2004

1. Write the electronic Schrödinger Equation for the Beryllium ($Z=4$, $N=4$) atom, showing the dependence on \hbar , e , ϵ_0 , and m_e . (Do not use atomic units. You may use the Born-Oppenheimer approximation.)

$$\sum_{i=1}^4 \left(\frac{-\hbar^2}{2m_e} \nabla_i^2 - \frac{Ze^2}{4\pi\epsilon_0 r_i} \right) + \sum_{i=1}^3 \sum_{j=i+1}^4 \frac{e^2}{4\pi\epsilon_0 |\mathbf{r}_i - \mathbf{r}_j|}$$

2. Write a Slater determinant wave function for the $1s^2 4s^1$ excited state of the Lithium atom. Do not use the “shorthand” notation. Denote the spatial part of the $1s$ orbital with $\psi_{1s}(\mathbf{r})$ and the spatial part of the $2s$ orbital with $\psi_{4s}(\mathbf{r})$.

$$\Phi_{1s^2 4s^1}(\mathbf{r}_1, s_1; \mathbf{r}_2, s_2; \mathbf{r}_3, s_3) = \frac{1}{\sqrt{3!}} \begin{vmatrix} \psi_{1s}(\mathbf{r}_1)\alpha(1) & \psi_{1s}(\mathbf{r}_1)\beta(1) & \psi_{4s}(\mathbf{r}_1)\alpha(1) \\ \psi_{1s}(\mathbf{r}_2)\alpha(2) & \psi_{1s}(\mathbf{r}_2)\beta(2) & \psi_{4s}(\mathbf{r}_2)\alpha(2) \\ \psi_{1s}(\mathbf{r}_3)\alpha(3) & \psi_{1s}(\mathbf{r}_3)\beta(3) & \psi_{4s}(\mathbf{r}_3)\alpha(3) \end{vmatrix}$$

Using separation of variables, the wave function for the hydrogen atom can be written as

$$\Psi_{n,l,m}(\mathbf{r}) = R_{n,l}(r) Y_l^m(\theta, \phi).$$

3. Write the Schrödinger equation for radial wave function, $R_{n,l}(r)$, in the hydrogen atom. You may use atomic units.

$$\left(-\frac{1}{2r^2} \frac{d}{dr} r^2 \frac{d}{dr} + \frac{l(l+1)}{2r^2} - \frac{Z}{r} \right) R_{n,l}(r) = E_{n,l} R_{n,l}(r)$$

4. Is $\Psi(\mathbf{r}_1, \sigma(1), \mathbf{r}_2, \sigma(2)) \propto e^{-\zeta_1 r_1} e^{-\zeta_2 r_2} (1 + b|\mathbf{r}_1 - \mathbf{r}_2| + c|\mathbf{r}_1 - \mathbf{r}_2|^2)(\alpha(1)\beta(2) - \alpha(2)\beta(1))$ an acceptable approximate wave function for the Helium atom?

(a) yes

(b) no

5. Which of the following operators commute with the electronic Hamiltonian for an atom. (There may be more than one answer.)

(a) \hat{L}_x

(d) \hat{L}^2

(g) \hat{J}_x

(b) \hat{L}_y

(e) \hat{S}^2

(h) \hat{S}_y

(c) \hat{L}_z

(f) \hat{J}^2

(i) \hat{S}_x

Name:

6. Which of the following operators commute with \hat{L}_x . (There may be more than one answer.)

- | | | |
|-----------------|-----------------|-----------------|
| (a) \hat{L}_x | (e) \hat{J}_y | (g) \hat{S}_x |
| (b) \hat{L}_y | (f) \hat{L}^2 | (h) \hat{S}_y |
| (c) \hat{L}_z | (g) \hat{S}^2 | (i) \hat{S}_z |
| (d) \hat{J}_x | (h) \hat{J}^2 | |

7. Which of the following operators commute with \hat{S}_x . (There may be more than one answer.)

- | | | |
|-----------------|-----------------|-----------------|
| (a) \hat{L}_x | (e) \hat{J}_y | (g) \hat{S}_x |
| (b) \hat{L}_y | (f) \hat{L}^2 | (h) \hat{S}_y |
| (c) \hat{L}_z | (g) \hat{S}^2 | (i) \hat{S}_z |
| (d) \hat{J}_x | (h) \hat{J}^2 | |

Below, there is a table with three term symbols.

8. What are the possible values of J ?

9. What are the possible values for M_S ?

10. What are the possible values of M_L ?

	Term Symbol		
Values	6D	3G	1S
J	$ L - S = \frac{1}{2} \leq J \leq \frac{9}{2} = L + S$ $J = \frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \frac{7}{2}, \frac{9}{2}$	$ L - S = 3 \leq J \leq 5 = L + S$ $J = 3, 4, 5$	$ L - S = 0 \leq J \leq 0 = L + S$ $J = 0$
M_S	2S+1=6, so S=$\frac{5}{2}$. Thus $M_S = -\frac{5}{2}, -\frac{3}{2}, \dots, \frac{5}{2}$	2S+1=3, so S=1. Thus $M_S = -1, 0, 1$	2S+1=0, so S=0. Thus $M_S = 0$
M_J	L=2 so $M_L = -2, -1, 0, 1, 2$	L=4 so $M_L = -4, -3, -2, -1, 0, 1, 2, 3, 4$	L=0 so $M_L = 0$