

Oh

Pauline

A thesis presented for the degree of
Bachelor of Science (Honours) (Physics)

School of Physics
Department of Science
The University of Sydney
Australia
February 10, 2026

Supervisor: Prof. Jane Professor

Co-Supervisor: Dr. Jack Supervisor

Co-Supervisor: Mr. June Supervisor

Tous les fichiers svg :

Dedication

For those who hate looking at a template with 500 lines of code and an extra 300 lines commented out.

Declaration

Acknowledgements

Contents

1 Introduction

| | | |
|---------|--|---|
| 1.1 | Fermi gas preparation (+ Bose gas ?) | 1 |
| 1.1.1 | TC | 1 |
| 1.1.2 | Zeeman cooling | 1 |
| 1.1.3 | Blue MOT | 1 |
| 1.1.3.1 | The physics | 1 |
| 1.1.3.2 | How to optimize the superposition with the repumper | 1 |
| 1.1.3.3 | Comment on the hyperfine states (+boson 88) | 1 |
| 1.1.3.4 | Optical setup (blue + repump) | 1 |
| 1.1.4 | Repumper | 1 |
| 1.1.5 | BB MOT | 1 |
| 1.1.5.1 | First step | 1 |
| 1.1.5.2 | Second step | 1 |
| 1.1.6 | Stir | 1 |
| 1.1.7 | Narrow MOT | 1 |
| 1.1.7.1 | Optimization of the narrow MOT (intensity, frequency, effect on the size of the cloud) | 1 |
| 1.1.7.2 | Optical setup | 1 |
| 1.1.8 | ODT and evaporation | 1 |
| 1.1.8.1 | Charging the crossing | 1 |
| 1.1.8.2 | Optimization of the evaporation ramps : Dimple + reservoir, just reservoir, parameter to optimize (number of atoms, temperature) | 1 |
| 1.1.8.3 | Optical setup | 1 |
| 1.1.9 | Optical pumping | 1 |
| 1.2 | Spin measurement scheme | 1 |

2 Ramsey interferometers on qudit

| | | |
|---------|--|---|
| 2.1 | Preparation of arbitrary dimension Hilbert space | 2 |
| 2.1.1 | Raman process | 2 |
| 2.1.1.1 | $\delta m_F = \pm 1$ | 2 |
| 2.1.1.2 | $\delta m_F = \pm 2$ | 2 |
| 2.1.2 | Moglabs chain without cavity | 2 |
| 2.1.3 | Purification of the laser spectrum with a FP cavity | 2 |
| 2.2 | Interferometric sensing with multiple nuclear spin state | 2 |
| 2.2.1 | Driving long coherence time Rabi oscillations | 2 |
| 2.2.1.1 | Rabi oscillations | 2 |

| | | |
|----------|---|----------|
| 2.2.1.2 | Interferometer of su(2) symmetry | 3 |
| 2.2.1.3 | Discussion on inhomogeneites | 3 |
| 2.2.2 | Measuring two quantites at a time | 3 |
| 2.2.2.1 | Physical principle | 3 |
| 2.2.2.2 | Results | 3 |
| 2.2.3 | Measuring two non commuting observables | 3 |
| 2.2.3.1 | Principle | 3 |
| 2.3 | SU(N) symmetry (ce qu'il faudrait pr la tester e.g densité gaz, alimentation bobines - <i>i</i> comment faire mieux que les chiffres actuels) | 3 |
| 3 | Engineering highly entangled system of photoassociated ^{87}Sr atoms | 4 |
| 3.1 | Introduction on photoassociation | 5 |
| 3.1.1 | What is photoassociation | 5 |
| 3.1.2 | Molecular formalism/vocabulary (condon radius, optical length...) | 6 |
| 3.1.3 | Internal energy states | 6 |
| 3.1.3.1 | WKB approximation | 6 |
| 3.1.3.2 | | 7 |
| 3.1.4 | External energy states | 7 |
| 3.2 | About photoassociation on other species | 7 |
| 3.2.1 | Mass scaling (^{88}Sr) | 7 |
| 3.2.2 | Ytterbium: hfs | 8 |
| 3.3 | Experimental setup | 8 |
| 3.4 | ^{88}Sr Results | 8 |
| 3.4.1 | Technical issues of inhabilitation of photoassociation | 9 |
| 3.4.1.1 | Laser width | 9 |
| 3.5 | ^{87}Sr molecules | 9 |
| 3.5.1 | Physical sources of inhabilitation of photoassociation | 9 |
| 3.5.1.1 | On $F = 9/2$: predissociation | 9 |
| 3.5.1.2 | Coupling to more energetic state from the IR | 9 |
| 3.5.1.3 | Node of wavefunction for some vibrational states | 9 |
| 3.5.2 | Energy landscape of $^{87}\text{Sr}-^{87}\text{Sr}$ molecules | 9 |

List of Figures

List of Tables

Chapter 1

Introduction

1.1 Fermi gas preparation (+ Bose gas ?)

1.1.1 TC

Comment on increasing power in TC (thésard marc chesnais)

1.1.2 Zeeman cooling

1.1.3 Blue MOT

1.1.3.1 The physics

1.1.3.2 How to optimize the superposition with the repumper

1.1.3.3 Comment on the hyperfine states (+boson 88)

1.1.3.4 Optical setup (blue + repump)

1.1.4 Repumper




Figure 1.1: Caption

1.1.5 BB MOT




Figure 1.2: Caption

1.1.5.1 First step

1.1.5.2 Second step

1.1.6 Stir

Need a stir because :




Figure 1.3: Caption




Figure 1.4: Caption

1.1.7 Narrow MOT

cf p.43 S.Stellmer thesis

1.1.7.1 Optimization of the narrow MOT (intensity, frequency, effect on the size of the cloud)

Include images of the cloud for different I and detuning ?

1.1.7.2 Optical setup

1.1.8 ODT and evaporation

1.1.8.1 Charging the crossing




Figure 1.5: Caption

1.1.8.2 Optimization of the evaporation ramps : Dimple + reservoir, just reservoir, parameter to optimize (number of atoms, temperature)

Comment on the LS it does to each state

1.1.8.3 Optical setup

1.1.9 Optical pumping




Figure 1.6: Caption

1.2 Spin measurement scheme




Figure 1.7: Caption

Chapter 2

Ramsey interferometers on qudit

2.1 Preparation of arbitrary dimension Hilbert space

2.1.1 Raman process

2.1.1.1 $\delta m_F = \pm 1$

2.1.1.2 $\delta m_F = \pm 2$

2.1.2 Moglabs chain without cavity

2.1.3 Purification of the laser spectrum with a FP cavity

blablablagtg

2.2 Interferometric sensing with multiple nuclear spin state

2.2.1 Driving long coherence time Rabi oscillations

2.2.1.1 Rabi oscillations

Comment on what the FP could add as a longer coherence time of the qubit

- 2.2.1.2 Interferometer of su(2) symmetry
- 2.2.1.3 Discussion on inhomogeneites
- 2.2.2 Measuring two quantites at a time
 - 2.2.2.1 Physical principle
 - 2.2.2.2 Results
- 2.2.3 Measuring two non commuting observables
 - 2.2.3.1 Principle
- 2.3 SU(N) symmetry (ce qu'il faudrait pr la tester
e.g densité gaz, alimentation bobines -; comment faire mieux que les chiffres actuels)

Chapter 3

Engineering highly entangled system of photoassociated ^{87}Sr atoms

Engineering Dicke states

3.1 Introduction on photoassociation

3.1.1 What is photoassociation




Figure 3.1: Caption

Photoassociating two atoms consists in bounding two colliding atoms with light that occurs mostly with two-body and three body losses. Each species has a specific cross section for which it can collide with another particle, which is defined as

$$\sigma = 4\pi a^2 \quad (3.1)$$

As visible in From a non-bound state of a two-atoms system we couple them at resonance with a molecular vibrationnal state. In the case of spectroscopy we usually start with the least bounded state to have the atomic state reference. From that we sweep the atom frequency to address the molecular states red detuned from the atomic line.

In ultracold gases, only s-wave atoms can collide because they do not have enough kinetic energy to pass the barrier potential of higher angular momentum states. It imposes that for fermions -by parity of the total wavefunction being antisymmetric- only atoms with even angular momentum can collide meaning their spin is in a singlet state.

3.1.2 Molecular formalism/vocabulary (condon radius, optical length...)

3.1.3 Internal energy states

3.1.3.1 WKB approximation




FIGURE 2.1. Schematic diagram for behavior of wave function $u_E(x)$ in potential well $V(x)$ with turning points x_1 and x_2 .

Figure 3.2: Caption

3.1.3.2

3.1.4 External energy states

3.2 About photoassociation on other species

3.2.1 Mass scaling (^{88}Sr)




Figure 3.3: Caption

3.2.2 Ytterbium: hfs




FIG. 2. Adiabatic molecular potentials for a $^{173}\text{Yb}_2$ dimer in the $^1\text{S}_0 + ^3\text{P}_1$ channel as functions of the interatomic separation r . The molecular potentials for 205 different (T, F, R) configurations are displayed, which are accessible via PA from the initial s -wave colliding atoms in the $^1\text{S}_0 + ^1\text{S}_0$ channel. At large r , the potentials converge to three asymptotic branches which correspond to excited atomic states with hyperfine numbers of $f'_1 = 3/2, 5/2$, and $7/2$. Some of the potentials have a local minimum (inset), possibly hosting purely long-range bound states [14]. The energy offset is adjusted to the $f'_1 = 7/2$ asymptote. The shaded region indicates the spectral range of our measurements.

Figure 3.4: Caption

3.3 Experimental setup

3.4 88Sr Results

Lopt, power broadening, thermal broadening...




Figure 3.5: Caption

3.4.1 Technical issues of inhabilitation of photoassociation

3.4.1.1 Laser width

3.5 ^{87}Sr molecules

Lopt questions sur nb quantique / choix de pompage optique

3.5.1 Physical sources of inhabilitation of photoassociation

3.5.1.1 On $F = 9/2$: predissociation

3.5.1.2 Coupling to more energetic state from the IR

3.5.1.3 Node of wavefunction for some vibrational states

3.5.2 Energy landscape of $^{87}\text{Sr}-^{87}\text{Sr}$ molecules

Conclusion

Bibliography

Appendix A

Algorithms