Oh

Pauline

A thesis presented for the degree of
Bachelor of Science (Honours) (Physics)

School of Physics
Department of Science
The University of Sydney
Australia
March 16, 2026

Supervisor: Prof. Jane Professor

Co-Supervisor: Dr. Jack Supervisor

Co-Supervisor: Mr. June Supervisor

Tous les fichiers svg :

Dedication

For those who hate looking at a template with 500 lines of code and an extra 300 lines commented out.

Declaration

Acknowledgements

Contents

1 Introduction

1.1 Fermi gas preparation (+ Bose gas ?) . . . . . .

1.1.1 TC . . . . . .

1.1.2 Zeeman cooling . . . . . .

1.1.3 Blue MOT . . . . . .

1.1.3.1 The physics . . . . . .

1.1.3.2 How to optimize the superposition with the repumper

1.1.3.3 Comment on the hyperfine states (+boson 88) . . . .

1.1.3.4 Optical setup (blue + repump) . . . . . .

1.1.4 Repumper . . . . . .

1.1.5 BB MOT . . . . . .

1.1.5.1 First step . . . . . .

1.1.5.2 Second step . . . . . .

1.1.6 Stir . . . . . .

1.1.7 Narrow MOT . . . . . .

1.1.7.1 Optimization of the narrow MOT (intensity, frequency, effect on the size of the cloud) . . . . . .

1.1.7.2 Optical setup . . . . . .

1.1.8 ODT and evaporation 1

1.1.8.1 Charging the crossing 1

1.1.8.2 Optimization of the evaporation ramps : Dimple + reservoir, just reservoir, parameter to optimize (number of atoms, temperature) 1

1.1.8.3 Optical setup 1

1.1.9 Optical pumping 1

1.2 Spin measurement scheme 2

2 Ramsey interferometers on qudit **3**

2.1 Preparation of arbitrary dimension Hilbert space 3

2.1.1 Raman process 3

2.1.1.1 $\delta m_F = \pm 1$ 3

2.1.1.2 $\delta m_F = \pm 2$ 3

2.1.2 Moglabs chain without cavity 3

2.1.3 Purification of the laser spectrum with a FP cavity 3

2.2 Interferometric sensing with multiple nuclear spin state 3

2.2.1 Driving long coherence time Rabi oscillations 3

2.2.1.1 Rabi oscillations 3

2.2.1.2 Interferometer of su(2) symmetry 4
2.2.1.3 Discussion on inhomogeneites 4
2.2.2 Measuring two quantites at a time 4
2.2.2.1 Physical principle 4
2.2.2.2 Results 4
2.2.3 Measuring two non commuting observables 4
2.2.3.1 Principle 4
2.3 SU(N) symmetry (ce qu'il faudrait pr la tester e.g densité gaz, alimentation bobines -¿ comment faire mieux que les chiffres actuels) . . 4

3 Engineering highly entangled system of photoassociated 87Sr atoms 5
3.1 Introduction on photoassociation 5
3.1.1 Preamble on scattering theory 5
3.1.2 What is photoassociation 8
3.1.3 Parameters for PA study 10
3.1.4 Internal energy states 14
3.1.4.1 WKB approximation 14
3.1.4.2 Leroy-Bernstein theory 15
3.1.5 External energy states 15
3.2 About photoassociation on other species 16
3.2.1 Mass scaling (^{88}Sr) 16
3.2.2 Ytterbium: hfs 17
3.3 88Sr Results 19
3.3.1 Protocol 19
3.3.2 Technical issues of inhabilitation of photoassociation 20
3.3.2.1 Laser width 20
3.4 87Sr molecules 20
3.4.1 Physical sources of inhabilitation of photoassociation 20
3.4.1.1 On F = 9/2 : predissociation 20
3.4.1.2 Coupling to more energetic state from the IR 20
3.4.1.3 Node of wavefunction for some vibrational states . . 20
3.4.2 Energy landscape of 87Sr-87Sr molecules 20

A Algorithms ii

List of Figures

1.1 Caption . . . . . .
1.2 Caption . . . . . .
1.3 Caption . . . . . .
1.4 Caption . . . . . .
1.5 Caption . . . . . .
1.6 Caption . . . . . .
1.7 Caption . . . . . .
1.8 Caption 1
1.9 Caption 1
1.10 Caption 2

3.1 Caption 6
3.2 Caption 9
3.3 u-g symmetry 11
3.4 hund 13
3.5 Caption 14
3.6 Caption 17
3.7 Caption 18
3.8 Caption 19

List of Tables

Chapter 1

Introduction

1.1 Fermi gas preparation (+ Bose gas ?)

1.1.1 TC

Comment on increasing power in TC (thésard marc chesnais)

1.1.2 Zeeman cooling

1.1.3 Blue MOT

1.1.3.1 The physics

1.1.3.2 How to optimize the superposition with the repumper



Figure 1.1: Caption

1.1.3.3 Comment on the hyperfine states (+boson 88)

1.1.3.4 Optical setup (blue + repump)



Figure 1.2: Caption

1.1.4 Repumper



Figure 1.3: Caption

1.1.5 BB MOT



Figure 1.4: Caption

1.1.5.1 First step

1.1.5.2 Second step

1.1.6 Stir

Need a stir because :



Figure 1.5: Caption



Figure 1.6: Caption

1.1.7 Narrow MOT

cf p.43 S.Stellmer thesis

1.1.7.1 Optimization of the narrow MOT (intensity, frequency, effect on the size of the cloud)

Include images of the cloud for different I and detuning ?

1.1.7.2 Optical setup



FIGURE 14 – Préparation des différents lasers rouges

Figure 1.7: Caption

1.1.8 ODT and evaporation

1.1.8.1 Charging the crossing



Figure 1.8: Caption

1.1.8.2 Optimization of the evaporation ramps : Dimple + reservoir, just reservoir, parameter to optimize (number of atoms, temperature)

Comment on the LS it does to each state

1.1.8.3 Optical setup

1.1.9 Optical pumping



Figure 1.9: Caption

1.2 Spin measurement scheme



Figure 1.10: Caption

Chapter 2

Ramsey interferometers on qudit

2.1 Preparation of arbitrary dimension Hilbert space

2.1.1 Raman process

2.1.1.1 $\delta m_F = \pm 1$

2.1.1.2 $\delta m_F = \pm 2$

2.1.2 Moglabs chain without cavity

2.1.3 Purification of the laser spectrum with a FP cavity

blablablagtg

2.2 Interferometric sensing with multiple nuclear spin state

2.2.1 Driving long coherence time Rabi oscillations

2.2.1.1 Rabi oscillations

Comment on what the FP could add as a longer coherence time of the qubit

2.2.1.2 Interferometer of su(2) symmetry

2.2.1.3 Discussion on inhomogeneites

2.2.2 Measuring two quantites at a time

2.2.2.1 Physical principle

2.2.2.2 Results

2.2.3 Measuring two non commuting observables

2.2.3.1 Principle

2.3 SU(N) symmetry (ce qu'il faudrait pr la tester e.g densité gaz, alimentation bobines -¿ comment faire mieux que les chiffres actuels)

Chapter 3

Engineering highly entangled system of photoassociated 87Sr atoms

In metrology the major error source of measuring a parameter ϕ is the quantum shot noise $\delta\phi \simeq \frac{1}{\sqrt{N}}$. In the second chapter of this thesis we presented Ramsey interferometers for which a applied generators for qubit enhancement. The phase ϕ from this pulse was limited by the shot noise.

Engineering Dicke state would make possible to go beyond this limitation. For such system variance express as $\delta\phi \simeq \frac{1}{N}$ which is called the Heisenberg limit being the best accuracy we afford in atomic science. ^{87}Sr large Hilbert space enables to improve even more the accuracy of the measurement because this phase depend directly on the maximum value of J

3.1 Introduction on photoassociation

3.1.1 Preamble on scattering theory

It is convenient in the case of the ^{87}Sr to approximate the electrons following the movement of the nuclei because of the mass ratio. In a Born-Oppenheimer picture we neglect the coupled terms between the kinetic energy of the electrons and the kinetic energy of the nuclei. In this manner we can study separately the radial and the angular part of the electronic movement.
As in the LANDAU-ZENER we will find the solutions of the 1D radial Schrödinger equation of the electrons

$$-\frac{\hbar^2}{2\mu}\left(\frac{1}{r^2}\frac{d^2\Psi_l}{dr^2}r^2 - \frac{l(l+1)}{r^2} + \hat{V}(r)\right)\Psi = E\Psi \tag{3.1}$$

- Ψ the eigenfunction - by approximating the scattering of the atoms as a sum of an incoming plane wave with a spherical scattered one - considered as plane wave at long distance 3.1

$$\Psi = e^{i\mathbf{k}\cdot\mathbf{z}} + f(\theta)\frac{e^{ikr}}{r} \tag{3.2}$$



Figure 3.1: Caption

$f(\theta)$ a function that will tell the absorption of the scattered wave if the collision is elastic or inelastic. θ is the angle between the vector wore by the incoming wave with the vector worn by the scattered wave.

The solution at asymptotes contains the scattering part of the phenomenon studied on this subsection. To solve this hamiltonian we will take the results of [1] that changes the solution basis to find a general solution of the hamiltonian 3.1.

$$\Psi = \frac{a_l}{r} sin(kr - \frac{l\pi}{2} + \delta_l) \tag{3.3}$$

with a_l a normalization factor and δ_l the phase acquired by the atoms at the distance at which they scatter.

Continuity of the wavefunction in every region should be respected. It gives

$$\delta_l \propto k^{2l+1} \tag{3.4}$$

For s-waves ($l = 0$), this express as

$$a = -\lim_{k \to 0} \delta_0 / k \tag{3.5}$$

To understand how the atoms scatter we are going to express the scattered part of 3.2 as $\Psi - e^{ikz}$ in a spherical basis where the solutions are a sum on all the partial waves.

$$\Psi \simeq \frac{i}{2kr} \sum_{l=0}^{\infty} (2l+1) \left((-1)^l e^{-ikr} - S_l e^{ikr} \right) \tag{3.6}$$

with $S_l = e^{2i\delta_l}$ the scattering amplitude.

For a **conservative hamiltonian** $|S_l| = 1$ meaning $\delta_l \in \mathbb{R}$. In this case the wavefunction is well a sum of incoming and outcoming waves of same amplitude.

The elastic cross section representing the surface where the scope of the potential is effective writes as

$$\sigma_{el} = \frac{\pi}{k^2} \sum_{l=0}^{\infty} (2l+1) |1 - S_l|^2 \tag{3.7}$$

But we see in equation 3.6 that $|S_l|$ needs to be < 1 to allow dissipation of the wavefunction. Defining δ_l as a complex number satisfies this condition.

In this case the inelastic collisions cross section reads as

$$\sigma_{inel} = \frac{\pi}{k^2} \sum_{l=0}^{\infty} (2l+1)(1 - |S_l|^2) \tag{3.8}$$

In ultracold gases, only s-wave atoms can collide because they do not have enough kinetic energy to pass the potential barrier of higher angular momentum states so $l = 0$ and the scattering amplitude is

$$S_0 = e^{2i\delta_0} \simeq 1 + 2i\delta_0 \tag{3.9}$$

By using the result of 3.5

$$1 + 2i\delta_0 \simeq 1 - 2ika \tag{3.10}$$

Because δ_0 is complex, a direct result of these equations is that we can write a general scattering length as

$$a = a_{bg} + ia" \tag{3.11}$$

At low temperatures we can re-express equations 3.7 and 3.8 as

$$\sigma_{el} = 4\pi |a|^2 \tag{3.12}$$

$$\sigma_{inel} = 4\pi |a"|^2 \tag{3.13}$$

A usual case of the elastic collisions is atoms in the ground state. For inelasticity photoassociation is a good example of dissipative process where two atoms in the ground state are coupled to a molecular state by a laser.

3.1.2 What is photoassociation

Photoassociating two atoms consists in bounding two colliding atoms with light that occurs mostly with two-body and three body losses : it means adding a dissipative part on the interaction.

In a first step we can express the equation evolution of a pair of atoms without dissipation in the basis $\{|\psi_g\rangle, |\psi_e\rangle\}$ of a non coupled closed channel of free atoms $\psi_g = |g, g\rangle$ and their first molecular state virtual channel $\psi_e = |g, e\rangle$

$$i\hbar \frac{d}{dt} \begin{pmatrix} \psi_g \\ \psi_e \end{pmatrix} = \begin{pmatrix} -\frac{\hbar^2}{2\mu}\nabla + V_g & 0 \\ 0 & -\frac{\hbar^2}{2\mu}\nabla + V_e \end{pmatrix} \begin{pmatrix} \psi_g \\ \psi_e \end{pmatrix}. \tag{3.14}$$

In the virtual channel we see in 3.2 the presence of the internal states of the molecule. It is clear that the atoms get more bounded with the increase of the potential amplitude.



Figure 3.2: Caption

Then at resonance $\Delta = \omega_{at} - \omega_L = 0$ with a bounded molecular state a dissipative term appears on the evolution of the system because of the molecular spontaneous emission linewidth γ_{mol}.
The evolution of the atom number evolves

$$\frac{dN}{dt} = -\beta N n - \Gamma N \tag{3.15}$$

where the 2-body losses express as $\beta \propto \gamma_{mol}$ (we are going to comment more in the next section). In comparison with [2]), $\beta = \frac{K_2 V_e}{2}$, where V_e is the effective volume of the cloud $(2\pi)^{3/2}\sigma_x\sigma_y\sigma_z$ for a density Gaussian distribution and Γ the one-body losses rate. This latter is limited by the vacuum quality and the lifetime of the atoms in the trap.
Adding the inelastic collisions of the PA process gives the following effective evolution equation

$$i\hbar\frac{d}{dt}\begin{pmatrix}\psi_g\\ \psi_e\end{pmatrix} = \begin{pmatrix}-\frac{\hbar^2}{2\mu}\nabla + V_g & \Omega/2\\ \Omega/2 & -\frac{\hbar^2}{2\mu}\nabla + V_e + \hbar\Delta - i\frac{\gamma_{mol}}{2}\end{pmatrix}\begin{pmatrix}\psi_g\\ \psi_e\end{pmatrix}. \tag{3.16}$$

The eigenvector of the ground state $|\psi_g\rangle$ is then modified by this coupling. It also means that the phase δ_l (see 3.3) of the free state wavefunction is modified so the scattering length. The real part of a encodes two-body losses rate whereas its imaginary part changes the collisions properties. In this case we can write an effective scattering length enclosing the background scattering length a_{bg} standing for elastic collisions without any coupling, δa a modification of it because of the coupling and still the inelastic part of it

$$a = a_{bg} + \delta a + ia" \tag{3.17}$$

In the case of spectroscopy we usually start with the least bounded state to have the atomic state reference. From that we sweep the atom frequency to adress the molecular states red detuned from the atomic line. The quantity that will testify there is two-body losses is the atom number remaining in the trap after a photoassociation pulse. The molecule leaves the trap by acquiring kinetic energy from the binding energy.

PA is mostly used in Feschbach resonance field because determining the exact position of the vibrational states enables by changing the magnetic field to tune the scattering length of the atoms and thus the interactions in the system.

3.1.3 Parameters for PA study

In this subsection we are going to admit the expression of the two-body losses rate we introduced above but we will take a look on its dependencies.
This rate writes as

$$K_2 = \frac{1}{k}\frac{\frac{1}{4}\gamma_{stim}\gamma_{mol}}{\Delta^2 + (\frac{1}{2}(\gamma_{stim} + \gamma_{mol})^2)} \tag{3.18}$$

γ_{stim} is linear with the intensity of the PA laser, proportionnal to the 3D density n, and the probability to transit from ψ_g to ψ_e that we call **Franck-Condon factor**

$$F_{FC} = \left|\int_0^\infty \psi_g(r)\psi_e(r)dr\right|^2 \tag{3.19}$$

It depends directly with the overlap of the wavefunctions of the initial and final states. **The Condon radius** is the distance between the atoms for which this factor is maximum which also means that in a classical approach the atoms spend the most time at this position.

In the molecular community it is conventional to characterize the strength of the photoassociation rate with the **optical length**

$$K_2(I) = \frac{4\pi\hbar}{\mu} l_{opt}(I) \tag{3.20}$$

μ the reduced mass of the two atoms. From this formula we can say that the major experimental changes enabling the increase of the PA rate is the density of the cloud and the intensity if the. PA beam as we are going to test in the experimental results.

A first element we need to introduce to describe molecules is the symmetry of the orbital wavefunction. For an even wavefunction the exchange of the two atoms A and B does not change the sign of the wavefunction, as in left figures 3.3. The potential is then qualified **gerade**. The probability of finding electrons in between the two atoms is high compared to the **ungerade** potential where the odd wavefunction cancels the probability of finding electrons in between.



Figure 8.5 : Evolution le long de l'axe internucléaire Oz ($x=0, y=0$) des fonctions d'onde liante $|\psi_-\rangle$ et antiliante $|\psi_+\rangle$ et des densités de probabilité de présence de l'électron associées $|\psi_-|^2$ et $|\psi_+|^2$. La distance internucléaire est fixée à $R = 1{,}25$ Å (minimum de E_-).

Figure 3.3: u-g symmetry

Exciting an atom from a ground state to a single excited state is straightforward and quite easy to do precisely. It is not as easy for molecules because there exist a big number molecular potentials and internal states for each potential. These internal

and external accesible states of the molecule are presented in the following via the introduction of the molecular quantum numbers.

The good quantum numbers
We write the total angular momentum of the molecule $T = R + F = R + I + J$, F being the total spin of the two atoms $f_1 + f_2$, $J = j_1 + j_2$ is the angular momentum of the molecule that describes the spin-orbit coupling, and R the rotational angular momentum of the molecule.

We define the projection of the different angular momentum on the inter-nuclear axis as

$$\Omega = \Lambda + \Sigma \tag{3.21}$$

with Λ the projection of the orbital momentum of the molecule on the internuclear axis and Σ the projection of the momentum spin.

M_T is the projection of the total angular momentum T onto a defined quantization axis.

For the rotational quantum number there is a parity of the total wavefunction that needs to be fulfilled. In addition there are only s-wave collisions in cold gases then the part of the wavefunction should be odd the spin one antisymetric

$$|\Psi\rangle = \left(|\phi_e\rangle_A |\phi_g\rangle_B + (-1)^R |\phi_e\rangle_B |\phi_g\rangle_A\right) \otimes |\chi\rangle \tag{3.22}$$

$|\phi_e\rangle$ and $|\phi_g\rangle$ are the electronic wavefunctions of the two atoms A and B in ground and excited state respectively, $|\chi\rangle$ is the spin wavefunction. It gives accessible values $R = 0, 2$ or 4...

For $^1S_0$ - $^3P_1$ molecular state there is a strong spin-orbit coupling that couples its related momentum to the internuclear axis (3.4), quantified by Ω and described as Hund's case (c)



Fig. 11.4 A vector diagram for Hund's case (c), in which the spin–orbit coupling is very strong.

Figure 3.4: hund

The experiment presented in the following section is photoassociation in the $F = 11/2$ hyperfine state. We will focus in this state to simplify the discussion.
Say we have one atome in the $^1S_0$ state and one in the $^3P_1$ state: we can have a total spin of $J = 0, 1$ or 2. It gives possible values of total atomic angular momentum F:

$$\text{For } J = 0, \quad F = \frac{9}{2} \tag{3.23}$$

$$\text{For } J = 1, \quad F = \frac{7}{2}, \frac{9}{2}, \frac{11}{2} \tag{3.24}$$

$$\text{For } J = 2, \quad F = \frac{5}{2}, \frac{7}{2}, \frac{9}{2}, \frac{11}{2}, \frac{13}{2} \tag{3.25}$$

We have one atom with $f_1 = 9/2$ from the $^1S_0$ state and one in the $^3P_1$ state with $f_2 = 11/2$. The possible values of total angular momentum of the molecule are

$$\text{For } R = 0, \quad T = 1, 2, 3...10 \tag{3.26}$$

$$\text{For } R = 2, \quad T = 1, 2...12 \tag{3.27}$$

$$\text{For } R = 4, \quad T = 1, 2...14 \tag{3.28}$$

The initial $T^i = i_1 + i_2$ with $R = 0$ because we are in s-wave collisions. In respect with the parity $p = +1$ the possible values of T^i are $T^i = 0, 2, 4....$
As for atoms there are selection rules for transitions from a free state to a molecular state fixed by the angular momentum of the electrons. $\Delta J = 0, 1$ gives $\Delta T = 0, 1$ with $T = 0 \rightarrow T = 0$ forbidden.
In the end the accessible transitions in the first molecular excited state of the $f_1 =$

11/2 are

$$\text{For } R = 0, \quad T^f = 1, 2, 3...11 \tag{3.29}$$
$$\text{For } R = 2, \quad T^f = 1, 2...13 \tag{3.30}$$
$$\text{For } R = 4, \quad T^f = 1, 2...15 \tag{3.31}$$
$$... \tag{3.32}$$

There are 981 combinations of accessible (R, T^f) molecular external states and for each potential a numerous internal state we are going to demonstrate briefly in the following.

3.1.4 Internal energy states

3.1.4.1 WKB approximation

WKB approximation is a semi-classical approach where plane waves $\psi(x) = \frac{C}{\sqrt{(k(x))}} e^{i \int_0^\infty k(x)dx}$ are solutions of the 1D Schrödinger equation for two particles of eigenenergie E in a potential $V(x)$

$$\frac{-\hbar^2}{2\mu} \frac{d^2\psi}{dx^2} + V(x)\psi = E\psi \tag{3.33}$$

giving the expression of the wavevector k

$$k = \frac{p}{\hbar} = \sqrt{2\mu(E - V(x))} \tag{3.34}$$

The condition of validity $\frac{d\lambda}{dx} \ll 1$ tells that the atoms must follow adiabatically the potential to adapt their movement to its variation. It means at really small x of region I of figure 3.5 this approximation is not valid.



IGURE 2.1. Schematic diagram for behavior of wave function $u_E(x)$ in potential well $V(x)$ ith turning points x_1 and x_2.

Figure 3.5: Caption

At position $x = x_1$ and x_2 the momentum diverges which means that the plane wave approximation is not valid anymore. In the region I and III the wave is evanescent with a phase $k = i\sqrt{2\mu(|E - V(x)|)}$. It is then usual the average the potential as linear with x around these turning points.

By reflection with the left barrier of the potential at $x = x_1$ the stationnary wave must come back with a 2π at x_2. By continuity of the solutions in $[0, +\infty[$ the phase tends to

$$\int_{x_1}^{x_2} k(x)dx = (n + \frac{1}{2})\pi \tag{3.35}$$

Then this formula enables to identify the solutions E that satisfies the condition $(n + \frac{1}{2})\pi$. These values are what we call the vibrationnal states of a molecular potential.

3.1.4.2 Leroy-Bernstein theory

Leroy-Bernstein theory enables to identify the position of the less bounded states without knowing the behaviour of the potential that could be complicated to estimate and solve numerically. This model is valid only close to the dissociation limit occuring at long distance. Then we can approximate the potential as

$$V(r) \xrightarrow[r\to\infty]{} D_e - \frac{C_n}{r^n} \tag{3.36}$$

with D_e the dissociation energy. For a vibrational energy state ν of energy E_ν we express the energy difference $\epsilon = D_e - E_\nu$ to inject it in 3.35 and find solutions of

$$\int_{x_1}^{x_2} \sqrt{2\mu(\frac{C_n}{r^n} - \epsilon)}dr = (\nu_D - \nu + \frac{1}{2})\hbar\pi \tag{3.37}$$

with ν_D the dissociation number. After few calculations we finally obtain

$$\epsilon = \left((\nu_D - \nu)\frac{\sqrt{\pi}\hbar\nu}{\sqrt{2\mu C_n^{2/n}}}\frac{\Gamma(\frac{3}{2} - \frac{1}{\nu})}{\Gamma(1 - \frac{1}{\nu})} \right)^{\frac{2\nu}{\nu-2}} \tag{3.38}$$

Γ being the Beta Euler functions. This energy also writes as

$$\frac{E_\nu - D_e}{\nu_D - \nu} = \left(\frac{\sqrt{\pi}\hbar\nu}{\sqrt{2\mu C_n^{2/n}}}\frac{\Gamma(\frac{3}{2} - \frac{1}{\nu})}{\Gamma(1 - \frac{1}{\nu})} \right)^{\frac{2\nu}{\nu-2}} \tag{3.39}$$

from which we determine the energy difference between the dissociation limit and one of the least bounded states ν of the molecule.

3.1.5 External energy states

As seen in the subsection 3.1.1 in a molecular picture the hamiltonian express as

$$\hat{H} = \frac{\hat{p_r}^2}{2\mu} + \frac{\hbar^2}{2\mu r^2} R\,(R+1) + \hat{V}_{BO} + \hat{H}_{HF} \tag{3.40}$$

It is possible to neglect some orders of electronic interactions depending on the Hund's case and the distance region between the atoms. In our case we approximate the potential to be as for ^{173}Yb in [2]

$$\hat{V}_{BO} = \frac{-C_6}{r^6}\left(1 - \frac{\sigma^6}{r^6}\right) - s\frac{C_3^{\Omega}}{r^3} \tag{3.41}$$

$$\hat{H}_{\text{hf}} = A\,(\mathbf{i}_1 \cdot \mathbf{j}_1) + B\,\frac{3(\mathbf{i}_1 \cdot \mathbf{j}_1)^2 + \frac{3}{2}(\mathbf{i}_1 \cdot \mathbf{j}_1) - i_1(i_1+1)\,j_1(j_1+1)}{2i_1(2i_1-1)\,2j_1(2j_1-1)} \tag{3.42}$$

$s = \pm 1$ for gerade and ungerade potentials respectively. The hyperfine term of the atom in $^1S_0$ is zero because its electronic angular momentum is zero. Only the hyperfine interaction of the atom with i_1, j_1 in $^3P_1$ counts.

3.2 About photoassociation on other species

3.2.1 Mass scaling (^{88}Sr)

^{88}Sr is a bosonic isotope of strontium with no hyperfine structure. The photoassociation spectrum is then simpler than the fermionic ones : only two potentials in the first excited molecular state 0_u and 1_u. Results have been presented in [3] where their fitting model gives the following potential expression

$$\hat{V}_{BO} \sim \frac{-C_{6,\Lambda}}{r^6} - \frac{C_3}{r^3} + \frac{C_J}{r^2} \tag{3.43}$$

for the two accessible potentials $\Lambda = 0, 1$.

Mass scaling method presented in [4] enables to predict the position of the vibrational states of the different isotopes of strontium. The authors consider the potential is the same for all the isotopes because we can validate the Born-Oppenheimer estimation for all isotopes. It is actually wrong to make this approximation in the case of 87^Sr because the hyperfine structure modifies the expression of the potential. It is still giving an average position of the vibrational states. Then for this method only the reduced mass in the integral

$$\int_{x_1}^{x_2} \sqrt{2\mu(E - V(x))}dx = (n + \frac{1}{2})\pi \tag{3.44}$$

will change upon the isotope. Then with respect with the condition on the phase the energies E will be modified. The experimental positions of the first vibrationnal states of the O_u^+ potential for ^{84}Sr, ^{86}Sr and ^{88}Sr are represented by the red crosses in figure 3.6 and enables to adjust $V(r)$ shape. In blue lines the fitted functions are plotted from this model : it is then possible to deduct the position of the ^{87}Sr vibrationnal states thanks to the other isotopes lines.
We see discontinuites at the 84 isotope: these are avoided crossing coming from a

coupling with the ground state potential.



Figure 3.6: Caption

3.2.2 Ytterbium: hfs

Whereas the Strontium bosonic isotope ^{88}Sr has only two molecular channel 0_u and 1_u on the intercombination line it is more convenient to refer to species with a hyperfine structure such as ^{173}Yb. Its molecular potential landscape is more complex (see figure 3.7) than many species that have been studied and the density of each momentum projection $0_u, 1_g, 1_u, 0_g$ comes from the hyperfine structure. It is indeed expected to count 205 different configurations of (T, F, R) for the excited molecular states on intercombination line with an atomic potential written as 3.41.

Around 80 lines were measured. Atom loss spectra as a function of PA laser detuning reveal the position of the bound states relative to the atomic resonance $|^1S_0, F = 5/2\rangle \rightarrow |^3P_1, F = 7/2\rangle$. Near this resonance atoms are illuminated for 30ms at low intensity $I = 37\text{mW/cm}^2$ to avoid power broadening, allowing the nearest bound state to the dissociation to be resolved. At larger red detunings the intensity is increased and the illumination time optimized to maximize contrast.



FIG. 2. Adiabatic molecular potentials for a $^{173}Yb_2$ dimer in the $^1S_0 + {}^3P_1$ channel as functions of the interatomic separation r. The molecular potentials for 205 different (T, F, R) configurations are displayed, which are accessible via PA from the initial s-wave colliding atoms in the $^1S_0 + {}^1S_0$ channel. At large r, the potentials converge to three asymptotic branches which correspond to excited atomic states with hyperfine numbers of $f_1' = 3/2,\ 5/2$, and $7/2$. Some of the potentials have a local minimum (inset), possibly hosting purely long-range bound states [14]. The energy offset is adjusted to the $f_1' = 7/2$ asymptote. The shaded region indicates the spectral range of our measurements.

Figure 3.7: Caption

3.3 88Sr Results

Dicke states are particularly interesting to engineer for metrology because it is possible to go beyond quantum shot-noise limit on a specific parameter ϕ. The strategy we performed to attempt this high precision is to photoassociate atoms to induce two-body losses resulting in single atoms entangled to the $N-1$ other. The aim is to induce Dicke state on a gas of fermions with a precision that goes beyond quantum shot-noise limit.
As introduced in the preamble on scattering theory only s-wave collisions occur in ultracold gases. It enforce that for fermions -by parity of the total wavefunction being antisymetric- only atoms with even angular momentum can collide meaning for PA process their spin state must be antisymetric so is the case of ^{87}Sr. For bosons it is the contrary: only atoms in triplet states will be photoassociated by symetry of the total wavefunction.

3.3.1 Protocol

Prior to the main experiment on ^{87}Sr we did calibrated our protocol on isotope ^{88}Sr well refered in ref [5], [4], [3].



Figure 3.8: Caption

For PA process on strontium 88 the beam line currently used for strontium 87 red MOT and 11/2 hyperfine state was modified. Figure 3.8 shows in red the frequency detunings for adressing the hyperfine state $^3P_1, F = 11/2$ used on the red MOT

stages and the optical pumping. The beam for these two is tuned via a first double pass AOM called "Inj slave 1" at center frequency $f_{InjSlave1} = \pm 300$MHz and a second one at $f_{11/2} = \pm 200$: "AOM PompOpt" in a case, "AOM MOT 87" in the other case both in double pass configuration.

3.3.2 Technical issues of inhabilitation of photoassociation

3.3.2.1 Laser width

3.4 87Sr molecules

Lopt questions sur nb quantique / choix de pompage optique

3.4.1 Physical sources of inhabilitation of photoassociation

3.4.1.1 On F = 9/2 : predissociation

3.4.1.2 Coupling to more energetic state from the IR

3.4.1.3 Node of wavefunction for some vibrational states

3.4.2 Energy landscape of 87Sr-87Sr molecules

Conclusion

Bibliography

[1] J. Dalibard, "Les interactions entre atomes dans les gaz quantiques,"

[2] J. H. Han, J. H. Kang, M. Lee, and Y.-i. Shin, "Photoassociation spectroscopy of ultracold ^{173}Yb atoms near the intercombination line," *Physical Review A*, vol. 97, p. 013401, Jan. 2018. arXiv:1710.03072 [physics].

[3] T. Zelevinsky, M. M. Boyd, A. D. Ludlow, T. Ido, J. Ye, R. Ciurylo, P. Naidon, and P. S. Julienne, "Narrow Line Photoassociation in an Optical Lattice," *Physical Review Letters*, vol. 96, p. 203201, May 2006. Number: 20 arXiv:physics/0602135.

[4] B. J. Reschovsky, B. P. Ruzic, H. Miyake, N. C. Pisenti, P. S. Julienne, and G. K. Campbell, "Narrow-line photoassociation spectroscopy and mass-scaling of bosonic strontium," Oct. 2018. arXiv:1808.06507 [physics].

[5] M. Borkowski, P. Morzyński, R. Ciuryło, P. S. Julienne, M. Yan, B. J. DeSalvo, and T. C. Killian, "Mass scaling and nonadiabatic effects in photoassociation spectroscopy of ultracold strontium atoms," *Physical Review A*, vol. 90, p. 032713, Sept. 2014.

Appendix A

Algorithms