Practice quiz on Exponents and Logarithms

TOTAL POINTS 12

1. Re write the number $784 = 2 \times 2 \times 2 \times 2 \times 7 \times 7$ using exponents.

1 / 1 point

- \bigcirc $(16^4)(49^2)$
- \bigcirc $(2 \times 7)^6$
- \bigcirc $(2^6)(7^6)$

✓ Correct

For this type of problem, count the number of times each relevant factor appears in the product. That number is the exponent for that factor.

2. What is $(x^2-5)^0$?

1/1 point

- 1
- \bigcirc -4
- $\bigcirc (x^2)$
- $\bigcirc (x^2) 5$

/ Correc

Any real number (except zero) raised to the "zeroith" power =1.

3. Simplify $((x-5)^2)^{-3}$

1 / 1 point

- $(x-5)^{-1}$
- $(x-5)^{-6}$
- $\bigcirc (x-5)$
- $(x-5)^{-5}$

✓ Correc

By Rule 2, "Power to a Power," multiply the exponents and get:

$$(x-5)^{(2\times-3)} = (x-5)^{-6}$$

By the definition of negative exponents, this is equal to $\frac{1}{(x-5)^6}$

Simplify $(\frac{8^2}{8^7})^2$

1 / 1 point

- $O 8^{-5}$
- \circ 8⁻¹

We can first simplify what is inside the parenthesis to 8^{-5} using the Division and Negative Powers Rule

Then apply division and negative powers-- the result is the same. $\frac{8^4}{8^{14}}=8^{-10}$

5. $\log 35 = \log 7 + \log x$

1/1 point

Solve for x

- 5
- O 28
- O 4
- O 7

✓ Correct

$$\log(x) = \log 35 - \log 7$$

$$\log(x) = \log \left(\frac{35}{7}\right)$$

By the Quotient Rule $\log x = \log 5$

6. $\log_2(x^2 + 5x + 7) = 0$

1 / 1 point

1 / 1 point

Solve for \boldsymbol{x}

- $\bigcirc x = 3$
- $\bigcirc x = 2$
- $\bigcirc \ \ x=2 \ {\rm or} \ \ x=3$

✓ Correct

We use the property that $b^{\log_b a} = a$

Use both sides as exponent for 2.

$$2^{\log_2 x^2 + 5x + 7} = 2^0$$

$$x^2 + 5x + 7 = 1$$

$$x^2 + 5x + 6 = 0$$

$$(x+3)(x+2) = 0$$

$$x = -3$$

$$x = -2$$

7. Simplify $\log_2 72 - \log_2 9$

3	
$\bigcirc \log_2 4$	Ł
$\bigcirc \log_2 6$	33
O 4	
✓ Co	rrect
Ву	the quotient rule, this is $\log_2 \frac{72}{9} = \log_2 2^3 = 3$

8. Simplify $\log_3 9 - \log_3 3 + \log_3 5$

- 0 8
- O 15
- \bigcirc $\log_3 15$
- $\bigcirc \log_3 8$

✓ Correct By the Quotient and Product Rules, this is $\log_3 \frac{9 \times 5}{3} = \log_3 15$

9. Simplify $\log_2(3^8 imes 5^7)$

- $\bigcirc \ 15 \times \log_2 56$
- $\bigcirc 56 \times \log_2 15$
- $\bigcirc \ (5 \times \log_2 3) + (8 \times \log_2 5)$

✓ Correct

We first apply the Product Rule to convert to the sum: $\log_2(3^8) + \log_2(5^7)$. Then apply the power and root rule.

10. If $\log_{10} y = 100$, what is $\log_2 y = ?$

- O 500
- 301.03
- 332.19
- O 20

Use the change of base formula, $\log_a b = \frac{\log_{\scriptscriptstyle X} b}{\log_{\scriptscriptstyle X} a}$

Where the "old" base is x and the "new" base is a.

So
$$\frac{100}{\log_{10}(2)} = \frac{100}{0.30103} = 332.19$$

11. A tree is growing taller at a continuous rate. In the past 12 years it has grown from 3 meters to 15 meters. What is its rate of growth per year?

- O 12.41%
- 0 10.41%
- O 11.41%

$$\sqrt{\frac{\ln \frac{15}{3}}{12}} = 0.1341$$

12. Bacteria can reproduce exponentially if not constrained. Assume a colony grows at a continually compounded rate of 400% per day. How many days before a colony with initial mass of 6.25×10^{-10} grams weights 1000 Kilograms?

1 / 1 point

- $\bigcirc 0.875 \, \mathrm{days}$
- O 87.5 days
- 8.75 days
- O 875 days

$$\checkmark$$
 correct
$$6.25 \times 10^{-10} \times e^{4t} = 10^{6}$$

$$4t = \ln \left(\frac{10^{6}}{(6.25 \times 10^{-10})} \right) = 35.00878$$

$$t = \ln \frac{10^6}{6.25 \times 10^{-10}} = 8.752195$$