



CENTER FOR
MACHINE PERCEPTION



CZECH TECHNICAL
UNIVERSITY IN PRAGUE

MASTER'S THESIS

Semidefinite Programming for Geometric Problems in Computer Vision

Pavel Trutman

pavel.trutman@fel.cvut.cz

April 11, 2017

Available at
<http://cmp.felk.cvut.cz/~trutmpav/master-thesis/thesis/thesis.pdf>

Thesis Advisor: Ing. Tomáš Pajdla, PhD.

Center for Machine Perception, Department of Cybernetics
Faculty of Electrical Engineering, Czech Technical University
Technická 2, 166 27 Prague 6, Czech Republic
fax +420 2 2435 7385, phone +420 2 2435 7637, www: <http://cmp.felk.cvut.cz>

Acknowledgements

Author's declaration

I declare that I have work out the presented thesis independently and that I have listed all information sources used in accordance with the Methodical Guidelines about Maintaining Ethical Principles for Writing Academic Theses.

Prohlášení autora práce

Prohlašuji, že jsem předloženou práci vypracoval samostatně a že jsem uvedl veškeré použité informační zdroje v souladu s Metodickým pokynem o dodržování etických principů při přípravě vysokoškolských závěrečných prací.

V Praze dne

.....

Podpis autora práce

Abstract

Keywords:

Abstrakt

Keywords:

Contents

1. Introduction	4
2. Semidefinite programming	5
2.1. Preliminaries on semidefinite programs	5
2.1.1. Symmetric matrices	5
2.1.2. Semidefinite programs	5
2.2. State of the art review	6
2.3. Nesterov's approach	6
2.3.1. Self-concordant functions	6
2.3.2. Self-concordant barriers	8
2.3.3. Barrier function for semidefinite programming	12
2.4. Implementation details	14
2.5. Comparison with the state of the art methods	14
3. Conclusion	16
A. Contents of the enclosed CD	17
Bibliography	18

List of Algorithms

1.	Newton method for minimization of self-concordant functions	8
2.	Damped Newton method for analytic centers	10
3.	Path following algorithm	12

List of Symbols and Abbreviations

$\text{cl}(S)$	Closure of the set S .
$\text{dom } f$	Domain of the function f .
$\text{Dom } f$	$\text{cl}(\text{dom } f)$.
$\text{int } S$	Interior of the set S .
\mathcal{P}^n	Cone of positive semidefinite $n \times n$ matrices.
\mathcal{S}^n	Space of $n \times n$ real symmetric matrices.

1. Introduction

[1]

2. Semidefinite programming

2.1. Preliminaries on semidefinite programs

We introduce here some notation and preliminaries about symmetric matrices and semidefinite programs. We will introduce further notation and preliminaries later on in the text when needed.

At the beginning, let us denote the inner product for two vectors $x, y \in \mathbb{R}^n$

$$\langle x, y \rangle = \sum_{i=1}^n x_i y_i \quad (2.1)$$

and the Frobenius inner product for two matrices $X, Y \in \mathbb{R}^{n \times m}$.

$$\langle X, Y \rangle = \sum_{i=1}^n \sum_{j=1}^m X_{ij} Y_{ij} \quad (2.2)$$

2.1.1. Symmetric matrices

Let \mathcal{S}^n denotes the space of $n \times n$ real symmetric matrices.

For matrix $M \in \mathcal{S}^n$, the notation $M \succeq 0$ means that M is positive semidefinite. $M \succeq 0$ if and only if any of the following equivalent properties holds.

1. $x^\top M x \geq 0$ for all $x \in \mathbb{R}^n$.
2. All eigenvalues of M are nonnegative.

For matrix $M \in \mathcal{S}^n$, the notation $M \succ 0$ means that M is positive definite. $M \succ 0$ if and only if any of the following equivalent properties holds.

1. $M \succeq 0$ and $\text{rank } M = n$.
2. $x^\top M x > 0$ for all $x \in \mathbb{R}^n$.
3. All eigenvalues of M are positive.

2.1.2. Semidefinite programs

The standard (primal) form of a semidefinite program in variable $X \in \mathcal{S}^n$ is defined as follows:

$$\begin{aligned} p^* &= \sup_{X \in \mathcal{S}^n} \langle C, X \rangle \\ \text{s.t.} \quad &\langle A_i, X \rangle = b_i \quad (i = 1, \dots, m) \\ &X \succeq 0 \end{aligned} \quad (2.3)$$

where $C, A_1, \dots, A_m \in \mathcal{S}^n$ and $b \in \mathbb{R}^m$ are given.

The dual form of the primal form is the following program in variable $y \in \mathbb{R}^m$.

$$\begin{aligned} d^* &= \inf_{y \in \mathbb{R}^m} b^\top y \\ \text{s.t.} \quad &\sum_{i=1}^m A_i y_i - C \succeq 0 \end{aligned} \quad (2.4)$$

2.2. State of the art review

2.3. Nesterov's approach

In this section, we will follow Chapter 4 of [2] by Y. Nesterov, which is devoted to convex minimization problems. We will extract from it only the minimum, just to be able to introduce an algorithm for SDP programs solving. We will present some basic definitions and theorems, but we will not prove them. For the proofs look into [2].

2.3.1. Self-concordant functions

Definition 2.1 (Self-concordant function in \mathbb{R}). A closed convex function $f : \mathbb{R} \mapsto \mathbb{R}$ is self-concordant if there exist a constant $M_f \geq 0$ such that the inequality

$$|f'''(x)| \leq M_f f''(x)^{3/2} \quad (2.5)$$

holds for all $x \in \text{dom } f$.

For better understanding of self-concordant functions, we provide several examples.

Example 2.1.

1. Linear and convex quadratic functions.

$$f'''(x) = 0 \text{ for all } x \quad (2.6)$$

Linear and convex quadratic functions are self-concordant with constant $M_f = 0$.

2. Negative logarithms.

$$f(x) = -\ln(x) \text{ for } x > 0 \quad (2.7)$$

$$f'(x) = -\frac{1}{x} \quad (2.8)$$

$$f''(x) = \frac{1}{x^2} \quad (2.9)$$

$$f'''(x) = -\frac{2}{x^3} \quad (2.10)$$

$$\frac{|f'''(x)|}{f''(x)^{3/2}} = 2 \quad (2.11)$$

Negative logarithms are self-concordant functions with constant $M_f = 2$.

3. Exponential functions.

$$f(x) = e^x \quad (2.12)$$

$$f''(x) = f'''(x) = e^x \quad (2.13)$$

$$\frac{|f'''(x)|}{f''(x)^{3/2}} = e^{-x/2} \rightarrow \infty \text{ as } x \rightarrow -\infty \quad (2.14)$$

Exponential functions are not self-concordant functions.

Definition 2.2 (Self-concordant function in \mathbb{R}^n). A closed convex function $f : \mathbb{R}^n \mapsto \mathbb{R}$ is self-concordant if function

$$g(t) = f(x + tv) \quad (2.15)$$

is self-concordant for all $x \in \text{dom } f$ and all $v \in \mathbb{R}^n$.

Now, let us focus on the main properties of self-concordant functions.

Theorem 2.1. Let functions f_i be self-concordant with constants M_i and let $\alpha_i > 0$, $i = 1, 2$. Then the function

$$f(x) = \alpha_1 f_1(x) + \alpha_2 f_2(x) \quad (2.16)$$

is self-concordant with constant

$$M_f = \max \left\{ \frac{1}{\sqrt{\alpha_1}} M_1, \frac{1}{\sqrt{\alpha_2}} M_2 \right\} \quad (2.17)$$

and

$$\text{dom } f = \text{dom } f_1 \cap \text{dom } f_2. \quad (2.18)$$

Corollary 2.1. Let function f be self-concordant with some constant M_f and $\alpha > 0$. Then the function $\phi(x) = \alpha f(x)$ is also self-concordant with the constant $M_\phi = \frac{1}{\sqrt{\alpha}} M_f$.

We call function $f(x)$ as the standard self-concordant function if $f(x)$ is some self-concordant function with the constant $M_f = 2$. Using Corollary 2.1, we can see that any self-concordant function can be transformed into the standard self-concordant function by scaling.

Theorem 2.2. Let function f be self-concordant. If $\text{dom } f$ contains no straight line, then the Hessian $f''(x)$ is nondegenerate at any x from $\text{dom } f$.

For some self-concordant function $f(x)$, for which we assume, that $\text{dom } f$ contains no straight line (which implies that all $f''(x)$ are nondegenerate, see Theorem 2.2), we denote two local norms as

$$\|u\|_x = \sqrt{u^\top f''(x) u} \quad (2.19)$$

$$\|u\|_x^* = \sqrt{u^\top f''(x)^{-1} u}. \quad (2.20)$$

Consider following minimization problem

$$x^* = \arg \min_{x \in \text{dom } f} f(x) \quad (2.21)$$

as the minimization of the self-concordant function $f(x)$. Algorithm 1 is describing the iterative process of solving optimization problem (2.21). The algorithm is divided into two stages by the value of $\|f'(x_k)\|_{x_k}^*$. The splitting parameter β guarantees quadratic convergence rate for the second part of the algorithm. The parameter β is chosen from interval $(0, \bar{\lambda})$, where

$$\bar{\lambda} = \frac{3 - \sqrt{5}}{2}, \quad (2.22)$$

which is a root of the equation

$$\frac{\lambda}{(1 - \lambda)^2} = 1. \quad (2.23)$$

The first while loop (lines 2 – 5) represents damped Newton method, where at each iteration we have

$$f(x_k) - f(x_{k+1}) \geq \beta - \ln(1 + \beta) \text{ for } k \geq 0, \quad (2.24)$$

2. Semidefinite programming

Algorithm 1 Newton method for minimization of self-concordant functions

Input:

f a self-concordant function to minimize
 $x_0 \in \text{dom } f$ a starting point
 $\beta \in (0, \bar{\lambda})$ a parameter of size of the region of quadratic convergence
 ε a precision

Output:

x^* an optimal solution to the minimization problem (2.21)

```

1:  $k \leftarrow 0$ 
2: while  $\|f'(x_k)\|_{x_k}^* \geq \beta$  do
3:    $x_{k+1} \leftarrow x_k - \frac{1}{1 + \|f'(x_k)\|_{x_k}^*} f''(x_k)^{-1} f'(x_k)$ 
4:    $k \leftarrow k + 1$ 
5: end while
6: while  $\|f'(x_k)\|_{x_k}^* > \varepsilon$  do
7:    $x_{k+1} \leftarrow x_k - f''(x_k)^{-1} f'(x_k)$ 
8:    $k \leftarrow k + 1$ 
9: end while
10: return  $x^* \leftarrow x_k$ 

```

where

$$\beta - \ln(1 + \beta) > 0 \text{ for } \beta > 0, \quad (2.25)$$

and therefore the global convergence of the algorithm is ensured. It can be shown that the local convergence rate of the damped Newton method is also quadratic, but the presented switching strategy is preferred as it gives better complexity bounds.

The second while loop of the algorithm (lines 6 – 9) is the standard Newton method with quadratic convergence rate.

The algorithm terminates when the required precision ε is reached.

2.3.2. Self-concordant barriers

To be able to introduce self-concordant barriers, let us denote $\text{Dom } f = \text{cl}(\text{dom } f)$.

Definition 2.3 (Self-concordant barrier). Let $F(x)$ be a standard self-concordant function. We call it a ν -self-concordant barrier for set $\text{Dom } F$, if

$$\sup_{u \in \mathbb{R}^n} (2u^\top F'(x) - u^\top F''(x)u) \leq \nu \quad (2.26)$$

for all $x \in \text{dom } F$. The value ν is called the parameter of the barrier.

The inequality (2.26) can be rewritten into the following equivalent matrix notation:

$$F''(x) \succeq \frac{1}{\nu} F'(x) F'(x)^\top. \quad (2.27)$$

In Definition 2.3, the hessian $F''(x)$ is not required to be nondegenerate. However, in case that $F''(x)$ is nondegenerate, the inequality (2.26) is equivalent to

$$F'^\top(x) F''(x)^{-1} F'(x) \leq \nu. \quad (2.28)$$

Let us explore, which basic functions are self-concordant barriers.

Example 2.2.

1. Linear functions.

$$F(x) = \alpha + a^\top x, \text{ dom } F = \mathbb{R}^n \quad (2.29)$$

$$F''(x) = 0 \quad (2.30)$$

From (2.27) and for $a \neq 0$ follows, that linear functions are not self-concordant barriers.

2. Convex quadratic functions.

For $A = A^\top \succ 0$:

$$F(x) = \alpha + a^\top x + \frac{1}{2}x^\top Ax, \text{ dom } F = \mathbb{R}^n \quad (2.31)$$

$$F'(x) = a + Ax \quad (2.32)$$

$$F''(x) = A \quad (2.33)$$

After substitution into (2.28) we obtain

$$(a + Ax)^\top A^{-1}(a + Ax) = a^\top A^{-1}a + 2a^\top x + x^\top Ax, \quad (2.34)$$

which is unbounded from above on \mathbb{R}^n . Therefore, quadratic functions are not self-concordant barriers.

3. Logarithmic barrier for a ray.

$$F(x) = -\ln x, \text{ dom } F = \{x \in \mathbb{R} \mid x > 0\} \quad (2.35)$$

$$F'(x) = -\frac{1}{x} \quad (2.36)$$

$$F''(x) = \frac{1}{x^2} \quad (2.37)$$

From (2.28), when $F'(x)$ and $F''(x)$ are both scalars, we get

$$\frac{F'(x)^2}{F''(x)} = \frac{x^2}{x^2} = 1. \quad (2.38)$$

Therefore, the logarithmic barrier for a ray is a self-concordant barrier with parameter $\nu = 1$ on domain $\{x \in \mathbb{R} \mid x > 0\}$.

Now, let us focus on the main properties of self-concordant barriers.

Theorem 2.3. Let $F(x)$ be a self-concordant barrier. Then the function $c^\top x + F(x)$ is a self-concordant function on $\text{dom } F$.

Theorem 2.4. Let F_i be ν_i -self-concordant barriers, $i = 1, 2$. Then the function

$$F(x) = F_1(x) + F_2(x) \quad (2.39)$$

is a self-concordant barrier for convex set

$$\text{Dom } F = \text{Dom } F_1 \cap \text{Dom } F_2 \quad (2.40)$$

with the parameter

$$\nu = \nu_1 + \nu_2. \quad (2.41)$$

2. Semidefinite programming

Theorem 2.5. Let $F(x)$ be a ν -self-concordant barrier. Then for any $x \in \text{Dom } F$ and $y \in \text{Dom } F$ such that

$$(y - x)^\top F'(x) \geq 0, \quad (2.42)$$

we have

$$\|y - x\|_x \leq \nu + 2\sqrt{\nu}. \quad (2.43)$$

There is one special point of convex set, which is important for solving convex minimization problem. It is called analytic center of convex set and we will focus on its properties.

Definition 2.4. Let $F(x)$ be a ν -self-concordant barrier for the set $\text{Dom } F$. The point

$$x_F^* = \arg \min_{x \in \text{Dom } F} F(x) \quad (2.44)$$

is called the analytic center of convex set $\text{Dom } F$, generated by the barrier $F(x)$.

Theorem 2.6. Assume that the analytic center of a ν -self-concordant barrier $F(x)$ exists. Then for any $x \in \text{Dom } F$ we have

$$\|x - x_F^*\|_{x_F^*} \leq \nu + 2\sqrt{\nu}. \quad (2.45)$$

This property clearly follows from Theorem 2.5 and the fact, that $F'(x_F^*) = 0$.

Thus, if $\text{Dom } F$ contains no straight line, then the existence of x_F^* (which leads to nondegenerate $F''(x_F^*)$) implies, that the set $\text{Dom } F$ is bounded.

Now, we describe the algorithm and its properties for obtaining an approximation to the analytic center. To find the analytic center, we need to solve the minimization problem (2.44). For that, we will use the standard implementation of the damped Newton method with termination condition

$$\|F'(x_k)\|_{x_k}^* \leq \beta, \text{ for } \beta \in (0, 1). \quad (2.46)$$

The pseudocode of the whole minimization process is shown in Algorithm 2.

Algorithm 2 Damped Newton method for analytic centers

Input:

- F a ν -self-concordant barrier
- $x_0 \in \text{Dom } F$ a starting point
- $\beta \in (0, 1)$ a centering parameter

Output:

- x_F^* an approximation of the analytic center of the set $\text{Dom } F$

```

1:  $k \leftarrow 0$ 
2: while  $\|F'(x_k)\|_{x_k}^* > \beta$  do
3:    $x_{k+1} \leftarrow x_k - \frac{1}{1 + \|F'(x_k)\|_{x_k}^*} F''(x_k)^{-1} F'(x_k)$ 
4:    $k \leftarrow k + 1$ 
5: end while
6: return  $x_F^* \leftarrow x_k$ 

```

Theorem 2.7. Algorithm 2 terminates no later than after N steps, where

$$N = \frac{1}{\beta - \ln(1 + \beta)} (F(x_0) - F(x_F^*)). \quad (2.47)$$

The knowledge of analytic center allows us to solve the standard minimization problem

$$x^* = \arg \min_{x \in Q} c^\top x \quad (2.48)$$

with bounded closed convex set $Q \equiv \text{Dom } F$, which has nonempty interior, and which is endowed with a ν -self-concordant barrier $F(x)$. Denote

$$f(t, x) = tc^\top x + F(x), \text{ for } t \geq 0 \quad (2.49)$$

as a parametric penalty function. Using Theorem 2.3 we can see, that $f(t, x)$ is self-concordant in x . Let us introduce new minimization problem using the parametric penalty function $f(t, x)$

$$x^*(t) = \arg \min_{x \in \text{dom } F} f(t, x). \quad (2.50)$$

This trajectory is called the central path of the problem (2.48). We will reach the solution $x^*(t) \rightarrow x^*$ as $t \rightarrow \infty$. Moreover, since the set Q is bounded, the analytic center x_F^* of this set exists and

$$x^*(0) = x_F^*. \quad (2.51)$$

From the first-order optimality condition, any point of the central path satisfies equation

$$tc + F'(x^*(t)) = 0 \quad (2.52)$$

Since the analytic center lies on the central path and can be found by Algorithm 2, all we have to do, to find the solution x^* , is to follow the central path. This enables us an approximate centering condition:

$$\|f'(t, x)\|_x^* = \|tc + F'(x)\|_x^* \leq \beta, \quad (2.53)$$

where the centering parameter β is small enough.

Assuming $x \in \text{dom } F$, one iteration of the path-following algorithm consists of two steps:

$$t_+ = t + \frac{\gamma}{\|c\|_x^*}, \quad (2.54)$$

$$x_+ = x - F''(x)^{-1}(t_+c + F'(x)). \quad (2.55)$$

Theorem 2.8. Let x satisfy the approximate centering condition (2.53)

$$\|tc + F'(x)\|_x^* \leq \beta \quad (2.56)$$

with $\beta < \bar{\lambda} = \frac{3-\sqrt{5}}{2}$. Then for γ , such that

$$|\gamma| \leq \frac{\sqrt{\beta}}{1 + \sqrt{\beta}} - \beta, \quad (2.57)$$

we have again

$$\|t_+c + F'(x_+)\|_{x_+}^* \leq \beta. \quad (2.58)$$

2. Semidefinite programming

Algorithm 3 Path following algorithm

Input:

F a ν -self-concordant barrier
 $x_0 \in \text{dom } F$ a starting point satisfying $\|F'(x_0)\|_{x_0}^* \leq \beta$, e.g. the analytic center x_F^* of the set $\text{Dom } F$
 $\beta \in (0, 1)$ a centering parameter
 γ a parameter satisfying $|\gamma| \leq \frac{\sqrt{\beta}}{1+\sqrt{\beta}} - \beta$
 $\epsilon > 0$ an accuracy

Output:

x^* an optimal solution to the minimization problem (2.48)

```

1:  $t_0 \leftarrow 0$ 
2:  $k \leftarrow 0$ 
3: while  $\epsilon t_k < \nu + \frac{(\beta + \sqrt{\nu})\beta}{1-\beta}$  do
4:    $t_{k+1} \leftarrow t_k + \frac{\gamma}{\|c\|_{x_k}^*}$ 
5:    $x_{k+1} \leftarrow x_k - F''(x_k)^{-1}(t_{k+1}c + F'(x_k))$ 
6:    $k \leftarrow k + 1$ 
7: end while
8: return  $x^* \leftarrow x_k$ 

```

This theorem ensures the correctness of the presented iteration of the path-following algorithm. For the whole description of the path-following algorithm, please see Algorithm 3.

Theorem 2.9. Algorithm 3 terminates no more than after N steps, where

$$N \leq \mathcal{O}\left(\sqrt{\nu} \ln \frac{\nu \|c\|_{x_F^*}^*}{\epsilon}\right). \quad (2.59)$$

The parameters β and γ in Algorithm 2 and Algorithm 3 can be fixed. The reasonable values are:

$$\beta = \frac{1}{9}, \quad (2.60)$$

$$\gamma = \frac{\sqrt{\beta}}{1 + \sqrt{\beta}} - \beta = \frac{5}{36}. \quad (2.61)$$

The union of Algorithm 2 and Algorithm 3 can be easily used to solve the standard minimization problem (2.48), supposing we have a feasible point $x_0 \in Q$.

2.3.3. Barrier function for semidefinite programming

In this section, we are going to show, how to find a self-concordant barrier for the semidefinite program (2.4), so that we can use Algorithm 2 and Algorithm 3 to solve it. For the purpose of this section, we are interested only in the constraints of the problem. The constraints are defining us the feasibility set Q :

$$Q = \left\{ y \in \mathbb{R}^m \mid A_0 + \sum_{i=1}^m A_i y_i \succeq 0 \right\}, \quad (2.62)$$

where $A_0, \dots, A_m \in \mathcal{S}^n$. Let us denote $X(y) = A_0 + \sum_{i=1}^m A_i y_i$. If the matrix $X(y)$ is block diagonal

$$X(y) = \begin{bmatrix} X_1(y) & 0 & \cdots & 0 \\ 0 & X_2(y) & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & X_k(y) \end{bmatrix} \quad (2.63)$$

with $X_j(y) \in \mathcal{S}^{n_j}$ for $j = 1, \dots, k$ and $\sum_{j=1}^k n_j = n$, then the feasibility set Q can be expressed as

$$Q = \{y \in \mathbb{R}^m \mid X_j(y) \succeq 0, j = 1, \dots, k\}. \quad (2.64)$$

This rule allows us to easily add or remove some constraints without touching the others and to keep the sizes of the used matrices small, which can significantly speed up the computation.

Instead of the set Q , which is parametrized by y , we can directly optimize over the set of positive semidefinite matrices. This set \mathcal{P}^n is defined as

$$\mathcal{P}^n = \{X \in \mathcal{S}^n \mid X \succeq 0\} \quad (2.65)$$

and it is called the cone of positive semidefinite $n \times n$ matrices. This cone is a closed convex set, which interior is formed by positive definite matrices and on its boundary lie matrices, which have at least one eigenvalue equal to zero.

Let us denote the function $F(X)$:

$$F(X) = -\ln \prod_{i=1}^n \lambda_i(X), \quad (2.66)$$

where $X \in \text{int } \mathcal{P}^n$ and $\{\lambda_i(X)\}_{i=1}^n$ is the set of eigenvalues of the matrix X . To avoid the computation of eigenvalues, the function $F(X)$ can be also expressed as:

$$F(X) = -\ln \det(X). \quad (2.67)$$

Theorem 2.10. Function $F(X)$ is an n -self-concordant barrier for \mathcal{P}^n .

Note, that $\text{Dom } F \supseteq \mathcal{P}^n$ because $\det(X) \geq 0$ when the number of negative eigenvalues of X is even. Therefore, the set $\text{Dom } F$ is made by separated subsets, which one of them is \mathcal{P}^n . As Algorithm 2 and Algorithm 3 are interior point algorithms, when the starting point is from $\text{int } \mathcal{P}^n$, then we never leave \mathcal{P}^n during the execution of the algorithms and the optimal solution is found.

Similarly, the self-concordant barrier function for the set Q is a function

$$F(y) = -\ln \det(X(y)). \quad (2.68)$$

Example 2.3. To make it clearer, what is the difference between the set Q and $\text{Dom } F(y)$, we have prepared this example. Let

$$X(y) = \begin{bmatrix} y_2 & y_1 \\ y_1 & y_2 \end{bmatrix}, \quad (2.69)$$

where $y = [y_1 \ y_2]^\top$. The equation

$$z = \det(X(y)) = y_2^2 - y_1^2 \quad (2.70)$$

2. Semidefinite programming

represents a hyperbolic paraboloid, which you can see in Figure 2.1. Therefore, the equation $z = 0$ is a slice of it, denoted by the purple color in Figure 2.2. The domain of the self-concordant barrier function is

$$\text{Dom } F(y) = \left\{ y \mid \det(X(y)) \geq 0 \right\} \quad (2.71)$$

and is shaded by the blue color. We can see, that the set $\text{Dom } F(y)$ consists of two disjoint parts. One of them is the set, where $X(y) \succeq 0$ (denoted by the orange color) and the second part is an area, where both eigenvalues of $X(y)$ are negative.




Figure 2.1. Hyperbolic paraboloid $z = y_2^2 - y_1^2$.

2.4. Implementation details

2.5. Comparison with the state of the art methods




Figure 2.2. Illustration of the sets $\text{Dom } F(y)$ and $\{y \mid \det(X(y)) \geq 0\}$.

3. Conclusion

A. Contents of the enclosed CD

```
/
└─ thesis
    └─ thesis.pdf .....digital copy of this thesis
```

Bibliography

- [1] David Cox, John Little, and Donald O'Shea. *Ideals, Varieties, and Algorithms : An Introduction to Computational Algebraic Geometry and Commutative Algebra*. Undergraduate Texts in Mathematics. Springer, New York, USA, 2nd edition, 1997. 4
- [2] Yurii Nesterov. *Introductory lectures on convex optimization : A basic course*. Springer, 2004. 6