

Comprehensive Cinnamon Agronomy & Diagnostic Knowledge Base

Guidelines for Fertilizer, Disease Morphology, Pests, Harvesting,
and Treatment Remedies

Cinnamon Advisor RAG System

August 31, 2026

Contents

1	Introduction	3
2	Environmental Requirements for Cinnamon Cultivation	4
2.1	Climatic Conditions and Stress Indicators	4
2.2	Soil Requirements and Edaphic Factors	5
3	Fertilizer Management and Nutrient Uptake	6
3.1	Soil pH and Dolomite Application	6
3.2	Inorganic Fertilizer Recommendations	6
4	Disease Management: Diagnostics and Remedies	8
4.1	Rough Bark Disease (<i>Pestalotiopsis</i> , <i>Phomopsis</i> , <i>Colletotrichum spp.</i>)	8
4.1.1	Visual and Morphological Symptoms (Diagnostic Features) .	8
4.1.2	Remedies and Treatment Protocols	8

4.2	Leaf Blight (<i>Colletotrichum spp.</i>)	9
4.2.1	Visual and Morphological Symptoms (Diagnostic Features) .	9
4.2.2	Remedies and Treatment Protocols	9
4.3	Leaf Spot (<i>Colletotrichum gloeosporioides, Pestalotiopsis spp.</i>) . . .	10
4.3.1	Visual and Morphological Symptoms (Diagnostic Features) .	10
4.3.2	Remedies and Treatment Protocols	10
4.4	Leaf Gall (<i>Eriophyes boisi</i> - Mites / <i>Trioza cinnamomi</i> - Jumping Plant Lice)	10
4.4.1	Visual and Morphological Symptoms (Diagnostic Features) .	10
4.4.2	Remedies and Treatment Protocols	11
4.5	Stripe Canker (<i>Phytophthora cinnamomi</i>)	11
4.5.1	Visual and Morphological Symptoms (Diagnostic Features) .	11
4.5.2	Remedies and Treatment Protocols	12
4.6	Root Diseases: White and Brown Root Disease	12
4.6.1	White Root Disease (<i>Rigidoporus microporus</i>)	12
4.6.2	Brown Root Disease (<i>Phellinus noxius</i>)	12
5	Pest Management: Identification and Eradication	14
5.1	Caterpillar / Cinnamon Butterfly (<i>Chilasa clytia</i>)	14
5.2	Stem Borer / Cinnamon Wood Borer (<i>Ichneumoniptera cinnamo- mumi</i>)	14
5.3	Thrips (<i>Helionothrips annosus</i>)	15
5.4	Leaf Eating Beetle (<i>Popillia spp., Singhala tenella</i>)	15
5.5	Grasshopper (<i>Orthoptera order</i>)	15
6	Harvesting, Processing, and Quality Standards	16
6.1	Harvesting Technique and Indicators	16
6.2	Post-Harvest Processing (Peeling and Curing)	16
6.3	Hygiene and Regulatory Quality Assurance	17

1 Introduction

This document serves as the foundational knowledge base for the Cinnamon Retrieval-Augmented Generation (RAG) pipeline. It aggregates critical agronomy data, primarily sourced from the Department of Cinnamon Development (DCD) Sri Lanka and global agricultural bodies. It provides detailed, up-to-date guidelines on cinnamon cultivation, fertilizer management, pest and disease control, explicit disease remedies, and harvesting. The extended focus of this document is to provide high-density morphological descriptions tailored for automated computer vision diagnostics, ensuring accurate spatial feature extraction for precision agriculture.

2 Environmental Requirements for Cinnamon Cultivation

Cinnamon (*Cinnamomum verum* J. Presl) is a tropical, evergreen perennial plant. While highly adaptable, commercial cultivation requires specific environmental conditions to optimize bark yield and volatile oil content. Understanding these parameters is critical for diagnosing environmental stress versus pathogenic infection.

2.1 Climatic Conditions and Stress Indicators

- **Temperature:** The optimal temperature range is between 25°C and 35°C . Temperatures exceeding 38°C can induce heat stress, presenting as marginal leaf scorch which must not be confused with pest damage.
- **Rainfall:** Commercial cultivation thrives in areas with an annual rainfall exceeding 1875 mm. In drier regions with rainfall below this threshold, cultivation is possible provided adequate irrigation is available. Drought stress manifests as systemic wilting and a dulling of the leaf cuticle.
- **Elevation:** Cinnamon can be commercially grown from sea level up to an elevation of 700 meters. Growth and yield tend to decline at higher elevations, often resulting in stunted internode lengths.
- **Humidity:** The optimal relative humidity is between 75% and 85%. Environments exceeding 90% humidity for prolonged periods exponentially increase the risk of fungal proliferation, specifically *Colletotrichum* and *Phytophthora* species.

2.2 Soil Requirements and Edaphic Factors

Cinnamon requires deep, well-drained soils.

- **Wet Zone:** Red-Yellow Podzolic soils, common in the rolling hills of the wet zone, are highly suitable. These soils are typically somewhat acidic, which facilitates optimal nutrient uptake for this specific genus.
- **Coastal Areas:** Flat coastal lands with sandy or sandy loam soils are excellent for cinnamon cultivation, provided organic matter is regularly amended to improve cation exchange capacity.
- **Unsuitable Conditions:** Avoid lands with a soil depth of less than 1 meter, rocky terrain, poor drainage, fluctuating water tables, or high salinity. Cinnamon roots are highly susceptible to root rot; prolonged waterlogging (over two weeks) will result in root death. Avoid lands with a slope exceeding 60% to prevent severe topsoil erosion.

3 Fertilizer Management and Nutrient Uptake

Proper nutrient management is critical for maximizing the yield of cinnamon bark and leaves. Deficiencies in macronutrients often present visual symptoms that mimic viral or fungal pathogens. The guidelines below must be strictly adhered to.

3.1 Soil pH and Dolomite Application

- **Optimal pH:** The optimal soil pH for cinnamon is between 5.5 and 6.5. Outside this range, essential micronutrients become chemically locked in the soil matrix.
- **Acidity Correction:** If the soil pH falls below 5.0, Dolomite must be applied. A general recommendation is 400 kg of Dolomite per acre for soils with a pH below 5.0.
- **Application Timing:** Dolomite should be applied during periods of soil moisture (onset or end of the rainy season) to ensure proper dissolution and soil integration.
- **Delay Rule:** Chemical fertilizers must not be applied simultaneously with Dolomite. A minimum waiting period of six weeks (1.5 months) is required after Dolomite application before inorganic NPK fertilizers can be applied.

3.2 Inorganic Fertilizer Recommendations

The standard inorganic fertilizer recommendation for cinnamon is an N:P₂O₅:K₂O ratio of 23:7:15. Nitrogen drives leaf flush, Phosphorus establishes root architecture, and Potassium is vital for bark development and stress tolerance.

- **Mature Plants (Year 3 onwards, 4x3 ft spacing):** Apply 50g Urea, 25g Muriate of Potash (MOP), and 25g Eppawala Rock Phosphate (ERP) per bush annually.
- **Application Schedule:** The annual dosage should be split into two equal applications, timed with the onset of the Yala and Maha monsoon rains.
- **Placement:** For mature bushes, apply fertilizer in a circular band approximately 30 cm (12 inches) away from the base of the plant. Surface broadcasting without incorporation leads to high volatilization losses.

4 Disease Management: Diagnostics and Remedies

Precise visual identification of diseases paired with immediate, targeted remedies is paramount for maintaining plantation health. The following profiles are enriched with specific morphological descriptors to aid in automated and manual diagnoses.

4.1 Rough Bark Disease (*Pestalotiopsis*, *Phomopsis*, *Colletotrichum spp.*)

4.1.1 Visual and Morphological Symptoms (Diagnostic Features)

- **Early Stage Detection:** Initially appears as small brown spots on immature green stems. These micro-lesions often present a slight halo effect under direct light.
- **Late Stage Pathology:** The bark becomes distinctly rough, dry, and cracked, severely impacting bark quality and peelability. The textural shift to a fissured, scab-like surface is a primary diagnostic indicator. In imaging systems, this presents as high-frequency textural noise and distinct color space shifts from healthy green/brown to necrotic gray/black.

4.1.2 Remedies and Treatment Protocols

- **Cultural Remedies:** Immediately prune and burn infected branches away from the plantation. Enforce proper shade control to drastically reduce humidity in the canopy. Practice timely harvesting every 4-6 months to prevent over-maturation of susceptible bark.
- **Chemical Remedies:** For early-stage infections, prepare and apply a 1% Bordeaux mixture. Alternatively, apply a systemic fungicide containing

Tebuconazole diluted at 10 ml per 10 Liters of water. Apply during dry weather to ensure the fungicide adheres to the bark and penetrates the vascular tissue.

4.2 Leaf Blight (*Colletotrichum spp.*)

4.2.1 Visual and Morphological Symptoms (Diagnostic Features)

- **Lesion Characteristics:** Characterized by small brown blister-like spots on leaves that rapidly enlarge into circular, dark necrotic patches. These patches often exhibit concentric zonation rings indicating fungal growth cycles.
- **Tissue Deterioration:** The central necrotic tissue dries and falls out, leaving distinct holes. Primarily affects nurseries and young plants during wet weather. The resulting "shot-hole" appearance dramatically reduces the active photosynthetic surface area.

4.2.2 Remedies and Treatment Protocols

- **Cultural Remedies:** Visually isolate and quarantine infected nursery plants. Remove excess overhead shade to increase air circulation and reduce leaf wetness periods. Avoid overhead sprinkler irrigation; use drip or base watering instead to keep the foliage dry.
- **Chemical Remedies:** Apply a 1% Bordeaux mixture. For severe nursery outbreaks, use systemic fungicides such as Hexaconazole or Tebuconazole to halt fungal replication within the leaf tissue. Spray at 7-10 day intervals until new flush emerges completely healthy.

4.3 Leaf Spot (*Colletotrichum gloeosporioides*, *Pestalotiopsis* spp.)

4.3.1 Visual and Morphological Symptoms (Diagnostic Features)

- Manifests as distinct, small, dark brown or grayish spots scattered across the leaf lamina. The margins of these spots are typically well-defined, distinguishing them from nutritional deficiencies which present as diffused chlorosis.
- Heavily infected leaves will turn yellow and prematurely drop. This abscission leads to thinning canopies and reduced vigor.

4.3.2 Remedies and Treatment Protocols

- **Cultural Remedies:** Ensure adequate spacing between bushes (e.g., 4x3 ft) for maximum aeration. Diligently collect and remove all fallen infected leaves from the base of the plant, as these serve as the primary inoculum source for future outbreaks.
- **Chemical Remedies:** Apply broad-spectrum protective fungicides such as Mancozeb or a copper-based fungicide during the onset of the rainy season to prevent spore germination. Ensure complete coverage of both the adaxial and abaxial leaf surfaces.

4.4 Leaf Gall (*Eriophyes boisi* - Mites / *Trioza cinnamomi* - Jumping Plant Lice)

4.4.1 Visual and Morphological Symptoms (Diagnostic Features)

- **Topography:** Highly distinctive raised, pimple-like bumps or warty growths (galls) on the upper and lower leaf surfaces. These represent cellular hyper-

trophy induced by the pests' salivary toxins.

- **Deformation:** Leaves become distorted, curled, thickened, and turn from light green to red or brown. The geometric disruption of the leaf plane is highly recognizable in 3D spatial mapping and 2D bounding box localization.

4.4.2 Remedies and Treatment Protocols

- **Cultural Remedies:** Prune and immediately destroy heavily galled shoots by burning them. Do not leave pruned galls on the soil surface, as mites can migrate back to the host plant.
- **Botanical Remedies:** Apply a 3-5% Neem oil emulsion directly to the foliage as an organic deterrent to disrupt the feeding and reproductive cycles of the mites and lice.
- **Chemical Remedies:** For widespread and severe mite infestations, apply a targeted miticide such as Abamectin or Spiromesifen according to local agricultural guidelines. Ensure full coverage of the underside of the leaves where mites congregate.

4.5 Stripe Canker (*Phytophthora cinnamomi*)

4.5.1 Visual and Morphological Symptoms (Diagnostic Features)

- Notable vertical, dark brown or black stripes appearing on the bark of the main stem. These linear necrotic tracks follow the vascular cambium, disrupting sap flow.
- The bark rots, and distinct exudation (gumming) is visible from the lesions, leading to dieback. The presence of resinous exudate is a definitive diagnostic marker for severe *Phytophthora* infection.

4.5.2 Remedies and Treatment Protocols

- **Cultural Remedies:** Stripe canker is strictly tied to waterlogging; therefore, improving soil drainage immediately via deep trenching is mandatory. Avoid causing mechanical damage to the bark with weeding tools, as wounds act as entry points for the pathogen.
- **Chemical Remedies:** For localized lesions, mechanically scrape away the infected, rotting bark down to the healthy wood. Immediately apply a fungicidal paste, such as Bordeaux paste or Metalaxyl, directly onto the wound to seal and sterilize the tissue.

4.6 Root Diseases: White and Brown Root Disease

4.6.1 White Root Disease (*Rigidoporus microporus*)

- **Foliar Symptoms:** Above-ground symptoms are delayed; eventually, leaves turn yellow and wilt rapidly, mimicking severe drought stress, leading to total plant death.
- **Remedies:** Uproot and burn infected plants completely. Expose the remaining roots (4 inches deep) and apply a Tebuconazole drench (10 ml per 10 L water). Isolate the infected area by digging deep isolation trenches. Apply 10g of Sulphur dust per planting hole one week prior to replanting in old rubber lands.

4.6.2 Brown Root Disease (*Phellinus noxius*)

- **Symptoms:** Roots become encrusted with a thick, hard mass of soil, sand, and brown fungal mycelium.
- **Remedies:** Improve soil drainage, aggressively remove excess shade, and apply the same Tebuconazole root drench protocol as recommended for White

Root Disease.

5 Pest Management: Identification and Eradication

Effective pest management relies on the early visual detection of insects or their highly specific damage patterns on foliage and stems.

5.1 Caterpillar / Cinnamon Butterfly (*Chilasa clytia*)

- **Visual Symptoms:** Complete defoliation of tender young leaves and flushes, leaving only the thick midribs. The larvae undergo dramatic color changes through instars, making visual identification complex without temporal data.
- **Remedies:** Manual handpicking of caterpillars and pupae for small attacks. For severe nursery infestations, apply biological control via *Bacillus thuringiensis* (Bt) sprays. Bt targets the lepidopteran gut without harming beneficial pollinators.

5.2 Stem Borer / Cinnamon Wood Borer (*Ichneumoniptera cinnamomumi*)

- **Visual Symptoms:** The larva bores into the base of mature stems; brown, powdery frass (sawdust-like excrement) accumulates on the ground around the collar of the plant.
- **Remedies:** Cultural control via earthing up (mounding soil) around the plant base to a height of 2 inches (use clay/mud if soil is loose). Execute timely harvesting of stems older than 3 years to remove prime breeding habitats.

5.3 Thrips (*Helionothrips annosus*)

- **Visual Symptoms:** Scorched, silvery-black tips on young leaves with microscopic shiny spots (excreta). The silverying is caused by air filling the emptied plant cells after the thrips have extracted the sap.
- **Remedies:** Pluck and destroy infected leaves early. Manage overhead shade to prevent dry, exposed conditions. Apply systemic insecticides like Imidacloprid (15 ml per 16 L water) for heavy infestations.

5.4 Leaf Eating Beetle (*Popillia spp.*, *Singhala tenella*)

- **Visual Symptoms:** Jagged, irregular holes chewed entirely through the mature leaf lamina. Damage often occurs nocturnally, meaning the pest is rarely observed during standard daytime scouting.
- **Remedies:** Deploy light traps to monitor and reduce adult populations. Maintain clean inter-rows to destroy daytime hiding spots. Chemical intervention is rarely required unless defoliation exceeds 30%.

5.5 Grasshopper (*Orthoptera order*)

- **Visual Symptoms:** Large, aggressive chunks missing from the margins of young seedlings.
- **Remedies:** Slash perimeter weeds to destroy habitats. Apply concentrated garlic or neem extracts to nursery boundaries as an olfactory deterrent.

6 Harvesting, Processing, and Quality Standards

The culmination of proper agronomic care is the harvest. Processing requires immense manual dexterity and adherence to strict thermodynamic and hygiene standards.

6.1 Harvesting Technique and Indicators

- **Timing Indicator:** Harvest when the outer bark (subula) is entirely brown and the plant exhibits active green leaf flush, indicating high sap flow for easy peeling.
- **Cutting Mechanics:** Cut the stem at a 45-degree angle, 1.5 to 2 inches above the ground level, using a sharp, clean knife. This precision angled cut prevents water pooling on the stump, mitigating fungal rot and encouraging vigorous outward coppicing.
- **Selective Harvesting:** Leave at least one mature stem per bush unharvested to sustain the plant's nutritional needs and root health.
- **Time of Day:** Harvest in the early morning to prevent the bark from drying out and adhering to the wood.

6.2 Post-Harvest Processing (Peeling and Curing)

- **Scraping and Rubbing:** The outer bark must be meticulously scraped off. The stem is then rubbed strictly with a brass rod to loosen the inner bark; iron tools cause severe oxidative blackening of the tissue.
- **Drying Thermodynamics:** The peeled bark (quills) must be dried in the shade. Exposure to direct sunlight or mechanical dryers exceeding 35°C

completely denatures the volatile oils, drastically lowering the aromatic profile and market value.

6.3 Hygiene and Regulatory Quality Assurance

- **Sanitation Parameters:** Processing areas must remain sterile. Bark must never contact the bare ground to prevent microbial contamination.
- **Regulatory Standards:** Sri Lanka Cinnamon must adhere to SLS 81:2021 and ISO 6539:2014 standards. Moisture content must not exceed 14%, and Sulphur dioxide limits must not exceed 150 *mg/kg*.