

Comprehensive Cinnamon Agronomy & Diagnostic Knowledge Base

Extended Guidelines for Fertilizer, Disease Morphology, Pests, and Harvesting

Cinnamon Advisor RAG System

August 30, 2026

Contents

1	Introduction	3
2	Environmental Requirements for Cinnamon Cultivation	3
2.1	Climatic Conditions	3
2.2	Soil Requirements	3
3	Fertilizer Management	3
3.1	Soil pH and Dolomite Application	4
3.2	Inorganic Fertilizer Recommendations	4
3.3	Integrated Plant Nutrition Management (IPNM)	4
4	Disease Management	4
4.1	Rough Bark Disease (<i>Pestalotiopsis</i> , <i>Phomopsis</i> , <i>Colletotrichum spp.</i>)	5
4.1.1	Visual and Morphological Symptoms	5
4.1.2	Control Measures	5
4.2	Leaf Blight (<i>Colletotrichum spp.</i>)	5
4.2.1	Visual and Morphological Symptoms	5
4.2.2	Control Measures	5
4.3	Leaf Spot (<i>Colletotrichum gloeosporioides</i> , <i>Pestalotiopsis spp.</i>)	6
4.3.1	Visual and Morphological Symptoms	6
4.3.2	Control Measures	6
4.4	Leaf Gall (<i>Eriophyes boisi</i> - Mites / <i>Trioza cinnamomi</i> - Jumping Plant Lice) . .	6
4.4.1	Visual and Morphological Symptoms	6
4.4.2	Control Measures	6
4.5	Stripe Canker (<i>Phytophthora cinnamomi</i>)	6
4.5.1	Visual and Morphological Symptoms	6
4.5.2	Control Measures	7
4.6	Root Diseases: White and Brown Root Disease	7
4.6.1	White Root Disease (<i>Rigidoporus microporus</i>)	7

4.6.2	Brown Root Disease (<i>Phellinus noxius</i>)	7
5	Pest Management	7
5.1	Caterpillar / Cinnamon Butterfly (<i>Chilasa clytia</i>)	7
5.1.1	Visual Symptoms and Damage Patterns	7
5.1.2	Control Measures	8
5.2	Stem Borer / Cinnamon Wood Borer (<i>Ichneumoniptera cinnamomumi</i>)	8
5.2.1	Visual Symptoms and Damage Patterns	8
5.2.2	Control Measures	8
5.3	Thrips (<i>Helionothrips annosus</i>)	8
5.3.1	Visual Symptoms and Damage Patterns	8
5.3.2	Control Measures	8
5.4	Leaf Eating Beetle (<i>Popillia spp.</i> , <i>Singhala tenella</i>)	9
5.4.1	Visual Symptoms and Damage Patterns	9
5.4.2	Control Measures	9
5.5	Grasshopper (<i>Orthoptera order</i>)	9
5.5.1	Visual Symptoms and Damage Patterns	9
5.5.2	Control Measures	9
6	Harvesting and Processing (Post-Harvest Technology)	9
6.1	Harvesting Indicators	9
6.2	Harvesting Technique	10
6.3	Processing (Peeling and Drying)	10
6.4	Cinnamon Grading and Quality Standards	10
6.5	Hygiene and Quality Assurance	11

1 Introduction

This document serves as the foundational knowledge base for the Cinnamon Retrieval-Augmented Generation (RAG) pipeline[cite: 1, 2]. It aggregates critical agronomy data, primarily sourced from the Department of Cinnamon Development (DCD) Sri Lanka and global agricultural bodies[cite: 1]. It provides detailed, up-to-date guidelines on cinnamon cultivation, fertilizer management, pest and disease control, and harvesting[cite: 1, 2]. To support automated diagnostic systems, this version heavily emphasizes precise morphological descriptions, spatial distribution of symptoms, and visual boundary delineations for each pest and disease.

2 Environmental Requirements for Cinnamon Cultivation

Cinnamon (*Cinnamomum verum* J. Presl) is a tropical, evergreen perennial plant[cite: 1, 2]. While highly adaptable, commercial cultivation requires specific environmental conditions to optimize bark yield and volatile oil content[cite: 1, 2].

2.1 Climatic Conditions

- **Temperature:** The optimal temperature range is between 25°C and 35°C [cite: 1, 2].
- **Rainfall:** Commercial cultivation thrives in areas with an annual rainfall exceeding 1875 mm[cite: 1, 2]. In drier regions with rainfall below this threshold, cultivation is possible provided adequate irrigation is available[cite: 1, 2].
- **Elevation:** Cinnamon can be commercially grown from sea level up to an elevation of 700 meters[cite: 1, 2]. Growth and yield tend to decline at higher elevations[cite: 1, 2].
- **Humidity:** The optimal relative humidity is between 75% and 85%[cite: 1, 2].

2.2 Soil Requirements

Cinnamon requires deep, well-drained soils[cite: 1, 2].

- **Wet Zone:** Red-Yellow Podzolic soils, common in the rolling hills of the wet zone, are highly suitable[cite: 1, 2]. These soils are typically somewhat acidic[cite: 1, 2].
- **Coastal Areas:** Flat coastal lands with sandy or sandy loam soils are excellent for cinnamon cultivation[cite: 1, 2].
- **Unsuitable Conditions:** Avoid lands with a soil depth of less than 1 meter, rocky terrain, poor drainage, fluctuating water tables, or high salinity[cite: 1, 2]. Cinnamon roots are highly susceptible to root rot; prolonged waterlogging (over two weeks) will result in root death[cite: 1, 2]. Avoid lands with a slope exceeding 60%[cite: 1, 2].

3 Fertilizer Management

Proper nutrient management is critical for maximizing the yield of cinnamon bark and leaves[cite: 1, 2]. The guidelines below must be strictly adhered to[cite: 1].

3.1 Soil pH and Dolomite Application

- **Optimal pH:** The optimal soil pH for cinnamon is between 5.5 and 6.5[cite: 1, 2].
- **Acidity Correction:** If the soil pH falls below 5.0, Dolomite must be applied[cite: 1, 2]. A general recommendation is 400 kg of Dolomite per acre for soils with a pH below 5.0[cite: 1, 2].
- **Application Timing:** Dolomite should be applied during periods of soil moisture (onset or end of the rainy season)[cite: 1, 2].
- **Delay Rule:** Chemical fertilizers must not be applied simultaneously with Dolomite[cite: 1, 2]. A minimum waiting period of six weeks (1.5 months) is required after Dolomite application before inorganic NPK fertilizers can be applied[cite: 1, 2].

3.2 Inorganic Fertilizer Recommendations

The standard inorganic fertilizer recommendation for cinnamon is an N:P2O5:K2O ratio of 23:7:15[cite: 1, 2].

- **Mature Plants (Year 3 onwards, 4x3 ft spacing):** Apply 50g Urea, 25g Muriate of Potash (MOP), and 25g Eppawala Rock Phosphate (ERP) per bush annually[cite: 1, 2].
- **Application Schedule:** The annual dosage should be split into two equal applications, timed with the onset of the Yala and Maha monsoon rains[cite: 1, 2].
- **Placement:** For mature bushes, apply fertilizer in a circular band approximately 30 cm (12 inches) away from the base of the plant[cite: 1, 2].

3.3 Integrated Plant Nutrition Management (IPNM)

To maintain soil health, IPNM combines inorganic and organic fertilizers[cite: 1, 2].

- **Organic Integration:** Apply organic matter such as compost, Gliricidia leaves, or well-rotted cow dung[cite: 1, 2]. When using IPNM, the inorganic fertilizer dose can be reduced by 50%, supplemented with 1000g of organic manure per bush annually[cite: 1, 2].
- **Biochar:** Biochar produced from cinnamon wood waste can be incorporated to improve soil carbon and nutrient retention[cite: 1, 2].

4 Disease Management

Precise visual identification of diseases is paramount for maintaining plantation health[cite: 2]. The descriptions below detail specific morphological changes on the plant surface to aid in accurate diagnosis[cite: 2]. These detailed profiles outline the texture, color gradients, and geometric patterns of lesions, providing critical features for spatial analysis and pixel-level symptom masking.

4.1 Rough Bark Disease (*Pestalotiopsis*, *Phomopsis*, *Colletotrichum spp.*)

Rough Bark Disease is a severe fungal infection that drastically reduces the market value of cinnamon by destroying the peelability of the bark[cite: 1, 2].

4.1.1 Visual and Morphological Symptoms

- **Early Stage Detection:** Initially appears as small brown spots on immature green stems[cite: 1, 2]. These spots are often isolated but can cluster under high humidity.
- **Progression and Texture:** These develop into larger brown patches with distinct black margins[cite: 1, 2]. The central necrotic tissue begins to suberize prematurely.
- **Late Stage Pathology:** As the disease progresses, the bark becomes distinctly rough, dry, and cracked, severely impacting bark quality and peelability[cite: 1, 2]. The textural shift from smooth, healthy green/brown bark to a fissured, scab-like surface is a primary diagnostic indicator.

4.1.2 Control Measures

- **Cultural Control:** Prune and burn infected branches[cite: 1, 2]. Ensure proper shade control to reduce humidity in the canopy and practice timely harvesting (every 4-6 months)[cite: 1, 2].
- **Chemical Control:** Apply a 1% Bordeaux mixture or a fungicide containing Tebuconazole (10 ml per 10 L of water) to early-stage infections[cite: 1, 2].

4.2 Leaf Blight (*Colletotrichum spp.*)

Leaf Blight predominantly affects the photosynthetic capacity of the plant, being particularly destructive in nurseries.

4.2.1 Visual and Morphological Symptoms

- **Lesion Characteristics:** Characterized by small brown blister-like spots on leaves that rapidly enlarge into circular, dark necrotic patches[cite: 1, 2].
- **Tissue Deterioration:** As the lesions age, the central necrotic tissue may dry and fall out, leaving distinct holes in the leaf surface[cite: 1, 2].
- **Distribution:** Primarily affects nurseries and young plants during wet weather[cite: 1, 2]. The lesions often show concentric rings of fungal fruiting bodies under magnification.

4.2.2 Control Measures

- Remove excess shade to increase air circulation, visually isolate infected plants, and apply a 1% Bordeaux mixture, Hexaconazole, or Tebuconazole[cite: 1, 2].

4.3 Leaf Spot (*Colletotrichum gloeosporioides*, *Pestalotiopsis spp.*)

4.3.1 Visual and Morphological Symptoms

- Manifests as distinct, small, dark brown or grayish spots scattered across the leaf lamina[cite: 2].
- Unlike blight, these spots may remain smaller and more localized, but heavily infected leaves will turn yellow and prematurely drop, reducing the photosynthetic area[cite: 2]. The boundary between the infected spot and healthy tissue is typically sharp and well-defined.

4.3.2 Control Measures

- Ensure adequate spacing between bushes for aeration[cite: 2]. Apply protective fungicides such as Mancozeb or a copper-based fungicide during the onset of the rainy season[cite: 2]. Remove fallen infected leaves from the base of the plant[cite: 2].

4.4 Leaf Gall (*Eriophyes boisi* - Mites / *Trioza cinnamomi* - Jumping Plant Lice)

4.4.1 Visual and Morphological Symptoms

- **Topography:** Highly distinctive raised, pimple-like bumps or warty growths (galls) on the upper and lower surfaces of the leaves[cite: 2].
- **Deformation:** Heavily infested leaves become distorted, curled, and thickened[cite: 2].
- **Coloration:** Galls initially appear light green and turn red or brown as they mature[cite: 2]. This volumetric change disrupts the normal flat geometry of the leaf.

4.4.2 Control Measures

- Prune and immediately destroy heavily galled shoots[cite: 2]. For widespread mite infestations, apply a targeted miticide such as Abamectin or Spiromesifen according to local agricultural guidelines[cite: 2].

4.5 Stripe Canker (*Phytophthora cinnamomi*)

4.5.1 Visual and Morphological Symptoms

- Notable vertical, dark brown or black stripes appearing on the bark of the main stem[cite: 2].
- As the infection progresses, the bark rots, and a distinct exudation (gumming) may be visible from the lesions[cite: 2].
- Severe cases lead to dieback of the entire stem[cite: 2]. The vertical orientation of the lesions strongly contrasts with the horizontal peeling process.

4.5.2 Control Measures

- Highly dependent on soil moisture management[cite: 2]. Improve soil drainage immediately[cite: 2]. For localized lesions, scrape away the infected bark to the healthy wood and apply a fungicidal paste (e.g., Bordeaux paste or Metalaxyl)[cite: 2].

4.6 Root Diseases: White and Brown Root Disease

Root diseases are insidious as symptoms often appear on the foliage only after the root system is severely compromised.

4.6.1 White Root Disease (*Rigidoporus microporus*)

- **Foliar Symptoms:** Leaves turn yellow and wilt, leading to plant death[cite: 1, 2].
- **Root Symptoms:** White fungal threads (mycelium) are visible on the surface of the infected roots[cite: 1, 2]. Highly prevalent in old rubber lands or lands adjacent to rubber plantations[cite: 1, 2].
- **Control:** Uproot and burn infected plants[cite: 1, 2]. Expose the remaining roots (4 inches deep) and apply a Tebuconazole drench (10 ml per 10 L water)[cite: 1, 2]. Isolate the infected area by digging trenches[cite: 1]. When planting in old rubber lands, apply 10g of Sulphur dust per planting hole one week prior to planting[cite: 1].

4.6.2 Brown Root Disease (*Phellinus noxius*)

- **Symptoms:** Similar to White Root Disease, but roots turn brown and have soil/sand adhering tightly to them[cite: 1, 2]. Distinct brown lines are visible beneath the bark[cite: 1, 2].
- **Control:** Improve soil drainage, remove excess shade, and apply Tebuconazole drench as recommended for White Root Disease[cite: 1, 2].

5 Pest Management

Effective pest management relies on early visual detection of insects or their specific damage patterns on foliage and stems[cite: 2]. Recognizing distinct chewing patterns, excrement deposition, and tissue necrosis is essential for categorizing pest threats accurately.

5.1 Caterpillar / Cinnamon Butterfly (*Chilasa clytia*)

5.1.1 Visual Symptoms and Damage Patterns

- The larvae (caterpillars) aggressively feed on tender young leaves and flushes[cite: 2].
- They are heavy feeders and can strip a young shoot entirely, leaving only the thick midribs[cite: 2]. This specific damage pattern (complete defoliation sparing only structural veins) is highly characteristic.

- The caterpillars are distinctly colored, often mimicking bird droppings in early instars and later developing striking striped or spotted patterns[cite: 2].

5.1.2 Control Measures

- For small-scale attacks, manual handpicking of caterpillars and pupae is highly effective[cite: 2].
- In severe nursery infestations, the application of *Bacillus thuringiensis* (Bt) provides biological control without harming beneficial insects[cite: 2]. Chemical insecticides should be a last resort[cite: 2].

5.2 Stem Borer / Cinnamon Wood Borer (*Ichneumoniptera cinnamomumi*)

5.2.1 Visual Symptoms and Damage Patterns

- The larva bores into the base of mature stems, feeding on the tissue from the bark to the heartwood[cite: 1, 2].
- The most prominent diagnostic feature is the accumulation of brown, powdery frass (sawdust-like excrement) visible on the ground at the base of the infected plant[cite: 1, 2]. Infected stems may break and fall over[cite: 1].

5.2.2 Control Measures

- The most effective cultural control is earthing up (mounding soil) around the base of the plant to a height of about 2 inches, especially after the plant is 3 years old[cite: 1, 2]. If soil cannot be used, a clay/mud mixture can be applied[cite: 1]. Ensure timely harvesting, as older, thicker stems are more susceptible[cite: 2].

5.3 Thrips (*Helionothrips annosus*)

5.3.1 Visual Symptoms and Damage Patterns

- Thrips suck sap from the tender foliage[cite: 2]. Feeding causes the tips and margins of young leaves to appear burnt, scorched, or silvery-black[cite: 1, 2].
- Microscopic shiny spots (excreta) and small, elongated insects are visible on the underside of the leaves[cite: 1, 2]. Severe attacks can cause defoliation and dieback of shoots[cite: 1].

5.3.2 Control Measures

- Pluck and destroy infected leaves early[cite: 1, 2]. Ensure overhead shade is properly managed, as thrips thrive in dry, exposed conditions[cite: 2]. For severe attacks, apply systemic insecticides like Imidacloprid (15 ml per 16 L water)[cite: 1, 2].

5.4 Leaf Eating Beetle (*Popillia spp.*, *Singhala tenella*)

5.4.1 Visual Symptoms and Damage Patterns

- Adult chafer beetles feed on the foliage, often at night[cite: 2].
- The damage manifests as irregular, jagged holes chewed through the leaves, or feeding from the leaf margins inwards[cite: 2].
- Heavy infestations can cause skeletonization of the leaves[cite: 2]. The lack of frass on the leaves distinguishes this from caterpillar damage.

5.4.2 Control Measures

- Keep the plantation free of dense weed undergrowth where beetles hide during the day[cite: 2]. Use light traps at night to monitor and reduce adult beetle populations[cite: 2]. Neem oil applications can act as a feeding deterrent[cite: 2].

5.5 Grasshopper (*Orthoptera order*)

5.5.1 Visual Symptoms and Damage Patterns

- Grasshoppers are primarily a threat in the nursery stage or to very young transplants[cite: 2].
- They have strong mandibles and chew large, irregular chunks out of the leaf margins, sometimes consuming entire young seedlings[cite: 2]. The bite marks are generally larger and smoother than those left by beetles.

5.5.2 Control Measures

- Maintain a clean perimeter by slashing tall grass and weeds around nurseries[cite: 2]. Hand removal is feasible for small operations[cite: 2]. Botanical extracts (like localized garlic or neem sprays) can deter feeding on highly vulnerable seedlings[cite: 2].

6 Harvesting and Processing (Post-Harvest Technology)

The proper execution of harvesting and post-harvest processing directly determines the final export grade and economic value of the cinnamon[cite: 1, 2].

6.1 Harvesting Indicators

- **Timing:** Harvesting begins 2-3 years after planting[cite: 1, 2]. The optimal time to harvest is when the plant flushes new leaves and they turn green, as this indicates active sap flow, making peeling significantly easier[cite: 1, 2]. Avoid harvesting during flowering, fruiting, or extreme dry periods[cite: 1, 2].
- **Visual Cues:** The outer bark (subula) should be entirely brown, except for the very tips of the branches[cite: 1, 2].

- **Peeling Test:** The bark must easily and cleanly separate from the wood when tested with a knife[cite: 1, 2].

6.2 Harvesting Technique

- **Cutting:** Cut the stem at a 45-degree angle, 1.5 to 2 inches above the ground level, using a sharp, clean knife[cite: 1, 2]. This angle promotes the outward growth of new vegetative shoots[cite: 1, 2].
- **Selective Harvesting:** Leave at least one mature stem per bush unharvested (roughly 1 in 5 stems) to sustain the plant's nutritional needs and root health[cite: 1, 2].
- **Time of Day:** Harvest in the early morning[cite: 1, 2]. Prolonged exposure of cut stems to sunlight will cause them to dry out quickly, causing the bark to adhere tightly to the wood[cite: 1, 2].

6.3 Processing (Peeling and Drying)

- **Scraping (Kurutta Sirima):** The outer brown bark must be completely scraped off using a specialized curved knife (Kokaththa) to reveal the yellow inner bark[cite: 1, 2]. Failure to remove the outer bark results in a lower-grade product with a poor taste, unattractive color, and higher susceptibility to insect damage[cite: 1].
- **Rubbing:** Rub the scraped stem with a brass rod to loosen the inner bark[cite: 1, 2]. Iron rods must not be used, as iron reacts with the chemical compounds in cinnamon, turning the bark black[cite: 1, 2].
- **Drying:** Freshly peeled bark must be air-dried in the shade on clean nylon nets or racks until the edges curl inward[cite: 1, 2]. Never expose the bark to direct sunlight[cite: 1, 2]. If using mechanical dryers, the temperature must strictly not exceed 35°C to preserve volatile flavor compounds[cite: 1, 2].
- **Quill Formation:** The standard length for a high-quality cinnamon quill is 42 inches[cite: 1].

6.4 Cinnamon Grading and Quality Standards

Export-ready Ceylon Cinnamon is meticulously graded based on quill diameter, foxing (surface blemishes), and overall craftsmanship.

- **Alba:** The finest, rarest, and most expensive grade. Quills are pencil-thin (approx. 6 mm in diameter), pale gold, and delicately hand-rolled. It requires a minimum of 45 whole quills per kilogram.
- **Continental (C Grades):** High quality, golden-yellow quills spanning from C5 Special (7-9 mm diameter) down to C4. C5 is considered the premium everyday standard for culinary use.

- **Mexican (M Grades):** Slightly thicker, bolder quills (e.g., M5, M4) with a diameter up to 19 mm.
- **Hamburg (H Grades):** The thickest and strongest-flavored quills, primarily consisting of mature bark with darker patches. H1 is the highest subgrade within this category.
- **By-products:** Quillings (broken pieces), featherings, and chips represent the same authentic *Cinnamomum verum* species but fail the physical structure requirements for whole quills.

6.5 Hygiene and Quality Assurance

- **Sanitation:** Processing areas must be pristine[cite: 2]. Workers must wash hands, wear clean clothes, and refrain from eating, chewing betel, or spitting in the processing area[cite: 1, 2]. Cinnamon stems and peeled bark must never touch the ground[cite: 1].
- **Regulatory Standards:** Sri Lanka Cinnamon must adhere to SLS 81:2021 and ISO 6539:2014 standards[cite: 1, 2]. Moisture content must not exceed 14%, and Sulphur dioxide limits must not exceed 150 *mg/kg*[cite: 1, 2].