

Ve401 Probabilistic Methods in Engineering

Summer 2018 — Assignment 2

Date Due: 12:10 PM, Wednesday, the 30th of May 2018



JOINT INSTITUTE
交大密西根学院

This assignment has a total of (28 Marks).

Exercise 2.1 Discrete Uniform Distribution

A discrete random variable is said to be *uniformly distributed* if it assumes a finite number of values with each value occurring with the same probability. For example, in the generation of a single random digit taken from $\{0, 1, 2, 3, 4, 5, 6, 7, 8, 9\}$ the number generated is uniformly distributed with each possible digit occurring with probability $1/10$.

In general, the density for a uniformly distributed random variable $X: S \rightarrow \{x_1, \dots, x_n\} \subset \mathbb{R}$, $n \in \mathbb{N}$, is given by

$$f(x_k) = \frac{1}{n}, \quad k = 1, \dots, n.$$

- Find the moment-generating function for a discrete uniform random variable.
(2 Marks)
- Use the moment-generating function to find $E[X]$ and $\text{Var}[X]$.
(2 Marks)

Exercise 2.2 Uniqueness of Moment Generating Functions - Simple Case

Suppose that two discrete random variables (X, f_X) and (Y, f_Y) both take on values $\{0, \dots, n\}$, $n \in \mathbb{N}$. Suppose that the moment-generating functions are equal in some neighborhood of zero, i.e., there exists some $\varepsilon > 0$ such that

$$m_X(t) = m_Y(t) \quad \text{for all } t \in (-\varepsilon, \varepsilon).$$

Show that $f_X(x) = f_Y(x)$ for $x = 0, \dots, n$.
(4 Marks)

Exercise 2.3 Sums of Independent Discrete Random Variables

Two discrete random variables X and Y are said to be independent if

$$P[X = x \text{ and } Y = y] = P[X = x] \cdot P[Y = y] \quad \text{for any } x \in \text{ran } X \text{ and } y \in \text{ran } Y.$$

- Let $Z = X + Y$, where X and Y are assumed to be independent. Show that

$$P[Z = z] = \sum_{x+y=z} P[X = x] \cdot P[Y = y]$$

using the formula for total probability.

(3 Marks)

- Show that the sum of two independent and identical geometric random variables follows a Pascal distribution with $r = 2$.
(3 Marks)

Exercise 2.4 Drawing until First Success in the Hypergeometric Setting

- Let (X, f_X) be a discrete random variable taking on only positive values, i.e., $\text{ran } X \subset \mathbb{N}$. Show that

$$E[X] = \sum_{x=0}^{\infty} P[X > x].$$

(2 Marks)

- A box contains N balls, of which r are red and $N - r$ are black. Balls are drawn from the box until a red ball is drawn. Show that the expected number of draws is

$$\frac{N+1}{r+1}.$$

Hint: $\sum_{x=0}^r \binom{N-r+x}{N-r} = \binom{N+1}{N-r+1}$
(4 Marks)

Exercise 2.5 Density of the Poisson Approximation

The density $p_x(t)$, $x \in \mathbb{N}$, of the Poisson distribution is obtained iteratively from the following differential equations:

$$p'_0 = -\lambda p_0, \quad p'_x + \lambda p_x = \lambda p_{x-1}.$$

Use induction to prove that $p_x(t) = (\lambda t)^x e^{-\lambda t} / x!$. Justify the initial values that you apply in your derivation.
(4 Marks)

Exercise 2.6 Poisson Approximation to the Binomial Distribution

Consider the density f of the binomial distribution,

$$f(x) = \binom{n}{x} p^x (1-p)^{n-x}. \quad (*)$$

Let k be fixed so that $np = k$ and set $p = k/n$. Replace p by k/n everywhere in $(*)$ and then let $n \rightarrow \infty$. Show that for every x , $f(x) \rightarrow (k^x / x!) e^{-k}$, the density of the Poisson distribution with parameter k .
(4 Marks)